



Contents lists available at ScienceDirect

Journal of Ethnopharmacology

journal homepage: www.elsevier.com/locate/jep



Review

Plant-based remedies for wolf bites and rituals against wolves in the Iberian Peninsula: Therapeutic opportunities and cultural values for the conservation of biocultural diversity



José A. González^{a,*}, Ana Maria Carvalho^b, José Ramón Vallejo^c, Francisco Amich^a

^a Grupo de Investigación de Recursos Etnobiológicos del Duero-Douro (GRIRED), Facultad de Biología, Universidad de Salamanca, Salamanca E-37071, Spain

^b Mountain Research Centre (CIMO), School of Agriculture, Polytechnic Institute of Bragança, Campus Santa Apolónia, 5300-253 Bragança, Portugal

^c Área de Didáctica de Ciencias Experimentales, Equipo de Historia de la Ciencia y Antropología de la Salud, Salud, Facultad de Educación, Universidad de Extremadura, Badajoz, E-06006, Spain

ARTICLE INFO

Keywords:

Wound healing
Wolf injuries
Herbal remedies
Veterinary ethnopharmacology
Iberian Peninsula

ABSTRACT

Ethnopharmacological relevance: Combined approaches to local knowledge and folk plant use improve awareness and promote effective strategies for the conservation of significant biocultural patrimony. Moreover, the information reported might be the basis for further appropriate phytochemical and pharmacological research. Therefore we provide an insight into traditional herbal remedies and practices for healing bite injuries in humans and domestic animals caused by the Iberian wolf. Wolf bites are associated with inflammatory processes and rabies is a potential complication

Aims: This paper describes and summarises the medicinal-veterinary empirical and ritual uses of the Iberian flora for wolf injuries and reviews the ethnopharmacological data of specific plants that are already published. The Iberian wolf is a critically endangered subspecies of the grey wolf. Livestock attacks attributed to wolves are increasingly frequent in the Iberian Peninsula, resulting in serious social problems. Interesting strategies for Iberian wolf conservation might be related to traditional grazing practices that are deeply linked with empirical knowledge and local practices passed on by oral tradition, which are also vulnerable now.

Materials and methods: Based on documentary sources from the beginning of the twentieth century to the present, we systematically searched old monographs, regional documents, technical papers, project reports, as well as the international and national databases and the available scientific literature, without restrictions regarding the language of the publications consulted.

Results: A total of 39 remedies for healing wolf bite injuries in humans and domestic animals was reported, highlighting the medicinal use of 33 species of vascular plants, mostly wild herbs, belonging to 18 botanical families. The use of wood ashes was also reported. The number of use-reports found represents a very high number considering similar European studies. Leaves were the predominant plant part mentioned. Boiling plant materials in water for topical uses was the most frequent method of preparation found. Some traditional remedies combined two or more plant species in order to potentiate their effects. Moreover, some plant-based traditional practices and rituals to ward off wolves and to prevent wolf attacks were also documented. In these practices eleven other species (belonging to seven more families) were used.

Conclusions: Despite the decline of the Iberian wolf over the last few decades, wolves are still in the imaginary of rural communities that perceive this large carnivore as both a diabolic creature and a mythic and benign animal. Wolf-related cultural heritage is of great interest in terms of conservation strategies. This review emphasises the importance of local knowledge and provides useful information about several potential sources of phytochemicals and their claimed therapeutic effects, aiming at contributing to the conservation and appreciation of the Iberian biocultural heritage.

* Corresponding author.

E-mail addresses: ja.gonzalez@usal.es (J.A. González), anacarv@ipb.pt (A.M. Carvalho), joseramonvallejo@unex.es (J.R. Vallejo), amich@usal.es (F. Amich).



Fig. 1. A specimen of Iberian wolf showing the characteristic black marks on the front legs (photo by J. A. González). Current geographical distribution of the subspecies [Source: the authors based on national and regional censuses].

1. Introduction

For many centuries, the Iberian Peninsula (Portugal and Spain) was a region where wolves were abundant (Gragera, 1996; Grande del Brío, 1984, 2000; Massip, 2011; Pimenta et al., 2005). The Iberian wolf (*Canis lupus signatus* Cabrera, 1907), an endemic subspecies smaller than the other European wolves and exhibiting a reddish coat and characteristic dark signs –*signatus*– (Fig. 1) (Grande del Brío, 2000; Iglesias et al., 2017; Torres and Fonseca, 2016), is an important element of biocultural diversity (i.e. a rich patrimony combining wild resources, habitat and landscape, and cultural heritage) in the Iberian territory.

A long coexistence between man and wolves provided unique cultural manifestations such as myths, legends, superstitions, tales and peculiar ways to defend against attacks by wolves and to hunt these predators –e.g. particular and varied wolf traps known as *loberas*, *callejos*, *chorcos*, *fosos*, *fojos*, *cortellos*, *cousos*– (Álvares et al., 2000, 2011; Grande del Brío, 1984, 2000; Valverde and Teruelo, 2001). Emblematic Portuguese and Spanish literary works (e.g. Martínez-Barbeito, 1947; Ribeiro, 1918, 1958) and hundreds of tales and stories that have been passed down orally through generations (e.g. Riego Celada and Galhano Alves, 2006) highlight the common presence of wolves and how they are key elements of regional identity.

The range of the Iberian wolf declined throughout the last decades of the twentieth century due to human expansion and persecution, habitat loss, scarcity of natural prey, and forests fires. More recent estimates suggest that in the Iberian Peninsula there are no more than 2500 wolves (more or less 80% in Spain and 20% in Portugal), most of the individuals occurring in a large and continuous population in the north-western region, and some others in two isolated populations in Sierra Morena (Andalusia, southern Spain) and in the southern area of the Douro River (central Portugal) (see Fig. 1) (Blanco et al., 2007; MAPAMA, 2015; Pimenta et al., 2005; Sáenz de Buruaga et al., 2015; Torres and Fonseca, 2016).

Important conservation strategies improving recolonisation (e.g. Grupo de Trabajo del Lobo, 2006; MEDWOLF, 2016) and national and European regulations, some providing compensation programmes for damages by wolves, partly managed to control wolf numbers and their distribution (Eggermann et al., 2011; Fernández-Gil et al., 2016). However, this subspecies is still threatened as rural abandonment increases, and flora and fauna changes affecting wolves' food sources and conflicts with livestock producers and farmers are still occurring.

At the present time, livestock attacks attributed to the wolf are more and more frequent in the Iberian Peninsula (see De la Cal, 2016; Pimenta et al., 2017; Sereno, 2017), resulting in serious social problems. The non-profit organization *Fondo para la Protección de los Animales Salvajes* (FAPAS) argues that the main causes are the

fragmentation of wolf packs (unable to capture large wild prey), the significant decrease in wild ungulates populations and the indiscriminate elimination of wolf specimens without technical or scientific basis (see FAPAS, 2016).

In depopulated areas, with no human-wolf potential conflicts, wolves mainly predate wild ungulates, i.e. roe deer (*Capreolus capreolus* (Linnaeus, 1758)), red deer (*Cervus elaphus* Linnaeus, 1758) or wild boar (*Sus scrofa* Linnaeus, 1758), while domestic animals are consumed occasionally (e.g. Barja, 2009). However, wolves having some contact with human activities feed mainly on livestock, being stocks density (number of animals) the factor that better explain wolf occurrence and persistence. For instance, in Portugal, domestic goat (*Capra hircus* Linnaeus, 1758) predominates in wolf diet (> 60%), followed by cattle (*Bos taurus* Linnaeus, 1758) and sheep (*Ovis aries* Linnaeus, 1758) (Eggermann et al., 2011; Torres et al., 2015). Otherwise, in the Spanish Autonomous Communities of Asturias and Galicia large livestock species, horses (*Equus caballus* Linnaeus, 1758) and cattle (primarily foals and calves), are the dominant prey, due to important changes in environmental and agricultural policies affecting these regions (Lagos and Bárcena, 2015; Llaneza and López-Bao, 2015).

Livestock' availability near villages attracts wolves to the outskirts resulting in their adaptation to human presence and thus increasing predatory attacks (Eggermann et al., 2011; Llaneza et al., 2012).

According to data from the Spanish Ministry of Environment, in 2016 more than 4600 wolf attacks were registered for eight Spanish Autonomous Communities (De la Cal, 2016). Data from Castile and León and Asturias Communities are particularly noticeable, as 216 of the 297 wolf packs identified in Spain inhabit these areas. In Castile and León ca. 1800 attacks resulted in about 3300 dead animals having been reported. In Asturias the 37 packs registered caused 2126 attacks on livestock. Overall the number of farm animals affected by wolf damages in Asturias averaged annually ca. 3000 and compensation costs more or less 600,000€ (Fernández-Gil et al., 2016). Likewise, wolf attacks are growing in those territories that are being recolonised, such as the Sierra de Madrid. The number of cases recorded in this area increased over time from 91 attacks in 2015 to 209 in 2016, corresponding to 337 domestic animals killed and 89,625€ paid as compensations (De la Cal, 2016).

Considering the Portuguese territories, during 2012–2013 wolves were responsible for about 2725 attacks to all livestock species per year. The annual compensations paid for damages approximately correspond to 820,000€ (Pimenta et al., 2017). These authors also reported that 83% of the farmers affected (total of 260 farms) suffered between 1 and 3 attacks per year, while only 3.8% suffered ten or more attacks per year (Pimenta et al., 2017).

The number of animals compensated by depredations amounted to

< 1% of free-ranging livestock (Fernández-Gil et al., 2016). Even so, most problems are due to economic damage caused to domestic animals in isolated rural areas and to the lack of funding for damage compensation (ASCEL, 2016; Cerrillo, 2017; Rejón, 2016); several cases of fraud in relation to the payment system with public money have also been reported (Planelles, 2016a, 2016b).

Moreover wolves wounds on cattle are greatly feared by rural people because it is believed that these wounds “turn black” (undergo necrosis) and gangrene easily (e.g. Arízaga, 1915; Martínez González and Lastra, 2006). It is well known that wolf bite wounds may lead to serious infection. The microorganisms involved tend to originate from the oral cavity of the biting animal, which can also be influenced by the microbiome of their ingested prey and other foods, as well as the environment where the injury occurred. A variety of aerobic and anaerobic microorganisms have been isolated from large carnivore bite wounds, with infection ranging from localised cellulitis to systemic dissemination, leading to severe complications ranging from abscess to bone and joint infection, to endocarditis and brain abscess (Abrahamian and Goldstein, 2011; Brook, 2009; Thomas and Brook, 2011).

Frequently people receiving a relatively minor bite from domestic or feral dogs (*Canis lupus familiaris* Linnaeus, 1758) develop extremely serious complications (diverse infections, septicaemia, paralysis and renal failure) that might result in death (Linnell et al., 2002; Thomas and Brook, 2011). Apparently such medical conditions are also associated with wolf bite injuries.

Wolf biting has additional problems because rabies represents the most feared potential complication after wolf bites in many regions worldwide (e.g. Linnell et al., 2002; Simani et al., 2012; Türkmen et al., 2012). Rabies has not been present in the Iberian Peninsula for a long time. This viral zoonosis never became established in wildlife and was eradicated in domestic dogs by the 1970s within the territories of Portugal and Spain (Linnell et al., 2002).

In Spain, there are a number of reported cases where wolves have attacked or threatened adults in self-defence (defensive attacks by a wolf that was trapped or cornered, or a wolf protecting a den site with pups) and well documented episodes where predatory attacks by wolves have occurred on humans. In the latter case, all accidents happened in the 1950s and 1970s in Galicia, and were associated with specific circumstances increasing the probability of carnivores' undesirable behaviour. For instance, landscapes with fragmented habitats, low densities of wild prey, high level of wolf dependence on livestock and anthropogenic foods, great human densities experiencing poor rural lifestyles. Such episodes were mostly focused on defenceless children placed in vulnerable situations (Linnell et al., 2002; Valverde and Teruelo, 2001). Linnell and Alleau (2016) state that nowadays rural population generally do not carry out activities that would constitute potential hazards (e.g. using children as shepherds). This fact implies a very low risk for people, being the real social economic problem the damages on livestock.

According to several scientists and researchers, some interesting strategies for wolf conservation might be related to traditional livestock practices and new approaches to cattle grazing (Álvares and Blanco, 2014; MEDWOLF, 2016; Signatus.org, 2016; Torres and Fonseca, 2016; WOLF, 2014). For example, some projects conducted in Portugal and Spain are studying the advantages of having a donkey (*Equus asinus* Linnaeus, 1758) among the herd (e.g. ASZAL, 2014; Ramos, 2015). It seems donkeys are particularly aggressive against wolves, their behaviour warning of imminent attacks as well as driving back the pack (Mettler, 2014). Likewise, the presence of guardian dogs may also prevent livestock predators. These guardians are mastiff type dogs with stud collars protecting their neck from bites, traditionally known as *carlancea* or *carranca*, fitted with sharp points and metal studs. They bark loudly and are large enough to keep wolves away (MEDWOLF, 2016; Nieto Maceín, 2016; Rosa, 2013).

This paper assumes that medical and veterinary popular remedies

and practices used in the past, many of them largely forgotten and already abandoned, might contribute to Iberian wolf conservation strategies, as well as minimise the causes of livestock damage, reducing potential conflict and allowing traditional systems of grazing. Thus, the aims of the present review are as follows: (1) to document and analyse traditional knowledge about the medicinal and veterinary use of plants to treat wolf bites in the Iberian Peninsula; (2) to document traditional plant-based practices to prevent wolf attacks; (3) to make results available within the scientific community in order to provide information for future complementary research, for instance into the validation of the effects of some plants and practices, and (4) to contribute to the conservation of biocultural diversity.

2. Materials and methods

2.1. Data collection

To access the maximum number of documentary sources, a qualitative systematic review of international and national databases was conducted. The ISI Web of Science and Anthropology Plus and JSTOR III – Arts and Sciences international databases were consulted.

The Spanish resources referenced include the database of Ph.D. Theses, TESEO; the information system of the databases of the Spanish Research Council (Spanish acronym CSIC); the Dialnet bibliographic website; Google Scholar and the catalogue of Public State Libraries (BPE). Furthermore, all ethnobotanical studies included in the repository of the Spanish Inventory of Traditional Knowledge related to Biodiversity –IECTB for its acronym in Spanish– were checked (see Aceituno-Mata et al., 2014).

Most of the Portuguese documentary sources are the direct result of an intensive review of the literature using search engines and the repository of B-on (Biblioteca do Conhecimento Online). A few old ethnographic documents were retrieved in Bragança Public Libraries. Ph.D. and Master Theses were accessed online, as well as a number of academic and project reports, conference proceedings, published papers and other non-refereed publications, such as several ethnobotanical guides and books. Most of the sources found focused on the distribution of the Iberian wolf, its diet, behaviour, health and genetics. Quite a few works provided useful data about wild species used in the context of attacks by wolves and on the cultural dimensions of wolves. Nevertheless, Portuguese ethnobotanical studies exist, carried out in the areas of Iberian wolf distribution, and reporting on the use of different plants to heal injuries caused by animals, for example wild boars, feral dogs and wolves to humans, livestock and other domestic animals. At times, these use-reports have a global approach and do not specify the exact agent causing the injury. It was therefore assumed that these species and remedies were also used in the case of wolf bites and wounds.

The overall search pattern covered the title, abstract and key words concerning ethnobotany-related disciplines that have UNESCO codes (e.g. anthropology, the history of medicine, botany) and the terms wolf/wolves, wolf bite, wolf attack, folk medicine, folk veterinary medicine, folklore, pastoralism, ethnobiology, ethnobotany, ethnopharmacology, ethnomedicine and ethnoveterinary medicine, in conjunction with the Iberian geographical context. No restrictions regarding the language of the publications consulted were imposed (e.g. Spanish, Portuguese, Galician, English, etc.).

This review was carried out using data found in ca. 50 documentary sources from the beginning of the twentieth century to the present. As for the type of these documentary sources, we obtained empirical use-reports from 22 books, 10 within the scope of ethnobotany, seven concerning folklore or ethnography and only five directly related to ethnomedicine or ethnoveterinary medicine. It was also included, information from 10 journal papers, mostly published in seven journals in the field of folklore and ethnography, five theses, four of which focused on ethnobotanical studies carried out in particular geographi-

Table 1
Vascular plant species traditionally used in the Iberian Peninsula for healing wounds done by wolves.

Botanical families Species	Vernacular name (s)	Common English name	Status ^a	Use-category ^b	Part (s) used	Method of preparation	Geographical location	References
POLYPODIOPSIDA <i>Osmundaceae</i> <i>Osmunda regalis</i> L.	Llantosil, untosil, ventosil	Royal fern	W	EVM (all livestock)	Rhizome	Decoction	Piloña and Sobrescobio (Asturias), SPAIN	Álvarez Peña (2004)
MAGNOLIOPSIDA <i>Adoxaceae</i> <i>Sambucus nigra</i> L.	Sabugueiro, caneleiro	Elderberry	W, C	MED / EVM (all livestock)	Flowers Bark	Decoction Ointment (with olive oil, or with beeswax, red wine, sheep fat and wheat bran)	Mogadouro (Natural Park of Douro International), PORTUGAL Mogadouro (Natural Park of Douro International), PORTUGAL	Carvalho and Ramos (2012) Carvalho and Ramos (2012)
ARALIACEAE <i>Hedera helix</i> L. [and <i>H. hibernica</i> (G. Kirchn.) Carrière]	Edra, hera	Ivy	W, C	MED	Cordiform leaves from the sterile stems	Decoction Poultice (warm up)	Vinhais and Bragança (Natural Park of Montesinho), and Natural Park Serra da Estrela, PORTUGAL Vinhais and Bragança (Natural Park of Montesinho), and Natural Park Serra da Estrela, PORTUGAL	Carvalho (2010); Silva et al. (2011)
CISTACEAE <i>Cistus albidus</i> L.	Lobera, hierba lobera	Grey-leaved cistus	W	MED	Aerial parts	Decoction	South part of the province of Alava, SPAIN	Alarcón et al. (2015)
<i>Cistus populifolius</i> L.	Jara, jara de Salentinos	—	W	EVM (all livestock, mainly horses)	Aerial parts	Decoction	Palacios del Sil (León), SPAIN	García Jiménez (2007)
<i>Cistus salviifolius</i> L.	Lobera, hierba lobera, bedarra, iñerra zurie, árnica	Sage-leaved rock-rose	W	MED	Aerial part or leaves	Decoction	Northwest of the Basque Country (Biscay and Alava), SPAIN	Menéndez-Baceta et al. (2014); Alarcón et al. (2015)
<i>Tuberaria lignosa</i> (Sweet) Samp.	Erva-loba, yerba-lhoba, alcaria, erva-árcadia	—	W	MED / EVM (all livestock)	Whole plant, aerial parts	Decoction	Vinhais and Bragança (Natural Park of Montesinho), and Miranda do Douro (Natural Park of Douro International), PORTUGAL	Carvalho and Ramos (2012)
COMPOSTAE	Cardo de arzolla	—	W	EVM (all livestock)	Whole plant	Decoction	Comarca de Campoo (Cantabria), SPAIN	Pardo-de-Santayana (2008)
<i>Carthamus carduncellus</i> L. [syn. <i>Carduncellus carduncellus</i> (L.) Huth] <i>Carthamus mitissimus</i> L. [syn. <i>Carduncellus mitissimus</i> (L.) DC.]	Cardo de arzolla, cardo las zolla	—	W	MED / EVM (all livestock)	Whole plant	Decoction	Panzegos (La Rioja) and the Montaña Palentina district (Palencia), SPAIN	Arízaga (1915); Oria de Rueda et al. (1996)
<i>Centaurea lagasana</i> Graells	Arzolla	—	W	MED / EVM (all livestock)	Inflorescence or whole plant	Decoction	Comarca de Campoo (Cantabria), SPAIN	Pardo-de-Santayana (2004, 2008)
<i>Centaurea ornata</i> Willd.	Arzolla	—	W	MED / EVM (all livestock)	Whole plant (in flower) Root	Decoction	Cantabria, SPAIN	Pardo-de-Santayana (2004)
<i>Chamaemelum nobile</i> (L.) All. <i>Jurinea humilis</i> (Desf.) DC.	Manzanilla, manzanilla de campo, manzanillón	Roman chamomile	W	MED	Inflorescences (flower heads)	Infusion	Northwest of the Zamora province, SPAIN	Tallegón (2016)
	Arzolla	—	W	MED / EVM (all livestock, mainly sheep)	Whole plant	Poultice (boiled)	Comarca de Campoo (Cantabria), SPAIN	Pardo-de-Santayana (2008)
						Decoction	Natural Park of Fuentes Carrionas y Fuente Cobre (Palencia), SPAIN	Pascual Gil (2013)
GENTIANACEAE <i>Gentiana lutea</i> L.	Genciana, janciana,	Great yellow	W	EVM (cattle,	Rhizome	Decoction	Natural Park of Fuentes Carrionas y Fuente Cobre (Palencia), SPAIN	Pascual Gil (2013)
								Martínez González and (continued on next page)

Table 1 (continued)

Botanical families Species	Vernacular name (s)	Common English name	Status ^a	Use-category ^b	Part (s) used	Method of preparation	Geographical location	References
JUGLANDACEAE <i>Juglans regia</i> L.	xanquina, xancina gentian		EVM (sheep) equines	Rhizome, roots	Decoction	Cuevas del Sil (León), SPAIN	Lastra (2006)	
	Nogal, nozal, noguera	Common walnut	C	EVM (cattle)	Leaves	Decoction	Concejo de Aller (Asturias), SPAIN	García Jiménez (2007)
LAMIACEAE <i>Rosmarinus officinalis</i> L.	Romero	Rosemary	C	EVM (sheep)	Leaves (dried)	Decoction	Cuevas del Sil (León), SPAIN Vinhais and Bragança (Natural Park of Montesinho), PORTUGAL	Martínez González and Lastra (2006)
<i>Salvia officinalis</i> L.	Sálvia, salva	Sage	C	MED / EVM (cattle, sheep, goats)	Leaves	Decoction	Riofrío de Aliste (and other localities of Northwest of the Zamora province), SPAIN	García Jiménez (2007)
<i>Salvia sclarea</i> L.	Bálsamo, esclareja	Clary sage	W	MED	Leaves and flowers	Decoction	Vinhais and Bragança (Natural Park of Montesinho), PORTUGAL	Carvalho (2010)
<i>Teucrium scorodonia</i> L.	Hierba de lobo, hojas de lobo	Woodland germander	W	MED	Root	Decoction	Miranda do Douro (Natural Park of Douro International), PORTUGAL	Frazaõ-Moreira and Carvalho (2009); Carvalho and Ramos (2012)
LEGUMINOSAE <i>Genista tridentata</i> L. [syn. <i>Pterospartum tridentatum</i> (L.) Willk.]	Carqueja, carqueja	—	W	MED	Leaves	Poultice (warm up and with olive oil or pork lard)	Vinhais and Bragança (Natural Park of Montesinho), and Miranda do Douro (Natural Park of Douro International), PORTUGAL	Frazaõ-Moreira and Carvalho (2009); Carvalho and Ramos (2012)
<i>Trifolium angustifolium</i> L.	Rabo de zorra	Narrow clover	W	MED	Aerial part	Decoction	Northwest of the Basque Country (Biscay and Alava), SPAIN	Menéndez-Buceta et al. (2014)
LYTHRACEAE <i>Lythrum salicaria</i> L.	Salgueira, salgueirinha, salcária, erva-carapau	Purple loosestrife	W	MED / EVM (sheep)	Leaves	Decoction	Valle de Carranza (Biscay), SPAIN	Díaz García (2001)
MELANTHIACEAE <i>Veratrum album</i> L.	Vegambre, verdégambre	White hellebore	W	EVM (cattle, equines)	Flowers (dried)	Decoction	Bragança (Natural Park of Montesinho) and Miranda do Douro (Natural Park of Douro International), PORTUGAL	Frazaõ-Moreira and Carvalho (2009)
OLEACEAE <i>Prunus angustifolia</i> Vahl	Fresno	Narrow-leaved ash	W	MED	Aerial part	Decoction	Valle de Carranza (Biscay), SPAIN	Díaz García (2001)
PAPAVERACEAE <i>Chelidonium majus</i> L.	Cirigüeña, cirigóña	Greater celandine	W	EVM (cattle, equines)	Root	Decoction	Concejo de Aller (Asturias), SPAIN	Martínez González and Lastra (2006)
PLANTAGINACEAE <i>Digitalis purpurea</i> L.	Hoja de la lobera, digital	Foxglove	W	MED / EVM (all livestock)	Bark	Decoction	San Roque de Riomiera (Cantabria), SPAIN	Pardo-de-Santayana (2004)
<i>Plantago major</i> L.	Llantrén	Broadleaf	W	EVM (cattle)	Leaves	Poultice (fresh and crushed)	San Roque de Riomiera (Cantabria), SPAIN	Pardo-de-Santayana (2004)
						Decoction	Concejo de Aller (Asturias), SPAIN	Martínez González and Lastra (2006)

(continued on next page)

Table 1 (continued)

Botanical families Species	Vernacular name (s)	Common English name	Status ^a	Use-category ^b	Part (s) used	Method of preparation	Geographical location	References
RHAMNACEAE <i>Rhamnus cathartica</i> L.	Espino negro	Buckthorn	W	MED	Root Bark	Decoction Decoction	Valle de Carranza (Biscay), SPAIN Valle de Carranza (Biscay), SPAIN	Díaz García (2001) Díaz García (2001)
SANTALACEAE <i>Viseum album</i> L.	Almuertago	Mistletoe	W	MED	Whole plant	Decoction	Valle de Carranza (Biscay), SPAIN	Díaz García (2001)
SCROPHULARIACEAE <i>Serophularia canina</i> L.	Erva-da-quinta	Dog figwort	W	EVM (cattle, sheep, goats) MED / EVM (all livestock)	Leaves Whole plant	Decoction Decoction	Bragança (Natural Park of Montesinho), PORTUGAL El Cerro, Valdelamantanza and Valero (Salamanca), SPAIN	Carvalho (2010) Criado Coca (2010)
URTICACEAE <i>Urtica dioica</i> L.	Hierba lobuna, hierba del lobo, hierba lobera Ortiga, ortiga blanca, ortega, urtica, estruga, sapédiñas, herba do cego	Common nettle	W	EVM (cattle, equines)	Root, leaves	Decoction, poultice	Concejo de Aller (Asturias), SPAIN	Martínez González and Lastre (2006)
				EVM (cattle, goats)	Flowering tops	Decoction	As Pontes de García Rodríguez (La Coruña), SPAIN	Latorre (2008)

^a Status: W, wild; C, cultivated.^b Use-category: MED = ethnomedicine; EVM = ethnovenery medicine.

cal areas, and two communications presented in Conferences. On the other hand, most of the data related to the use of plants in rituals to ward off wolves and to prevent wolf attacks have been found in ethnographic works.

2.2. Plant taxonomy and nomenclature

To avoid ambiguities and errors in botanical scientific nomenclature, pharmacological or biomedical studies on plants require correct spellings, accepted names, author citations and current family designations (see Bennett and Balick, 2014; Rivera et al., 2014). In order to properly use botanical nomenclature we checked The Plant List (www.theplantlist.org/). Botanical families, local names and common English names, if available, are also provided.

3. Results and discussion

3.1. Quantitative ethnobotanical analyses

A total of 39 herbal remedies to treat wolf bites was recorded, all of which are based on the medicinal use of 33 species of vascular plants, belonging to 18 botanical families (Table 1). Moreover, 14 of the reported remedies were exclusively prescribed for the treatment of persons with wolf bites, 12 were used for sheep, goats, equines and cattle, and 13 applied to humans and domestic animals.

Most of the compiled remedies were found in documentary sources published between 2001 and 2016, as only two of all the references providing herbal remedies date from the beginning of the 20th century (Arizaga, 1915; Oria de Rueda et al., 1996).

Plants, preparations and uses mentioned have been registered within the current geographical areas of wolf distribution in the Iberian Peninsula (Fig. 2). The only exceptions are two remedies based on the use of ivy –*Hedera helix* L., *H. hibernica* (G. Kirchn.) Carrière–documented by Silva et al. (2011) in the Serra da Estrela (Portugal). The Estrela mountain range in the centre of Portugal used to be a traditional area of wolf occurrence. During the 1980s the subspecies almost disappeared due to landscape humanisation (Cândido and Petrucci-Fonseca, 2000; Pimenta et al., 2005).

Considering the biological types, most of the taxa cited are herbs (21). Shrubs represent about one-fifth of the useful plant species. Three trees and three climbers were also reported. The three most relevant botanical families regarding their contribution to treat wolf bites in the Iberian Peninsula are Compositae (7 species) and Cistaceae and Lamiaceae (4 species in both cases).

The taxa obtained are mainly wild; however, five plant species cultivated were also reported. According to the phytosociological category to which each plant species belonged, most plants were collected from forest, woodland, scrub vegetation and synanthropic vegetation zones, modified by human activity.

The total number of 39 remedies registered constitutes a very high number of use-reports, considering that among several recent ethnopharmacological studies, herbal medicines for wolf injuries are rarely mentioned. Four studies, carried out in Italy and Albania, were retrieved while searching the European literature on-line. They specifically report that infection from wolf bites in cows and sheep was prevented in Latium and Liguria (Italy) by a poultice made of olive oil and the boiled rhizome of *Veratrum lobelianum* Bernh. (Melanthiaceae) (Viegi et al., 2003), and in Central Italy (Marche, Abruzzo and Latium) by a poultice prepared from the boiled root of *V. nigrum* L. (Guarrera, 2005). To treat animals bitten by wolves, the cinders of aerial parts of *Pteridium aquilinum* (L.) Kuhn (Dennstaedtiaceae) were topically applied with olive oil in the Italian region of Basilicata (Guarrera et al., 2005). Fresh seeds of *Aesculus hippocastanum* L. (Sapindaceae) were crushed and rubbed on the necks of sheep to cure wolf bites in the Rraicë and Mokra areas (Albania) (Pieroni et al., 2015).




Fig. 2. Geographical location of the documented herbal remedies (black spots) and rituals (white spots); in red current geographical distribution of the Iberian wolf.

3.2. Plant parts used and methods of preparation and application

Leaves are the most frequently sought plant parts, accounting for 12 of the claimed medicinal plants. Of a total of nine species the whole plant and for seven the aerial part are used (flowered or not), and of other seven species, their flowers or inflorescences. Few were harvested for their bark, roots and/or rhizomes. The majority of the remedies were harvested for immediate use. Small proportions were reported to be stored for future use.

All herbal remedies documented are based on topical use, including direct application to injured surfaces of the human or animal body.

The predominant method of preparation is that commonly known as *cocimientos* or *cozimentos* (made by boiling plant materials in water for more than 15–20 min) (Table 1). These are water extractions (decoctions) that correspond to the method of preparation of 85% of all the remedies recorded (39). Another practice involves plant materials, such as branches of rosemary (*Rosmarinus officinalis* L.), boiled in wine. Likewise, infusion was the method employed in another registered remedy. In that case, the plant parts are not subjected to boiling water; extraction is accomplished by steeping the material in very hot water. The residual liquid of decoctions and infusions is used externally, soaked in pieces of cloth or directly applied to the skin and wounds.

Some interesting examples of species and practices were registered. For instance, *Carthamus mitissimus* L. was the most useful vulnerary plant documented in Campoo (Cantabria, Spain). The species was claimed to have wound-healing action, being a powerful disinfectant and known for cicatrising wounds in humans and animals (Pardo-de-Santayana, 2004). A method mentioned as both simple and highly effective against severe wounds (wolf bites and injuries) involved inflorescences and whole plant decoctions. These materials were boiled for a long time in order to obtain a solution with a high concentration, that was used to wash the injuries or was applied using wet pads (Pardo-de-Santayana, 2004, 2008). In the early 20th century, shepherds from Panzuegos (La Rioja, Spain) and the Montaña Palentina district (Palencia, Spain) also employed, in the same way, the decoctions of a similar species, *Carthamus carduncellus* L., for the treatment of humans and livestock injured by wolves. It is said that the healing process was short, only taking a few days (Arízaga, 1915; Oria de Rueda et al., 1996).

Poultices were a common technique for providing the plants' active principles to the injured area. Some herbal substances were prepared with soft leaves, previously heated and greased with olive oil, pig fat and/or beeswax; other plant parts were boiled and mashed, used raw and then crushed or ground, and applied as pastes. As many as six remedies are related to these methods of preparation (Table 1).

Some examples cited are the balsamic leaves of *Salvia sclarea* L.

used in north-eastern Portugal. Large and smooth leaves of clary sage (best harvested before the plant came into flower) were heated up in the fireplace, greased with olive oil or pig fat and applied directly to the injured skin (Carvalho, 2010; Carvalho and Ramos, 2012; Frazão-Moreira and Carvalho, 2009). In Mogadouro (NE Portugal), a semi-solid oily preparation was obtained with the bark of elderberry (*Sambucus nigra* L.) crushed with wheat bran and olive oil, or with beeswax, red wine or sheep fat. This ointment was kept in a small glass container and used for its antiseptic qualities, especially in the case of injuries caused by wolves and wild boars (Carvalho and Ramos, 2012).

3.3. Mixtures

Several herbal medicines for healing wolf injuries were prepared combining different species and parts of plants to enhance their potential therapeutic properties, which might be related to synergistic effects.

Gentiana lutea L. and *Juglans regia* L. are well known species and have a long history of use as herbal medicines for their antiseptic and anti-inflammatory characteristics (Font Quer, 1961). The dried underground parts of yellow gentian (i.e. roots and rhizome) and the leaves of the walnut tree were cooked together in water and the remaining liquid was used to clean the wounds in Cuevas del Sil (León, Spain) (García Jiménez, 2007).

In the Concejo de Aller (Asturias, Spain), wounds caused by the wolf on the legs of cows and mares were cleaned and disinfected with decoctions prepared using walnut leaves, broadleaf plantain leaves (*Plantago major* L.) and the roots of *Urtica dioica* L. and *Veratrum album* L. Another folk remedy consisted of the liquid resulting from the decoction of the rhizome of yellow gentian together with the aerial part of *Chelidonium majus* L. and the same two roots mentioned above. In both cases, after washing the wound carefully with these decoctions, a daily poultice of cooked nettles was applied to the wound (Martínez González and Lastra, 2006).

In Valle de Carranza (Biscay, Spain) a decoction was prepared combining three different plant materials: *hojas de lobo* (i.e. the leaves of *Teucrium scorodonia* L.), the bark of the common ash tree (*Fraxinus angustifolia* Vahl.) and *rabos de zorra* (i.e. the aerial parts and inflorescences of *Trifolium angustifolium* L.). The extract obtained was kept in a bottle for subsequent use. However, the solution was heated before being used with a pad or cloth to disinfect the wounds (Díaz García, 2001).

Use-reports from Bragança (Portugal) also highlight some traditional healing practices combining two different herbal remedies: for instance, the external use of clary sage leaves (*Salvia sclarea*) previously heated and greased combined with a decoction of dog figwort leaves (*Scrophularia canina* L.) (Carvalho, 2010). Clary sage leaves are vulnerary, relieving pain and sterilising wounds (Peana and Moretti, 2002; Gülcin et al., 2004a; Džamić et al., 2008; Sienkiewicz et al., 2015); the water extract of figwort leaves cleans, protects from re-infection and promotes cicatrisation (Santos Galíndez et al., 2002; Venditti et al., 2016).

3.4. Potential pharmacological uses

Wound healing is a continuous and complex pathophysiological process divided in well defined phases: haemostasis, inflammation, cellular migration and proliferation, protein synthesis and wound contraction (Dreifke et al., 2015; Velnar et al., 2009).

Herbal approaches to the treatment of wounds focus the use of healing species with antiseptic, anti-inflammatory and analgesic properties (Merino and Blanquer, 2016; Nagori and Solanki, 2011; Raina et al., 2008). There is currently a great interest in isolating and identifying active constituents from plant species and in screening those active compounds for pharmacological activities. Many ethnopharmacological studies of wound healing plants have been carried out

around the world (e.g. Adetutu et al., 2011; Kumar et al., 2007). Several papers on particular medicinal plant species controlling various phases of the wound healing process are also available in the scientific literature (e.g. Mukherjee et al., 2000; Rashed et al., 2003).

Medicinal plants beneficial to skin and wound healing concern antifungal, antiviral and antibacterial activities against infections and activity against inflammatory disorders affecting skin (Mantle et al., 2001; Nagori and Solanki, 2011; Pazyar et al., 2014). Moreover, wound healing protocols depend on each type of injury, considering the level of contamination, extension and size, as well as the cause and depth of the wound (Kasote et al., 2015; Pazyar et al., 2014; Pereira and Bártolo, 2016). In the case of clean wounds (i.e. injuries with a low level of contamination, where skin is likely to heal without complications), apart from suturing, washing with herbal preparations with the above-described properties may be sufficient. On the other hand, infected wounds (exhibiting clinical signs of infection, as well as the presence of pathogenic microorganisms) will need more effective treatments, such as poultices or compresses prepared with extracts of the same plants. Minor and small wounds may be treated with plants with antiseptic properties (antibacterial, antifungal and antiviral). Wounds requiring a healing process are treated with antiseptic species too, but plants and herbal preparations with powerful cicatrisation action are also fundamental to speed the rate of healing and reduce the risk of infection (Merino and Blanquer, 2016).

A number of secondary metabolites-active compounds isolated from plants and tested in animal models (in vivo) have been reported as active principles responsible for facilitating wound healing. Some of the most important ones are phenolic compounds, tannins, alkaloids (monoterpens, triterpenes), flavonoids and secoiridoid glycosides, among others (Dreifke et al., 2015; Kumar et al., 2007). For example, tannins have been also reported to have antiviral and antibacterial effects (Akiyama et al., 2001). There is a strong association between tannin content and the effects popularly attributed to wound healing and anti-inflammatory plants (Araújo et al., 2008). Tannins might be included in haemostatic formulations, not only to check bleeding by contracting the tissues but also to promote wound healing internally, as they have the ability to form a protective layer over the exposed tissue preventing reinfection (Ashok and Upadhyaya, 2012). Triterpenes proved to increase the rate of successful healing, as well as of epithelialisation and collagen deposition, because they are effective in accelerating re-epithelialisation and improving scar formation (Agra et al., 2015; Kindler et al., 2016). Plant flavonoids have antioxidant activity that destroy free radicals and favour wound healing, as described in several studies (e.g. Rasik and Shukla, 2001).

Most frequently, wolves' attacks on flocks and free-ranging livestock involve the combined effort of the group or pack, in a similar way they predate on large wild ungulates (Grande del Brío, 2000). Sometimes, they capture domestic animals by the neck or throat and many of these animals perish; other times hindquarters (including tail, groin and perineal regions, vulva and scrotum) are the main focus of the attempts; face, front legs and flanks are also potentially highly exposed to wolf attacks. Wolf nasty bites usually cause damage in the underlying tissues, frayed wounds and lacerations, crushed skull, broken spine, disembowelment and massive tissue damage. Injured animals experience blood loss, get in shock, lie down and eventually succumb. However, under certain conditions some more or less seriously wounded animals can survive (Acorn and Dorrance, 1990; Vantassel, 2012).

The literature systematically searched provided ethnopharmacological information about some of the reported vascular plant species used in the Iberian Peninsula to treat wolf bites. Table 2 summarises the known therapeutic activities of these species related to the wound healing process, based on laboratory and clinical evidence, according to the reviewed literature.

Therefore, considering the different stages of the wound healing process and the most relevant therapeutic properties and effects

promoting and speeding the cure, it seems that most of the documented plants and herbal preparations used in the Iberian Peninsula might control and reduce the inflammatory phase (antiseptic activity), and promote the onset of the proliferative phase, improving wound contraction, re-epithelialisation, angiogenesis and collagen deposition.

Several authors provided scientific evidence that many of the species listed in Table 2, or at least other species in the same genera, have analgesic activity. Nine of the reported plant species present antinociceptive effects (inhibits nociception, the sensation of pain). For instance, *Plantago major* (e.g. Gonçalves and Romano, 2016) and *Rosmarinus officinalis* (e.g. Rahbardar et al., 2017). Other works highlight important bioactive properties of some species and there are clinical data available for their medicinal and veterinary uses.

The Committee on Herbal Medicinal Products (HMPC) of the European Medicines Agency (EMA) develop monographs in accordance with European Union (EU) regulations, covering the therapeutic uses and safe conditions of well-established and/or traditional use for herbal substances and preparations (EMA, <http://www.ema.europa.eu/ema/>). Some of the reported species used within the Iberian Peninsula for healing wounds caused by wolves are included in EU Monographs and has a simplified registration procedure for traditional herbal medicinal products in EU Member States, considering the HMPC's scientific opinion on safety and efficacy data about such herbal substances and preparations. Moreover such monographs inform of interesting constituents and possible pharmacodynamic actions although they might not play a role for the already known traditional use. Examples are *Fraxinus angustifolia* (EMA, 2012a, 2012b), *Genciana lutea* (EMA, 2010c, 2010d), *Juglans regia* (EMA, 2014a, 2014b), *Salvia officinalis* L. (EMA, 2010a, 2010b) or *Rosmarinus officinalis* (EMA, 2011b).

Several studies and monographs consulted concern some of the species listed in Table 2 and show that such plants are used worldwide in the treatment of a number of diseases and particularly skin diseases. The chemical constituents of selected examples from Table 2 and their biological activities are described considering different authors findings.

Roman chamomile flowers (*Chamomillae romanae flos*) are commonly used within the Iberian territory. Preparations derived from the flower heads of *Chamaemelum nobile* (L.) All. are externally applied to treat skin inflammation and irritation; they have also emollient properties and are protective for cracks, abrasions, frostbites, chaps and insect bites (Edwards et al., 2015; EMA, 2012f). Besides many medicinal and veterinary purposes and different solid and liquid formulations for oral or topical application, the consumption of *C. nobile* flower heads infusion (capitula) is also widespread as recreational herbal tea or as an ingredient to prepare other traditional beverages (e.g. Carvalho, 2010; Pardo-de-Santayana, 2008). The species contains an essential oil (up to about 1.75%) with complex composition: esters of angelic and butyric acids, sesquiterpene lactones of the germacranolide type, mainly nobilin and 3-epinobilin. The dried flowers have many terpenoids and flavonoids, as well as 1–2% volatile oils including alpha-bisabolol, alpha-bisabolol oxides A and B, and matricin (Edwards et al., 2015; Srivastava et al., 2010). Major constituents of Roman chamomile are related with its medicinal properties, particularly with anti-inflammatory effects. Moreover, trials in human volunteers showed that such flavonoids and essential oils entered into the deeper skin layers, an important characteristic for a topical anti-inflammatory. Srivastava et al. (2010) reviewed the use of chamomile in traditional medicine and described its curative and preventive properties to enhance wound healing, as well as the results of some trials. The findings reported highlight the efficacy of chamomile extracts on wound drying and in speeding epithelialisation, in addition to antimicrobial activity against various microorganisms. According these authors, such effects validate the use of Roman chamomile in wound management (Srivastava et al., 2010).

Chelidonium majus, greater celandine, is commonly known for its

Table 2
Scientific evidence of medicinal properties of vascular plants used in the Iberian Peninsula to treat wolf bite injuries.

Plant species	Therapeutic activities ^a						References (chronologically arranged)
	WH	AS	AI	AT	AO	AA	
<i>Carlina hispanica</i> ^b	+	+	+	+	+	+	Ivancheva et al. (2006) ^b ; Hermann et al. (2011) ^b
<i>Carthamus (C. cardunculus and C. mitissimus)</i> ^b	+	+	+	+	+	+	Almeida et al. (2001) ^b ; Ivancheva et al. (2006) ^b ; Mothana et al. (2010) ^b ; Kueite et al. (2012) ^b ; Asgarpanah and Kazemivash (2013) ^b ; Turghaybayeva et al. (2014) ^b
<i>Centaura (C. lagasca and C. ornata)</i> ^b	+	+	+	+	+	+	Barnbour et al. (2004) ^b ; Borchardt et al., 2008 ^b ; Koca et al. (2009) ^b ; Khammar and Djeddi (2012) ^b ; Zater et al. (2016) ^b
<i>Chamaemelum nobile</i>	+	+	+	+	+	+	Chao et al. (2000); Bai et al. (2009); Srivastava et al. (2010); EMA (2012b); Zhao et al. (2014); Edwards et al. (2015); Kazenian et al. (2015); Al-Snafi (2016)
<i>Chelidonium majus</i>	+	+	+	+	+	+	Kokoska et al. (2002); Zuo et al. (2008); WHO (2010); EMA (2012e); Gynovian and Trchoumian (2017)
<i>Cistus albidus</i> ^b	+	+	+	+	+	+	Papaeftimioú et al. (2014) ^b
<i>Cistus populifolius</i>	+	+	+	+	+	+	De Andrés et al. (1999); Barragón-Catalán et al. (2010); Papaeftimioú et al. (2014); El Euchi et al. (2015); Habibi et al. (2015); Mahmoudi et al. (2016); Rebaya et al. (2016)
<i>Cistus salviifolius</i>	+	+	+	+	+	+	Abouzeed et al. (2013); Tomás-Menor et al. (2013); Papaeftimioú et al. (2013); Tomás-Menor et al. (2013); Papaeftimioú et al. (2013); Gynovian and Trchoumian (2017)
<i>Digitalis purpurea</i>	–	–	–	–	–	–	–
<i>Fraxinus angustifolia</i>	+	+	+	+	+	+	EMA (2012a) (2012b); Moulaoui et al. (2015)
<i>Genista tridentata</i>	+	+	+	+	+	+	Pinela et al. (2011); Martins et al., 2015; Ares et al. (2016)
<i>Gentiana lutea</i>	+	+	+	+	+	+	Mathew et al. (2004); Ortíz et al. (2006); WHO (2007); Singh (2008); Šavikin et al. (2009); EMA (2010c) (2010d); Azman et al. (2014); Edwards et al. (2015)
<i>Hedera helix</i>	+	+	+	+	+	+	Geddiremen et al. (2005); Lutsenko et al. (2010); Uddin et al. (2011); Rashed (2013); EMA (2015, 2016)
<i>Juglans regia</i>	+	+	+	+	+	+	Vanaclocha and Cañigueral (2003); Santos et al. (2013); Sharma et al. (2013); EMA (2014a) (2014b); Ponth et al. (2016)
<i>Jurinea humilis</i> ^b	+	+	+	+	+	+	Ortíz et al. (2011) ^b
<i>Lythrum salicaria</i>	+	+	+	+	+	+	Becker et al. (2005); Tunaliér et al. (2007); Borchardt et al. (2008a); (2008b); Humadi and Istidor (2009); Vafi et al. (2016)
<i>Osmunda regalis</i>	+	+	+	+	+	+	Banerjee and Sen (1980); Thomas (2011)
<i>Plantago major</i>	+	+	+	+	+	+	Samuelson (2000); Vanaclocha and Cañigueral (2003); Mahmood and Phipps (2006); Stanislavjević et al. (2008); WHO (2010); Gonçalves and Romano (2016)
<i>Rhamnus cathartica</i>	–	–	–	–	–	–	Borchardt et al. (2008a); Locatelli et al. (2011); Hamed et al. (2015)
<i>Rosmarinus officinalis</i>	+	+	+	+	+	+	Vanaclocha and Cañigueral (2003); Moreno et al. (2006); Bozin et al. (2007); Takaki et al. (2008); Martínez et al. (2009); WHO (2009); Abu-Al-Bassel (2010); EMA (2011b); Lai et al. (2012); Abuzeid et al. (2014); Pazvar et al. (2014); Edwards et al. (2015); Rahbardar et al. (2017)
<i>Salvia officinalis</i>	+	+	+	+	+	+	Baricic et al. (2001); Vanaclocha and Cañigueral (2003); Mitić-Culařić et al. (2005); Bozin et al. (2007); EMA (2010a) (2010b); WHO (2010); Abouzeed et al. (2013); Edwards et al. (2015); Kadhim et al. (2016); Karimzadeh and Farahpour (2017)
<i>Sativia sclarea</i>	+	+	+	+	+	+	Peana and Moretti (2002); Gülin et al. (2004a); Džanić et al. (2008); Sienkiewicz et al. (2015)
<i>Sambucus nigra</i>	+	+	+	+	+	+	WHO (2004); EMA (2008a); Hearst et al. (2010); Mogosanu et al. (2014); Edwards et al. (2015); Gynovian and Trchoumian (2017)
<i>Scrophularia canina</i> ^b	+	+	+	+	+	+	Santos Galindez et al. (2002); Venditti et al. (2016)
<i>Teneturia scorodonia</i> ^b	+	+	+	+	+	+	Alatiyakoli et al. (2012) ^b ; Tabatabaei Yazdi and Alizadeh Behbahani (2013) ^b
<i>Trifolium angustifolium</i> ^b	+	+	+	+	+	+	Kolodziejczyk-Czepas (2012) ^b ; Renda et al. (2013) ^b
<i>Tuberaria lignosa</i>	+	+	+	+	+	+	Pinela et al. (2012)
<i>Urtica dioica</i>	+	+	+	+	+	+	Almeida et al. (2001); Vanaclocha and Cañigueral (2003); Gülcin et al. (2004b); WHO (2004, 2010); EMA (2008b) (2011a) (2012c) (2012d); Modarresi-Chahardehi et al. (2012); Salehzadeh et al. (2014); Edwards et al. (2015); Babaei et al. (2017)
<i>Veratrum album</i>	+	+	+	+	+	+	Tosun et al. (2005); Režić and Ferrier (2011); Gynovian and Trchoumian (2017)
<i>Viscum album</i>	+	+	+	+	+	+	Ertlik et al. (2003); Orhan et al. (2006); Sengül et al. (2009); Hegde et al. (2011); EMA (2013); Gynovian and Trchoumian (2017)

^a Therapeutic activities: WH = wound-healing (vulnery, cicatrizing, skin regeneration, angiogenic); AS = antiseptic (antibiotic, antimicrobial, bacteriostatic, bacteriocidal, antifungal, fungistatic, antiviral); AI = anti-inflammatory; AT = astringent; AO = antioxidant; AA = analgesic (including antinociceptive effects); HS = haemostatic.

^b Information provided for other species of the same genus.

orange coloured latex which topical traditional use against warts, moles, callus and corns is frequent (EMA, 2012e). Major constituents of dried aerial parts (*Chelidonium herba*) are over 20 different alkaloids, belonging to the benzylisoquinoline group. The plant also contains flavonoids, saponins, organic acids, vitamins, carotenoids, and several hydroxycinnamic acid derivatives including caffeoylelmalic, chelidonic, malic and citric acids, as well proteolytic enzymes detected in the latex (EMA, 2012e; WHO, 2010). Experimental pharmacology describes anti-inflammatory activity in vitro during a test conducted to determine constituent's ability to inhibit enzymes in the inflammatory cascade (WHO, 2010).

Fraxinus angustifolia provides different traditional herbal medicinal products: the leaves are reported for minor joint pain (EMA, 2012a, 2012b), and the bark as diuretic and for gastrointestinal complaints in veterinary (Moulaoui et al., 2015; Viegi et al., 2003). The active components of leaves and bark ethanolic extracts and their content in polyphenols were assessed by HPLC to elucidate the polyphenolic composition of *F. angustifolia* extracts, and to evaluate their efficacy in wound healing. Quercetin, catechin, rutin and tannic acid were identified as the main components. *In vivo* results showed that the highest antioxidant and anti-inflammatory effects were provided by the polyphenolic phytocomplexes that consequently promote wound healing (Moulaoui et al., 2015).

Gentiana lutea is another species documented by WHO and EU monographs on selected medicinal plants (EMA, 2010c, 2010d; WHO, 2007). Plant materials correspond to dried underground organs, roots and rhizomes (*Gentianae radix*). Different types of extracts proved to have antimicrobial activity, i.e. inhibition of bacteria and fungi growth (WHO, 2007). It seems that gentian leaves and roots exhibit considerable antioxidant properties, expressed either by their capability to scavenge DPPH or superoxide radicals; however further research is needed to confirm these observations (EMA, 2010c, 2010d). According to Oztürk et al. (2006) and Singh (2008) active principles responsible for wound healing are: gentianine a monoterpenoid alkaloid with anti-inflammatory, analgesic and antibacterial properties; gentianadine, an alkaloid that possesses anti-inflammatory activity; and sweroside, a secoiridoid glycoside with antimicrobial and antifungal action. Savikin et al. (2009) point out that the methanolic extracts of flowers and leaves of *G. lutea* have a good antimicrobial effect on a variety of Gram-positive and Gram-negative bacteria as well as on the yeast *Candida albicans*, which is related with their secoiridoid glycoside content. Antioxidant activity has been attributed to the phenolic content of gentian extracts, especially gentiopicroside, which also exhibited analgesic properties (Edwards et al., 2015).

Juglans regia leaves (drug *Juglandis folium*) are usually boiled in water to make a decoction which is applied to mild and superficial skin inflammations. Walnut leaves contain approximately 10% tannins and about 3.4% of flavonoids (EMA, 2014a, 2014b). Santos et al. (2013) identified and quantified twenty-five phenolic compounds in walnut leaves decoctions prepared for topical applications, as also the presence of pro-cyanidins, taxifolin derivatives, and tocopherols. According to Panth et al. (2016), the volatile oil obtained from *J. regia* leaves and its key constituents (β -pinene, α -pinene, limonene, germacrene D and β -caryophyllene) exhibited strong activity against bacteria (primarily Gram-positive bacteria). Wynn and Fougère (2007) suggest veterinary indications for walnut leaves, applied topically for focal inflammation and skin lesions.

Plantago major (broadleaf plantain) and other species of the genus *Plantago* are plants which seeds, leaves and aerial parts have widespread medicinal uses. Plantain fresh leaves have been used externally for a long time as an antiseptic for the treatment of skin irritations and skin inflammations (Samuelson, 2000). WHO monographs (WHO, 2010) document the dried leaves of plantain as useful plant material with different therapeutic properties: antimicrobial due to the pectic polysaccharide fraction that protects against bacterial infections; antinociceptive and anti-inflammatory as ursolic and oleanolic acids

were isolated from a hexane extract of dried leaves using bioactivity-directed fractionation. Such components have important wound-healing activity. The pectin-type polysaccharide fraction (PTPF) isolated from the leaves of *P. major* is a potent complement activator with an activity of the same order of magnitude as that of aggregated human immunoglobulin G (WHO, 2010). *Plantago major* might have also veterinary interest being topically used in poultices for skin wounds (Wynn and Fougère, 2007).

Rosmarinus officinalis, known as rosemary, has great aromatic and medicinal potential. The most valuable plant materials are the leaves and the flowering aerial parts that are used in solid and liquid herbal preparations for skin affections, among other therapeutic purposes (EMA, 2011b). Both leaves and the essential oil obtained by steam distillation of the flowering aerial parts are considered by EMA (2011b) and WHO (2009) monographs. Experimental pharmacology shows antimicrobial and inflammatory activities as well as antioxidant and pro-oxidant properties both in vitro and in vivo (WHO, 2009; Edwards et al., 2015). Rosemary essential oil exhibit peripheral antinociceptive activity in animal models (Takaki et al., 2008). Besides the chief constituents of the essential oil (e.g. camphor, limonene and others) and phenolic compounds (e.g. flavonoids and phenolic acids), tricyclic diterpenes such as rosmarinidiphenol, carnosol and carnosic acid are interesting constituents. Carnosol presented potent antioxidant activity when compared with other compounds extracted from rosemary. Moreover, possible mechanisms for anti-inflammatory activity are related with carnosol that suppresses nitric oxide production and inducible nitric oxide synthase gene expression by inhibiting activation of nuclear factor-B (WHO, 2009; Edwards et al., 2015).

Salvia officinalis leaves (*Salviae folium*) are traditionally used as a vulnerary and an antiseptic. They contain thujone, cineol and camphor, and other constituents such as tannins, diterpene bitter principles, triterpenes, and flavonoids (such as luteolin and salvigenin). Several studies performed show positive antibacterial and anti-inflammatory effects of *Salvia officinalis* extracts. Specifically, flavonoids and phenolic acids such as rosmarinic acid found in sage may be responsible for some of the biological activities, as these compounds have known antiviral, antibacterial, anti-inflammatory and antioxidant properties (see Peterson and Simmonds, 2003). Sage oil has antimicrobial properties; the oil shows inhibitory activity against Gram-positive and Gram-negative bacteria and against a range of fungi. Most frequent dermatological therapeutic indications concern minor skin inflammation (EMA, 2010a, 2010b; WHO, 2010; Edwards et al., 2015). Karimzadeh and Farahpour' recent research (2017) reports that the hydroethanolic leaves extract has a high total flavonoid and phenolic content and antioxidant capacity. The topical application of ointments with different doses of hydroethanolic sage leaf extract improved wound healing process in rats, as significantly increased the percentage of wound contraction, and promoted the period of re-epithelialisation, and the new vessel formation and fibroblast distribution. Thus, it can be considered an appropriate potential compound for clinical application (Karimzadeh and Farahpour, 2017).

Urtica dioica dried leaves and rhizome, as well as fresh and dried aerial parts harvested during the flowering period (*Urticae folium* and *Urticae herba*), are reported for their traditional medicinal applications (EMA, 2008b, 2011a, 2012c, 2012d; WHO, 2004). Major constituents of underground organs (*Urticae radix*) are many compounds with different polarity and belonging to various chemical classes, such as triterpenes, phenylpropanes, lignans, coumarins, ceramides, sterols and lectins (WHO, 2004; Edwards et al., 2015). Principal components of *Urticae herba* are flavonoids, caffeic acid esters (mainly caffeoylelmalic acid, chlorogenic, neochlorogenic), chlorophyll degradation products and carotenoids, coumarins, triterpenes and sterols, leukotrienes (EMA, 2008b, 2012c; Edwards et al., 2015). Experimental pharmacological assays proved anti-inflammatory activity; an ethanol extract inhibited the activity of human leukocyte elastase and reduced the amount of the enzyme released by activated polymorphonuclear

granulocytes during the inflammatory response (WHO, 2004, 2010). Topical ointments prepared from the extracts of *Urtica dioica* and *Sambucus ebulus* L. and their combination showed strong wound healing properties. It is postulated that a synergistic effect may exist between the two extracts since the combination 2% showed better results than the sole extracts (Babaei et al., 2017).

Viscum album L. lipophilic extract contains pharmacologically active pentacyclic triterpenes that are known to exhibit wound healing activity. Preliminary clinical observations indicate that this extract was able to influence cutaneous wound healing in vivo. Data support the casual observation that it might modulate wound healing related processes in vivo (Kuonen et al., 2013).

Among the plant species for which there is no clinical data available regarding medicinal and veterinary uses, highlight *Cistus albidus* L., which chemical composition is one of the most studied within the genus *Cistus*. This perennial shrub has been highly reported for its traditional use in ethnomedicine (Alarcón et al., 2015; Gonçalves et al., 2013). Some of the compounds identified among their volatiles such as the phenolic monoterpenes thymol or carvacrol, are well known as antimicrobial compounds, exhibiting significant growth-inhibitory effects on bacteria (Maccioni et al., 2007). Gonçalves et al. (2013) experimental research showed that *C. albidus* extracts are exceptionally effective and potentially new sources of antioxidants, confirming the folk use of species, particularly for the prevention and treatment of diseases that are known to be caused or accelerated by oxidative stress.

On the other hand, the traditional use of the leaves of foxglove (*Digitalis purpurea* L.) or of woodland germander (*Teucrium scorodonia*) as a healing remedy for wolf bites and infected wounds seems to be an exception. For example, foxglove preparations were applied to humans and domestic animals bitten by wolves in different Spanish regions, as reported by Pardo-de-Santayana (2004), Verde et al. (2008) and Criado (2010), and woodland germander is one of the ingredients present in a patented homeopathic formulation (<https://www.google.com/patents/WO2013189908A1?cl=en>) for healing wounds. Moreover, in many Spanish regions, such as the Basque Country, the common name of woodland germander, *herba de lobo* (literally “wolf herb”) is linked with the carnivore presence or its injuries’ treatment (Díaz García, 2001; Menéndez-Baceta et al., 2014). Besides *T. scorodonia*, other five plants also used for healing wolf bites are locally known by popular names related with wolves, e.g. *lobera*, *herba lobera*, *herbalobuna*, *erva-loba* (see Table 1). However, assays to determine the antimicrobial activity of foxglove extracts (e.g. Dogruoz et al., 2008) or of woodland germander essential oils (e.g. Djabou et al., 2013) did not confirm bacterial inhibition. Hydroalcoholic extract of *Teucrium polium* L. has antimicrobial effect and decreases the duration of complete wound healing process in rats (Alahvakoli et al., 2012; Bahramikia and Yazdanparast, 2012; Tabatabaei Yazdi and Alizadeh Behbahani, 2013), emphasising the importance of ethnopharmacological approaches and the development of new studies.

3.5. Incertae sedis

A common practice of Spanish shepherds from Zafra (Badajoz) and Tierra de Cameros (La Rioja) consisted of making plasters with wood ashes that were then applied to wolf bites. It was claimed that such preparations dried and healed the injury (Elías and Muntián, 1989; Penco, 2005). Ashes, salt and water were the main ingredients to prepare a decoction that was used to cure bites in Asturias (Spain) (Castañón, 1976; Junceda Avello, 1987).

In Sardinia (Italy), Piluzza et al. (2015) have documented the use of ash in water, wrapped with a cloth around the wounds from red fox – *Vulpes vulpes* (Linnaeus, 1758)– or dog bites. In fact, several tests conducted with animal models have confirmed that the topical administration of ashes (corresponding to complete burning of woody materials) accelerates the wound healing process, enhances epithelialisation and reduces infection (Hamid and Shaikh, 2009; Shaikh and

Hamid, 2009). The analysis of different types of wood ashes revealed that they contain the appropriate combination of metals required to heal a wounded area faster (Hamid and Shaikh, 2009; Shaikh and Hamid, 2009).

3.6. Plant-based traditional practices to ward off wolves

Until recently and within the Iberian territories wolves were scared away by several peculiar methods such as: tying a red cloth ribbon around the lambs’ necks (Aguirre Sorondo, 1996; Penco, 2005); playing the zambomba, a traditional musical instrument similar to a friction drum (Barandarián and Manterola, 2000; Goñi, 1975); or reciting a set of prayers to St. Antony to get protection from wolves (e.g. Domínguez Moreno, 1994; Mariño Ferro, 1985; Prieto, 1949; Valverde and Teruelo, 2001; Vidal-González, 2013). Moreover, the literature also refers to the use of plant species in some practices to prevent attacks by wolves (see Fig. 2). In the Sierra de la Culebra (Zamora, Spain), livestock was kept temporarily in the mountain inside the so-called corralas or corrales. These safe enclosures were erected with stone walls and topped with a vegetal parapet made of heather species (*Erica arborea* L. and *E. australis* L., Ericaceae), gorse (*Genista* spp., Fabaceae) and/or elm-leaf blackberry (*Rubus ulmifolius* Schott., Rosaceae) (Dacosta, 2010). In the case of the Pallars region (Lérida, Spain), shepherds carved thick pointed branches of box wood (*Buxus sempervirens* L., Buxaceae) to use in enclosures to prevent or, at least, to hinder the carnivores (Agelet, 1999). In Valencia (Spain) shepherds believed that the sound produced by flutes made with the wood of *lledoner* (the honeyberry or nettle tree, *Celtis australis* L., Cannabaceae) would ward off wolves (Zurriaga, 2012). These last two use-reports are very interesting, as they are practices to ward off wolves documented in geographical areas where this large carnivore disappeared in the early twentieth century (see Fig. 2) (Grande del Brío, 1984, 2000; Massip, 2011).

3.7. Plant-based rituals for the prevention of the wolf attacks

Different magic-religious rituals, based on the use of certain very fragrant plant species, are among several traditional practices that were used to protect livestock and people from wolves. Thus, in Teverga (Asturias, Spain) bunches of fennel (*Foeniculum vulgare* Mill., Apiaceae) were left on the streets during the Feast of Corpus Christi. It was believed that the branches would be blessed as the priest wandered the streets carrying the custodia. Later, on the eve of St. John’s Day, these holy fennel bunches would feed domestic animals that would be protected against wolves (Castañón, 1966).

On the same day, St. John’s Eve, the inhabitants of Ahigal (Cáceres, Spain) implemented a *sahumerio* made of aromatic plants. This Spanish word, which is derived from the Latin *suffumāre*, is related to the combustion of aromatic biotic material which releases fragrant smoke and is used to purify. Bunches of *cantueso* (*Lavandula pedunculata* (Mill.) Cav., Lamiaceae) and rosemary, previously blessed during the procession at Corpus Christi, were burned and the smoke spread throughout the house, the barn and the stable. It was considered that being exposed to the blessed smoke would keep everybody and everything safe from wolf attacks. For the same purpose, domestic animals were passed over ashes at dawn to get protection against wolves (Grande del Brío, 1984).

The purifying power and the beneficial virtues of fire and smoke were also claimed to prevent wolf attacks in El Bierzo (León, Spain). Piles of burning fragrant plants were used to frighten the beasts, such as *ruda* (rue, *Ruta* spp., Rutaceae), *caléndula* (field marigold, *Calendula arvensis* M. Bieb., Compositae), *cantueso* or rosemary (Criado, 2012). Furthermore, in Tierra de Granadilla (Cáceres, Spain), making bonfires using rosemary near houses was a common practice during St. John’s Eve. The next day, the livestock would walk over the ashes and such cinders would be distributed over the sheepfolds (Domínguez Moreno, 1991, 1994).

4. Conclusions

Despite the decline of the Iberian wolf during over the last few decades, wolves are still in the imaginary of rural communities that perceive this large carnivore as both a diabolic creature and a mythic and benign animal. The wolf-related cultural dimension is of great interest in terms of conservation strategies and local knowledge is intrinsically bound up with cultural legacies. Thus, it has great significance to assess local knowledge about 39 plant-based remedies traditionally used for healing injuries from wolves in humans and domestic animals within the Iberian territory, including all the areas of wolf distribution, as well as to describe the practices and rituals for improving preventive methods and warding off wolves that involved eleven plants. Additionally, the gathered information may contribute to the conservation of the Iberian biocultural patrimony by recognising, understanding and valuing natural resources, as well as rural beliefs and practices.

The vascular plants reported are claimed to have analgesic and antiseptic properties, to reduce inflammation and improve wound contraction, re-epithelialisation, angiogenesis and collagen deposition. Some specific uses currently cannot be explained as they concern the magic-ritual domain.

The documented species are culturally important plants and may be potential sources of phytochemicals. Although it was found that certain species mentioned for healing injuries from wolves are effective in the wound-healing process, there is clearly a need for further phytochemical and pharmacological research and bioassays, in order to better understand the properties and effects of these interesting plant resources.

Plants and their extracts have great potential for the management and treatment of wounds in domestic animals. Phytomedicines for wound healing are not only less expensive and affordable but are also purportedly safe as hyper sensitive reactions are rarely encountered with the use of these agents. Nevertheless, there is a need for scientific validation, standardization and safety evaluation of the plants used in Ethnoveterinary Medicine (EVM) before they could be recommended for treatment of the wounds. Further clinical studies of phytoconstituents should be carried out to determine safety, efficacy and doses. The combination of traditional and modern knowledge can produce better drugs for wound healing with fewer side effects.

Acknowledgments

We wish to thank several anonymous contributors for their help in improving the manuscript. We would also like to show our gratitude to Rachel Fell for her professional English technical support.

References

- Abouzeed, Y.M., Elfahem, A., Zgheel, F., Ahmed, M.O., 2013. Antibacterial in-vitro activities of selected medicinal plants against methicillin resistant *Staphylococcus aureus* from Libyan environment. *J. Environ. Anal. Toxicol.* 3, 7.
- Abrahamian, F.M., Goldstein, E.J.C., 2011. Microbiology of animal bite wound infections. *Clin. Microbiol. Rev.* 24 (2), 231–246.
- Abu-Al-Basal, M.A., 2010. Healing potential of *Rosmarinus officinalis* L. on full-thickness excision cutaneous wounds in alloxan-induced-diabetic BALB/c mice. *J. Ethnopharmacol.* 131 (2), 443–450.
- Abuzeid, N., Kalsum, S., Koshy, R.J., Larsson, M., Glader, M., Andersson, H., et al., 2014. Antimycobacterial activity of selected medicinal plants traditionally used in Sudan to treat infectious diseases. *J. Ethnopharmacol.* 157, 134–139.
- Aceituno-Mata, L., et al., 2014. Metodología para la elaboración del inventario Español de los Conocimientos Tradicionales relativos a la Biodiversidad. In: Pardo-de-Santayana, M., Morales, R., Aceituno, L., Molina, M. (Eds.), *Inventario Español de los Conocimientos Tradicionales relativos a la Biodiversidad. Primera Fase: introducción, metodología y fichas*. Ministerio de Agricultura, Alimentación y Medio Ambiente, Madrid, 31–49.
- Acorn, R.C., Dorrance, M.J., 1990. Methods of investigating predation of livestock. Alberta Agriculture and Rural Development, Edmonton.
- Adetutu, A., Morgan, W.A., Corcoran, O., 2011. Ethnopharmacological survey and in vitro evaluation of wound-healing plants used in South-western Nigeria. *J. Ethnopharmacol.* 137, 50–56.
- Agelet, A., 1999. Estudis d'Etnobotànica farmacèutica al Pallars (Ph.D.thesis). Facultad de Farmacia, Universidad de Barcelona, Barcelona (in Catalan).
- Agra, L.C., Ferro, J.N.S., Barbosa, F.T., Barreto, E., 2015. Triterpenes with healing activity: a systematic review. *J. Dermatol. Treat.* 26 (5), 465–470.
- Aguirre Sorondo, A., 1996. Veterinaria popular vasca. Orain, Bilbao.
- Aires, A., Marrinhos, E., Carvalho, R., Dias, C., Saavedra, M.J., 2016. Phytochemical composition and antibacterial activity of hydroalcoholic extracts of *Pterospartum tridentatum* and *Mentha pulegium* against *Staphylococcus aureus* isolates. *BioMed. Res. Int.* 2016, article ID 5201879.
- Akiyama, H., Fujii, K., Yamasaki, O., Oono, T., Iwatsuki, K., 2001. Antibacterial action of several tannins against *Staphylococcus aureus*. *J. Antimicrob. Chemother.* 48 (4), 487–491.
- Alahtavakoli, M., Vazirinejad, R., Ansari Jaber, A., Negahban, T., Mashayekhi, H., Nazari, M., Ghoreshi, S., Nematollahi, F., 2012. Effect of *Teucrium polium* extract on skin wound healing in rat. *HJM* 16 (1), 17–24.
- Alarcón, R., Pardo-de-Santayana, M., Priestley, C., Morales, R., Heinrich, M., 2015. Medicinal and local food plants in the south of Alava (Basque Country, Spain). *J. Ethnopharmacol.* 176, 207–224.
- Almeida, R.N., Navarro, D.S., Barbosa-Filho, J.M., 2001. Plants with central analgesic activity. *Phytomed.* 8 (4), 310–322.
- Al-Snafi, A.E., 2016. Medical importance of *Anthemis nobilis* (*Chamaemelum nobile*)—a review. *AJPST* 6 (2), 89–95.
- Álvares, F., Blanco, J.C., 2014. Recovering traditional husbandry practices to reduce wolf predation on free-ranging cattle in Iberia. *CDP News* 10, 4–9.
- Álvares, F., Alonso, P., Sierra, P., Petrucci-Fonseca, F., 2000. Os fojos dos lobos na Península Ibérica. Sua inventariação, caracterização e conservação. *Galemys* 12, 57–77.
- Álvares, F., Domingues, J., Sierra, P., Primavera, P., 2011. The cultural dimension of wolves in the Iberian Peninsula: implications of ethnozoology in conservation biology. *Innov. Eur. J. Soc. Sci. Res.* 24 (3), 313–331.
- Álvarez Peña, A., 2004. Melecina máxico-tradicional n'Asturias. VTP Editorial, Gijón (in Asturian).
- Araújo, T.A.S., Alencar, N.L., Amorim, E.L.C., Albuquerque, U.P., 2008. A new approach to study medicinal plants with tannins and flavonoids contents from the local knowledge. *J. Ethnopharmacol.* 120 (1), 72–80.
- Arizaga, J., 1915. Itinerarios botánicos. Diputación Provincial de Álava, Vitoria.
- ASCEL, 2016. (Asociación para la Conservación y Estudio del Lobo Ibérico). <<http://loboiberico.com/>> (accessed 24 October 2016).
- Asgarpanah, J., Kazemivash, N., 2013. Phytochemistry, pharmacology and medicinal properties of *Carthamus tinctorius* L. *Chin. J. Integr. Med.* 19 (2), 153–159.
- Ashok, P.K., Upadhyaya, K., 2012. Tannins are astringent. *J. Pharmacogn. Phytochem.* 1 (3), 45–50.
- ASZAL, 2014. (Asociación Nacional de Criadores de la Raza Asnal Zamorano-Leonesa). Proyecto Burros y lobos en Zamora <<http://www.aszal.com/paginas/proyectos.php?I=6>> (accessed 21 October 2016).
- Azman, N.A.M., Segovia, F., Martínez-Farré, X., Gil, E., Almajano, M.P., 2014. Screening of antioxidant activity of *Gentian lutea* root and its application in oil-in-water emulsions. *Antioxidants* 3, 455–471.
- Babaei, E., Asghari, M.H., Mehdihani, F., Moloudizargari, M., Ghobadi, E., Pouya, S.R.H., 2017. The healing effects of herbal preparations from *Sambucus ebulus* and *Urtica dioica* in full-thickness wound models. *Asian Pac. J. Trop. Biomed.* 7 (5), 421–427.
- Bahramikia, S., Yazdanparast, R., 2012. Phytochemistry and medicinal properties of *Teucrium polium* L. (Lamiaceae). *Phytother. Res.* 26 (11), 1581–1593.
- Bail, S., Buchbauer, G., Jirovec, L., Denkova, Z., Slavchev, A., Stoyanova, A., Schmidt, E., Geissler, M., 2009. Antimicrobial activities of Roman chamomile oil from France and its main compounds. *J. Essent. Oil Res.* 21 (3), 283–286.
- Banerjee, R.D., Sen, S.P., 1980. Antibiotic activity of pteridophytes. *Econ. Bot.* 34 (3), 284–298.
- Barandiarán, J.M., Manterola, A.(Dirs.), 2000. Ganadería y pastoreo en Vasconia. Etniker Euskalerria – Eusko Jaurlaritza – Gobierno de Navarra, Bilbao.
- Barbour, E.K., Al Sharif, M., Sagherian, V.K., Habre, A.N., Talhouk, R.S., Talhouk, S.N., 2004. Screening of selected indigenous plants of Lebanon for antimicrobial activity. *J. Ethnopharmacol.* 93 (1), 1–7.
- Baricevic, D., Sosa, S., Della Loggia, R., Tubaro, A., Simonovska, B., Krasna, A., Zupancic, A., 2001. Topical anti-inflammatory activity of *Salvia officinalis* L. leaves: the relevance of ursolic acid. *J. Ethnopharmacol.* 75 (2–3), 125–132.
- Barja, I., 2009. Prey and prey-age preference by the Iberian wolf *Canis lupus signatus* in a multiple-prey ecosystem. *Wildl. Biol.* 15, 147–154.
- Barrajón-Catalán, E., Fernández-Arroyo, S., Saura, D., Guillén, E., Fernández-Gutiérrez, A., Segura-Carretero, A., Micol, V., 2010. Cistaceae aqueous extracts containing ellagitannins show antioxidant and antimicrobial capacity, and cytotoxic activity against human cancer cells. *Food Chem. Toxicol.* 48, 2273–2282.
- Becker, H., Scher, J.M., Speakman, J.B., Zapp, J., 2005. Bioactivity guided isolation of antimicrobial compounds from *Lythrum salicaria*. *Fitoterapia* 76 (6), 580–584.
- Bennett, B.C., Balick, M.J., 2014. Does the name really matter? The importance of botanical nomenclature and plant taxonomy in biomedical research. *J. Ethnopharmacol.* 152 (3), 387–392.
- Blanco, J.C., Sáenz de Buruaga, M., Llaneza, L., 2007. *Canis lupus Linnaeus, 1758*. In: Palomo, L.J., Gisbert, J., Blanco, J.C. (Eds.), *Atlas y Libro Rojo de los Mamíferos Terrestres de España. Dirección General para la Biodiversidad–SECEM–SECEMU*, Madrid, pp. 272–276.
- Bozin, B., Mimica-Dukic, M., Samoilik, I., Jovin, E., 2007. Antimicrobial and antioxidant properties of rosemary and sage (*Rosmarinus officinalis* L. and *Salvia officinalis* L., Lamiaceae) essential oils. *J. Agric. Food Chem.* 55 (19), 7879–7885.
- Borchardt, J.R., Wyse, D.L., Sheaffer, C.C., Kauppi, K.L., Fulcher, R.G., Ehlke, N.J.,

- Biesboer, D.D., Bey, R.F., 2008a. Antioxidant and antimicrobial activity of seed from plants of the Mississippi river basin. *J. Med. Plant. Res.* 2 (4), 81–93.
- Borchardt, J.R., Wyse, D.L., Sheaffer, C.C., Kauppi, K.L., Fulcher, R.G., Ehlke, N.J., Biesboer, D.D., Bey, R.F., 2008b. Antimicrobial activity of native and naturalized plants of Minnesota and Wisconsin. *J. Med. Plant. Res.* 2 (5), 98–110.
- Brook, I., 2009. Management of human and animal bite wound infection: an overview. *Curr. Infect. Dis. Rep.* 11 (5), 389–395.
- Cândido, A.T., Petrucci-Fonseca, F.R., 2000. O lobo da Serra da Estrela: passado, presente e futuro. *Galemys* 12 (special issue), 209–222.
- Carvalho, A.M., 2010. Plantas y sabiduría popular del Parque Natural de Montesinho. Un estudio etnobotánico en Portugal. CSIC, Madrid.
- Carvalho, A.M., Ramos, M.T., 2012. Guia da etnoflora da Terra de Miranda. FRAUGA & IPB, Bragança.
- Castañón, L., 1966. Detalles y características de la noche sanjuanera en Asturias. *Bol. Inst. Estud. Astur.* 59, 19–56.
- Castañón, L., 1976. Supersticiones y creencias de Asturias. Ayalga, Salinas (Asturias).
- Cerrillo, A., 2017. La política se mete en la boca del lobo. *La Vanguardia* 17.05.2017 < <http://www.lavanguardia.com/natural/20170517/422634548801/politica-boca-lobo.html> >;
- Chao, S.C., Young, D.G., Oberg, C.J., 2000. Screening for inhibitory activity of essential oils on selected bacteria, fungi and viruses. *J. Essent. Oil Res.* 12 (5), 639–649.
- Criado, J., 2010. Plantas medicinales y tóxicas. In: Velasco, J.M., Criado, J., Blanco, E. (Eds.), *Usos tradicionales de las plantas en la provincia de Salamanca*. Instituto de las Identidades. Diputación de Salamanca, Salamanca, 119–215.
- Criado, T., 2012. Lobos por El Bierzo: realidad, tradición y leyenda. Ediciones del Lobo Sapiens, León.
- Dacosta, A., 2010. Una mirada a la tradición. La arquitectura popular en Aliste, Tábara y Alba. Centro de Estudios Cervantinos, Alcalá de Henares (Madrid).
- De Andrés, A.I., Gómez-Serranillos, M.P., Iglesias, I., Villar, A.M., 1999. Effects of extract of *Cistus populifolius* L. on the central nervous system. *Phytother. Res.* 13, 575–579.
- De la Cal, L., 2016. Los 4.650 ataques en el año del lobo feroz. *El Mundo* 22.12.2016 < <http://www.elmundo.es/cronica/2016/12/22/585443ace2704eda6e8b45be.html> >;
- Delamare, A.P.L., Moschen-Pistorello, I.T., Artico, L., Atti-Serfani, L., Echeverrigaray, S., 2007. Antibacterial activity of the essential oils of *Salvia officinalis* L. and *Salvia triloba* L. cultivated in South Brazil. *Food Chem.* 100 (2), 603–608.
- Díaz García, J., 2001. La ganadería a lo largo del siglo XX en el Valle de Carranza (Bizkaia). Anu. Eusko-Folk 43, 131–182.
- Djabou, N., Lorenzi, V., Guinoiseau, E., Andreani, S., Giuliani, M.C., Desjober, J.M., Bolla, J.M., Costa, J., Berti, L., Luciani, A., Muselli, A., 2013. Phytochemical composition of Corsican *Teucrium* essential oils and antibacterial activity against foodborne or toxin-infectious pathogens. *Food Control* 30 (1), 354–363.
- Dogruoz, N., Zeybek, Z., Karagöz, A., 2008. Antibacterial activity of some plant extracts. *IUFS J. Biol.* 67 (1), 17–21.
- Dominguez Moreno, J.M., 1991. La fiesta del lobo en Extremadura. *Rev. Folk.* 132, 183–187.
- Domínguez Moreno, J.M., 1994. La etnoveterinaria en Extremadura: el tratamiento del ganado lanar. *Rev. Folk.* 160, 111–121.
- Dreifke, M.B., Jayasuriya, A.A., Jayasuriya, A.C., 2015. Current wound healing procedures and potential care. *Mater. Sci. Eng. C. Mater. Biol. Appl.* 48, 651–662.
- Džanić, A., Soković, M., Ristić, M., Gruijić-Jovanović, S., Vučkojević, J., Marin, P.D., 2008. Chemical composition and antifungal activity of *Salvia sclarea* (Lamiaceae) essential oil. *Arch. Biol. Sci.* 60 (2), 233–237.
- Edwards, S.E., da Costa Rocha, I., Williamson, E.M., Heinrich, M., 2015. *Phytopharmacy: an evidence-based guide to herbal medicinal products*. John Wiley & Sons Ltd, Chichester.
- Eggermann, J., Ferrão da Costa, G., Guerra, A.M., Kirchner, W.H., Petrucci-Fonseca, F., 2011. Presence of Iberian wolf (*Canis lupus signatus*) in relation to land cover, livestock and human influence in Portugal. *Mamm. Biol.* 76 (2), 217–221.
- El Euch, S.K., Bouajila, J., Bouzouita, N., 2015. Chemical composition, biological and cytotoxic activities of *Cistus salviifolius* flower buds and leaves extracts. *Ind. Crops Prod.* 76, 1100–1105.
- Elías, L.V., Muntión, C., 1989. *Los Pastores de Cameros. Gobierno de La Rioja, Logroño*.
- EMA, 2008a. (European Medicines Agency). Community herbal monograph on *Sambucus nigra* L., flos. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2008b. (European Medicines Agency). Assessment report on *Urtica dioica* L., and *Urtica urens* L., herba. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2010a. (European Medicines Agency). Assessment report on *Salvia officinalis* L., folium and *Salvia officinalis* L., aetheroleum. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2010b. (European Medicines Agency). Community herbal monograph on *Salvia officinalis* L., folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2010c. (European Medicines Agency). Assessment report on *Gentiana lutea* L., radix. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2010d. (European Medicines Agency). Community herbal monograph on *Gentiana lutea* L., radix. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2011a. (European Medicines Agency). Assessment report on *Urtica dioica* L., and *Urtica urens* L., folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2011b. (European Medicines Agency). Assessment report on *Rosmarinus officinalis* L., aetheroleum and *Rosmarinus officinalis* L., folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2012a. (European Medicines Agency). Assessment report on *Fraxinus excelsior* L. or *Fraxinus angustifolia* Vahl, folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2012b. (European Medicines Agency). Community herbal monograph on *Fraxinus excelsior* L. or *Fraxinus angustifolia* Vahl, folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2012c. (European Medicines Agency). Assessment report on *Urtica dioica* L., *Urtica urens* L., their hybrids or their mixtures, radix. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2012d. (European Medicines Agency). Community herbal monograph on *Urtica dioica* L., *Urtica urens* L., their hybrids or their mixtures, radix. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2012e. (European Medicines Agency). Assessment report on *Chelidonium majus* L., herba. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2012f. (European Medicines Agency). Assessment report on *Chamaemelum nobilis* (L.) All., flos. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2013. (European Medicines Agency). Assessment report on *Viscum album* L., herba. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2014a. (European Medicines Agency). Assessment report on *Juglans regia* L., folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2014b. (European Medicines Agency). Community herbal monograph on *Juglans regia* L., folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2015. (European Medicines Agency). Assessment report on *Hedera helix* L., folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- EMA, 2016. (European Medicines Agency). European Union herbal monograph on *Hedera helix* L., folium. Committee on Herbal Medicinal Products (HMPC). Available at <www.ema.europa.eu> .
- Ertürk, Ö., Kati, H., Yayı, N., Demirbağ, Z., 2003. Antimicrobial activity of *Viscum album* L. subsp. *abietis* (Wiesb.). *Turk. J. Biol.* 27, 255–258.
- Fernández-Gil, A., Naves, J., Ordiz, A., Quevedo, M., Revilla, E., Delibes, M., 2016. Conflict misleads large carnivore management and conservation: brown bears and wolves in Spain. *PLoS ONE* 11 (3), e0151541.
- Font Quer, P., 1961. *Plantas Medicinales: el Dioscórides renovado*. Labor, Barcelona.
- Frazão-Moreira, A., Carvalho, A.M., 2009. Etnobotánica do Nordeste Português: Saberes, plantas e usos. Relatório final do Projeto POCI/ANT/59395/2004. Centro em Rede de Investigação em Antropologia (CRIA) e Centro de Investigação de Montanha (CIMO), Bragança.
- Gallego, E., 2009. Estudio etnobotánico del occidente asturiano. Instituto de Estudios Zamoranos Florián de Ocampo, Zamora.
- García Jiménez, R., 2007. *Etnobotánica leonesa: Municipio de Palacios del Sil* (Ph.D. thesis) Facultad de Biología. Universidad Complutense de Madrid, Madrid.
- Gepdiremen, A., Mshvidladze, V., Süleymanı, H., Elias, R., 2005. Acute anti-inflammatory activity of four saponins isolated from ivy: alpha-hederin, hederasaponin-C, hederacolchiside-E and hederacolchiside-F in carrageenan-induced rat paw edema. *Phytomedicine* 12 (6–7), 440–444.
- Gonçalves, S., Romano, A., 2016. The medicinal potential of plants from the genus *Plantago* (Plantaginaceae). *Ind. Crops Prod.* 83, 213–226.
- Gonçalves, S., Gomes, D., Costa, P., Romano, A., 2013. The phenolic content and antioxidant activity of infusions from Mediterranean medicinal plants. *Ind. Crops Prod.* 43, 465–471.
- Goñi, K., 1975. Estudio etnográfico del pueblo de Zerain. Guipúzcoa. Anu. Eusko-Folk 26, 255–442.
- Gragera, F., 1996. *El lobo Ibérico en la Baja Extremadura*. Universitas Editorial, Badajoz.
- Grande del Brío, R., 1984. *El lobo Ibérico: Biología y Mitología*. Hermann Blume, Madrid.
- Grande del Brío, R., 2000. *El lobo Ibérico: Biología, Ecología y Comportamiento*. Amarí Ediciones, Salamanca.
- Grupo de Trabajo del Lobo, 2006. *Estrategia Para la Conservación del lobo en España*. Ministerio de Medio Ambiente, Madrid.
- Guarrera, P.M., 2005. Traditional Phytotherapy in Central Italy (Marche, Abruzzo, and Latium). *J. Ethnopharmacol.* 76 (1), 1–25.
- Guarrera, P.M., Salerno, G., Caneva, G., 2005. Folk Phytotherapeutic Plants from Maratea Area (Basilicata, Italy). *J. Ethnopharmacol.* 99 (3), 367–378.
- Gülçin, I., Uğuz, M.T., Oktay, M., Beydemir, S., Kürevioglu, O.I., 2004a. Evaluation of the antioxidant and antimicrobial activities of clary sage (*Salvia sclarea* L.). *Turk. J. Agric. For.* 28, 25–33.
- Gülçin, I., Kürevioglu, O.I., Oktay, M., Büyükköroğlu, M.E., 2004b. Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.). *J. Ethnopharmacol.* 90 (2–3), 205–215.
- Gynovian, M.M., Trchounian, A.H., 2017. Screening of some plant materials used in Armenian traditional medicine for their antimicrobial activity. *Proc. Yerevan St. Univ. Chem. Biol.* 51 (1), 44–53.
- Habibi, A.A., Zubek, S.A., Abushihhi, M.A., Ahmed, M.O., El-Khodery, S.A., Osman, H.Y., Bennour, E.M., 2015. Antibacterial activity of selected Libyan medicinal plants against *Pseudomonas aeruginosa* and *Escherichia coli*. *J. Pharmacogn. Phytochem.* 3 (6), 197–201.
- Hamed, M.M., Refahy, L.A., Abdel-Aziz, M.S., 2015. Evaluation of antimicrobial activity of some compounds isolated from *Rhamnus cathartica* L. *Orient. J. Chem.* 31 (2), 1133–1140.

- Hamid, Z.S., Shaikh, D.M., 2009. Wood ash and charcoal ash, an instrument to skin tissue repair in acute injury in rabbit model. *Pak. J. Physiol.* 5 (1), 68–72.
- Hearst, C., McCollum, G., Nelson, D., Ballard, L.M., Millar, B.C., Goldsmith, C.E., Rooney, P.J., Loughrey, A., Moore, J.E., Rao, J.R., 2010. Antibacterial activity of elder (*Sambucus nigra* L.) flower or berry against hospital pathogens. *J. Med. Plant. Res.* 4 (17), 1805–1809.
- Hegde, P., Maddur, M.S., Friboulet, A., Bayry, J., Kaveri, S.V., 2011. *Viscum album* exerts anti-inflammatory effect by selectively inhibiting cytokine-induced expression of cyclooxygenase-2. *PLoS ONE* 6 (10), e26312.
- Herrmann, F., Hamoud, R., Sporer, F., Tahraní, A., Wink, M., 2011. Carlina oxide—a natural polyacetylene from *Carlina acaulis* (Asteraceae) with potent antitypanosomal and antimicrobial properties. *Planta Med.* 77, 1905–1911.
- Humadi, S.S., Istudor, V., 2009. *Lythrum salicaria* (purple loosestrife). Medicinal use, extraction and identification of its total phenolic compounds. *Farmacia* 57 (2), 192–200.
- Iglesias, A., España, A.J., España, J., 2017. Lobos ibéricos: anatomía, ecología y conservación. Náyade, Medina del Campo (Valladolid).
- Ivancheva, S., Nikolova, M., Tsvetkova, R., 2006. In: Imperato, F. (Ed.), *Pharmacological Activities and Biologically Active Compounds of Bulgarian Medicinal Plants. Phytochemistry: Advances in Research*, Trivandrum, Kerala, 87–103.
- Junceda Avello, E., 1987. *Medicina Popular en Asturias*. Instituto de Estudios Asturianos, Oviedo.
- Kadhim, S.M., Mohammed, M.T., Ahmed, O.M., Jassimand, A.M.N., 2016. Study of some *Salvia officinalis* L. (sage) components and effect of their aqueous extract on antioxidant. *Int. J. Chem. Sci.* 14 (2), 711–719.
- Karimzadeh, S., Farahpour, M.R., 2017. Topical application of *Salvia officinalis* hydroethanolic leaf extract improves wound healing process. *Indian J. Exp. Biol.* 55, 98–106.
- Kasote, D., Ahmad, A., Viljoen, A., 2015. Proangiogenic potential of medicinal plants in wound healing. In: Mukherjee, P.K. (Ed.), *Evidence-Based Validation of Herbal Medicine*. Elservier, Amsterdam, 149–164.
- Kazemian, H., Ghafourian, S., Heidari, H., Amiri, P., Yamchi, J.K., Shavalipour, A., Hourri, H., Maleki, A., Sadeghifard, N., 2015. Antibacterial, anti-swarming and anti-biofilm formation activities of *Chamaelium nobilis* against *Pseudomonas aeruginosa*. *Rev. Soc. Bras. Med. Trop.* 48 (4), 432–436.
- Khammar, A., Djeddi, S., 2012. Pharmacological and biological properties of some *Centaurea* species. *Eur. J. Sci. Res.* 84 (3), 398–416.
- Kindler, S., Schuster, M., Seebauer, C., Rutkowski, R., et al., 2016. Triterpenes for well-balanced scar formation in superficial wounds. *Molecules* 21, 1129.
- Koca, U., Süntar, I.P., Keles, H., Yesilada, E., Akkol, E.K., 2009. In vivo anti-inflammatory and wound healing activities of *Centaurea iberica* Trev. ex Spreng. *J. Ethnopharmacol.* 126 (3), 551–556.
- Kokosa, L., Polesny, Z., Rada, V., Nepovim, A., Vanek, T., 2002. Screening of some Siberian medicinal plants for antimicrobial activity. *J. Ethnopharmacol.* 82, 51–53.
- Kolodziejczyk-Czepas, J., 2012. *Trifolium* species-derived substances and extracts—biological activity and prospects for medicinal applications. *J. Ethnopharmacol.* 143 (1), 14–23.
- Kuete, V., Wiench, B., Hegazy, M.E.F., Mohamed, T.A., Fankam, A.G., Shahat, A.A., Efferth, T., 2012. Antibacterial activity and cytotoxicity of selected Egyptian medicinal plants. *Planta Med.* 78, 193–199.
- Kumar, B., Vijayakumar, M., Govindarajan, R., Pushpangadan, P., 2007. Ethnopharmacological approaches to wound healing—exploring medicinal plants of India. *J. Ethnopharmacol.* 114, 103–113.
- Kuonen, R., Weissenstein, U., Urech, K., Kunz, M., Hostanska, K., Estko, M., Heusser, P., Baumgartner, S., 2013. Effects of lipophilic extract of *Viscum album* L. and oleanolic acid on migratory activity of NIH/3T3 fibroblasts and on HaCat keratinocytes. *Evid. Based Complement. Altern. Med.* 2013, 718105.
- Lai, B., Teixeira, G., Moreira, I., Correia, A.I., Duarte, A., Madureira, A.M., 2012. Evaluation of the antimicrobial activity in species of a Portuguese “Montado” ecosystem against multidrug resistant pathogens. *J. Med. Plant. Res.* 6 (12), 2381–2387.
- Lagos, L., Bárcena, F., 2015. EU sanitary regulation on livestock disposal: implications for the diet of wolves. *Environ. Manag.* 56 (4), 890–902.
- Latorre, J.A., 2008. Estudio Etnobotánico de la Provincia de La Coruña Ph.D. Thesis. Facultad de Farmacia, Universitat de València, Valencia.
- Linnell, J.D.C., Allee, J., 2016. Predators that kill humans: myth, reality, context and the politics of wolf attacks on people. In: Angelici, F.M. (Ed.), *Problematic Wildlife: A Cross-disciplinary Approach*. Springer, Heidelberg, (375–371).
- Linnell, J.D.C., Andersen, R., Andersone, Z., Balciuskas, L., Blanco, J.C., Boitani, L., Brainerd, S., Breitenmoser, U., Kojola, I., Liberg, O., Loe, J., Okarma, H., Pedersen, H.C., Promberger, C., Sand, H., Solberg, E.J., Valdmann, H., Wabakken, P., 2002. The fear of wolves: a review of wolf attacks on humans. *NINA Oppdragsmedd.* 731, 1–65.
- Llaneza, L., López-Bao, J.V., 2015. Indirect effects of changes in environmental and agricultural policies on the diet of wolves. *Eur. J. Wildl. Res.* 61, 895–902.
- Llaneza, L., López-Bao, J.V., Sazatornil, V., 2012. Insights into wolf presence in human-dominated landscapes: the relative role of food availability, humans and landscape attributes. *Divers. Distrib.* 18, 459–469.
- Locatelli, M., Epifano, F., Genovese, S., Carlucci, G., Koncić, M.Z., Kosalec, I., Kremer, D., 2011. Anthraquinone profile, antioxidant and antimicrobial properties of bark extracts of *Rhamnus catharticus* and *R. orbiculatus*. *Nat. Prod. Commun.* 6 (9), 1275–1280.
- Lutsenko, Y., Bylka, W., Matlawska, I., Darmohray, R., 2010. *Hedera helix* as a medicinal plant. *Herba Pol.* 56 (1), 83–96.
- Maccioni, S., Baldini, R., Cioni, P.L., Tebano, M., Flamini, G., 2007. *In vivo* volatiles emission and essential oils from different organs and pollen of *Cistus albidus* from Caprione (Eastern Liguria, Italy). *Flavour Fragr. J.* 22, 61–65.
- Mahmood, A.A., Phipps, M.E., 2006. Wound healing activities of *Plantago major* leaf extract in rats. *Int. J. Trop. Med.* 1 (1), 33–35.
- Mahmoudi, H., Aouadhi, C., Kadour, R., Gruber, M., Zargouni, H., et al., 2016. Comparison of antioxidant and antimicrobial activities of two cultivated *Cistus* species from Tunisia. *Biosci. J. Überländia* 32 (1), 226–237.
- Mantle, D., Gok, M.A., Lennard, T.W., 2001. Adverse and beneficial effects of plant extracts on skin and skin disorders. *Advers. Drug React. Toxicol. Rev.* 20 (2), 89–103.
- MAPAMA, 2015. (Ministerio de Agricultura, Alimentación y Medio Ambiente). Censo 2012–2014 de lobo ibérico (*Canis lupus*, Linnaeus, 1758) en España. <http://www.mapama.gob.es/es/biodiversidad/temas/inventarios-nacionales/censo_lobo_espana_2012_14_tcm7-414048.pdf>.
- Maríño Ferro, X.R., 1985. *La medicina popular interpretada*. Ediciones Xerais de Galicia, Vigo.
- Martínez, A.L., González-Trujano, M.E., Pellicer, F., López-Muñoz, F.J., Navarrete, A., 2009. Antinociceptive effect and GC/MS analysis of *Rosmarinus officinalis* L. essential oil from its aerial parts. *Planta Med.* 75, 508–511.
- Martínez-Barbeito, C., 1947. *El bosque de Ancines*. Ayma, Barcelona.
- Martínez González, J., Lastra, J.J., 2006. Aportaciones al conocimiento de la medicina tradicional en el Concejo de Aller (Asturias, España). In: *Actas del V Congreso Internacional de Fitoterapia y Técnicas Afines Ciudad de Oviedo*. Sociedad Española de Fitoterapia, Oviedo, pp. 131–144.
- Massip, J.M., 2011. *El llop i els humans. Passat i present a Catalunya*. Arola, Tarragona (in Catalan).
- Mathew, A., Taranalli, A.D., Torgal, S.S., 2004. Evaluation of anti-inflammatory and wound healing activity of *Gentiana lutea* rhizome extracts in animals. *Pharm. Biol.* 42 (1), 8–12.
- MEDWOLF, 2016. LIFE project MED-WOLF: Best practice actions for wolf conservation in Mediterranean type-areas. <<http://www.medwolf.eu/>> (accessed 21 October 2016).
- Menéndez-Baceta, G., Aceituno-Mata, L., Molina, M., Reyes-García, V., Tardío, J., Pardo-de-Santayana, M., 2014. Medicinal plants traditionally used in the northwest of the Basque Country (Biscay and Alava), Iberian Peninsula. *J. Ethnopharmacol.* 152, 113–134.
- Merino, V., Blanquer, A., 2016. Fitoterapia para las afecciones dermatológicas. In: Castillo García, E., Martínez Solís, I. (Eds.), *Manual de Fitoterapia* 2nd ed. Elsevier, Barcelona, 411–428.
- Mettler, D., 2014. Alternatives to livestock guarding dogs. Llamas, donkeys, fence-systems and aversive conditioning: possibilities and limits. *CDP News* 10, 18–20.
- Mitić-Čulafić, D., Vuković-Gačić, B., Knežević-Vukčević, J., Stanković, S., Simić, D., 2005. Comparative study on the antibacterial activity of volatiles from sage (*Salvia officinalis* L.). *Arch. Biol. Sci.* 57 (3), 173–178.
- Modarresi-Chahardehi, A., Ibrahim, D., Fariza-Sulaiman, S., Mousavi, L., 2012. Screening antimicrobial activity of various extracts of *Urtica dioica*. *Rev. Biol. Trop.* 60 (4), 1567–1576.
- Mogoșanu, G.D., Popescu, F.C., Busuioc, C.J., Pop, O.T., Mogoanta, L., Parvanescu, H., Rau, G., Lascăr, I., 2014. Effects of a topical preparation containing Sambuci Folium extract in experimental model of thermal skin burns on rats. *Farmacia* 62 (4), 693–703.
- Moreno, S., Scheyer, T., Romano, C.S., Vojnov, A.A., 2006. Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. *Free Radic. Res* 40 (2), 223–231.
- Mothana, R.A.A., Abdo, S.A.A., Hasson, S., Althawab, F.M.N., Alaghbari, S.A.Z., Lindequist, U., 2010. Antimicrobial, antioxidant and cytotoxic activities and phytochemical screening of some Yemeni medicinal plants. *Evid. Based Complement. Altern. Med.* 7 (3), 323–330.
- Moulaoui, K., Caddeo, C., Manca, M.L., Castangia, I., Valentí, D., Escrivano, E., Atmani, D., Fadda, A.M., Manconi, M., 2015. Identification and nanoentrainment of polyphenolic phytocomplex from *Fraxinus angustifolia*: *In vitro* and *in vivo* wound healing potential. *Eur. J. Med. Chem.* 89 (7), 179–188.
- Mukherjee, P.K., Verpoorte, R., Suresh, B., 2000. Evaluation of *in-vivo* wound healing activity of *Hypericum patulum* (Family: hypericaceae) leaf extract on different wound model in rats. *J. Ethnopharmacol.* 70, 315–321.
- Nagori, P.B., Solanki, R., 2011. Role of medicinal plants in wound healing. *Res. J. Med. Plant* 5 (4), 392–405.
- Nieto Macén, D., 2016. Cómo proteger el ganado con perros. Tundra, Almenara (Castellón).
- Orhan, D.D., Küpeli, E., Yesilada, E., Ergun, F., 2006. Anti-inflammatory and antinociceptive activity of flavonoids isolated from *Viscum album* ssp. *album*. *Z. Nat. C.* 61, 26–30.
- Oria de Rueda, J.A., Diez, J., Rodriguez, M., 1996. *Guía de las Plantas Silvestres de Palencia*. Cáalamo, Palencia.
- Oztürk, H., Kolak, U., Meric, C., 2011. Antioxidant, anticholinesterase and antibacterial activities of *Jurinea consanguinea* DC. *Rec. Nat. Prod.* 5 (1), 43–51.
- Oztürk, N., Korkmaz, S., Oztürk, Y., Baser, K.H.C., 2006. Effects of gentiopicroside, sweroside and swertiafamarine, secoiridoids from gentian (*Gentiana lutea* ssp. *sympphyandra*), on cultured chicken embryonic fibroblasts. *Planta Med.* 72, 289–294.
- Panth, N., Paudel, K.R., Karki, R., 2016. Phytochemical profile and biological activity of *Juglans regia*. *J. Integr. Med.* 14 (5), 359–373.
- Papaefthimiou, D., Papanikolaou, A., Falara, V., Givanoudi, S., Kostas, S., Kanellis, A.K., 2014. Genus *Cistus*: a model for exploring labdane-type diterpenes' biosynthesis and a natural source of high value products with biological, aromatic, and pharmacological properties. *Front Chem.* 2, 35.
- Pardo-de-Santayana, M., 2004. *Guía de las Plantas Medicinales de Cantabria*. Librerías Estudio, Santander.

- Pardo-de-Santayana, M., 2008. Estudios Etnobotánicos en Campoo (Cantabria): Conocimiento y uso Tradicional de Plantas. CSIC, Madrid.
- Pascual Gil, J.C., 2013. Etnobotánica de La Pernía, Polentinos y Valle de Castillería (Palencia) (Degree thesis). ETS Ingenierías Agrarias. Universidad de Valladolid, Palencia.
- Pazyar, N., Yaghoobi, R., Rafiee, E., Mehrabian, A., Feily, A., 2014. Skin wound healing and phytotherapy: a review. *Skin. Pharmacol. Physiol.* 27 (6), 303–310.
- Peana, A.T., Moretti, M.D.L., 2002. Pharmacological activities and applications of *Salvia sclarea* and *Salvia desoleana* essential oils. *Stud. Nat. Prod. Chem.* 26, 391–423.
- Penco, A.D., 2005. Medicina Popular Veterinaria en la Comarca de Zafra. Facultad de Medicina. Universidad de Extremadura, Badajoz.
- Pereira, R.F., Bartolo, P.J., 2016. Traditional therapies for skin wound healing. *Adv. Wound Care* 5 (5), 208–229.
- Peterson, M., Simmonds, M.S., 2003. Rosmarinic acid. *Phytochemistry* 62 (2), 121–125.
- Pieroni, A., Ibraliiu, A., Abbasi, A.M., Papajani-Toska, V., 2015. An ethnobotanical study among Albanians and Aromanians living in the Rracaë and Mokra areas of Eastern Albania. *Genet. Resour. Crop Evol.* 62, 477–500.
- Piluzza, G., Virdis, S., Serralutzi, F., Bullitta, S., 2015. Uses of plants, animal and mineral substances in Mediterranean ethno-veterinary practices for the care of small ruminants. *J. Ethnopharmacol.* 168, 87–99.
- Pimenta, V., Barroso, I., Álvares, F., Correia, J., Ferrão da Costa, G., Moreira, L., Nascimento, J., Petrucci-Fonseca, F., Roque, S., Santos, E., 2005. Situação Populacional do lobo em Portugal: Resultados do Censo Nacional 2002/2003. Relatório Técnico. ICNF, Lisboa.
- Pimenta, V., Barroso, I., Boitani, L., Beja, P., 2017. Wolf predation on cattle in Portugal: assessing the effects of hunting systems. *Biol. Conserv.* 207, 17–26.
- Pinela, J., Barros, L., Carvalho, A.M., Ferreira, I.C.F.R., 2011. Influence of the drying method in the antioxidant potential and chemical composition of four shrubby flowering plants from the tribe Genistae (Fabaceae). *Food Chem. Toxicol.* 49 (11), 2983–2989.
- Pinela, J., Barros, L., Dueñas, M., Carvalho, A.M., Santos-Buelga, C., Ferreira, I.C.F.R., 2012. Antioxidant activity, ascorbic acid, phenolic compounds and sugars of wild and commercial *Tuberaria lignosa* samples: effects of drying and oral preparation methods. *Food Chem.* 135, 1028–1035.
- Planelles, M., 2016a. El negocio sucio que rodea al lobo. *El País* 03.07.2016 <http://politica.elpais.com/politica/2016/07/01/actualidad/1467394364_030402.html?Rel=mas&Rel=mas>.
- Planelles, M., 2016b. La Guardia Civil cifra en 200.000 euros el fraude por ataques de lobo en Asturias. *El País* 16.07.2016 <http://politica.elpais.com/politica/2016/07/16/actualidad/1468687494_357490.html>.
- Prieto, L., 1949. Notas etnográficas sobre animales domésticos y salvajes de La Gudiña (Ourense). Douro Litoral 5, 50–67.
- Rahbaran, M.G., Amin, B., Mehri, S., Mirnajafi-Zadeh, S.J., Hosseinzadeh, H., 2017. Anti-inflammatory effects of ethanolic extract of *Rosmarinus officinalis* L. and rosmarinic acid in a rat model of neuropathic pain. *Biomed. Pharmacother.* 86, 441–449.
- Raina, R., Parwez, S., Verma, P.K., Pankaj, N.K., 2008. Medicinal plants and their role in wound healing. *Vet. Scan* 3 (1), article 21.
- Ramos, S., 2015. Burros pastores contra el lobo. *La Opinión de Zamora* 21.03.2015 <<http://www.laopiniondezamora.es/agroalimentacion/2015/03/19/proyecto-comprobara-eficacia-burros-proteger/830155.html>>.
- Rashed, K.N.Z., 2013. Antioxidant activity of *Hedera helix* L. extracts and the main phytoconstituents. *Int. J. Allied Med. Sci. Clin. Res.* 1 (2), 62–64.
- Rashed, A.N., Afifi, F.U., Disi, A.M., 2003. Simple evaluation of the wound healing activity of a crude extract of *Portulaca oleracea* L. (growing in Jordan) in *Mus musculus* JVI-1. *J. Ethnopharmacol.* 88, 131–136.
- Rasik, A.M., Shukla, A., 2001. Antioxidant status in delayed healing type of wounds. *Int. J. Exp. Path.* 81, 257–263.
- Rebaya, A., Belghith, S.I., Hammrouni, S., Maaroufi, A., Ayadi, M.T., Chérif, J.K., 2016. Antibacterial and antifungal activities of ethanol extracts of *Halimium halimifolium*, *Cistus salviifolius* and *Cistus monspeliensis*. *Int. J. Pharm. Clin. Res.* 8 (4), 243–247.
- Redzic, S., Ferrier, J., 2011. Ethnobotany and ethnopharmacology of the genus *Veratrum* L. in veterinary medicine (Bosnia and Herzegovina, W. Balkan). *Planta Med.* 77 (12), 1320–1321.
- Rejón, R., 2016. Los ataques de lobos al ganado afectan a menos del 1% de la cabaña española. *ElDiario.es* 30.07.2016 <http://www.eldiario.es/sociedad/ataques-ganado-suponen-cabana-espanola_0_541045973.html>.
- Renda, G., Yalçın, F.N., Nemutlu, E., Akkol, E.K., Süntar, I., Keleş, H., Ina, H., Çalış, I., Ersöz, T., 2013. Comparative assessment of dermal wound healing potentials of various *Trifolium* L. extracts and determination of their isoflavone contents as potential active ingredients. *J. Ethnopharmacol.* 148 (2), 423–432.
- Ribeiro, A., 1918. Terras do Demo. Bertrand Editora, Lisboa.
- Ribeiro, A., 1958. Quando os lobos Uivam. Bertrand Editora, Lisboa.
- Riego Celada, L., Galhano Alves, J.P., 2006. Contos y cuentos de lobos. Recorrido por los relatos loberos de Sanabria y Tras os Montes. Ayuntamiento de Puebla de Sanabria, Puebla de Sanabria (Zamora).
- Rivera, D., Allkin, R., Obón, C., Alcaraz, F., Verpoorte, R., Heinrich, M., 2014. What is in a name? The need for accurate scientific nomenclature for plants. *J. Ethnopharmacol.* 152 (3), 393–402.
- Rosa, J.L., 2013. Cães de gado, os grandes guardiães dos rebanhos. In: Rodrigues, J.B. (Ed.), *Entre lobos e pastores*. Aldeia-CoraNe, Bragança, 35–41.
- Sáenz de Buruaga, M., Canales, F., Campos, M.A., Noriega, A., Muñoz, F.J., Navamuel, N., 2015. Censo regional de lobo (*Canis lupus*) en Castilla y León. Consultora de Recursos Naturales S. L. – TRAGSATEC, Valladolid.
- Salehzadeh, A., Asadpour, L., Naeemi, A.S., Houshangd, E., 2014. Antimicrobial activity of methanolic extracts of *Sambucus ebulus* and *Urtica dioica* against clinical isolates of methicillin resistant *Staphylococcus aureus*. *Afr. J. Tradit. Complement. Altern. Med* 11 (5), 38–40.
- Samuelson, A.B., 2000. The traditional uses, chemical constituents and biological activities of *Plantago major* L.: a review. *J. Ethnopharmacol.* 71, 1–21.
- Santos, A., Barros, L., Calhelha, R.C., Dueñas, M., Carvalho, A.M., Santos-Buelga, C., Ferreira, I.C.F.R., 2013. Leaves and decoction of *Juglans regia* L.: different performances regarding bioactive compounds and in vitro antioxidant and antitumor effects. *Ind. Crops Prod.* 51, 430–436.
- Santos Galíndez, J., Díaz Lanza, A.M., Fernández Matellano, L., 2002. Biologically active substances from the genus *Scrophularia*. *Pharm. Biol.* 40 (1), 45–59.
- Šavikin, K., Menković, N., Zdunić, G., Stević, T., Radanović, D., Janković, T., 2009. Antimicrobial activity of *Gentiana lutea* L. extracts. *Z. Nat. C.* 64, 339–342.
- Sengul, M., Yıldız, H., Gungor, N., Cetin, B., Eser, Z., Ercisli, S., 2009. Total phenolic content, antioxidant and antimicrobial activities of some medicinal plants. *Pak. J. Pharm. Sci.* 22 (1), 102–106.
- Sereno, E., 2017. Los ataques de lobo a la ganadería se intensifican: más de 7.500 animales muertos. *elEconomista.es* 30.01.2017 <<http://www.eleconomista.es/empresas-finanzas/agro/noticias/8118852/01/17/Los-ataques-de-lobo-a-la-ganaderia-se-intensifican-mas-de-7500-animales-muertos.html>>.
- Shaikh, D.M., Hamid, Z.S., 2009. Ash as a unique natural medicine for wound healing. *Isra Med. J.* 1 (3), 72–78.
- Sharma, P., Ravikumar, G., Kalaiselvi, M., Gomathi, D., Uma, C., 2013. In vitro antibacterial and free radical scavenging activity of green hull of *Juglans regia*. *J. Pharm. Anal.* 3 (4), 298–302.
- Sienkiewicz, M., Głowiak, A., Pożnańska-Kurowska, K., Kaszuba, A., Urbaniak, A., Kowalczyk, E., 2015. The effect of clary sage oil on staphylococci responsible for wound infections. *Post. Dermatol. Alergol.* 32 (1), 21–26.
- Signatus.org, 2016. Medidas de defensa del ganado frente a ataques de lobo. Signatus.org: en defensa del lobo ibérico y sus hábitats <<https://www.signatus.org/defensa-ganado>> (accessed 21 October 2016).
- Silva, A., Meireles, C., Dias, C., Sales, F., Conde, J., Salgueiro, L., Batista, T., 2011. Plantas aromáticas e medicinais do Parque Natural da Serra da Estrela – guia etnobotânico. CISE, Seia.
- Simani, S., Fayaz, A., Rahimi, P., Eslami, N., Howeizi, N., Biglari, P., 2012. Six fatal cases of classical rabies virus without biting incidents, Iran 1990–2010. *J. Clin. Virol.* 54 (3), 251–254.
- Singh, A., 2008. Phytochemicals of Gentianaceae: a review of pharmaceutical properties. *Int. J. Pharm. Sci. Nanotechnol.* 1 (1), 33–36.
- Srivastava, J.K., Shankar, E., Gupta, S., 2010. Chamomile: a herbal medicine of the past with bright future. *Mol. Med. Report.* 3 (6), 895–901.
- Stanislavljević, I.T., Stojčević, S.S., Veljković, D.T., Lazić, M.L., Veljković, V.B., 2008. Screening the antioxidant and antimicrobial properties of the extracts from plantain (*Plantago major* L.) leaves. *Spec. Sci. Tech.* 43 (14), 3652–3662.
- Tabatabaei Yazdi, F., Alizadeh Behbahani, B., 2013. Antimicrobial effect of aqueous and ethanolic extracts *Teucrium polium* L. on *Streptococcus pyogenes*, *Pseudomonas aeruginosa* and *Staphylococcus epidermidis*. *Sci. J. Biol. Sci.* 2 (3), 54–61.
- Takaki, I., Bersani-Amado, L.E., Vendruscolo, A., Sartoretto, S.M., Diniz, S.P., Bersani-Amado, C.A., Cuman, R.K., 2008. Anti-inflammatory and antinociceptive effects of *Rosmarinus officinalis* L. essential oil in experimental animal models. *J. Med. Food* 11 (4), 741–746.
- Talegón, J., 2016. El lobo en la cultural rural del noroeste de Zamora. Algunos datos preliminares. XIV Jornadas de Cultura tradicional sobre la cultura loba en el medio rural del noroeste de la provincia de Zamora, Zamora. <<http://www.furmientu.org/01Documentos/999999999998LlobosHomes.pdf>>.
- Thomas, T., 2011. Preliminary antibacterial and phytochemical assessment of *Osmunda regalis* L. *IJPBA* 2 (1), 559–562.
- Thomas, N., Brook, I., 2011. Animal bite-associated infections: microbiology and treatment. *Expert Rev. Anti Infect. Ther.* 9 (2), 15–26.
- Tomás-Menor, L., Morales-Soto, A., Barragón-Catalán, E., Roldán-Segura, C., Segura-Carretero, A., Micó, V., 2013. Correlation between the antibacterial activity and the composition of extracts derived from various Spanish *Cistus* species. *Food Chem. Toxicol.* 55, 313–322.
- Torres, R.T., Fonseca, C., 2016. Perspectives on the Iberian wolf in Portugal: population trends and conservation threats. *Biodivers. Conserv.* 25 (3), 411–425.
- Torres, R.T., Silva, N., Brotas, G., Fonseca, C., 2015. To eat or not to eat? The diet of the endangered Iberian wolf (*Canis lupus signatus*) in a human-dominated landscape in central Portugal. *PLoS One* 10 (6), e0129379.
- Tosun, F., Akyüz Kızılıy, Ç., Şener, B., Vural, M., 2005. The evaluation of plants from Turkey for in vitro antimycobacterial activity. *Pharm. Biol.* 43 (1), 58–63.
- Tunalier, Z., Koşar, M., Küpeli, E., Çalış, İ., Can Başer, K.H., 2007. Antioxidant, anti-inflammatory, anti-nociceptive activities and composition of *Lythrum salicaria* L. extracts. *J. Ethnopharmacol.* 110 (3), 539–547.
- Turgumbayeva, A.A., Ustanova, G.O., Stabayeva, G.C., Ross, S.A., 2014. Phytochemical screening and biological activities of the Camel Thorn (*Alhagi kirghisorum*) and Safflower Flowers (*Carthamus tinctorius* L.) grown in Kazakhstan. *Am.-Eurasian J. Agric. Environ. Sci.* 14 (12), 1487–1491.
- Türkmen, S., Sahin, A., Gunaydin, M., Tatlı, O., Karaca, Y., Turedi, S., Gunduz, A., 2012. A wild wolf attack and its unfortunate outcome: rabies and death. *Wilderness Environ. Med.* 23 (3), 248–250.
- Uddin, G., Rauf, A., Qaisar, M., Rehman, T.U., Latif, A., Ali, M., 2011. Preliminary phytochemical screening and antimicrobial activity of *Hedera helix* L. *Middle East J. Sci. Res.* 8 (1), 198–202.
- Vafi, F., Bahrami-Soltani, R., Abdollahi, M., et al., 2016. Burn wound healing activity of *Lythrum salicaria* L. and *Hypericum scabrum* L. *Wounds* 28 (12), 448–458.
- Valverde, J.A., Teruelo, S., 2001. Los lobos de Morla. Al Andalus Ediciones, Sevilla.
- Vanaclocha, B., Cañigueral, S., 2003. Fitoterapia. Vademécum de Prescripción 4th ed.

- Masson, Barcelona.
- Vantassel, S.M., 2012. The Wildlife Damage Inspection Handbook: A Guide to Identifying Vertebrate Damage to Structures, Landscapes and Livestock 3rd ed. Wildlife Control Consultant, Lincoln.
- Velnar, T., Bailey, T., Smrkolj, V., 2009. The wound healing process: an overview of the cellular and molecular mechanisms. *J. Int. Med. Res.* 37 (5), 1528–1542.
- Venditti, A., Frezza, C., Riccardelli, M., Foddai, S., Nicoletti, M., Serafini, M., Bianco, A., 2016. Secondary metabolites from *Scrophularia canina* L. *Nat. Prod. Res.* 30 (14), 1665–1669.
- Verde, A., Rivera, D., Fajardo, J., Obón, C., Cebrián, F., 2008. Guía de las plantas medicinales de Castilla-La Mancha (y otros recursos medicinales de uso tradicional). Altabán Ediciones, Albacete.
- Viapiana, A., Wesolowski, M., 2017. The phenolic contents and antioxidant activities of infusions of *Sambucus nigra* L. *Plant Foods Hum. Nutr.* 72 (1), 82–87.
- Vidal-González, P., 2013. Sacred rituals and popular religiousness amongst transhumant shepherds of Teruel region, Spain. *Pastoralism* 3, 24.
- Viegi, L., Pieroni, A., Guarnera, P.M., Vangelisti, R., 2003. A review of plants used in folk veterinary medicine in Italy as basis for a databank. *J. Ethnopharmacol.* 89, 221–244.
- WHO (World Health Organization), 2004. WHO Monographs on Selected Medicinal Plants 2. WHO Press, Geneva.
- WHO (World Health Organization), 2007. WHO Monographs on Selected Medicinal Plants 3. WHO Press, Geneva.
- WHO (World Health Organization), 2009. WHO Monographs on Selected Medicinal Plants 4. WHO Press, Geneva.
- WHO (World Health Organization), 2010. WHO Monographs on Medicinal Plants Commonly used in the Newly Independent States (NIS). WHO Press, Geneva.
- WOLF, 2014. Good practice guide for co-existence wolf – livestock farming. WOLF, wild life & farmers Cooperation project. <http://www.wolf-project.com/lang/publicaciones/index_d.shtml?Refbol=publicaciones&refsec=publicaciones-y-documenta&idarticulo=125307> (accessed 21 October 2016).
- Wynn, S.G., Fougère, B.J., 2007. *Materia Medica*. In: Wynn, S.G., Fougère, B.J. (Eds.), *Veterinary Herbal Medicine*. Mosby-Elsevier, St. Louis, 459–672.
- Zater, H., Huet, J., Fontaine, V., Benayache, S., Stévigny, C., Duez, P., Benayache, F., 2016. Chemical constituents, cytotoxic, antifungal and antimicrobial properties of *Centaurea diluta* Ait. subsp. *algeriensis* (Coss. & Dur.) Maire. *Asian Pac. J. Trop. Med.* 9 (6), 554–561.
- Zhao, J., Khan, S.I., Wang, M., Vasquez, Y., Yang, M.H., Avula, B., Wang, Y.H., Avonto, C., Smillie, T.J., Khan, I.A., 2014. Octulosonic acid derivatives from Roman chamomile (*Chamaemelum nobile*) with activities against inflammation and metabolic disorder. *J. Nat. Prod.* 77 (3), 509–515.
- Zuo, G.Y., Meng, F.Y., Hao, X.Y., Zhang, Y.L., Wang, G.C., Xu, G.L., 2008. Antibacterial alkaloids from *Chelidonium majus* Linn (Papaveraceae) against clinical isolates of methicillin-resistant *Staphylococcus aureus*. *J. Pharm. Pharm. Sci.* 11 (4), 90–94.
- Zurriaga, F., 2012. El lledoner: l'arbre de les forques. In: Climent, D., Zurriaga, F. (Eds.), *Herbari: Viure amb les Plantes. Monografies Mètode 6*, Valencia, 245–257, (in Catalan).