

Shyamal K. Hui; Richard S. Lemence; Pradip Mandal

Wintgen inequalities on Legendrian submanifolds of generalized Sasakian-space-forms

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 61 (2020), No. 1, 105–117

Persistent URL: <http://dml.cz/dmlcz/148079>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2020

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## Wintgen inequalities on Legendrian submanifolds of generalized Sasakian-space-forms

SHYAMAL K. HUI, RICHARD S. LEMENCE, PRADIP MANDAL

*Abstract.* A submanifold  $M^m$  of a generalized Sasakian-space-form  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  is said to be  $C$ -totally real submanifold if  $\xi \in \Gamma(T^\perp M)$  and  $\varphi X \in \Gamma(T^\perp M)$  for all  $X \in \Gamma(TM)$ . In particular, if  $m = n$ , then  $M^n$  is called Legendrian submanifold. Here, we derive Wintgen inequalities on Legendrian submanifolds of generalized Sasakian-space-forms with respect to different connections; namely, quarter symmetric metric connection, Schouten–van Kampen connection and Tanaka–Webster connection.

*Keywords:* generalized Sasakian-space-form; Legendrian submanifold

*Classification:* 53C25, 53C15

### 1. Introduction

A generalized Sasakian-space-form is an almost contact metric manifold  $\overline{M}(\varphi, \xi, \eta, g)$  whose curvature tensor  $\overline{R}$  is of the form, see [1],

$$\begin{aligned}
 \overline{R}(X, Y)Z &= f_1\{g(Y, Z)X - g(X, Z)Y\} \\
 &\quad + f_2\{g(X, \varphi Z)\varphi Y - g(Y, \varphi Z)\varphi X + 2g(X, \varphi Y)\varphi Z\} \\
 &\quad + f_3[\eta(Z)\{\eta(X)Y - \eta(Y)X\} \\
 &\quad + \{g(X, Z)\eta(Y) - g(Y, Z)\eta(X)\}\xi]
 \end{aligned}
 \tag{1.1}$$

for all vector fields  $X, Y, Z$  on  $\overline{M}$ , where  $f_i \in C^\infty(\overline{M})$ ,  $i = 1, 2, 3$ . Such a manifold of dimension  $(2n + 1)$ ,  $n > 1$ , is denoted by  $\overline{M}^{2n+1}(f_1, f_2, f_3)$ .

In particular, if  $f_1 = (c + 3)/4$ ,  $f_2 = f_3 = (c - 1)/4$  then  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  reduces to the notion of Sasakian-space-forms. Many authors studied  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  in different context such as ([2]–[5], and references therein).

Beside the Riemannian connection, there exist some other connections on smooth manifolds. In 1975, S. Golab in [6] introduced the idea of quarter symmetric connection. The quarter symmetric connection is called metric connection if the covariant derivative of such connection is zero.

---

DOI 10.14712/1213-7243.2020.006

The third author gratefully acknowledges to the CSIR(File No. 09/025(0221)/2017-EMR-1), Government of India for the award of Junior Research Fellowship.

The Schouten–van Kampen connection (SVKC) introduced for the study of non-holomorphic manifolds, see [11]. In 2006, A. Bejancu in [3] studied SVKC connection on foliated manifolds. Recently Z. Olszak in [10] studied SVKC on almost (para) contact metric structure.

The Tanaka–Webster connection (TWC), see [12], [14], is the canonical affine connection defined on a non-degenerate pseudo-Hermitian CR-manifold. S. Tanno in [13] defined the TWC for contact metric manifolds. Here we denote quarter symmetric metric connection (QSMC), SVKC and TWC on  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  by  $\widetilde{\nabla}$ ,  $\widehat{\nabla}$ ,  $\overset{*}{\nabla}$ , respectively.

After introducing Wintgen inequality in [15], I. Mihai derived Wintgen inequality for submanifolds of complex-space-form, see [8], and Sasakian-space-form, see [9]. In this paper we derive Wintgen inequality for Legendrian submanifolds of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\widetilde{\nabla}$ ,  $\widehat{\nabla}$  and  $\overset{*}{\nabla}$ .

### 2. Preliminaries

On an almost contact metric manifold  $\overline{M}(\varphi, \xi, \eta, g)$ , we have in [4]

$$(2.1) \quad \varphi^2(X) = -X + \eta(X)\xi, \quad \varphi\xi = 0,$$

$$(2.2) \quad \eta(\xi) = 1, \quad g(X, \xi) = \eta(X), \quad \eta(\varphi X) = 0,$$

$$(2.3) \quad g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y),$$

$$(2.4) \quad g(\varphi X, Y) = -g(X, \varphi Y).$$

On  $\overline{M}^{2n+1}(f_1, f_2, f_3)$ , we have in [1]

$$(2.5) \quad (\overline{\nabla}_X \varphi)(Y) = (f_1 - f_3)[g(X, Y)\xi - \eta(Y)X].$$

The relations of  $\widetilde{\nabla}$ ,  $\widehat{\nabla}$  and  $\overset{*}{\nabla}$  with  $\overline{\nabla}$  on  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  are

$$(2.6) \quad \widetilde{\nabla}_X Y = \overline{\nabla}_X Y + \eta(Y)\varphi X - g(\varphi X, Y)\xi,$$

$$(2.7) \quad \widehat{\nabla}_X Y = \overline{\nabla}_X Y + (f_1 - f_3)\eta(Y)\varphi X - (f_1 - f_3)g(\varphi X, Y)\xi$$

and

$$(2.8) \quad \overset{*}{\nabla}_X Y = \overline{\nabla}_X Y + \eta(X)\varphi Y + (f_1 - f_3)\eta(Y)\varphi X - (f_1 - f_3)g(\varphi X, Y)\xi.$$

Let  $\widetilde{\widetilde{R}}$  (or  $\widehat{\widehat{R}}, \widehat{\widehat{R}}^*$ ) be the curvature tensor of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\widetilde{\nabla}$  ( $\widehat{\nabla}, \widehat{\nabla}^*$ , respectively). Then

$$(2.9) \quad \begin{aligned} \widetilde{\widetilde{R}}(X, Y, Z, W) = & \overline{R}(X, Y, Z, W) + g(\varphi X, Z)g(\varphi Y, W) \\ & - g(\varphi Y, Z)g(\varphi X, W) + (f_1 - f_3) [\{\eta(X)g(Y, W) \\ & - \eta(Y)g(X, W)\}\eta(Z) + \{g(X, Z)\eta(Y) \\ & - g(Y, Z)\eta(X)\}\eta(W)]. \end{aligned}$$

Also, we have

$$(2.10) \quad \begin{aligned} \widehat{\widehat{R}}(X, Y, Z, W) = & f_1\{g(Y, Z)g(X, W) - g(X, Z)g(Y, W)\} \\ & + f_2\{g(X, \varphi Z)g(\varphi Y, W) - g(Y, \varphi Z)g(\varphi X, W) \\ & + 2g(X, \varphi Y)g(\varphi Z, W)\} + \{f_3 + (f_1 - f_3)^2\} [\eta(X)\eta(Z)g(Y, W) \\ & - \eta(Y)\eta(Z)g(X, W) + \{g(X, Z)\eta(Y) - g(Y, Z)\eta(X)\}\eta(W)] \\ & + (f_1 - f_3)^2 [g(X, \varphi Z)g(\varphi Y, W) - g(Y, \varphi Z)g(\varphi X, W)], \end{aligned}$$

$$(2.11) \quad \begin{aligned} \widehat{\widehat{R}}^*(X, Y, Z, W) = & f_1\{g(Y, Z)g(X, W) - g(X, Z)g(Y, W)\} \\ & + f_2\{g(X, \varphi Z)g(\varphi Y, W) - g(Y, \varphi Z)g(\varphi X, W) \\ & + 2g(X, \varphi Y)g(\varphi Z, W)\} + \{f_3 + (f_1 - f_3)^2\} \\ & \times [\eta(X)\eta(Z)g(Y, W) - \eta(Y)\eta(Z)g(X, W) \\ & + \{g(X, Z)\eta(Y) - g(Y, Z)\eta(X)\}\eta(W)] \\ & + (f_1 - f_3)^2 [g(X, \varphi Z)g(\varphi Y, W) - g(Y, \varphi Z)g(\varphi X, W)] \\ & + 2(f_1 - f_3)g(X, \varphi Y)g(\varphi Z, W), \end{aligned}$$

where  $(f_1 - f_3)$  is a constant function.

Let  $M$  be a submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$ . If  $\nabla$  and  $\nabla^\perp$  are the induced connections on  $\Gamma(TM)$  and  $\Gamma(T^\perp M)$ , respectively, then the Gauss and Weingarten formulas are given by [17]

$$(2.12) \quad \widetilde{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad \widetilde{\nabla}_X V = -A_V X + \nabla_X^\perp V$$

for all  $X, Y \in \Gamma(TM)$  and  $V \in \Gamma(T^\perp M)$ , where  $h$  and  $A_V$  are second fundamental form and shape operator respectively and they are related by [17]  $g(h(X, Y), V) = g(A_V X, Y)$ .

Let  $\widetilde{R}$  (or  $\widehat{R}, \widehat{R}^*$ ) be the curvature tensor of  $M$  for the induced connection  $\nabla$  ( $\widetilde{\nabla}, \widehat{\nabla}, \widehat{\nabla}^*$ , respectively) and  $\widetilde{h}$  (or  $\widehat{h}, \widehat{h}^*$ ) be the second fundamental forms and  $\widetilde{A}_V$

(or  $\widehat{A}_V, \widehat{A}_V^*$ ) shape operators with respect to the induced connection  $\widetilde{\nabla}$  ( $\widehat{\nabla}, \widehat{\nabla}^*$ , respectively).

From (2.12), we have the Gauss and Ricci equations as

$$(2.13) \quad \begin{aligned} \overline{R}(X, Y, Z, W) &= R(X, Y, Z, W) - g(h(X, W), h(Y, Z)) \\ &\quad + g(h(X, Z), h(Y, W)) \end{aligned}$$

and

$$(2.14) \quad R^\perp(X, Y, \mu, \nu) = \overline{R}(X, Y, \mu, \nu) + g([A_\mu, A_\nu]X, Y)$$

where  $\mu, \nu \in \Gamma(T^\perp M)$ . In a similar way, we have

$$(2.15) \quad \begin{aligned} \widetilde{R}(X, Y, Z, W) &= \widetilde{\overline{R}}(X, Y, Z, W) + g(\widetilde{h}(X, W), \widetilde{h}(Y, Z)) \\ &\quad - g(\widetilde{h}(X, Z), \widetilde{h}(Y, W)), \end{aligned}$$

$$(2.16) \quad \widetilde{R}^\perp(X, Y, \mu, \nu) = \widetilde{\overline{R}}(X, Y, \mu, \nu) + g([\widetilde{A}_\mu, \widetilde{A}_\nu]X, Y),$$

$$(2.17) \quad \begin{aligned} \widehat{R}(X, Y, Z, W) &= \widehat{\overline{R}}(X, Y, Z, W) + g(\widehat{h}(X, W), \widehat{h}(Y, Z)) \\ &\quad - g(\widehat{h}(X, Z), \widehat{h}(Y, W)), \end{aligned}$$

$$(2.18) \quad \widehat{R}^\perp(X, Y, \mu, \nu) = \widehat{\overline{R}}(X, Y, \mu, \nu) + g([\widehat{A}_\mu, \widehat{A}_\nu]X, Y),$$

$$(2.19) \quad \begin{aligned} \overline{R}^*(X, Y, Z, W) &= \overline{R}^*(X, Y, Z, W) - g(h(X, W), h(Y, Z)) \\ &\quad + g(h^*(X, Z), h^*(Y, W)), \end{aligned}$$

$$(2.20) \quad \overline{R}^{\perp*}(X, Y, \mu, \nu) = \overline{R}^*(X, Y, \mu, \nu) + g([\overline{A}_\mu^*, \overline{A}_\nu^*]X, Y).$$

Let  $p \in M^m$  and  $\{e_1, \dots, e_m\}$  be an orthonormal basis of  $T_p M$  and  $\{e_{m+1}, \dots, e_{2n}, e_{2n+1} = \xi\}$  be an orthonormal basis of  $T^\perp M^m$ . We define the mean curvature vector as

$$H(p) = \frac{1}{m} \sum_{i=1}^m h(e_i, e_i).$$

Following [16], we define

$$(2.21) \quad K_N = -\frac{1}{4} \sum_{r,s=1}^{2n-m+1} \text{Tr} [A_r, A_s]^2,$$

where  $A_r = A_{e_{n+r}}$ ,  $r \in \{1, \dots, 2n-m+1\}$  and call it the scalar normal curvature of  $M^m$ . The normalized scalar normal curvature is given by  $\varrho_N = \frac{2}{m(m-1)}\sqrt{K_N}$ .

Since  $A_\xi = 0$ , it follows that

$$(2.22) \quad \begin{aligned} K_N &= -\frac{1}{2} \sum_{1 \leq r < s \leq 2n-m} \text{Tr} [A_r, A_s]^2 \\ &= \sum_{1 \leq r < s \leq 2n-m} \sum_{1 \leq i < j \leq m} (g([A_r, A_s]e_i, e_j))^2. \end{aligned}$$

Also we can express  $K_N$  as

$$(2.23) \quad K_N = \sum_{1 \leq r < s \leq 2n-m} \sum_{1 \leq i < j \leq m} (h_{jk}^r h_{ik}^s - h_{ik}^r h_{jk}^s)^2.$$

Again we define

$$(2.24) \quad \varrho_N = \frac{2}{m(m-1)} \left[ \sum_{1 \leq r < s \leq 2n-m} \sum_{1 \leq i < j \leq m} \left( \sum_{k=1}^m (h_{jk}^r h_{ik}^s - h_{ik}^r h_{jk}^s) \right)^2 \right]^{1/2}.$$

The normalized scalar curvature is given by

$$(2.25) \quad \varrho = \frac{2\tau}{m(m-1)} = \sum_{1 \leq i < j \leq m} \frac{2}{m(m-1)} R(e_i, e_j, e_j, e_i),$$

where  $\tau$  is the scalar curvature and  $\{e_i : i = 1, 2, \dots, m\}$  is an orthonormal basis of  $TM^m$ .

The normalized normal scalar curvature is given by

$$(2.26) \quad \varrho^\perp = \frac{2\tau^\perp}{m(m-1)} = \frac{2}{m(m-1)} \sqrt{\sum_{1 \leq i < j \leq m} \sum_{1 \leq \alpha < \beta \leq m} (R^\perp(e_i, e_j, u_\alpha, u_\beta))^2},$$

where  $R$  and  $R^\perp$  are the curvature tensor and normal curvature of  $M^m$ .

In similar of (2.25) and (2.26) we can define  $\tilde{\varrho}$ ,  $\tilde{\varrho}^\perp$ ;  $\hat{\varrho}$ ,  $\hat{\varrho}^\perp$  and  $\tilde{\varrho}^*$ ,  $\tilde{\varrho}^{\perp*}$  with respect to  $\tilde{\nabla}$ ;  $\hat{\nabla}$  and  $\tilde{\nabla}^*$  as

$$(2.27) \quad \tilde{\varrho} = \frac{2\tilde{\tau}}{m(m-1)} = \sum_{1 \leq i < j \leq m} \frac{2}{m(m-1)} \tilde{R}(e_i, e_j, e_j, e_i),$$

$$(2.28) \quad \tilde{\varrho}^\perp = \frac{2\tilde{\tau}^\perp}{m(m-1)} = \frac{2}{m(m-1)} \sqrt{\sum_{1 \leq i < j \leq m} \sum_{1 \leq \alpha < \beta \leq m} (\tilde{R}^\perp(e_i, e_j, u_\alpha, u_\beta))^2},$$

$$(2.29) \quad \hat{\varrho} = \frac{2\hat{\tau}}{m(m-1)} = \sum_{1 \leq i < j \leq m} \frac{2}{m(m-1)} \hat{R}(e_i, e_j, e_j, e_i),$$

$$(2.30) \quad \hat{\varrho}^\perp = \frac{2\hat{\tau}^\perp}{m(m-1)} = \frac{2}{m(m-1)} \sqrt{\sum_{1 \leq i < j \leq m} \sum_{1 \leq \alpha < \beta \leq m} (\hat{R}^\perp(e_i, e_j, u_\alpha, u_\beta))^2},$$

$$(2.31) \quad \varrho^* = \frac{2\tau^*}{m(m-1)} = \sum_{1 \leq i < j \leq m} \frac{2}{m(m-1)} R^*(e_i, e_j, e_j, e_i),$$

$$(2.32) \quad \varrho^{*\perp} = \frac{2\tau^{*\perp}}{m(m-1)} = \frac{2}{m(m-1)} \sqrt{\sum_{1 \leq i < j \leq m} \sum_{1 \leq \alpha < \beta \leq m} (\tilde{R}^\perp(e_i, e_j, u_\alpha, u_\beta))^2}.$$

A submanifold  $M^m$  of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  is said to be  $C$ -totally real submanifold if  $\xi \in \Gamma(T^\perp M)$  and  $\varphi X \in \Gamma(T^\perp M)$  for all  $X \in \Gamma(TM)$ . In particular, if  $m = n$ , then  $M^n$  is called Legendrian submanifold.

### 3. Some basic results

**Proposition 3.1.** *Let  $M$  be a  $C$ -totally real submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\tilde{\nabla}$ . Then following relations hold on  $M$ :*

- (i)  $\tilde{h}(X, Y) = h(X, Y)$ ,  $\tilde{H} = H$ ;
- (ii)  $\tilde{A}_V X = A_V X$ .

PROOF: From (2.12), we have

$$(3.1) \quad \tilde{\nabla}_X Y = \tilde{\nabla}_X Y + \tilde{h}(X, Y)$$

and

$$(3.2) \quad \tilde{\nabla}_X V = \tilde{\nabla}_X^\perp V - \tilde{A}_V X.$$

From (2.6), (2.12) and (3.1) we have

$$(3.3) \quad \tilde{\nabla}_X Y + \tilde{h}(X, Y) = \nabla_X Y + h(X, Y) + \eta(Y)\varphi X - g(\varphi X, Y)\xi$$

for any  $X, Y \in \Gamma(TM)$ .

Since  $\xi \in \Gamma(T^\perp M)$  and  $\varphi X \in \Gamma(T^\perp M)$  for all  $X$ , then from (3.3) we have

$$\tilde{\nabla}_X Y = \nabla_X Y \quad \text{and} \quad \tilde{h}(X, Y) = h(X, Y).$$

Again from (2.6), (2.12) and (3.2)

$$(3.4) \quad \tilde{\nabla}^{\perp}_X V - \tilde{A}_V X = \nabla^{\perp}_X V - A_V X + \eta(V)\varphi X - g(\varphi X, V)\xi.$$

Equating tangential and normal part of (3.4) we have

$$\tilde{A}_V X = A_V X, \quad \tilde{\nabla}^{\perp}_X V = \nabla^{\perp}_X V + \eta(V)\varphi X - g(\varphi X, V)\xi.$$

□

**Proposition 3.2.** *Let  $M$  be a  $C$ -totally real submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\tilde{\nabla}$ . Then following relations hold on  $M$ :*

- (i)  $\hat{h}(X, Y) = h(X, Y), \hat{H} = H;$
- (ii)  $\hat{A}_V X = A_V X.$

PROOF: The proof is similar to the proof of Proposition 3.1. □

**Proposition 3.3.** *Let  $M$  be a  $C$ -totally real submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\tilde{\nabla}^*$ . Then following relations hold on  $M$ :*

- (i)  $\overset{*}{h}(X, Y) = h(X, Y), \overset{*}{H} = H;$
- (ii)  $\overset{*}{A}_V X = A_V X.$

PROOF: The proof is similar to the proof of Proposition 3.1. □

#### 4. Wintgen inequality on Legendrian submanifolds of $\overline{M}^{2n+1}(f_1, f_2, f_3)$ with respect to $\tilde{\nabla}$

**Proposition 4.1.** *Let  $M^m$  be a  $C$ -totally real submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\tilde{\nabla}$ . Then*

$$(4.1) \quad \|H\|^2 + f_1 \geq \tilde{q} + \varrho_N.$$

PROOF: We see that

$$(4.2) \quad m^2 \|H\|^2 = \sum_{r=1}^{2n-m} \left( \sum_{i=1}^m h_{ii}^r \right)^2 = \frac{1}{m-1} \sum_{r=1}^{2n-m} \sum_{1 \leq i < j \leq m} (h_{ii}^r - h_{jj}^r)^2 + \frac{2m}{m-1} \sum_{r=1}^{2n-m} \sum_{1 \leq i < j \leq m} h_{ii}^r h_{jj}^r.$$



From [7] we have the inequality

$$(4.3) \quad \sum_{r=1}^{2n-m} \sum_{1 \leq i < j \leq m} (h_{ii}^r - h_{jj}^r)^2 + 2 \sum_{r=1}^{2n-m} \sum_{1 \leq i < j \leq m} h_{ij}^r h_{ij}^r \geq \left[ \sum_{1 \leq r < s \leq 2n-m} \sum_{1 \leq i < j \leq m} \left( \sum_{k=1}^m (h_{jk}^r h_{ik}^s - h_{ik}^r h_{jk}^s) \right)^2 \right]^{1/2}.$$

From (2.24), (4.2) and (4.3), we get

$$(4.4) \quad m^2 \|H\|^2 - m^2 \varrho_N \geq \frac{2m}{m-1} \sum_{r=1}^{2n-m} \sum_{1 \leq i < j \leq m} [h_{ii}^r h_{jj}^r - (h_{ij}^r)^2].$$

Now from (2.9) and (2.15) we have

$$(4.5) \quad \tilde{\tau} = \tilde{R}(e_i, e_j, e_j, e_i) = \frac{m(m-1)}{2} f_1 + \sum_{r=1}^{2n-m} \sum_{1 \leq i < j \leq m} [h_{ii}^r h_{jj}^r - (h_{ij}^r)^2].$$

Substituting (4.5) in (4.4) we get (4.1). □

**Theorem 4.1.** *Let  $M^n$  be a Legendrian submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\tilde{\nabla}$ . Then*

$$(4.6) \quad (\tilde{\varrho}^\perp)^2 \leq (\|H\|^2 - \tilde{\varrho} + f_1) + \frac{2}{n(n-1)}(f_2 - 1)^2 + \frac{4}{n(n-1)}(f_2 - 1)(\tilde{\varrho} - f_1).$$

PROOF: Let us consider  $\{e_1, \dots, e_n\}$  be an orthonormal basis of  $TM^n$  and  $\{e_{n+1} = \varphi e_1, \dots, e_{2n} = \varphi e_n, e_{2n+1} = \xi\}$  be orthonormal basis of  $T^\perp M^n$ . Now from (2.9) and (2.16) we have

$$(4.7) \quad \begin{aligned} g(\tilde{R}^\perp(e_i, e_j)e_{n+r}, e_{n+s}) &= f_2[g(\varphi e_i, e_{n+s})g(\varphi e_j, e_{n+r}) - g(\varphi e_i, e_{n+r})g(\varphi e_j, e_{n+s})] \\ &\quad + \{g(\varphi e_i, e_{n+r})g(\varphi e_j, e_{n+s}) - g(\varphi e_j, e_{n+r})g(\varphi e_i, e_{n+s})\} \\ &\quad + g([A_r, A_s]e_i, e_j) \\ &= (f_2 - 1)(\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js}) + g([A_r, A_s]e_i, e_j). \end{aligned}$$

Contracting (4.7) and using (2.28) we have

$$\begin{aligned}
 (\tilde{\tau}^\perp)^2 &= \sum_{1 \leq r < s \leq n} \sum_{1 \leq i < j \leq n} g^2(\tilde{R}^\perp(e_i, e_j)e_{n+r}, e_{n+s}) \\
 &= \sum_{1 \leq r < s \leq n} \sum_{1 \leq i < j \leq n} [(f_2 - 1)(\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js}) + g([A_r, A_s]e_i, e_j)]^2 \\
 (4.8) \quad &= \sum_{1 \leq r < s \leq n} \sum_{1 \leq i < j \leq n} [g^2([A_r, A_s]e_i, e_j) + (f_2 - 1)^2(\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js})^2 \\
 &\quad + 2(f_2 - 1)(\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js})g([A_r, A_s]e_i, e_j)] \\
 &= \frac{n^2(n-1)^2}{4} \varrho_N^2 + \frac{n(n-1)(f_2-1)^2}{2} - (f_2-1)\|h\|^2 \\
 &\quad + (f_2-1)n^2\|H\|^2.
 \end{aligned}$$

From (2.9), (2.15) and (2.27) we have

$$2\tilde{\tau} = n^2\|H\|^2 - \|h\|^2 + n(n-1)f_1$$

or equivalently,

$$(4.9) \quad n^2\|H\|^2 - \|h\|^2 = n(n-1)(\tilde{\varrho} - f_1).$$

Substituting (4.9) in (4.8) and using (2.28) we get

$$(4.10) \quad (\tilde{\varrho}^\perp)^2 \leq \varrho_N^2 + \frac{4}{n(n-1)}(\tilde{\varrho} - f_1)(f_2 - 1)\frac{2(f_2 - 1)^2}{n(n-1)}.$$

By virtue of Proposition 4.1 and (4.10), we obtain the inequality (4.6).  $\square$

## 5. Wintgen inequality on Legendrian submanifolds of $\overline{M}^{2n+1}(f_1, f_2, f_3)$ with respect to $\widehat{\nabla}$

**Proposition 5.1.** *Let  $M^m$  be a  $C$ -totally real submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\widehat{\nabla}$ . Then*

$$(5.1) \quad \|H\|^2 + f_1 \geq \hat{\varrho} + \varrho_N.$$

PROOF: The proof is similar to the proof of Proposition 4.1  $\square$

**Theorem 5.1.** *Let  $M^n$  be a Legendrian submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\widehat{\nabla}$ . Then*

$$(5.2) \quad \begin{aligned} (\widehat{\varrho}^\perp)^2 &\leq (\|H\|^2 - \widehat{\varrho} + f_1) + \frac{2}{n(n-1)}(f_2 + (f_1 - f_3)^2)^2 \\ &\quad + \frac{4}{n(n-1)}(f_2 + (f_1 - f_3)^2)(\widehat{\varrho} - f_1). \end{aligned}$$

PROOF: Now from (2.10) and (2.18) we have

$$(5.3) \quad \begin{aligned} &g(\widehat{R}^\perp(e_i, e_j)e_{n+r}, e_{n+s}) \\ &= f_2[g(\varphi e_i, e_{n+s})g(\varphi e_j, e_{n+r}) - g(\varphi e_i, e_{n+r})g(\varphi e_j, e_{n+s})] \\ &\quad + (f_1 - f_3)^2\{g(\varphi e_j, e_{n+r})g(\varphi e_i, e_{n+s}) \\ &\quad - g(\varphi e_i, e_{n+r})g(\varphi e_j, e_{n+s})\} + g([A_r, A_s]e_i, e_j) \\ &= (f_2 + (f_1 - f_3)^2)(\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js}) + g([A_r, A_s]e_i, e_j). \end{aligned}$$

From (2.30) and (5.3) we have

$$(5.4) \quad \begin{aligned} (\widehat{\tau}^\perp)^2 &= \sum_{1 \leq r < s \leq n} \sum_{1 \leq i < j \leq n} g(\widehat{R}^\perp(e_i, e_j)e_{n+r}, e_{n+s})^2 \\ &= \sum_{1 \leq r < s \leq n} \sum_{1 \leq i < j \leq n} [(f_2 + (f_1 - f_3)^2)(\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js}) \\ &\quad + g([A_r, A_s]e_i, e_j)]^2 \\ &= \sum_{1 \leq r < s \leq n} \sum_{1 \leq i < j \leq n} [g^2([A_r, A_s]e_i, e_j) + (f_2 + (f_1 - f_3)^2)^2 \\ &\quad \times (\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js})^2 + 2(f_2 + (f_1 - f_3)^2)(\delta_{is}\delta_{jr} - \delta_{ir}\delta_{js}) \\ &\quad \times g([A_r, A_s]e_i, e_j)] \\ &= \frac{n^2(n-1)^2}{4} \varrho_N^2 + \frac{n(n-1)(f_2 + (f_1 - f_3)^2)^2}{2} \\ &\quad - (f_2 + (f_1 - f_3)^2)\|h\|^2 + f_2 n^2 \|H\|^2. \end{aligned}$$

From (2.10) and (2.17) we have

$$2\widehat{\tau} = n^2\|H\|^2 - \|h\|^2 + n(n-1)f_1$$

or equivalently,

$$(5.5) \quad n^2\|H\|^2 - \|h\|^2 = n(n-1)(\widehat{\varrho} - f_1).$$

Substituting (5.5) in (5.4) and using (2.30) we get

$$(5.6) \quad (\hat{\varrho}^\perp)^2 \leq \varrho_N^2 + \frac{4}{n(n-1)}(\hat{\varrho} - f_1)(f_2 + (f_1 - f_3)^2) + \frac{2(f_2 + (f_1 - f_3)^2)^2}{n(n-1)}.$$

By virtue of Proposition 5.1 and (5.6) we have the inequality (5.2). □

**6. Wintgen inequality on Legendrian submanifolds of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\overset{*}{\nabla}$**

**Proposition 6.1.** *Let  $M^m$  be a  $C$ -totally real submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\overset{*}{\nabla}$ . Then*

$$(6.1) \quad \|H\|^2 + f_1 \geq \overset{*}{\varrho} + \varrho_N.$$

PROOF: The proof is similar to the proof of Proposition 4.1. □

**Theorem 6.1.** *Let  $M^n$  be a Legendrian submanifold of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to  $\overset{*}{\nabla}$ . Then*

$$(6.2) \quad \begin{aligned} (\overset{*}{\varrho}^\perp)^2 &\leq (\|H\|^2 - \hat{\varrho} + f_1) + \frac{2}{n(n-1)}(f_2 + (f_1 - f_3)^2)^2 \\ &\quad + \frac{4}{n(n-1)}(f_2 + (f_1 - f_3)^2)(\hat{\varrho} - f_1). \end{aligned}$$

PROOF: The proof is similar to the proof of Theorem 5.1. □

**7. Summary**

Here, we present a summary of the results obtained on submanifolds of  $\overline{M}^{2n+1}(f_1, f_2, f_3)$  with respect to the three connections considered; namely, quarter symmetric metric connection  $\overset{\sim}{\nabla}$ , Schouten–van Kampen connection  $\overset{\wedge}{\nabla}$  and Tanaka–Webster connection  $\overset{*}{\nabla}$ .

Connection	$C$ -totally inequality	Wintgen inequality
$\tilde{\nabla}$	$\ H\ ^2 + f_1 \geq \tilde{\varrho} + \varrho_N$	$(\tilde{\varrho}^\perp)^2 \leq (\ H\ ^2 - \tilde{\varrho} + f_1)$ $+ \frac{2}{n(n-1)}(f_2 - 1)^2$ $+ \frac{4}{n(n-1)}(f_2 - 1)(\tilde{\varrho} - f_1)$
$\hat{\nabla}$	$\ H\ ^2 + f_1 \geq \hat{\varrho} + \varrho_N$	$(\hat{\varrho}^\perp)^2 \leq (\ H\ ^2 - \hat{\varrho} + f_1)$ $+ \frac{2}{n(n-1)}(f_2 + (f_1 - f_3)^2)^2$ $+ \frac{4}{n(n-1)}(f_2 + (f_1 - f_3)^2)(\hat{\varrho} - f_1)$
$\overset{*}{\nabla}$	$\ H\ ^2 + f_1 \geq \overset{*}{\varrho} + \varrho_N$	$(\overset{*}{\varrho}^\perp)^2 \leq (\ H\ ^2 - \overset{*}{\varrho} + f_1)$ $+ \frac{2}{n(n-1)}(f_2 + (f_1 - f_3)^2)^2$ $+ \frac{4}{n(n-1)}(f_2 + (f_1 - f_3)^2)(\overset{*}{\varrho} - f_1)$

REFERENCES

- [1] Alegre P., Blair D. E., Carriazo A., *Generalized Sasakian-space-forms*, Israel J. Math. **141** (2004), 157–183.
- [2] Alegre P., Carriazo A., *Structures on generalized Sasakian-space-forms*, Differential Geom. Appl. **26** (2008), no. 6, 656–666.
- [3] Bejancu A., *Schouten–van Kampen and Vranceanu connections on foliated manifolds*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.) **52** (2006), no. 1, 37–60.
- [4] Blair D. E., *Contact Manifolds in Riemannian Geometry*, Lecture Notes in Mathematics, 509, Springer, Berlin, 1976.
- [5] Carriazo A., *On generalized Sasakian-space-forms*, Proc. of the Ninth International Workshop on Differential Geometry, Kyungpook Nat. Univ., Taegu, (2005), 31–39.
- [6] Gołąb S., *On semi-symmetric and quarter-symmetric linear connections*, Tensor (N.S.) **29** (1975), no. 3, 249–254.
- [7] Lu Z., *Normal scalar curvature conjecture and its applications*, J. Funct. Anal. **261** (2011), no. 5, 1248–1308.
- [8] Mihai I., *On the generalized Wintgen inequality for Lagrangian submanifolds in complex space forms*, Nonlinear Anal. **95** (2014), 714–720.
- [9] Mihai I., *On the generalized Wintgen inequality for Legendrian submanifolds in Sasakian space forms*, Tohoku Math. J. (2) **69** (2017), no. 1, 43–53.
- [10] Olszak Z., *The Schouten van-Kampen affine connection adapted to an almost (para) contact metric structure*, Publ. Inst. Math. (Beograd) (N.S.) **94(108)** (2013), 31–42.
- [11] Schouten J. A., van Kampen E. R., *Zur Einbettungs- und Krümmungstheorie nicht-holonomer Gebilde*, Math. Ann. **103** (1930), 752–783 (German).
- [12] Tanaka, N., *On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections*, Japan J. Math. (N.S.) **2** (1976), no. 1, 131–190.
- [13] Tanno, S., *Variational problems on contact Riemannian manifolds*, Trans. Amer. Math. Soc. **314** (1989), no. 1, 349–379.
- [14] Webster S. M., *Pseudo-Hermitian structures on a real hypersurface*, J. Differential Geometry **13** (1978), no. 1, 25–41.
- [15] Wintgen P., *Sur l'inégalité de Chen-Willmore*, C. R. Acad. Sci. Paris Sér. A-B **288** (1979), no. 21, A993–A995 (French. English summary).

- [16] Yano K., Kon M., *Anti-invariant Submanifolds*, Lecture Notes in Pure and Applied Mathematics, 21, Marcel Dekker, New York, 1976.
- [17] Yano K., Kon M., *Structures on Manifolds*, Series in Pure Mathematics, 3, World Scientific Publishing, Singapore, 1984.

S. K. Hui (corresponding author):

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF BURDWAN, GOLAPBAG,  
BURDWAN 713104, WEST BENGAL, INDIA

*E-mail:* skhui@math.buruniv.ac.in

R. S. Lemence:

DEPARTMENT OF MATHEMATICS, INSTITUTE OF ARTS AND SCIENCES,  
FAR EASTERN UNIVERSITY, NICANOR REYES ST, SAMPALOC, MANILA 1008,  
PHILIPPINES

*E-mail:* rlemence@feu.edu.ph

P. Mandal:

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF BURDWAN, GOLAPBAG,  
BURDWAN 713104, WEST BENGAL, INDIA

*E-mail:* pm2621994@gmail.com

(Received September 11, 2018, revised January 14, 2019)