

The probabilistic approach to limited packings in graphs

Andrei Gagarin^{a,*}, Vadim Zverovich^b

^a*Department of Computer Science, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, UK*

^b*University of the West of England, Bristol, BS16 1QY, UK*

Abstract

We consider (closed neighbourhood) packings and their generalization in graphs. A vertex set X in a graph G is a k -limited packing if for every vertex $v \in V(G)$, $|N[v] \cap X| \leq k$, where $N[v]$ is the closed neighbourhood of v . The k -limited packing number $L_k(G)$ of a graph G is the largest size of a k -limited packing in G . Limited packing problems can be considered as secure facility location problems in networks.

In this paper, we develop a new application of the probabilistic method to limited packings in graphs, resulting in lower bounds for the k -limited packing number and a randomized algorithm to find k -limited packings satisfying the bounds. In particular, we prove that for any graph G of order n with maximum vertex degree Δ ,

$$L_k(G) \geq \frac{kn}{(k+1) \sqrt[k]{\binom{\Delta}{k}} (\Delta+1)}.$$

Also, some other upper and lower bounds for $L_k(G)$ are given.

Keywords: k -Limited packings, The probabilistic method, Lower and upper bounds, Randomized algorithm

1. Introduction

We consider simple undirected graphs. If not specified otherwise, standard graph-theoretic terminology and notations are used (e.g., see [1, 2]). We are interested in the classical packings and packing numbers of graphs as introduced in [9], and their generalization, called limited packings and limited packing numbers, respectively, as presented in [6]. In the literature, the classical packings are often referred to under different names: for example, as (distance) 2-packings [9, 13], closed neighborhood packings [11] or strong stable sets [8]. They can also be considered as generalizations of independent (stable) sets which, following the terminology of [9], would be (distance) 1-packings.

*Corresponding author, fax: +44 (0) 1784 439786, phone: +44 (0) 1784 443437

Email addresses: andrei.gagarin@rhul.ac.uk (Andrei Gagarin),
vadim.zverovich@uwe.ac.uk (Vadim Zverovich)

Formally, a vertex set X in a graph G is a *k-limited packing* if for every vertex $v \in V(G)$,

$$|N[v] \cap X| \leq k,$$

where $N[v]$ is the closed neighbourhood of v . The *k-limited packing number* $L_k(G)$ of a graph G is the maximum size of a *k-limited packing* in G . In these terms, the classical (distance) 2-packings are 1-limited packings, and hence $\rho(G) = L_1(G)$, where $\rho(G)$ is the 2-packing number.

The problem of finding a 2-packing (1-limited packing) of maximum size is shown to be *NP*-complete by Hochbaum and Schmoys [8]. In [4], it is shown that the problem of finding a maximum size *k-limited packing* is *NP*-complete even for the classes of split and bipartite graphs.

Graphs usually serve as underlying models for networks. A number of interesting application scenarios of limited packings are described in [6], including network security, market saturation, and codes. These and others can be summarized as secure location or distribution of facilities in a network. In a more general sense, these problems can be viewed as (maximization) facility location problems to place/distribute in a given network as many resources as possible subject to some (security) constraints.

2-Packings (1-limited packings) are well-studied in the literature from the structural and algorithmic point of view (e.g., see [8, 9, 11, 12]) and in connection with other graph parameters (e.g., see [3, 7, 9, 11, 13]). In particular, several papers discuss connections between packings and dominating sets in graphs (e.g., see [3, 4, 6, 7, 11]). Although the formal definitions for packings and dominating sets may appear to be similar, the problems have a very different nature: one of the problems is a maximization problem not to break some (security) constraints, and the other is a minimization problem to satisfy some reliability requirements. For example, given a graph G , the definitions imply a simple inequality $\rho(G) \leq \gamma(G)$, where $\gamma(G)$ is the domination number of G (e.g., see [11]). However, the difference between $\rho(G)$ and $\gamma(G)$ can be arbitrarily large as illustrated in [3]: $\rho(K_n \times K_n) = 1$ for the Cartesian product of complete graphs, but $\gamma(K_n \times K_n) = n$.

In this paper, we develop an application of the probabilistic method to *k-limited packings* in general and to 2-packings (1-limited packings) in particular. In Section 2 we present the probabilistic construction and use it to derive two lower bounds for the *k-limited packing number* $L_k(G)$. Also, using a greedy algorithm approach, we provide an improved lower bound for the 2-packing (1-limited packing) number $\rho(G) = L_1(G)$. The probabilistic construction implies a randomized algorithm to find *k-limited packings* satisfying the lower bounds. The algorithm and its analysis are presented in Section 3. Section 4 shows that the main lower bound is asymptotically sharp, and discusses the improvement for 1-limited packings from the greedy algorithm approach. Finally, Section 5 provides upper bounds for $L_k(G)$, e.g. in terms of the *k-tuple domination number* $\gamma_{\times k}(G)$.

Notice that the probabilistic construction and approach are different from the well-known probabilistic constructions used for independent sets (e.g., see [1], p.27–28). In terms of packings, an independent set in a graph G is a distance 1-packing: for any two vertices in an independent set, the distance between them in G is greater

than 1. To the best of our knowledge, the proposed application of the probabilistic method is a new approach to work with packings and related maximization problems.

2. The probabilistic construction and lower bounds

Let $\Delta = \Delta(G)$ denote the maximum vertex degree in a graph G . Notice that $L_k(G) = n$ when $k \geq \Delta + 1$. We define

$$c_t = c_t(G) = \binom{\Delta}{t} \quad \text{and} \quad \tilde{c}_t = \tilde{c}_t(G) = \binom{\Delta + 1}{t}.$$

In what follows, we put $\binom{a}{b} = 0$ if $b > a$.

The following theorem gives a new lower bound for the k -limited packing number. It may be pointed out that the probabilistic construction used in the proof of Theorem 1 implies a randomized algorithm for finding a k -limited packing set, whose size satisfies the bound of Theorem 1 with a positive probability (see Algorithm 1 in Section 3).

Theorem 1. *For any graph G of order n with $\Delta \geq k \geq 1$,*

$$L_k(G) \geq \frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}}. \quad (1)$$

PROOF. Let A be a set formed by an independent choice of vertices of G , where each vertex is selected with the probability

$$p = \left(\frac{1}{\tilde{c}_{k+1} (1+k)} \right)^{1/k}. \quad (2)$$

For $m = k, \dots, \Delta$, we denote

$$A_m = \{v \in A : |N(v) \cap A| = m\}.$$

For each set A_m , we form a set A'_m in the following way. For every vertex $v \in A_m$, we take $m - (k - 1)$ neighbours from $N(v) \cap A$ and add them to A'_m . Such neighbours always exist because $m \geq k$. It is obvious that

$$|A'_m| \leq (m - k + 1)|A_m|.$$

For $m = k + 1, \dots, \Delta$, let us denote

$$B_m = \{v \in V(G) - A : |N(v) \cap A| = m\}.$$

For each set B_m , we form a set B'_m by taking $m - k$ neighbours from $N(v) \cap A$ for every vertex $v \in B_m$. We have

$$|B'_m| \leq (m - k)|B_m|.$$

Let us construct the set X as follows:

$$X = A - \left(\bigcup_{m=k}^{\Delta} A'_m \right) - \left(\bigcup_{m=k+1}^{\Delta} B'_m \right).$$

It is easy to see that X is a k -limited packing in G . The expectation of $|X|$ is

$$\begin{aligned} \mathbf{E}[|X|] &\geq \mathbf{E} \left[|A| - \sum_{m=k}^{\Delta} |A'_m| - \sum_{m=k+1}^{\Delta} |B'_m| \right] \\ &\geq \mathbf{E} \left[|A| - \sum_{m=k}^{\Delta} (m-k+1)|A_m| - \sum_{m=k+1}^{\Delta} (m-k)|B_m| \right] \\ &= pn - \sum_{m=k}^{\Delta} (m-k+1)\mathbf{E}[|A_m|] - \sum_{m=k+1}^{\Delta} (m-k)\mathbf{E}[|B_m|]. \end{aligned}$$

Let us denote the vertices of G by v_1, v_2, \dots, v_n and the corresponding vertex degrees by d_1, d_2, \dots, d_n . We will need the following lemma:

Lemma 2. *If $p = \left(\frac{1}{\tilde{c}_{k+1}(1+k)} \right)^{1/k}$, then, for any vertex $v_i \in V(G)$,*

$$\binom{d_i}{m} (1-p)^{d_i-m} \leq \binom{\Delta}{m} (1-p)^{\Delta-m}, \quad m \geq k. \quad (3)$$

PROOF. The inequality (3) holds if $d_i = \Delta$. It is also true if $d_i < m$ because in this case $\binom{d_i}{m} = 0$. Thus, we may assume that

$$m \leq d_i < \Delta.$$

Now, it is easy to see that inequality (3) is equivalent to the following:

$$(1-p)^{\Delta-d_i} \geq \binom{d_i}{m} / \binom{\Delta}{m} = \frac{(\Delta-m)!/(d_i-m)!}{\Delta!/d_i!} = \prod_{i=0}^{\Delta-d_i-1} \frac{\Delta-m-i}{\Delta-i}. \quad (4)$$

Further, $\Delta \geq k$ implies $\frac{\Delta}{k} \leq \frac{\Delta-i}{k-i}$, where $0 \leq i \leq k-1$. Taking into account that $\Delta > 0$, we obtain

$$\left(\frac{\Delta}{k} \right)^k \leq \prod_{i=0}^{k-1} \frac{\Delta-i}{k-i} = c_k < \tilde{c}_{k+1}(1+k)$$

or

$$\frac{1}{\tilde{c}_{k+1}(1+k)} < \left(\frac{k}{\Delta} \right)^k.$$

Thus,

$$p^k < \left(\frac{k}{\Delta} \right)^k \quad \text{or} \quad p < \frac{k}{\Delta} \leq \frac{m}{\Delta}.$$

We have $p < \frac{m}{\Delta}$, which is equivalent to $1 - p > \frac{\Delta - m}{\Delta}$. Therefore,

$$(1 - p)^{\Delta - d_i} > \left(\frac{\Delta - m}{\Delta} \right)^{\Delta - d_i} \geq \prod_{i=0}^{\Delta - d_i - 1} \frac{\Delta - m - i}{\Delta - i},$$

as required in (4). \square

Now we go on with the proof of Theorem 1. By Lemma 2,

$$\begin{aligned} \mathbf{E}[|A_m|] &= \sum_{i=1}^n \mathbf{P}[v_i \in A_m] \\ &= \sum_{i=1}^n p \binom{d_i}{m} p^m (1 - p)^{d_i - m} \\ &\leq p^{m+1} \sum_{i=1}^n \binom{\Delta}{m} (1 - p)^{\Delta - m} \\ &= p^{m+1} (1 - p)^{\Delta - m} c_m n, \end{aligned}$$

where $p \binom{d_i}{m} p^m (1 - p)^{d_i - m}$ is the probability of having vertex v_i , $i = 1, \dots, n$, in the set A_m , $m = k, \dots, \Delta$. Again, by Lemma 2,

$$\begin{aligned} \mathbf{E}[|B_m|] &= \sum_{i=1}^n \mathbf{P}[v_i \in B_m] \\ &= \sum_{i=1}^n (1 - p) \binom{d_i}{m} p^m (1 - p)^{d_i - m} \\ &\leq p^m \sum_{i=1}^n \binom{\Delta}{m} (1 - p)^{\Delta - m + 1} \\ &= p^m (1 - p)^{\Delta - m + 1} c_m n, \end{aligned}$$

where $(1 - p) \binom{d_i}{m} p^m (1 - p)^{d_i - m}$ is the probability of having vertex v_i , $i = 1, \dots, n$, in the set B_m , $m = k + 1, \dots, \Delta$.

Taking into account that $c_{\Delta+1} = \binom{\Delta}{\Delta+1} = 0$, we obtain

$$\begin{aligned} \mathbf{E}[|X|] &\geq pn - \sum_{m=k}^{\Delta} (m - k + 1) p^{m+1} (1 - p)^{\Delta - m} c_m n - \sum_{m=k+1}^{\Delta+1} (m - k) p^m (1 - p)^{\Delta - m + 1} c_m n \\ &= pn - \sum_{m=0}^{\Delta - k} (m + 1) p^{m+k+1} (1 - p)^{\Delta - m - k} c_{m+k} n \\ &\quad - \sum_{m=0}^{\Delta - k} (m + 1) p^{m+k+1} (1 - p)^{\Delta - m - k} c_{m+k+1} n \end{aligned}$$

$$\begin{aligned}
&= pn - \sum_{m=0}^{\Delta-k} (m+1)p^{m+k+1}(1-p)^{\Delta-m-k} n (c_{m+k} + c_{m+k+1}) \\
&= pn - p^{k+1}n \sum_{m=0}^{\Delta-k} (m+1)\tilde{c}_{m+k+1}p^m(1-p)^{\Delta-k-m}.
\end{aligned}$$

Furthermore,

$$\begin{aligned}
(m+1)\tilde{c}_{m+k+1} &= \binom{\Delta-k}{m} \frac{(m+1)!(\Delta+1)!}{(m+k+1)!(\Delta-k)!} \\
&\leq \binom{\Delta-k}{m} \frac{(\Delta+1)!}{(k+1)!(\Delta-k)!} = \binom{\Delta-k}{m} \tilde{c}_{k+1}.
\end{aligned}$$

We obtain, by the binomial theorem,

$$\begin{aligned}
\mathbf{E}[|X|] &\geq pn - p^{k+1}n \sum_{m=0}^{\Delta-k} \binom{\Delta-k}{m} \tilde{c}_{k+1}p^m(1-p)^{\Delta-k-m} \\
&= pn - p^{k+1}n\tilde{c}_{k+1} \\
&= pn(1 - p^k\tilde{c}_{k+1}) \\
&= \frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}}.
\end{aligned}$$

Since the expectation is an average value, there exists a particular k -limited packing of size at least $\frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}}$, as required. The proof of the theorem is complete. \square

The lower bound of Theorem 1 can be written in a simpler but weaker form as follows:

Corollary 3. *For any graph G of order n ,*

$$L_k(G) > \frac{kn}{e(1+\Delta)^{1+1/k}}.$$

PROOF. It is not difficult to see that

$$\tilde{c}_{k+1} \leq \frac{(\Delta+1)^{k+1}}{(k+1)!}$$

and, using Stirling's formula,

$$(k!)^{1/k} > \left(\sqrt{2\pi k} \left(\frac{k}{e} \right)^k \right)^{1/k} = \sqrt[2k]{2\pi k} \frac{k}{e}.$$

By Theorem 1,

$$L_k(G) \geq \frac{kn((k+1)!)^{1/k}}{(\Delta+1)^{1+1/k}(1+k)^{1+1/k}} > \frac{kn}{e(1+\Delta)^{1+1/k}} \times \frac{\sqrt[2k]{2\pi k} k}{1+k} > \frac{kn}{e(1+\Delta)^{1+1/k}}.$$

Note that $\frac{2^k \sqrt{2\pi k}}{1+k} = \frac{2^k \sqrt{2\pi k}}{1+1/k} > 1$. The last inequality is obviously true for $k = 1$, while for $k \geq 2$ it can be rewritten in the equivalent form: $2\pi k > (1 + 1/k)^{2k} = e^2 - o(1)$. \square

In the case $k = 1$, Theorem 1 gives the following lower bound for the 2-packing (1-limited packing) number:

Corollary 4. *For any graph G of order n with $\Delta \geq 1$,*

$$\rho(G) = L_1(G) \geq \frac{n}{2\Delta(\Delta + 1)}. \quad (5)$$

Let $\delta = \delta(G)$ denote the minimum vertex degree in a graph G . The lower bound of Corollary 4 can be improved as follows:

Theorem 5. *For any graph G of order n ,*

$$\rho(G) = L_1(G) \geq \frac{n + \Delta(\Delta - \delta)}{\Delta^2 + 1} \geq \frac{n}{\Delta^2 + 1}. \quad (6)$$

PROOF. Choose any vertex $v \in V(G)$ of the minimum degree δ in G . Then add v to a set X and remove vertices of $N[N[v]]$ from the graph to obtain $G' = G - N[N[v]]$, where $N[N[v]] = \{w : w \in N[u] \text{ for some } u \in N[v]\}$ is the so-called *second closed neighbourhood* of v in G . Recursively apply the same procedure to the remaining graph G' until it is empty. It is not difficult to see that X is a 1-limited packing (distance 2-packing) of size at least $\left\lceil \frac{n + \Delta(\Delta - \delta)}{\Delta^2 + 1} \right\rceil$: we remove at most $1 + \Delta + \Delta(\Delta - 1) = 1 + \Delta^2$ vertices at each iteration, but at most $1 + \delta + \delta(\Delta - 1) = 1 + \delta\Delta$ vertices at the first iteration, and $(1 + \Delta^2) - (1 + \delta\Delta) = \Delta(\Delta - \delta)$. \square

The proof of Theorem 5 provides a greedy algorithm to find a distance 2-packing (1-limited packing) satisfying bound (6). We explain later in Section 4 why the lower bound of Theorem 5 is as good as lower bound (5) of Corollary 4 for almost all graphs.

3. Randomized algorithm

A pseudocode presented in Algorithm 1 explicitly describes a randomized algorithm to find a k -limited packing set, whose size satisfies bound (1) with a positive probability. Notice that Algorithm 1 constructs a (preliminary) k -limited packing X' by recursively removing unwanted vertices from a randomly generated set A . This is different from the probabilistic construction used in the proof of Theorem 1. The recursive removal of vertices from the set A may be more effective and efficient, especially if one tries to remove overall as few vertices as possible from A by maximizing intersections of the sets A'_m ($m = k, \dots, \Delta$) and B'_m ($m = k+1, \dots, \Delta$).

At the final stage, Algorithm 1 does a (greedy) extension of the preliminary k -limited packing X' derived from the randomly generated set A . Our experimental tests with randomly generated problem instances show the following: although

the randomized part of Algorithm 1 may eventually return a preliminary k -limited packing set slightly smaller than lower bound (1), the extension of this set to a maximal k -limited packing always satisfies (1). This is of no surprise, because the expectation of the size of randomly formed set A in Algorithm 1 is $\mathbf{E}[|A|] = pn$, where $p = \left(\binom{\Delta}{k} (\Delta + 1) \right)^{-1/k}$, while the expression for lower bound in (1) yields a smaller value:

$$\frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}} = \frac{k}{k+1} pn = \frac{k}{k+1} \mathbf{E}[|A|] < \mathbf{E}[|A|].$$

From the experimental tests, an initially formed set A may contain only few redundant vertices to be removed to obtain the preliminary k -limited packing X' . As a result, the preliminary k -limited packing X' in many cases satisfies lower bound (1), and the extension of X' to a maximal k -limited packing X seems to always satisfy (1). In our view, since the problem is NP -hard, Algorithm 1 constitutes a simple efficient approach to tackle the problem in practice and, hopefully, can be useful to solve some hard instances of the problem.

Algorithm 1: Randomized k -limited packing

Input: Graph G and integer k , $1 \leq k \leq \Delta$.

Output: k -Limited packing X in G .

begin

```

    Compute  $p = \left( \frac{1}{\tilde{c}_{k+1} (1+k)} \right)^{1/k}$  ;
    Initialize  $A = \emptyset$ ;                                /* Form a set  $A \subseteq V(G)$  */
    foreach vertex  $v \in V(G)$  do
    | with the probability  $p$ , decide whether  $v \in A$  or  $v \notin A$ ;
    end
    /* Recursively remove redundant vertices from  $A$  */
    foreach vertex  $v \in V(G)$  do
    | Compute  $r = |N(v) \cap A|$ ;
    | if  $v \in A$  and  $r \geq k$  then
    | | remove any  $r - k + 1$  vertices of  $N(v) \cap A$  from  $A$ ;
    | end
    | if  $v \notin A$  and  $r > k$  then
    | | remove any  $r - k$  vertices of  $N(v) \cap A$  from  $A$ ;
    | end
    end
    Put  $X' = A$ ;                                        /*  $X'$  is a  $k$ -limited packing */
    Extend  $X'$  to a maximal  $k$ -limited packing  $X$ ;
    return  $X$ ;

```

end

Algorithm 1 can be implemented to run in $O(n^2)$ time. To compute the probability $p = \left(\binom{\Delta}{k} (\Delta + 1) \right)^{-1/k}$, the binomial coefficient $\binom{\Delta}{k}$ can be computed

by using the dynamic programming and Pascal's triangle in $O(k\Delta) = O(\Delta^2)$ time using $O(k) = O(\Delta)$ memory. The maximum vertex degree Δ of G can be computed in $O(m)$ time, where m is the number of edges in G . Then p can be computed in $O(m + \Delta^2) = O(n^2)$ steps. It takes $O(n)$ time to find the initial set A . Computing the intersection numbers $r = |N(v) \cap A|$ and removing unwanted vertices of $N(v) \cap A$'s from A can be done in $O(n + m)$ steps. Finally, checking whether X' is maximal or extending X' to a maximal k -limited packing X can be done in $O(n + m)$ time: try to add vertices of $V(G) - X'$ to X' recursively one by one, and check whether the addition of a new vertex $v \in V(G) - X'$ to X' violates the conditions of a k -limited packing for v or at least one of its neighbours in G with respect to $X' \cup \{v\}$. Thus, overall Algorithm 1 takes $O(n^2)$ time, and, since $m = O(n^2)$ in general, it is linear in the graph size $(m + n)$ when $m = \theta(n^2)$.

Also, this randomized algorithm for finding k -limited packings in a graph G can be implemented in parallel or as a local distributed algorithm. As explained in [5], this kind of algorithms are especially important, e.g. in the context of ad hoc and wireless sensor networks. We hope that this approach can be also extended to design self-stabilizing or on-line algorithms for k -limited packings. For example, a self-stabilizing algorithm searching for maximal 2-packings in a distributed network system is presented in [12]. Notice that self-stabilizing algorithms are distributed and fault-tolerant, and use the fact that each node has only a local view/knowledge of the distributed network system. This provides another motivation for efficient distributed search and algorithms to find k -limited packings in graphs and networks.

4. Sharpness of the lower bounds

We now show that the lower bound of Theorem 1 is asymptotically best possible for some values of k . The bound of Theorem 1 can be rewritten in the following form for $\Delta \geq k$:

$$L_k(G) \geq \frac{kn}{(k+1)^k \sqrt[k]{\binom{\Delta}{k}} (\Delta+1)}.$$

Combining this bound with the upper bound of Lemma 8 from [6], we obtain that for any connected graph G of order n with minimum degree $\delta(G) \geq k$,

$$\frac{1}{\sqrt[k]{\binom{\Delta}{k}} (\Delta+1)} \times \frac{k}{k+1}n \leq L_k(G) \leq \frac{k}{k+1}n. \quad (7)$$

Notice that the upper bound in the inequality (7) is sharp (see [6]), so these bounds provide an interval of values for $L_k(G)$ in terms of k and Δ when $k \leq \delta$. For regular graphs, $\delta = \Delta$, and, when $k = \Delta$, we have

$$\frac{1}{\sqrt[k]{\binom{\Delta}{k}} (\Delta+1)} = \frac{1}{(k+1)^{1/k}} \rightarrow 1 \quad \text{as } k \rightarrow \infty.$$

Therefore, the bound of Theorem 1 is asymptotically sharp for regular connected graphs in the case $k = \Delta$. In other words, there are graphs whose k -limited packing number is arbitrarily close to the bound of Theorem 1. Thus, the following result holds:

Theorem 6. *When n is large, there exist graphs G such that*

$$L_k(G) \leq \frac{kn}{\tilde{c}_{k+1}^{1/k} (1+k)^{1+1/k}} (1 + o(1)). \quad (8)$$

As shown above, the graphs satisfying Theorem 6 contain regular connected ones for $k = \Delta$. This class of graphs can be extended, because it is possible to prove that the bound of Theorem 1 is asymptotically sharp for connected graphs with $k = \Delta(1 - o(1))$, $\delta(G) \geq k$.

Notice that, for regular graphs, the condition $k = \Delta$ and Lemma 5 from [6] imply $L_k(G) = n - \gamma(G)$. Then the classical upper bound (9) for $\gamma(G)$ gives a weaker lower bound for $L_k(G)$ than Theorem 1.

As shown in Theorem 5, in contrast to the situation for relatively ‘large’ values of k , bound (1) of Theorem 1 (see Corollary 4) can be improved for distance 2-packings (1-limited packings), i.e. when $k = 1$. However, this improvement is irrelevant for almost all graphs. A 1-limited packing set X in G has a very strong property that any two vertices in X are at distance at least 3 in G . It is well known that almost every graph has diameter equal to 2 (e.g., see [10]). Therefore, $\rho(G) = L_1(G) = 1$ for almost all graphs. Thus, in the case $k = 1$, Theorem 1 yields a lower bound of 1 for almost all graphs and is as good as Theorem 5. Notice that the bound of Theorem 5 is sharp, for example, for any number of disjoint copies of the Petersen graph. In the other cases, when G has a diameter larger than 2, one is encouraged to use the greedy algorithm and lower bound (6) provided by Theorem 5, because it improves bound (5) of Corollary 4 by a factor of $2 + o(1)$.

5. Upper bounds

As mentioned earlier, $\rho(G) = L_1(G) \leq \gamma(G)$. In [6], the authors provide several upper bounds for $L_k(G)$, e.g. $L_k(G) \leq k\gamma(G)$ for any graph G . Using the well-known bound (see e.g. [1])

$$\gamma(G) \leq \frac{\ln(\delta + 1) + 1}{\delta + 1} n, \quad (9)$$

we obtain

$$L_k(G) \leq \frac{\ln(\delta + 1) + 1}{\delta + 1} kn. \quad (10)$$

Even though this bound does not work well when k is ‘close’ to δ , it is very reasonable for small values of k .

We now prove an upper bound for the k -limited packing number in terms of the k -tuple domination number. A set X is called a k -tuple dominating set of G if for every vertex $v \in V(G)$, $|N[v] \cap X| \geq k$. The minimum cardinality of a k -tuple dominating set of G is the k -tuple domination number $\gamma_{\times k}(G)$. The k -tuple domination number is only defined for graphs with $\delta \geq k - 1$.

Theorem 7. For any graph G of order n with $\delta \geq k - 1$,

$$L_k(G) \leq \gamma_{\times k}(G). \quad (11)$$

PROOF. We prove inequality (11) by contradiction. Let X be a maximum k -limited packing in G of size $L_k(G)$, and let Y be a minimum k -tuple dominating set in G of size $\gamma_{\times k}(G)$. We denote $B = X \cap Y$, i.e. $X = A \cup B$ and $Y = B \cup C$, where A and C are disjoint. Assume to the contrary that $L_k(G) > \gamma_{\times k}(G)$, thus $|A| > |C|$.

Since Y is k -tuple dominating set, each vertex of A is adjacent to at least k vertices of Y . Hence the number of edges between A and $B \cup C$ is as follows:

$$e(A, B \cup C) \geq k|A|.$$

Now, every vertex of C is adjacent to at most k vertices of X , because X is a k -limited packing set. Therefore, the number of edges between C and $A \cup B$ satisfies

$$e(C, A \cup B) \leq k|C|.$$

We obtain

$$e(C, A \cup B) \leq k|C| < k|A| \leq e(A, B \cup C),$$

i.e. $e(C, A \cup B) < e(A, B \cup C)$. By eliminating the edges between A and C , we conclude that

$$e(C, B) < e(A, B).$$

Now, let us consider an arbitrary vertex $b \in B$ and denote $s = |N(b) \cap A|$. Since $X = A \cup B$ is a k -limited packing set, we obtain $|N(b) \cap X| \leq k - 1$, and hence $|N(b) \cap B| \leq k - s - 1$. On the other hand, $Y = B \cup C$ is k -tuple dominating set, so $|N(b) \cap Y| \geq k - 1$. Therefore, $|N(b) \cap C| \geq s$. Thus, $|N(b) \cap C| \geq |N(b) \cap A|$ for any vertex $b \in B$. We obtain,

$$e(C, B) \geq e(A, B),$$

a contradiction. We conclude that $L_k(G) \leq \gamma_{\times k}(G)$. \square

Notice that it is possible to have $k = \Delta + 1$ in the statement of Theorem 7, which is not covered by Theorem 1. Then $\delta = \Delta$, which implies the graph is regular. However, $L_k(G) = \gamma_{\times k}(G) = n$ for $k = \delta + 1 = \Delta + 1$. In non-regular graphs, $\delta + 1 \leq \Delta$, and $k \leq \Delta$ to satisfy the conditions of Theorem 1 as well.

For $t \leq \delta$, we define

$$\delta' = \delta - k + 1 \quad \text{and} \quad \tilde{b}_t = \tilde{b}_t(G) = \binom{\delta + 1}{t}.$$

Using the upper bound for the k -tuple domination number from [5], we obtain:

Corollary 8. For any graph G with $\delta \geq k$,

$$L_k(G) \leq \left(1 - \frac{\delta'}{\tilde{b}_{k-1}^{1/\delta'} (1 + \delta')^{1+1/\delta'}} \right) n. \quad (12)$$

In some cases, Theorem 1 and Corollary 8 simultaneously provide good bounds for the k -limited packing number. For example, for a 40-regular graph G :

$$0.312n < L_{25}(G) < 0.843n.$$

Acknowledgement

The authors are grateful to the anonymous referee for helpful comments and suggestions.

References

- [1] N. Alon, J.H. Spencer, *The Probabilistic Method*, 2nd ed., John Wiley & Sons Inc., New York, 2000.
- [2] B. Bollobás, *Graph Theory: An Introductory Course*, Springer-Verlag, New York, 1979.
- [3] A.P. Burger, M.A. Henning, J.H. van Vuuren, On the ratios between packing and domination parameters of a graph, *Discrete Math.* 309 (2009) 2473–2478.
- [4] M.P. Dobson, V. Leoni, G. Nasini, The multiple domination and limited packing problems in graphs, *Inform. Process. Lett.* 111 (2011) 1108–1113.
- [5] A. Gagarin, A. Poghosyan, V.E. Zverovich, Randomized algorithms and upper bounds for multiple domination in graphs and networks, *Discrete Appl. Math.* 161 (2013) 604–611.
- [6] R. Gallant, G. Gunther, B. Hartnell, D. Rall, Limited packings in graphs, *Discrete Appl. Math.* 158(12) (2010) 1357–1364.
- [7] M.A. Henning, C. Löwenstein, D. Rautenbach, Dominating sets, packings, and the maximum degree, *Discrete Math.* 311 (2011) 2031–2036.
- [8] D.S. Hochbaum, D.B. Shmoys, A best possible heuristic for the k -center problem, *Math. Oper. Res.* 10(2) (1985) 180–184.
- [9] A. Meir, J.W. Moon, Relations between packing and covering numbers of a tree, *Pacific J. Math.* 61 (1975) 225–233.
- [10] J.W. Moon, L. Moser, Almost all $(0,1)$ matrices are primitive, *Studia Sci. Math. Hungar.* 1 (1966) 153–156.
- [11] R.R. Rubalcaba, A. Schneider, P.J. Slater, A survey on graphs which have equal domination and closed neighborhood packing numbers, *AKCE J. Graphs. Combin.* 3(2) (2006) 93–114.
- [12] Z. Shi, A self-stabilizing algorithm to maximal 2-packing with improved complexity, *Inform. Process. Lett.* 112 (2012) 525–531.
- [13] J. Topp, L. Volkmann, On packing and covering numbers of graphs, *Discrete Math.* 96 (1991) 229–238.