

# BMJ Open

## Validity and effectiveness of paediatric early warning systems and track and trigger tools for identifying and reducing clinical deterioration in hospitalised children: a systematic review

Rob Trubey,<sup>1</sup> Chao Huang,<sup>1</sup> Fiona V Lugg-Widger,<sup>1</sup> Kerensa Hood,<sup>1</sup> Davina Allen,<sup>2</sup> Dawn Edwards,<sup>3</sup> David Lacy,<sup>4</sup> Amy Lloyd,<sup>1</sup> Mala Mann,<sup>5</sup> Brendan Mason,<sup>6</sup> Alison Oliver,<sup>7</sup> Damian Roland,<sup>8</sup> G Sefton,<sup>9</sup> Richard Skone,<sup>7</sup> Emma Thomas-Jones,<sup>1</sup> Lyvonne N Tume,<sup>10</sup> Colin Powell<sup>11,12</sup>

**To cite:** Trubey R, Huang C, Lugg-Widger FV, *et al.* Validity and effectiveness of paediatric early warning systems and track and trigger tools for identifying and reducing clinical deterioration in hospitalised children: a systematic review. *BMJ Open* 2019;0:e022105. doi:10.1136/bmjopen-2018-022105

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-022105>).

Received 6 February 2018  
Revised 7 March 2019  
Accepted 8 March 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

### Correspondence to

Dr Rob Trubey;  
trubeyrj@cf.ac.uk

### ABSTRACT

**Objective** To assess (1) how well validated existing paediatric track and trigger tools (PTTT) are for predicting adverse outcomes in hospitalised children, and (2) how effective broader paediatric early warning systems are at reducing adverse outcomes in hospitalised children.

**Design** Systematic review.

**Data sources** British Nursing Index, Cumulative Index of Nursing and Allied Health Literature, Cochrane Central Register of Controlled Trials, Database of Abstracts of Reviews of Effectiveness, EMBASE, Health Management Information Centre, Medline, Medline in Process, Scopus and Web of Knowledge searched through May 2018.

**Eligibility criteria** We included (1) papers reporting on the development or validation of a PTTT or (2) the implementation of a broader early warning system in paediatric units (age 0–18 years), where adverse outcome metrics were reported. Several study designs were considered.

**Data extraction and synthesis** Data extraction was conducted by two independent reviewers using template forms. Studies were quality assessed using a modified Downs and Black rating scale.

**Results** 36 validation studies and 30 effectiveness studies were included, with 27 unique PTTT identified. Validation studies were largely retrospective case-control studies or chart reviews, while effectiveness studies were predominantly uncontrolled before-after studies. Metrics of adverse outcomes varied considerably. Some PTTT demonstrated good diagnostic accuracy in retrospective case-control studies (primarily for predicting paediatric intensive care unit transfers), but positive predictive value was consistently low, suggesting potential for alarm fatigue. A small number of effectiveness studies reported significant decreases in mortality, arrests or code calls, but were limited by methodological concerns. Overall, there was limited evidence of paediatric early warning system interventions leading to reductions in deterioration.

**Conclusion** There are several fundamental methodological limitations in the PTTT literature, and the predominance of single-site studies carried out in specialist centres greatly limits

### Strengths and limitations of this study

- Paediatric early warning systems and paediatric track and trigger tools (PTTT) are increasingly used by paediatric units across Europe, North America, Australia and elsewhere—this study is a timely review of the evidence for their validity and effectiveness.
- A comprehensive search was carried out across multiple databases and included published as well as grey literature.
- The review highlights methodological weaknesses and gaps in the current evidence base and makes suggestions for future research.
- Heterogeneity in study populations, study designs and outcome measures make it difficult to compare and synthesise findings across the wide range of early warning systems and PTTT being used in practice.
- The review is limited in scope to quantitative validation and effectiveness studies, so must be considered alongside wider literature reflecting on potential secondary benefits of early warning systems and PTTT for communication, teamwork and empowerment.

generalisability. With limited evidence of effectiveness, calls to make PTTT mandatory across all paediatric units are not supported by the evidence base.

**PROSPERO registration number** CRD42015015326

### BACKGROUND

Failure to recognise and respond to clinical deterioration in hospitalised children is a major safety concern in healthcare. The underlying causes of this problem are

clearly multifactorial,<sup>1-3</sup> but paediatric 'early warning systems' have been strongly advocated as one approach to improving recognition of deterioration in paediatric units.<sup>1,2,4</sup>

A paediatric 'early warning system' can be considered any patient safety initiative or programme which aims to monitor, detect and respond to signs of deterioration in hospitalised children in order to avert adverse outcomes and premature death. Such systems are often multifaceted and may include the use of rapid response teams (RRT) or medical emergency teams (MET), education or training to improve clinical staff's ability to identify deterioration or strategies aimed at improving staff communication and situational awareness.

An increasingly commonplace paediatric 'early warning system' initiative is the use of a 'track and trigger tool': these tools, also commonly used in adult care, provide a formal framework for evaluating routine physiological, clinical and observational data for early indicators of patient deterioration. They are typically integrated into routine observation charts or electronic health records and compare patient observations with predefined 'normal' thresholds. When one or more observation is considered abnormal, staff are directed to various clinical actions, including but not limited to altered frequency of observations, review by senior staff or more appropriate treatment or management. Tools may be paper based or electronic and monitoring may be automated or manually undertaken by staff.

These tools have been referred to in the literature using a number of different terms: paediatric early warning scores (PEWS); paediatric early warning tools (PEWT), track and trigger tools (TTT) and many others. Here, we refer to the tools themselves using the term 'paediatric track and trigger tools' (PTTT). A variety of PTTT have been developed, typically by teams based in specialist paediatric centres and often used as a means of triggering a dedicated response team. Their advocacy has recently led to widespread uptake across a variety of different paediatric units, including many non-specialist centres where patient populations and resources may differ. In the UK, a recent cross-sectional survey found that 85% of paediatric units were using some form of PTTT, most of which were non-specialist centres without a dedicated response team.<sup>5</sup> Despite their widespread use, recent reviews have questioned the evidence base for their effectiveness in improving patient outcomes.<sup>6,7</sup> The current review aimed to build on this work, assessing in depth the evidence base for both the validity of PTTT for predicting in-patient deterioration and the effectiveness of broader 'early warning systems' at reducing instances of mortality and morbidity in paediatric settings:

- ▶ Question 1: how well validated are existing PTTT and their component parts for predicting inpatient deterioration?
- ▶ Question 2: how effective are paediatric early warning systems (with or without a PTTT) at reducing mortality and critical events?

## METHODS

This systematic review is reported in accordance with the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) guidelines.<sup>8</sup> Our review protocol is registered with the PROSPERO database CRD42015015326.

### Search strategy

A comprehensive search was conducted across a range of databases to identify relevant studies in the English language. Published and unpublished literature was considered where publicly available, as were studies in press. The following databases were searched through May 2018: British Nursing Index, Cumulative Index of Nursing and Allied Health Literature, Cochrane Central Register of Controlled Trials, Database of Abstracts of Reviews of Effectiveness, EMBASE, Health Management Information Centre, Medline, Medline in Process, Scopus and Web of Knowledge (Science Citation Indexes). To identify additional papers, published, unpublished or research reported in the grey literature, a range of relevant websites and trial registers were searched including ClinicalTrials.gov. To identify published papers that had not yet been catalogued in the electronic databases, recent editions of key journals were hand-searched. The search terms included 'early warning scores', 'alert criteria', 'rapid response', 'track and trigger' and 'early medical intervention' (see online supplementary table 1).

### Eligibility screening and study selection

PICOS parameters guided inclusion criteria for the validation and effectiveness studies (see online supplementary table 2). Papers reporting development of validation of a PTTT were included for question 1, whereas papers reporting the implementation of any broader 'paediatric early warning system' (with or without a PTTT) were eligible for question 2. Both research questions were limited to studies that involved inpatients aged 0–18 years. Outcome measures considered were mortality and critical events, including: unplanned admission to a higher level of care, cardiac arrest, respiratory arrest, medical emergencies requiring immediate assistance, children reviewed by paediatric intensive care unit (PICU) staff on the ward (in specialist centres) or reviewed by external PICU staff (for non-specialist centres), acuity at PICU admission and PICU outcomes. A range of study designs were considered for both questions.

Two of the review authors independently screened the titles and abstracts yielded in the search. Full texts were reviewed independently by six reviewers against the above eligibility criteria and were assigned to the relevant review question if included. Reasons for exclusion were recorded. Separate data extraction forms were developed for validation and effectiveness studies. The forms had common elements (study design, country, setting, study population, description of the PTTT or early warning system, statistical techniques used, outcomes assessed). Additional data items for validation studies included

the items in the PTTT, modifications to the PTTT from previous versions, predictive ability of individual items and the overall tool, sensitivity and specificity and inter-rater and intra-rater reliability. Effectiveness studies included an assessment of outcomes in terms of mortality and various morbidity variables. Data extraction was carried out by two reviewers and discrepancies were resolved by discussion. For effectiveness studies, effect sizes and 95% CIs were calculated or reported as risk ratios (RR) or ORs as appropriate, with p values reported to assess statistical significance. Data analysis was conducted using an online medical statistics tool.

### Quality appraisal

Methodological quality and risk of bias was assessed for each included study using a modified version of the Downs and Black rating scale<sup>9</sup> (templates shown in online supplementary table 3).

### Patient and public involvement

This review was conducted as part of a larger mixed-methods study (ISRCTN94228292), which used a formal, facilitated parental advisory group. The group comprised parents of children who had experienced an unexpected adverse event in a paediatric unit and provided input which helped to shape the broader research questions and outcome measures. The results of the review will be disseminated to parents through this group.

## RESULTS

Figure 1 shows the PRISMA flow diagram for both research questions.

### Study characteristics

Table 1 summarises the study characteristics of validation and effectiveness papers in the review.

### Types of PTTS and components

Across 66 studies, we identified 27 unique PTTT (table 2). Twenty PTTTs were based on one of four different tools: Monaghan's Brighton PEWS,<sup>10</sup> the Bedside PEWS,<sup>11</sup> the Bristol PEWT<sup>12</sup> and the Melbourne Activation Criteria (MAC).<sup>13</sup> Other PTTT described in the literature included the National Health Service Institute for Innovation and Improvement (NHS III) PEWS<sup>14</sup> (the second most commonly used PTTT in UK paediatric settings<sup>5</sup>), RRT and MET activation criteria<sup>15–18</sup> and one prediction algorithm developed from a large dataset of electronic health data.<sup>19</sup>

Table 2 illustrates the range of physiological and behavioural parameters underpinning PTTT. Common parameters included heart rate (present in 26 out of 27 PTTT), respiratory rate (24), respiratory effort (24) and level of consciousness or behavioural state (24). All PTTT required at least six different parameters to be collected.

### Question 1: how well validated are PTTT and component parts for predicting inpatient deterioration?

Nine validation papers meeting inclusion criteria were excluded from analysis: eight did not report any

performance characteristics of the PTTT for predicting deterioration<sup>20–27</sup> and one study calculated incorrect sensitivity/specificity outcomes<sup>12</sup> (see online supplementary table 4). The remaining 27 validation studies, evaluating the performance of 18 unique PTTT, are described in table 3. Four studies evaluated multiple PTTTs<sup>3 19 28 29</sup> and one paper described three separate studies of the same PTTT.<sup>30</sup>

Five cohort studies were included,<sup>14 31–34</sup> three based on the same dataset. All other studies were either case-control or chart reviews. Thirteen papers implemented the PTTT in practice,<sup>23 30 31 34–43</sup> while the remaining studies 'bench tested' the PTTT—researchers retrospectively calculated the score based on data abstracted from medical charts and records. All studies were conducted in specialist centres with only one multicentre study reported.<sup>44</sup>

### Outcome measures

PTTT were evaluated for their ability to predict a wide range of clinical outcomes. Composite measures were used in 8 studies,<sup>14 23 29 32 33 37 45 46</sup> cardiac/respiratory arrest or a 'code call' was used (singularly or part of a composite outcome) in 6 studies,<sup>23 28 29 37 45 47</sup> while 22 studies used transfer to a PICU or paediatric high-dependency unit as the main outcome.<sup>3 11 19 23 28–34 36 37 39 41–44 46 48 49</sup>

### Predictive ability of individual PTTT components


Three validation papers reported on the performance characteristics of individual components of the tool for predicting adverse outcomes.<sup>11 33 42</sup> Parshuram *et al*, for instance, reported area under the receiver operating characteristic curve (AUROC) values for individual PTTT items of a pilot version of the Bedside PEWS: ranging from 0.54 (bolus fluid) to 0.81 (heart rate), compared with 0.91 for the overall PTTT.<sup>11</sup> All other studies reported outcomes for the PTTT as a whole.

### PEWS score

The predictive ability of the 16-item PEWS score was assessed by one internal<sup>47</sup> (AUROC=0.90) and two external case-control studies<sup>28 29</sup> (AUROC range=0.82–0.88) with a range of outcome measures and scoring thresholds. One case-control study used an observed prevalence rate to calculate a positive predictive value (PPV) of 4.2% for the tool in predicting code calls<sup>47</sup> (for every 1000 patients triggering the PTTT, 42 would be expected to deteriorate).

### Bedside PEWS and derivatives

The Bedside PEWS was evaluated in one internal<sup>11</sup> (AUROC=0.91) and five external case-control studies<sup>19 28 29 44 46</sup> (AUROC range=0.73–0.90) for a range of different outcome measures and at different scoring thresholds. One case-control study calculated a PPV of 2.1% for identifying children requiring urgent PICU transfer within 24 hours of admission, based on locally observed prevalence rates.<sup>19</sup> A modified version of the Bedside PEWS (with temperature added) demonstrated an AUROC of 0.86 in an external case-control study with



**Figure 1** Preferred Reporting Items for Systematic Review and Meta-Analyses flow diagram of study inclusion. PEWS, paediatric early warning scores.

a composite outcome of death, arrest or unplanned PICU transfer.<sup>29</sup>

### Brighton PEWS and derivatives

Six different PTTT based on the original Brighton PEWS were evaluated across 11 studies.<sup>19 29 31 37 39-42 45 48 50</sup> The Modified Brighton PEWS (a) was evaluated for its ability to predict PICU transfers in one large prospective cohort study (AUROC=0.92, PPV=5.8%),<sup>31</sup> and an external case-control study tested the same score for predicting

urgent PICU transfers within 24 hours of admission (AUROC=0.74, PPV=2.1%).<sup>19</sup>

An external case-control study used a composite measure of death, arrest or PICU transfer to evaluate the Modified Brighton PEWS (b) (AUROC=0.79) and the Modified Brighton PEWS (d) (AUROC=0.74).<sup>29</sup> The latter tool was evaluated in a further internal case-control study for predicting PICU transfer (AUROC=0.82).<sup>48</sup>



**Table 1** Summary of the study characteristics of the 36 validation (question 1) and 30 effectiveness (question 2) papers included in the review

Validation studies (n=36)	N (%)	Effectiveness studies (n=30)	n (%)
Type		Type	
Full text	22 (61.1)	Full text	21 (70.0)
Abstract	14 (38.9)	Abstract	9 (30.0)
Country		Country	
USA	15 (41.7)	USA	18 (60.0)
UK	12 (33.3)	UK	3 (10.0)
Canada	2 (5.5)	Canada	2 (6.7)
Australia	0 (0.0)	Australia	3 (10.0)
Other	5 (13.9)	Other	3 (10.0)
Multiple	1 (2.8)	Multiple	1 (3.3)
Unclear	1 (2.8)	Unclear	0 (0.0)
Year of study		Year of study	
Pre-2012	10 (27.8)	Pre-2012	15 (50.0)
2012	3 (8.3)	2012	1 (3.3)
2013	6 (16.7)	2013	2 (6.7)
2014	5 (13.9)	2014	6 (20.0)
2015	7 (19.4)	2015	0 (0.0)
2016	2 (5.6)	2016	2 (6.7)
2017	3 (8.3)	2017	1 (3.3)
2018	0 (0.0)	2018	3 (10.0)
Setting		Setting	
Specialist/tertiary	33 (91.7)	Specialist/tertiary	29 (96.7)
Non-specialist/community	0 (0.0)	Non-specialist/community	1 (3.3)
Unclear	3 (8.3)	Unclear	0 (0.0)
Single-centre/multicentre		Single-centre/multicentre	
Single-centre	35 (97.2)	Single-centre	28 (93.3)
Multicentre	1 (2.8)	Multicentre	2 (6.7)
Study population		Study population	
General inpatients	23 (63.9)	General inpatients	20 (66.6)
Specialist population	11 (30.6)	Specialist population	5 (16.7)
Unclear	2 (5.6)	Unclear	5 (16.7)
Study design		Study design	
Case-control	18 (50.0)	Uncontrolled before-after	26 (86.7)
Case/chart review	10 (27.8)	Controlled before-after	1 (3.3)
Cohort	7 (19.4)	Interrupted time series	2 (6.7)
Pilot study	1 (2.8)	Cluster randomised trial	1 (3.3)

The Children's Hospital Early Warning Score (CHEWS) had a reported AUROC of 0.90 for predicting PICU transfers or arrests in a large internal case-control study.<sup>50</sup> A modification for cardiac patients, the Cardiac CHEWS (C-CHEWS) was evaluated by one internal study on a cardiac unit<sup>37</sup> (AUROC=0.90) looking at arrests or unplanned PICU transfers, and two external studies of oncology/haematology units<sup>41 42</sup> for the same outcome (AUROC=0.95). Finally, the Children's Hospital

Los Angeles PEWS was evaluated by in a small internal case-control study for prediction of re-admission to PICU after initial PICU discharge<sup>40</sup> (AUROC=0.71).

#### MAC and derivatives

The MAC was assessed by one external case-control study with an outcome of death, arrest or unplanned PICU transfer<sup>29</sup> (AUROC=0.71) and a large external cohort study with an outcome of death or unplanned

Table 2 Summary of PTTTs

PTT name (references)	Development/modification details	Score/trigger	Choice of thresholds/parameters	Age-dependent thresholds	No. of items in the tool*	PTTT parameters															
						Respiratory rate	Heart rate	Respiratory effort/distress	LOC/behaviour	Oxygen saturation	Capillary refill time	Oxygen therapy	Systolic blood pressure	Pain	Staff concern	Skin colour	Airway problems	Temperature	Pulses	Family concern	Other items
<b>Paediatric early warning system (PEWS) score and derivatives</b>																					
PEWS score <sup>38-47</sup>	Developed for use in Canadian tertiary centre. <sup>47</sup> Nurse-generated candidate items reduced by focus groups/Delphi and evaluation with clinical dataset (code blue calls, n=587; controls, n=528). Development and validation datasets not independent.	Score	Expert opinion	Yes	16	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Bolus fluid, medications, home oxygen, any previous admission to an ICU, central venous line in situ, transplant recipient, severe cerebral palsy, gastrostomy tube, greater than three medical specialities involved in care.
Bedside PEWS <sup>11-19, 23, 27, 28, 44, 46, 59-64, 65, 70</sup>	Developed for use in US tertiary centres. <sup>11</sup> Routinely collected items assessed for discriminatory ability, using clinical dataset (PICU admission, n=60; controls, n=120). Development and validation set not independent.	Score	Expert opinion	Yes	7	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
Modified Bedside PEWS (a) <sup>30</sup>	Modification to Bedside PEWS for use in Dutch tertiary centre. Added temperature; modified wording of respiratory effort and oxygen therapy items.	Score	Expert opinion	Yes	8	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
Modified Bedside PEWS (b) <sup>49</sup>	Modification to Bedside PEWS for use in US tertiary centre. Changed normal thresholds for HR and RR based on analysis of local clinical data.	Score	HR/RR data driven	Yes	7	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
<b>Brighton PEWS and derivatives</b>																					
Brighton PEWS <sup>53, 64</sup>	Initial development for use in UK tertiary centre. Adapted from existing adult scores, but amended based on local clinical consensus. Small audit of patients (n=30) described but no formal validation.	Score	Expert opinion	No	5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	¼ hourly nebulisers, persistent vomiting postsurgery.
Modified Brighton PEWS (a) <sup>19, 21, 39</sup>	Modification of Brighton PEWS for use in general medical ward of a US tertiary centre. Altered thresholds for oxygen therapy; changed wording for respiratory effort; modified escalation algorithm.	Score	Expert opinion	No	5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	¼ hourly nebulisers, persistent vomiting postsurgery.
Modified Brighton PEWS (b) <sup>45, 72</sup>	Modification of Brighton PEWS for use in US tertiary centre. Added age-dependent thresholds for HR and RR.	Score	Expert opinion	Yes	5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	¼ hourly nebulisers, persistent vomiting postsurgery.

Continued

**Table 2** Continued

PTT name (references)	Development/ modification details	Score/ trigger	Choice of thresholds/ parameters	Age- dependent thresholds	No. of items in the tool*	PTT parameters																
						Respiratory rate	Heart rate	Respiratory effort/ distress	LOC/ behaviour	Oxygen saturation	Capillary refill time	Oxygen therapy	Systolic blood pressure	Pain	Staff concern	Skin colour	Airway problems	Temperature	Pulses	Family concern	Other items	
Modified Brighton PEWS (c) <sup>25</sup>	Modification of Brighton PEWS for use in a US haematology/oncology unit. Altered thresholds; changed respiratory effort wording; modified escalation algorithm; added and removed items. No formal validation study reported.	Score	Expert opinion	No	3	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		
Modified Brighton PEWS (d) <sup>26</sup>	Modification of Brighton PEWS for use in a US tertiary centre. Modified wording of behaviour component; added age-dependent thresholds for HR and RR; removed nebulisers and persistent vomiting.	Score	Expert opinion	Yes	3	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		
Modified Brighton PEWS (e) <sup>27</sup>	Modification of Brighton PEWS for use in a US tertiary centre. Modified wording of behaviour and respiratory effort items; altered thresholds for O <sub>2</sub> therapy; removed nebulisers and persistent vomiting items. No formal validation study reported.	Score	Expert opinion	No	3	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		
Texas Children's Hospital PEWS <sup>28</sup>	Modification of Brighton PEWS for use in a US tertiary centre. Modified wording of behaviour category; added scoring items to respiratory and cardiovascular categories; changed O <sub>2</sub> therapy thresholds; modified escalation algorithm.	Score	Expert opinion	No	5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Hourly respiratory treatments; persistent vomiting postsurgery.	
Children's Hospital Early Warning Score <sup>30</sup>	Modification of Brighton PEWS for use in a US tertiary centre. Altered thresholds for O <sub>2</sub> therapy; changed wording for behaviour and respiratory categories; added staff and family concern; removed nebulisers and vomiting; modified escalation algorithm.	Score	Expert opinion	No	5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
Children's Hospital Cardiac Early Warning Score <sup>31,42,67</sup>	Modification of Brighton PEWS for cardiac ward of a US tertiary centre. Altered O <sub>2</sub> therapy thresholds; added items to behaviour, respiratory and cardiovascular categories; added family and staff concern; added age-related thresholds; removed nebulisers and vomiting items; modified escalation algorithm.	Score	Expert opinion	Yes	5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

Continued

AUTHOR PROOF

AUTHOR PROOF

Table 2 Continued

PTTT name (references)	Development/ modification details	Score/ trigger	Choice of thresholds/ parameters		Age-dependent thresholds	No. of items in the tool*	PTTT parameters										Other items		
			Expert opinion	No			Respiratory rate	Heart rate	Respiratory effort/ distress	LOC/behaviour	Oxygen saturation	Capillary refill time	Oxygen therapy	Systolic blood pressure	Pain	Staff concern		Skin colour	Airway problems
Burn-specific PEWS <sup>34</sup>	Modification of Brighton PEWS, for use in a specialist Burn Centre of a US tertiary centre. Added temperature; added intake and output scoring items; added skin component.	Score	Expert opinion	No	6	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Intake; outputs; skin.
Children's Hospital Los Angeles PEWS <sup>37</sup>	Modification of Brighton PEWS for use in a US tertiary centre. Added medical history scoring item; added single ventricle physiology scoring item; changed O <sub>2</sub> therapy thresholds; added items to respiratory category.	Score	Expert opinion		4	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	RRT, code blue or transfer from/ to PICU in past 2 weeks; single ventricle physiology; any assisted ventilation.
<b>Melbourne Activation Criteria (MAC) and derivatives</b>																			
MAC <sup>13,33,32</sup>	Initial development for use in an Australian tertiary centre to activate MET. Adapted from adult MET calling criteria, using age-appropriate thresholds. No formal validation study reported.	Trigger	Expert opinion	Yes	9	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Cardiac or respiratory arrest.
Modified MAC <sup>33</sup>	Modification of MAC for use in a Canadian tertiary centre, to activate an RRS. Removed cardiac/respiratory arrest outcome. No formal validation study reported.	Trigger	Expert opinion	Yes	8	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
Cardiff and Vale PEWS <sup>33</sup>	Modification of MAC for evaluation in a UK tertiary centre. Removed cardiac/respiratory arrest outcome; altered thresholds of some items; evaluated as aggregate score rather than single-item trigger.	Score	Expert opinion	Yes	8	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
<b>Bristol paediatric early warning tool (PEWT) and derivatives</b>																			
Bristol PEWT <sup>1,2,28,34,35</sup>	Initial development for use in a UK tertiary centre. Initial candidate items drawn from unvalidated Plymouth tool—retrospectively evaluated for ability to predict adverse events among cases (n=350, HDU or PICU transfers). Development and validation dataset not independent.	Trigger	APLS values	Yes	14	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Required nebulised epinephrine; hyperkalaemia; suspected meningococcus; diabetic ketoacidosis; persistent convulsion.
Modified Bristol PEWT (a) <sup>38</sup>	Modification of Bristol PEWT for a UK tertiary centre. Adjusted wording of Airway parameters; added respiratory items; added APU evaluation; removed suspected meningococcus and diabetic ketoacidosis; added pH <7.2 and unresolved pain. No formal validation study reported.	Trigger	APLS values	Yes	14	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Required nebulised epinephrine or no improvement after nebulisers; pH <7.2; unresolved pain or current analgesic therapy; fitting.

Continued



**Table 2** Continued

PTTT name (references)	Development/ modification details	Score/ trigger	Choice of thresholds/ dependent parameters	Age- dependent thresholds	No. of items in the tool*	PTTT parameters													
						Respiratory rate	Heart rate	Respiratory effort/ distress	LOC/ behaviour	Oxygen saturation	Capillary refill time	Oxygen therapy	Systolic blood pressure	Pain	Staff concern	Skin colour	Airway problems	Temperature	Pulses
Modified Bristol PEWT (b) <sup>3</sup>	Modification of Bristol PEWT for cardiac ward of a UK tertiary centre. Amended HR and RR thresholds. Adjusted wording of airway parameters; added respiratory items; added APU evaluation; removed suspected meningococcus and diabetic ketoacidosis; added pH < 7.2 and unresolved pain.	Trigger	HR/RR data driven	Yes	14	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Required nebulised ephedrine or no improvement after nebulisers; pH < 7.2; unresolved pain or current analgesic therapy; fitting.
<b>Other PTTT</b>																			
NHS Institute for Innovation and Improvement PEWS <sup>4</sup>	Designed as part of a NHS Institute fellowship project. Adapted from adult scores and Brighton PEWS. No formal development or internal validation study published.	Score	APLS values	Yes	6	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Worsening retractions; cyanosis.	
Paediatric medical emergency team (PMET) triggering criteria (a)	Initial development for use in a US tertiary centre to activate a MET. Retrospective chart review of case patients (n=44, code calls) used to generate candidate items. Clinical judgement used to select final items. No formal validation of final tool reported.	Trigger	Expert opinion	No	4	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Worsening retractions; cyanosis.	
PMET triggering criteria (b)	Initial development for use in a US tertiary centre to activate a MET. Minimal description of tool development—authors deliberately chose broad criteria and categories of illness rather than specific vital signs. No formal validation study reported.	Trigger	Expert opinion	Unclear	12	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Cardiac or respiratory arrest; seizures with apnoea; progressive lethargy; circulatory compromise/acute shock syndrome.	
Paediatric rapid response team (PRRT) triggering criteria (a)	Initial development for use in a US tertiary centre, to activate an RRT. Triggering items elected through expert consensus locally—reference to similarity to MAC and PMET triggering criteria (a). No formal validation study reported.	Trigger	Expert opinion	No	6	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Convulsion.	
PRRT triggering criteria (b)	Initial development for use in calling RRT team in a tertiary centre in Pakistan. Minimal explanation for selection of calling criteria. No formal validation study reported in the literature.	Trigger	Unclear	Yes	8	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Convulsion.	

Continued

AUTHOR PROOF

AUTHOR PROOF

Table 2 Continued

PTTT name (reference)	Development/ modification details	Score/ trigger	Choice of thresholds/ parameters	Age-dependent thresholds	No. of items in the tool*	PTTT parameters															
						Respiratory rate	Respiratory effort/ distress	LOC/ behaviour	Oxygen saturation	Capillary refill time	Oxygen therapy	Systolic blood pressure	Pain	Staff concern	Skin colour	Airway problems	Temperature	Pulses	Family concern	Other items	
Logistic regression algorithm <sup>6</sup>	Initial development based on data mining of electronic health records in US tertiary centre. Extracted 24 hours of clinical data from inpatients (n=6722 controls, 526 PICU transfers) and used logistic regression model to select 29 item tool. Validation performed on subset of development dataset.	Score	Expert opinion	Yes	29	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Acuity level (local measure); tissue perfusion and oxygenation.

\*Multiple parameters are often required to be collected for each scoring item/category, eg, scoring the 'cardiovascular' category in the Brighton PEWS requires collection/evaluation of HR, skin colour and capillary refill time.

†Denotes a study included in the effectiveness review.

APLS, advanced paediatric life support; AWPV, alert, voice, pain, unresponsive; HDU, high-dependency unit; HR, heart rate; LOC, level of consciousness; NHS, National Health Services; PICU, paediatric intensive care unit; PTTs, paediatric track and trigger tool; RR, respiratory rate.

PICU or HDU transfer<sup>33</sup> (AUROC=0.79, PPV=3.6%). A derivative of the MAC using an aggregate score, the Cardiff and Vale PEWS (C&VPEWS), was tested using the same cohort and outcome measures in an earlier external study (AUROC=0.86, PPV=5.9%)<sup>32</sup> and was the best performing PTTT in an external case-control study evaluating multiple PTTT<sup>29</sup> (AUROC=0.89).

**Bristol PEWT**

The Bristol PEWT was evaluated by five external validation studies: two chart review studies<sup>3, 35</sup> (no AUROC), one small cohort study of PICU transfers<sup>34</sup> (AUROC=0.91, PPV=11%), and two case-control studies looking at code calls<sup>28</sup> (AUROC=0.75) and a composite of death, arrests and PICU transfers<sup>29</sup> (AUROC=0.62).

**Other PTTT**

The NHS III PEWS was tested by one external cohort study looking at a composite of death or unplanned transfers to PICU or HDU<sup>14</sup> (AUROC=0.88, PPV=4.3%) and one external case-control study looking at a composite of death, arrests and PICU transfers<sup>29</sup> (AUROC=0.82). Zhai *et al* developed and retrospectively evaluated a logistic regression algorithm in an internal case-control study looking at urgent PICU transfers in the first 24 hours of admission<sup>19</sup> (AUROC=0.91, PPV=4.8%).

Across PTTT, studies reporting performance characteristics of a tool at a range of different scoring thresholds demonstrate the expected interaction and trade-off between sensitivity and specificity—at lower triggering thresholds, sensitivity is high but specificity is low; at higher thresholds, the opposite is true.

**Inter-rater reliability and completeness of data**

Accurate assessment of the ability of a PTTT to predict clinical deterioration is contingent on accuracy and reliability of tool scoring (whether by bedside nurses in practice or by researchers abstracting data) and the availability of underpinning observations. Only five papers made reference to accuracy or reliability of scoring,<sup>28 31 37 42 45</sup> with mixed results: for example, two nurses separately scoring a subset of patients on the Modified Brighton PEWS (a) achieved an intra-class coefficient of 0.92,<sup>31</sup> but a study nurse and bedside nurse achieved only 67% agreement in scoring the C-CHEWS tool.<sup>37</sup> Completeness of data was reported in 11 studies.<sup>11 14 19 29 30 32 33 34 44 45 47</sup> An evaluation of the Modified Bedside PEWS (a) reported that 'the PEWS was correctly performed and could be used for inclusion in the study' in 59% of cases,<sup>30</sup> a prospective study bench-testing the C&VPEWS found an average completeness rate of 44% for the seven different parameters in daily practice,<sup>32</sup> while a multicentre study of the Bedside PEWS reported that 'only 5.1% (of observation sets) had measurements on all seven items'.<sup>44</sup>

**Table 3** Summary of PTTT validation study outcomes

PTTT	First author, year	Country	Study population	Study design	Number of centres	PTTT used in practice?	Internal/external validation study?	Outcome measures	Sample size	Score or trigger?	Score tested/maximum score	Which score used (frequency of scoring)?	AUROC	Sensitivity	Specificity	PPV	NPV	Notes on accuracy/reliability of scoring and missing data	Quality score (max=24)
Paediatric early warning system (PEWS) score	Duncan 2006 <sup>28</sup>	Canada	All inpatients	Case-control study (retrospective)	1	No	Int	Code blue call for actual or suspected cardiopulmonary arrest	215 (87 cases)	S	5/26	Max 24 hours before event (hourly)	0.90	78.0	95.0	4.2†		No details on data abstraction. 100% of eligible cases and 83% of eligible controls excluded due to incomplete clinical data.	14
	Robson 2013 <sup>29</sup>	USA	All inpatients	Case-control study (retrospective)	1	No	Ext	Code blue call	192 (86 cases)	S	5/32	Max 24 hours before event (six hourly)	0.85	86.6	72.2			Four researchers scored PTTT from 20 charts. Inter-rater reliability of 0.95. No details on extent of missing data.	8
	Chapman 2017 <sup>29</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or unplanned PICU transfer	608 (297 cases)	S	7/32	Max 48 hours before event (per usual practice)	0.82	70.0	75.0	72.6	72.0	Data abstraction by single researcher. 36% of observation sets contained HR, RR, O2 Sats, systolic BP, temperature and assessment of consciousness.	17
Bedside PEWS	Praeshuram 2009 <sup>1</sup>	Canada	All inpatients	Case-control study (retrospective)	1	No	Int	Urgent PICU transfer (without code blue call)	180 (60 cases)	S	8/26	Max 24 hours before event (hourly)	0.91	82.0	93.0			Availability of scoring items in medical records varied from 27% (cap refill time) to 95% (oxygen therapy).	21
	Praeshuram 2011 <sup>44</sup>	Canada and UK	All inpatients	Case-control study (prospective)	4	No	Ext	Urgent PICU transfer or immediate call to resuscitation team	2074 (686 cases)	S	7/26	Max 24 hours before event (hourly)	0.87	64.0	91.0			PTTT scores calculated electronically after abstraction by research nurse. 5.1% of records had all seven items recorded, 31% had at least five items.	22
	Robson 2013 <sup>29</sup>	USA	All inpatients	Case-control study (retrospective)	1	No	Ext	Code blue call	192 (86 cases)	S	7/26	Max 24 hours before event (six hourly)	0.73	56.3	78.1			See above.	8
	Zhai 2014 <sup>49</sup>	USA	All inpatients	Case-control study (retrospective)	1	No	Ext	Urgent PCU transfer within 24 hours of admission	6362 (63 cases)	S	7/26	Max 24 hours before event (hourly)	0.82	73.6	71.7	2.1†		Data extracted from electronic health records. Excluded two items of Bedside PEWS (oxygen therapy and respiratory effort) due to difficulty abstracting.	17
Modified Bedside PEWS (a)	Gawronski 2016 <sup>30</sup>	Italy	Stem Cell Transplant Unit	Case-control study (retrospective)	1	No	Ext	Unexpected death, urgent consult with HRT or urgent PICU transfer	99 (19 cases)	S	6/26	Score 4 hours before event	0.90	79.0	97.5			Data abstracted by research nurse. No details on extent of missing data. Conflicting missing observations were identified in interviews with clinical staff.	15
	Chapman 2017 <sup>29</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	S	6/26	Max 48 hours before event (per usual practice)	0.88	72.0	89.0	86.0	77.0	See above.	17
	Fulkschot 2015 <sup>30</sup> (study 1)	The Netherlands	Oncology ward	Case-cohort study (retrospective)	1	Yes	Int	Emergency medical intervention or reviewed by PICU staff or staff concern	118 (15 cases)	S	8/28	Unclear (minimum eight hourly)			73.0		41% of admissions excluded from study due to incomplete PTTT scores.	10	
Modified Bedside PEWS (b)	Fulkschot 2015 <sup>30</sup> (study 2)	The Netherlands	All inpatients	Case-cohort study (retrospective)	1	Yes	Int	PICU transfer	Unclear (24 cases)	S	8/28	Score 2-6 hours before event (minimum eight hourly)		66.6				High rate of exclusions reported due to missing data.	10
	Fulkschot 2015 <sup>30</sup> (study 3)	The Netherlands	All inpatients	Case-cohort study (prospective)	1	Yes	Int	Emergency medical intervention	Unclear (14 cases)	S	8/28	Unclear (minimum eight hourly)		100				No details on missing data.	10
	Chapman 2017 <sup>29</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	S	7/28	Max 48 hours before event (per usual practice)	0.87	69.0	91.0	87.9	79.0	See above.	17
Modified Bedside PEWS (b)	Ross 2015 <sup>49</sup>	USA	All inpatients	Case-control study (retrospective)	1	No	Int	Urgent PICU transfer	4628 (848 cases)	S	8/26	Max during admission		70.0	84.0			No details on data abstraction. Respiratory effort category excluded due to difficulty abstracting. No details on missing data.	9

Continued

AUTHOR PROOF

Table 3 Continued

PTTT	First author, year	Country	Study population	Study design	Number of centres	PTTT used in practice?	Internal/external validation in practice?	Outcome measures	Sample size	Score or trigger?	Score tested/maximum score	Which score used (frequency of scoring)?	AUROC	Sensitivity	Specificity	PPV	NPV	Notes on accuracy/reliability of scoring and missing data	Quality score (max=24)
Modified Brighton PEWS (a)	Tucker 2009 <sup>11</sup>	USA	General medical unit	Cohort study (prospective)	1	Yes	Int	PICU transfer	2979 (51 cases)	S	3/11	Max during admission (four hourly)	0.89	90.2	74.4	5.8	99.8	Intraclass coefficient of 0.92 reported for two bedside nurses scoring 35 patients. No details on missing data.	14
	Zhai 2014 <sup>13</sup>	USA	All inpatients	Case-control study (retrospective)	1	No	Ext	Urgent PCU transfer within 24 hours of admission	6352 (53 cases)	S	2/11	Max 24 hours before event (hourly)	0.74	68.4	81.6	2.3	Data extracted from electronic health records. Only included records with complete PEWS score: 64% of eligible cases and 51% of eligible controls excluded.	17	
	Fenix 2015 <sup>28</sup>	USA	PICU transfers among all inpatients (trauma, haematology, oncology, surgical and cardiac wards)	Case-control study (retrospective)	1	Yes	Ext	Non-elective PICU transfer followed by deterioration event	97 PICU transfers followed by deterioration event	S	3/11	Max during admission	80.0	43.0	61.0	67.0	No details on missing data.	15	
Modified Brighton PEWS (b)	Akre 2010 <sup>15</sup>	USA	All inpatients	Chart review study (retrospective)	1	No	Int	Rapid response (170 RRT calls, 16 code blue call)	186 cases	S	4/13	Max 24 hours before event (minimum four hourly)	85.5	85.5			Scores abstracted from charts by single nurse, having calibrated with advanced nurse practitioner. Charts were scored missing if any items missing. 25% of charts missing behavioural state, 28% cardiovascular colour.	14	
	Chapman 2017 <sup>29</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	S	4/13	Max 48 hours before event (per usual practice)	0.79	61.0	84.0	78.4	69.0	See above.	17
Modified Brighton PEWS (c)	Skalicky 2012 <sup>24</sup>	USA	Medical surgical wards	Case-control study (retrospective)	1	No	Int	PICU transfer	350 (100 cases)	S	2.5/9	Max 48 hours before event (four hourly)	0.81	62.0	89.0		Data abstracted from medical charts and notes. Researcher blinded to PICU status. No details on missing data.	15	
	Chapman 2017 <sup>29</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	S	4/9	Max 48 hours before event (per usual practice)	0.74	46.0	90.0	81.3	63.0	See above.	17
Children's Hospital Early Warning Score	McLellan 2014 <sup>30</sup>	USA	All inpatients	Case-control study (retrospective)	1	Yes	Int	Arrest or unplanned PICU transfer	1136 (380 cases)	S	4/12	Max in admission (four hourly)	0.90	84.2	80.9		No details on missing data.	10	
Children's Hospital Los Angeles PEWS Score	McLellan 2013 <sup>30</sup>	USA	Cardiovascular unit	Case-control study (retrospective)	1	Yes	Int	Arrest or unplanned PICU transfer	312 (64 cases)	S	3/12	Max 18 hours before event (four hourly)	0.86	95.3	76.2	50.8	98.4	Study nurse and bedside nurse abstracted vital signs for 37 patients, 67% agreement. No details on missing data.	9
	Agulnik 2016 <sup>41</sup>	USA	Oncology unit	Case-control study (retrospective)	1	Yes	Ext	Unplanned PICU transfer	330 (110 cases)	S	4/12	Max 24 hours before event (hourly)	0.96	86.0	95.0		PTTT scores abstracted by researcher. Did not abstract if vital signs were present but no PTTT score calculated by nurse. No details on missing data.	14	
Children's Hospital Los Angeles PEWS Score	Agulnik 2017 <sup>42</sup>	Guatemala	Oncology unit	Case-control study (retrospective)	1	Yes	Ext	Unplanned PICU transfer	258 (129 cases)	S	4/12	Max 24 hours before event (three hourly)		91.0	88.0		Researcher excluded charts with related scores, reporting 14% error rate (PTTT score calculated incorrectly) and 3% omission rate (vital signs recorded but no PTTT score calculated). One out of 130 cases excluded due to missing PTTT documentation.	16	
Children's Hospital Los Angeles PEWS Score	Mandell 2015 <sup>40</sup>	USA	Inpatients discharged from PICU to ward	Case-control study (retrospective)	1	Yes	Int	Early unplanned re-admission to PICU (within 48 hours of discharge from PICU to ward)	189 (68 cases)	S	2/10	First score assigned on ward, post-PICU discharge	0.71	76.0	56.0		No details on missing data.	12	

Continued

**Table 3** Continued

PTTT	First author, year	Country	Study population	Study design	Number of centres	PTTT used in practice?	Internal/external validation study?	Outcome measures	Sample size	Score or trigger?	Score test/minimum score	Which score used (frequency of scoring)?	AUROC	Sensitivity	Specificity	PPV	NPV	Notes on accuracy/reliability of scoring and missing data	Quality score (max=24)
Melbourne Action Criteria	Tume 2007 <sup>23</sup>	UK	Inpatients with unplanned PICU transfer	Chart review study (retrospective)	1	No	Ext	Unplanned PICU transfer	33 cases	T	N/A	Unclear	0.78	87.8				Data abstracted by two reviewers. Due to a large number of missing records and observation charts'. See above.	11
	Tume 2007 <sup>23</sup>	UK	Inpatients with an unplanned PHDU transfer	Chart review study (retrospective)	1	No	Ext	Unplanned PHDU transfer	32 cases	T	N/A	Unclear	0.79	87.5	83.2	3.6	99.7	Observation charts altered to include all PTTT parameters. 96% of records missing at least one component. Missing data assumed to be normal.	11
	Edwards 2011 <sup>24</sup>	UK	All inpatients	Cohort study (retrospective)	1	No	Ext	Death or unplanned PICU or HDU transfer	1000 (16 cases)	T	N/A	Any trigger over admission (per usual practice)	0.82	68.3	83.2	3.6	99.7	Observation charts altered to include all PTTT parameters. 96% of records missing at least one component. Missing data assumed to be normal.	17
	Chapman 2017 <sup>25</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	T	N/A	Max 48 hours before event (per usual practice)	0.71	93.0	49.0	64.0	88.0	See above.	17
Cardiff and Vale PEWS	Edwards 2009 <sup>26</sup>	UK	All inpatients	Cohort study (prospective)	1	No	Int	Death or unplanned PICU or HDU transfer	1000 (16 cases)	S	2/8	Max score during admission (per usual practice)	0.86	69.5	89.9	5.9	99.7	Observation charts altered to include all PTTT parameters. 96% of records missing at least one component. Missing data assumed to be normal.	18
	Chapman 2017 <sup>25</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	S	3/8	Max 48 hours before event (per usual practice)	0.89	80.0	86.0	84.0	82.0	See above.	17
Bristol paediatric early warning tool (PEWT)	Tume 2007 <sup>23</sup>	UK	Inpatients with an unplanned PICU transfer	Chart review (retrospective)	1	No	Ext	Unplanned PICU transfer	33 cases	T	N/A	Unclear	0.86	87.8				See above.	11
	Tume 2007 <sup>23</sup>	UK	Inpatients with an unplanned PHDU transfer	Chart review (retrospective)	1	No	Ext	Unplanned PHDU transfer	32 cases	T	N/A	Unclear	0.84	84.4				See above.	11
	Wright 2011 <sup>28</sup>	UK	All inpatients	Chart review (retrospective)	1	Yes	Ext	Cardiac arrest	55 cases	T	N/A	If triggered 24 hours before event	0.91	49.1				One case excluded due to missing notes. No details on missing data.	11
	O'Loughlin 2012 <sup>24</sup>	UK	All inpatients	Cohort study (prospective)	1	Yes	Ext	PICU transfer	331 (7 cases)	T	N/A	Triggered during admission (12 hourly)	0.91	100	81.0	11.0		No details on missing data.	6
	Robson 2013 <sup>28</sup>	USA	All inpatients	Case-control study (retrospective)	1	No	Ext	Code blue call	192 (86 cases)	T	N/A	Triggered 24 hours before event (6 hourly)	0.75	76.3	61.5			See above.	8
	Chapman 2017 <sup>25</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	T	N/A	If triggered 48 hours before event (per usual practice)	0.82	96.0	28.0	56.0	88.0	See above.	17
Modified Bristol PEWT (b)	Clayson 2014 <sup>28</sup>	UK	Cardiac ward	Cohort study (prospective)	1	Yes	Int	'A deteriorating patient'	126 (unclear number of cases)	T	N/A	Unclear		12.5	97.0			No details on missing data.	5
NHS Institute for Innovation and Improvement PEWS	Mason 2016 <sup>4</sup>	UK	All inpatients	Cohort study (retrospective)	1	No	Ext	Death or unplanned PICU or HDU transfer	1000 (16 cases)	S	2/7	Max score over admission (per usual practice)	0.88	80.0	81.0	4.3	99.7	Observation charts altered to include all PTTT parameters. 96% of records missing at least one component. Missing data assumed to be normal.	15
	Chapman 2017 <sup>25</sup>	UK	All inpatients	Case-control study (retrospective)	1	No	Ext	Death, arrest or PICU transfer	608 (297 cases)	S	2/7	Max 48 hours before event (per usual practice)	0.82	83.0	65.0	69.6	80.0	See above.	17
Logistic regression algorithm	Zhai 2014 <sup>19</sup>	USA	All inpatients	Case-control study (retrospective)	1	No	Ext	Urgent PICU transfer within 24 hours of admission	6552 (53 cases)	S	>0.5	Max 24 hours before event (hourly)	0.91	84.9	85.9	4.8		Data extracted from electronic health records. No details on extent of missing data. Authors report that 'missing data were a major cause of incorrect prediction'.	17

Continued

AUTHOR PROOF

AUTHOR PROOF

Table 3 Continued

PTTT	First author, year	Country	Study population	Study design	Number of centres	PTTT used in practice?	Internal/external validation study?	Outcome measures	Sample size	Score or trigger?	Score tested/maximum score	Which score used (frequency of scoring)?*	AUROC	Sensitivity	Specificity	PPV	NPV	Notes on accuracy/reliability of scoring and missing data	Quality score (max=24)
Burton Paediatric Early Warning Score	Ahmed 2012 <sup>38</sup>	UK	PICU admissions only	Chart review (retrospective)	1	Yes	Int	PICU admission	23	S	4/19	Max 24 hours before event (unclear)	93.0					Data extracted from case notes by two reviewers. No details on missing data.	4
Between the Flags' PEWS	Blackstone 2017 <sup>33</sup>	UK	Urgent PICU admissions only	Chart review (retrospective)	1	Yes	Ext	Urgent PICU admission	100	T	N/A	Unclear	91.0					Data extracted from health records. No details on missing data.	8

All studies conducted in a specialist/tertiary centre.

PPV and NPV values in italics represent results from case-control studies—these values are misleading in isolation because they assume that the wider prevalence rate of the adverse event is equal to the case to control ratio used in the research study (eg, if the researchers studied 300 cases and 300 controls, the prevalence rate of adverse events for the calculation of PPV is 50%). As per the cohort studies, prevalence rates of critical events are typically far lower among hospitalised paediatric populations than the case-control ratios used in studies, and so PPV values would be considerably lower in clinical practice.

Studies classified as internal validation if the setting for the study was the same hospital and same research team as those who developed the score. Studies classified as external validation if the score was tested in a different centre and by a different research team to those who developed it.

\*Typically, study researchers collected or abstracted multiple PTTT scores for each patient at different time points, but can only use one score per patient for the analysis of the tool's predictive ability. This column specifies which score the researchers used. In most cases, the study team used the maximum PTTT score recorded for each patient in a given study window, eg, 24 hours prior to a critical event for case patients. The text in parentheses describes the frequency with which scores were assessed or abstracted for each patient, if this information was described in the paper.

†Case-control study, but PPV value calculated based on clinical prevalence of event as measured at local centre during the study.

AUROC, area under the receiver operating characteristic curve; Ext, external validation; HFNC, high flow nasal cannula; Int, internal validation; Max, maximum; N/A, not applicable; NPV, negative predictive value; PHUO, paediatric high-dependency unit; PICU, paediatric intensive care unit; PPV, positive predictive value; PTTT, paediatric track and trigger tool; RRT, rapid response team; S, score; T, trigger.

## Question 2: how effective are early warning systems at reducing mortality and critical events in hospitalised children?

Eleven papers meeting inclusion criteria were excluded from analysis for providing insufficient statistical information (eg, denominator data, absolute numbers of events) to calculate effect sizes.<sup>39 51–59</sup> Further details on papers excluded from analysis are provided in online supplementary table 5. Findings from the 19 studies included in the analysis are summarised in table 4.

### Type of early warning system interventions

Seventeen interventions involved the introduction of a new PTTT,<sup>13 15–18 60–72</sup> one intervention introduced a mandatory triggering element to an existing PTTT<sup>71</sup> and one study reported a large, multicentre analysis of MET introduction with no details on PTTT use.<sup>73</sup> Twelve interventions included the introduction of a new MET or RRT,<sup>13 15–18 60–65 69</sup> while four further interventions introduced a new PTTT in a hospital with an existing MET or RRT. Only three studies therefore evaluated a PTTT in the absence of a dedicated response team.<sup>67 68 70</sup> A staff education programme was explicitly described in 10 interventions.<sup>13 15 17 61 62 64 67 68 70 72</sup>

Of the 18 studies that used a PTTT, only 7 used a tool that had been formally evaluated for validity: 3 used the Bedside PEWS,<sup>64 65 70</sup> 2 used the MAC,<sup>13 62</sup> 1 used the Modified Brighton PEWS (b)<sup>72</sup> and 1 used the C-CHEWS.<sup>67</sup> One study did not report the PTTT used,<sup>61</sup> while 10 studies used a variety of calling criteria and local modifications to validated tools that had not been evaluated for validity.<sup>15–18 60 63 66 68 69 71</sup>

### Mortality (ward or hospital wide)

Two uncontrolled before-after studies (both with MET/RRT) reported significant mortality rate reductions postintervention: one in hospital wide deaths per 100 discharges<sup>17</sup> (RR=0.82, 95% CI 0.70 to 0.95) and one in total hospital deaths per 1000 admissions (RR=0.65, 95% CI 0.57 to 0.75) and deaths on the ward ('unexpected deaths') per 1000 admissions<sup>62</sup> (RR=0.35, 95% CI 0.13 to 0.92). Seven studies found no reductions in mortality, including two high-quality multicentre studies.<sup>13 15 60 63–65 73</sup> Parshuram *et al* conducted a cluster randomised trial and found no difference in all-cause hospital mortality rates between 10 hospitals randomly selected to receive an intervention centred around use of the Bedside PEWS and 11 usual care hospitals, 1-year postintervention (OR=1.01, 95% CI 0.61 to 1.69).<sup>64</sup> Kutty *et al*<sup>73</sup> assessed the impact of MET implementation in 38 US paediatric hospitals with an interrupted time series study, and reported no difference in the slope of hospital mortality rates 5 years postintervention and the expected slope based on preimplementation trends (OR=0.94, 95% CI 0.93 to 0.95).

### PICU mortality

Two uncontrolled before-after studies (both with MET/RRT) reported a significant postintervention reduction in



**Table 4** Summary of early warning system effectiveness study outcome

Outcome	Intervention				Country	Number of centres	Specialist unit?	Existing RRT/MET?	Population	Study design	Study duration in months	Events before, n (rate)	Events after, n (rate)	Effect size (95% CI)	P value	Quality score (max=26)
	First author, year	Implemented a new PTTT	Implemented a new RRT/MET	Modified escalation process												
<b>Mortality</b>																
Deaths on ward (per 1000 admissions)	Tibballs 2005 <sup>13</sup>	✓	✓	✓	Australia	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	53 (41 before, 12 after)	13 (0.12)	2 (0.06)	RR=0.45 (0.10 to 1.99)†	0.29	10
Hospital-wide deaths (per 100 discharges)	Shank 2007 <sup>17</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	64 (67 before, 17 after)	547 (1.01)	158 (0.83)	RR=0.82 (0.70 to 0.95)	<b>0.007</b>	15
Hospital-wide deaths, excluding neonate ICU and ED (per 1000 discharges)	Zenker 2007 <sup>20</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	34 (23 before, 11 after)	97 (4.30)	52 (4.45)	RR=1.04 (0.74 to 1.45)†	0.57	12
Deaths outside ICU (per 1000 non-ICU patient days)	Brill 2007 <sup>15</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	27 (15 before, 12 after)	9 (0.10)	2 (0.04)	RR=0.39 (0.09 to 1.80)†	0.13	14
Ward death rate (per 1000 ward admissions)	Hanson 2010 <sup>61</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (retrospective)	36 (24 before, 12 after)	13 (1.50)	2 (0.45)	RR=0.30 (0.07 to 1.31)†	0.07	18
Total hospital deaths (per 1000 admissions)	Tibballs 2009 <sup>62</sup>	✓	✓	✓	Australia	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	89 (41 before, 48 after)	469 (4.36)	398 (2.67)	RR=0.65 (0.37 to 0.75)	<b>&lt;0.0001</b>	15
Deaths on ward (per 1000 admissions)	Tibballs 2009 <sup>62</sup>	✓	✓	✓	Australia	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	89 (41 before, 48 after)	13 (0.12)	6 (0.04)	RR=0.35 (0.13 to 0.92)	<b>0.03</b>	15
All-cause hospital mortality (per 1000 admissions)	Kotsalis 2011 <sup>63</sup>	✓	✓	✓	Canada	4	Y	N	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	553 (9.97)	540 (9.65)	RR=0.97 (0.831 to 1.12)	0.65	18
All-cause hospital mortality (per 1000 discharges)	Prabhram 2018 <sup>64</sup>	✓	✓	✓	Belgium, Ireland, The Netherlands, England, Italy, Canada, New Zealand	21	Y	N	All inpatients	Cluster randomised trial (prospective)	18 (6 pre, 12 post)	61 (1.31)	147 (1.56)	OR=1.01 (0.61 to 1.68)	0.96	23
Hospital mortality (per 1000 admissions)	Kutty 2016 <sup>23</sup>	✓	✓	✓	USA	38	Y	N	All inpatients	Interrupted time series (retrospective)	180 (60 before, 120 after)	N/A	N/A	OR=0.94 (0.93 to 0.95)	0.98	20
<b>PICU mortality</b>																
PICU mortality after ICU admission from ward (per PICU admission)	Anwarul-Haque 2010 <sup>65</sup>	✓	✓	✓	Pakistan	1	Y	N	All inpatients	Uncontrolled before-after study (retrospective)	18 (9 before, 9 after)	23 (51.1)	5 (15.65)	RR=0.31 (0.13 to 0.72)†	<b>0.0071</b>	6
PICU mortality after ICU readmission within 48 hours of discharge (per 1000 admissions)	Kotsalis 2011 <sup>63</sup>	✓	✓	✓	Canada	4	Y	N	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	16 (0.29)	7 (0.13)	RR=0.43 (0.17 to 0.99)	<b>&lt;0.05</b>	18
PICU mortality after urgent PICU admission from ward (per 1000 admissions)	Kotsalis 2011 <sup>63</sup>	✓	✓	✓	Canada	4	Y	N	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	70 (1.3)	61 (1.1)	RR=0.80 (0.70 to 1.00)	0.25	18
Death prior to discharge (per unplanned PICU transfer)	Boravida 2014 <sup>66</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Interrupted time series study (prospective)	59 (32 before, 27 after)	51 (6.3)	56 (6.5)	RR=1.03 (0.72 to 1.49)†	0.99	23
PICU mortality (per PICU admission)	Duns 2014 <sup>68</sup>	✓	✓	✓	Australia	1	Y	Y	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	30 (6.57)	20 (5.49)	RR=0.64 (0.37 to 1.11)†	0.14	7
Death in PICU (per 1000 patient-days)	Agnihit 2017 <sup>27</sup>	✓	✓	✓	Guatemala	1	Y	N	Oncology unit	Uncontrolled before-after study (retrospective)	24 (12 before, 12 after)	21 (1.25)	22 (1.10)	RR=0.89 (0.49 to 1.61)†	0.76	19
Death in PICU (per emergency PICU admission)	Sifton 2015 <sup>68</sup>	✓	✓	✓	UK	1	Y	N	All PICU admissions	Controlled before-after study (retrospective)	24 (12 before, 12 after)	17 (10.8)	14 (8.4)	RR=0.78 (0.40 to 1.53)†	0.47	16
Deaths in PICU (per unplanned PICU admission)	Kolovec 2018 <sup>69</sup>	✓	✓	✓	USA	1	Y	N	All unplanned PICU admissions	Uncontrolled before-after study (retrospective)	78 (42 before, 36 after)	54† (4.9)	40† (3.8)	RR=0.77 (0.32 to 1.15)†	0.20†	12
PICU mortality (per 1000 discharges)	Pashuram 2018 <sup>74</sup>	✓	✓	✓	Belgium, Ireland, The Netherlands, Canada, New Zealand	21	Y	N	All inpatients	Cluster randomised trial (prospective)	18 (6 pre, 12 post)	34 (0.73)	91 (0.96)	OR=0.85 (0.48 to 1.86)	0.88	23
<b>Cardiac arrest</b>																
Cardiac arrests on ward (per 1000 admissions)	Tibballs 2005 <sup>13</sup>	✓	✓	✓	Australia	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	53 (41 before, 12 after)	20 (0.19)	4 (0.11)	RR=0.58 (0.20 to 1.70)	0.33	10
Cardiopulmonary arrests (per 1000 non-ICU patient-days)	Brill 2007 <sup>15</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	27 (15 before, 12 after)	7 (0.09)	2 (0.04)	RR=0.50 (0.10 to 2.42)†	0.11	14

Continued

AUTHOR PROOF

AUTHOR PROOF

Table 4 Continued

Outcome	First author, year	Intervention			Country	Number of centres	Specialist unit?	Existing RRT/MET?	Population	Study design	Study duration in months	Events before, n (rate)	Events after, n (rate)	Effect size (95% CI)	P value	Quality score (max:26)
		Implemented a new PTTT	Implemented new RRT/MET	Staff training/education												
Ward cardiac arrest rate (per 1000 ward admissions)	Hanson 2010 <sup>41</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (retrospective)	38 (24 before, 12 after)	11 (1.27)	2 (0.45)	RR=0.35 (0.08 to 1.58)†	0.13	18
Ward cardiopulmonary arrests (per 1000 patient-days)	Hunt 2008 <sup>16</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	24 (12 before, 12 after)	5 (0.10)	5 (0.10)	RR=0.38 (0.22 to 0.64)	0.97	17
Preventable cardiac arrests (per 1000 ward admissions)	Tibballs 2009 <sup>42</sup>	✓	✓	✓	Australia	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	89 (41 before, 48 after)	17 (0.16)	10 (0.07)	RR=0.45 (0.20 to 0.97)	<b>0.04</b>	15
Unexpected cardiac arrests (per 1000 ward admissions)	Tibballs 2009 <sup>42</sup>	✓	✓	✓	Australia	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	89 (41 before, 48 after)	20 (0.19)	24 (0.17)	RR=0.31 (0.19 to 0.64)	0.75	15
Actual cardiopulmonary arrests (per 1000 ward admissions)	Kotsakis 2011 <sup>43</sup>	✓	✓	✓	Canada	4	Y	N	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	69 (1.9)	66 (1.8)	RR=0.95 (0.76 to 1.98)	0.68	18
Near cardiopulmonary arrests (per 1000 ward admissions)	Kotsakis 2011 <sup>43</sup>	✓	✓	✓	Canada	4	Y	N	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	123 (3.4)	67 (1.9)	RR=0.54 (0.52 to 0.57)	<0.001	18
Cardiac arrests on ward (per 1000 non-ICU patient-days)	Bonafide 2014 <sup>44</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Interrupted time series study (prospective)	69 (32 before, 27 after)	61 (0.03)	21 (0.01)	RR=0.36 (0.07 to 1.78)†	0.21	23
Cardiac arrests (per 1000 patient-days)	Parekh 2018 <sup>45</sup>	✓	✓	✓	Belgium, Ireland, The Netherlands, England, Italy, Canada, New Zealand	21	Y	N	All inpatients	Cluster randomised trial (prospective)	18 (six pre, 12 post)	18 (0.11)	32 (0.10)	RR=1.02 (0.65 to 1.62)	0.92	23
Respiratory arrest	Hunt 2008 <sup>16</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	24 (12 before, 12 after)	11 (0.23)	3 (0.06)	RR=0.27 (0.07 to 0.95)	<b>0.04</b>	17
Cardiac or respiratory arrest	Zenker 2007 <sup>46</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	34 (23 before, 11 after)	180 (7.98)	60 (5.13)	RR=0.64 (0.48 to 0.86)†	0.19	12
Code calls (per 1000 non-ICU patient-days)	Brill 2007 <sup>15</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	27 (15 before, 12 after)	25 (0.27)	6 (0.11)	RR=0.42 (0.17 to 1.03)†	0.08†	14
Code calls (per 1000 non-ICU patient-days)	Shank 2007 <sup>17</sup>	✓	✓	✓	USA	1	Y	N	All inpatients	Uncontrolled before-after study (prospective)	84 (67 before, 17 after)	53 (0.52)	5 (0.15)	RR=0.29 (0.10 to 0.65)	<b>0.008</b>	15
Code calls (per 1000 admissions)	Anwar-Haque 2010 <sup>47</sup>	✓	✓	✓	Pakistan	1	Y	N	All inpatients	Uncontrolled before-after study (retrospective)	18 (9 before, 9 after)	26 (5.25)	12 (2.79)	RR=0.52 (0.28 to 1.03)	0.06	6
Calls for urgent review/assistance	Parekh 2014 <sup>48</sup>	✓	✓	✓	Canada	1	N	N	All inpatients	Uncontrolled before-after study (prospective)	8 (3 before, 5 after)	8 (9.5)	8 (3.4)	RR=0.36 (0.13 to 0.95)†	<b>0.04†</b>	23
Urgent calls to paediatrician (per 1000 patient-days)	Parekh 2014 <sup>48</sup>	✓	✓	✓	Canada	1	N	N	All inpatients	Uncontrolled before-after study (prospective)	8 (3 before, 5 after)	19 (22.6)	12 (5.1)	RR=0.23 (0.11 to 0.46)†	<0.0001	23
Code blue calls on the ward (per 1000 admissions)	Kotsakis 2011 <sup>43</sup>	✓	✓	✓	Canada	4	Y	N	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	210 (3.75)	150 (2.79)	RR=0.71 (0.51 to 0.93)	<0.0001	18
Urgent calls to outreach team (per 1000 admissions)	Duns 2014 <sup>48</sup>	✓	✓	✓	Australia	1	Y	Y	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	1058 (39.5)	2120 (76.0)	RR=1.92 (1.78 to 2.07)†	<b>0.02</b>	7
RRT calls (per 1000 patient-days)	Panesar 2014 <sup>41</sup>	✓	✓	✓	USA	1	Y	Y	All inpatients	Uncontrolled before-after study (retrospective)	42 (18 before, 24 after)	44 (3.14)	69 (4.22)	RR=1.35 (0.92 to 1.96)†	0.11	15
RRT calls (per 1000 patient days)	Douglas 2016 <sup>42</sup>	✓	✓	✓	USA	1	Y	Y	All inpatients	Uncontrolled before-after study (retrospective)	24 (12 before, 12 after)	184 (6.17)	292 (9.88)	RR=1.59 (1.33 to 1.90)†	<0.001	12
Code calls (per 1000 patient days)	Douglas 2016 <sup>42</sup>	✓	✓	✓	USA	1	Y	Y	All inpatients	Uncontrolled before-after study (retrospective)	24 (12 before, 12 after)	31 (0.98)	20 (0.67)	RR=0.68 (0.39 to 1.19)†	0.21	12
ICU transfers	Parekh 2014 <sup>48</sup>	✓	✓	✓	Canada	1	N	N	All inpatients	Uncontrolled before-after study (prospective)	8 (3 before, 5 after)	5 (5.9)	19 (8.1)	RR=1.37 (0.51 to 3.63)†	0.54†	23
Clinical deterioration events on ward prior to transfer to specialist unit (per 1000 patient-days)	Parekh 2011 <sup>10</sup>	✓	✓	✓	Canada	1	N	N	All inpatients	Uncontrolled before-after study (prospective)	8 (3 before, 5 after)	2 (2.4)	1 (0.43)	RR=0.18 (0.02 to 1.97)†	0.16†	23
ICU transfers (per 1000 admissions)	Duns 2014 <sup>48</sup>	✓	✓	✓	Australia	1	Y	Y	All inpatients	Uncontrolled before-after study (prospective)	48 (24 before, 24 after)	350 (13.1)	364 (13.1)	RR=1.00 (0.88 to 1.16)†	0.98	7

Continued

Table 4 Continued

Outcome	First author, year	Intervention			Country	Number of centres	Specialist unit?	Excluding MET?	Study design	Study duration in months	Events (n, n (rate))	Events (n, n (rate))	Effect size (95% CI)	P value	Quality score (max:26)
		Implemented a new PTTT	Implemented a MET	Staff training/education											
Unplanned PICU transfers from ward (per 1000 non-ICU patient-days)	Bonafide 2014 <sup>46</sup>	✓	✓	✓	USA	1	Y	N	All inpatients interrupted time series study (prospective)	59 (32 before, 27 after)	874 (4,54)	936 (5,26)	IRR=0.73 (0.46 to 1.14)	0.16	23
Unplanned transfers to PICU from ward (per 1000 patient-days)	Agulnik 2017 <sup>47</sup>	✓	✓	✓	Guatemala	1	Y	N	Uncontrolled before-after study (retrospective)	24 (12 before, 12 after)	157 (9,3)	130 (6,5)	RR=0.70 (0.56 to 0.88)†	0.003	19
Urgent PICU admissions (per 1000 patient-days)	Pashuram 2018 <sup>48</sup>	✓	✓	✓	Belgium, Netherlands, England, Italy, Canada, New Zealand	21	Y	N	Cluster randomised trial (prospective)	18 (6 pre, 12 post)	Con: (4,01) Int: (3,48)	Con: (4,01) Int: (3,48)	RR=0.35 (0.62 to 1.09)	0.45	23
<b>PICU outcomes</b>															
Critical deterioration events after PICU transfer (per 1000 non-ICU patient-days)	Bonafide 2014 <sup>46</sup>	✓	✓	✓	USA	1	Y	N	Interrupted time series study (prospective)	59 (32 before, 27 after)	260† (1,35)	282† (1,58)	IRR=0.38 (0.20 to 0.75)	0.01	23
Mechanical ventilation within 1 hour of unplanned PICU transfer (per unplanned transfer to PICU)	Bonafide 2014 <sup>46</sup>	✓	✓	✓	USA	1	Y	N	Interrupted time series study (prospective)	59 (32 before, 27 after)	45 (5,1)	42 (4,3)	RR=0.87 (0.58 to 1.31)†	0.51	23
Mechanical ventilation within 12 hours of unplanned PICU transfer (per unplanned transfer to PICU)	Bonafide 2014 <sup>46</sup>	✓	✓	✓	USA	1	Y	N	Interrupted time series study (prospective)	59 (32 before, 27 after)	112 (12,8)	103 (11,0)	IRR=0.17 (0.07 to 0.44)	<0.001	23
Vasopressors within 1 hour of unplanned PICU transfer (per unplanned transfer to PICU)	Bonafide 2014 <sup>46</sup>	✓	✓	✓	USA	1	Y	N	Interrupted time series study (prospective)	59 (32 before, 27 after)	41 (4,7)	16 (1,7)	RR=0.36 (0.21 to 0.64)†	<0.001	23
Vasopressors within 12 hours of unplanned PICU transfer (per unplanned transfer to PICU)	Bonafide 2014 <sup>46</sup>	✓	✓	✓	USA	1	Y	N	Interrupted time series study (prospective)	59 (32 before, 27 after)	71 (8,1)	57 (6,1)	IRR=0.20 (0.06 to 0.62)	0.006	23
Invasive ventilation in PICU (per emergency PICU admission)	Sefton 2015 <sup>49</sup>	✓	✓	✓	UK	1	Y	N	Controlled before-after study (retrospective)	24 (12 before, 12 after)	118 (75,2)	104 (62,7)	RR=0.83 (0.72 to 0.97)†	0.002	16
Intubates in PICU (per emergency PICU admission)	Sefton 2015 <sup>49</sup>	✓	✓	✓	UK	1	Y	N	Controlled before-after study (retrospective)	24 (12 before, 12 after)	50 (31,8)	40 (24,1)	RR=0.76 (0.53 to 1.08)†	0.12	16
Intubation within 24 hours of PICU admission (per 1000 patient-days)	Agulnik 2017 <sup>47</sup>	✓	✓	✓	Guatemala	1	Y	N	Uncontrolled before-after study (retrospective)	24 (12 before, 12 after)	11 (0,65)	18 (0,90)	RR=1.38 (0.65 to 2.92)†	0.46	19
Vasopressors within 24 hours of PICU admission (per 1000 patient-days)	Agulnik 2017 <sup>47</sup>	✓	✓	✓	Guatemala	1	Y	N	Uncontrolled before-after study (retrospective)	24 (12 before, 12 after)	29 (1,72)	37 (1,86)	RR=1.08 (0.69 to 1.75)†	0.60	19
Mechanical ventilation during PICU admission (per PICU admission)	Kolovos 2018 <sup>48</sup>	✓	✓	✓	USA	1	Y	N	RRT activation criteria* Uncontrolled before-after study (retrospective)	78 (42 before, 36 after)	285 (25,88)	233 (22,09)	RR=0.85 (0.73 to 0.99)†	0.03†	12
Intubation within 1 hour of PICU admission (per PICU admission)	Kolovos 2018 <sup>48</sup>	✓	✓	✓	USA	1	Y	N	RRT activation criteria* Uncontrolled before-after study (retrospective)	78 (42 before, 36 after)	49 (4,47)	88 (8,94)	RR=1.87 (1.33 to 2.62)	0.0003	12
Significant critical deterioration events (per 1000 patient-days)	Pashuram 2018 <sup>48</sup>	✓	✓	✓	Belgium, Ireland, The Netherlands, England, Canada, New Zealand	21	Y	N	Cluster randomised trial (prospective)	18 (6 pre, 12 post)	Con: (1,44) Int: (0,89)	Con: (1,44) Int: (0,84)	RR=0.77 (0.61 to 0.97)	0.03	23

P, values in bold denote statistical significance (<0.05).  
 †Critical deterioration event is defined as transfer to the ICU followed by non-invasive or invasive mechanical ventilation or vasopressor infusion within 12 hours.  
 ‡Indicates a PTTT not described or validated in the published literature.  
 †Data calculated by research team, based on data presented in the journal article. All data calculated via <https://www.medcalc.org>.  
 Con, control group; ED, emergency department; ICU, intensive care unit; Int, intervention group; IRR, incident risk ratio; MET, medical emergency team; N/A, not available; PICU, paediatric intensive care unit; PTTT, paediatric track and trigger tool; RR, relative risk; RRT, rapid response team.

AUTHOR PROOF

rates of PICU mortality among ward transfers (RR=0.31, 95% CI 0.13 to 0.72),<sup>18</sup> and PICU mortality rates among patients readmitted within 48 hours (RR=0.43, 95% CI 0.17 to 0.99).<sup>63</sup> Six studies (including a high-quality cluster randomised trial and interrupted time series study) reported no postintervention change in PICU mortality using a variety of metrics.<sup>64–69</sup>

### Cardiac and respiratory arrests

Two uncontrolled before-after studies (both with RRT/MET) reported significant postintervention rate reductions in subcategories of cardiac arrests: one in 'near cardiopulmonary arrests'<sup>63</sup> (RR=0.54, 95% CI 0.52 to 0.57) but not 'actual cardiopulmonary arrests' and one in 'preventable cardiac arrests'<sup>62</sup> (RR=0.45, 95% CI 0.20 to 0.97) but not 'unexpected cardiac arrests'. One uncontrolled before-after study (with RRT/MET) reported a significant postintervention reduction in rates of ward respiratory arrests per 1000 patient-days<sup>16</sup> (RR=0.27, 95% CI 0.07 to 0.95). Seven studies (including one high-quality cluster randomised trial and one high-quality interrupted time series study) found no change in cardiac arrest rates using a variety of metrics<sup>13 15 16 61 64 65</sup> or cardiac and respiratory arrests combined.<sup>60</sup>

### Calls for urgent review/assistance

Two uncontrolled before-after studies (all with RRT/MET) reported significant postintervention reductions in rates of code calls<sup>17 63</sup> (RR=0.29, 95% CI 0.10 to 0.65; RR=0.71, 95% CI 0.61 to 0.83) while three studies found no change in rates of code calls.<sup>15 18 72</sup> One uncontrolled before-after study in a community hospital (without RRT/MET) found significant postintervention reductions in rates of urgent calls to the in-house paediatrician (RR=0.23, 95% CI 0.11 to 0.46) and respiratory therapist<sup>70</sup> (RR=0.36, 95% CI 0.13 to 0.95). Two uncontrolled before-after studies (with RRT/MET) found increases in rates of RRT calls<sup>72</sup> (RR=1.59, 95% CI 0.33 to 1.90) and outreach team calls<sup>66</sup> (RR=1.92, 95% CI 1.79 to 2.07). One study found no change in rates of RRT calls.<sup>71</sup>

### PICU transfers

One uncontrolled before-after study (without RRT/MET) found a significant postintervention decrease in the rate of unplanned PICU transfers per 1000 patient-days<sup>67</sup> (RR=0.70, 95% CI 0.56 to 0.88). Four studies (including one high-quality cluster randomised trial and one high-quality interrupted time series study) found no change in rates of PICU admissions postintervention.<sup>64–66 70</sup>

### PICU outcomes

Two studies, one interrupted time series and one multi-centre cluster randomised trial (both with RRT/MET), found significant reductions in rates of 'critical deterioration events' (life-sustaining interventions administered within 12 hours of PICU admission) relative to preimplementation trends and relative to control hospitals, respectively (IRR=0.38, 95% CI 0.20 to 0.75; OR=0.77, 95% CI 0.61 to 0.97).<sup>64 65</sup> One controlled before-after

study (without RRT/MET) reported a significant reduction in rates of invasive ventilation given to emergency PICU admissions postintervention (RR=0.83, 95% CI 0.72 to 0.97), with no significant change observed in a control group of patients admitted to PICU from outside of the hospital.<sup>68</sup> One uncontrolled before-after study reported a significant postintervention decrease in rates of PICU admissions receiving mechanical ventilation (RR=0.85, 95% CI 0.73 to 0.99), but an increase in rates of early intubation (RR=1.87, 95% CI 1.33 to 2.62).<sup>69</sup>

### Implementation outcomes

Only three studies reported outcomes relating to the quality of implementation of the intervention. One study reported 99% of audited observation sets of the Bedside PEWS had at least five vital signs present postintervention, up from 76% preintervention (no change in control hospitals).<sup>64</sup> A previous study of the same PTTT reported 3% of audited cases had used the incorrect age chart but reported an intraclass coefficient of 0.90 for agreement between bedside nurses scoring the PTTT in practice and research nurses retrospectively assigned scores.<sup>70</sup> Finally, error rates in C-CHEWS scoring were reported to have reduced from an initial 47% to below 10% by the end of the study.<sup>67</sup>

## DISCUSSION

This paper reviewed the published PTTT and early warning system literature in order to assess the validity of PTTT for predicting inpatient deterioration (question 1) and the effectiveness of early warning system interventions (with or without PTTT) for reducing mortality and morbidity outcomes in hospitalised children (question 2). We believe that the consideration of broader 'early warning systems' differentiates this paper from previous reviews, as does the inclusion of two recently published high-quality effectiveness studies.<sup>64 73</sup>

### How well validated are existing tools for predicting inpatient deterioration?

Given a growing understanding and emphasis on the importance of local context in healthcare interventions, it is perhaps not surprising that such a wide range of PTTT have been developed and evaluated internationally, and modifications to existing PTTT are common. The result, however, is that a large number of different PTTT have been narrowly validated, but none has been broadly validated across a variety of different settings and populations. With only one exception,<sup>44</sup> all studies evaluating the validity of PTTT have been single-centre reports from specialist units, greatly limiting the generalisability of the findings.

PTTT such as the Bedside PEWS, C&VPEWS, NHS III PEWS and C-CHEWS have demonstrated very good (AUROC  $\geq$  0.80) or excellent (AUROC  $\geq$  0.90) diagnostic accuracy, typically for predicting PICU transfers, in internal and external validation studies.<sup>11 14 19 29 32 37 42 44</sup>

However, methodological issues common to the validation studies mean that such results need to be interpreted with a degree of caution. First, each of the studies was conducted in a clinical setting where paediatric inpatients are subject to various forms of routine clinical intervention throughout their admission. There are numerous statistical modelling techniques which can account for co-occurrence of clinical interventions and the longitudinal nature of the predictors,<sup>74 75</sup> but none of these were used in the validation studies and so estimates of predictive ability are likely to be distorted. Indeed, the majority of outcomes used in the validation studies are clinical interventions themselves (eg, PICU transfer). Second, while it is understandable that a majority of studies ‘bench-tested’ the PTTT rather than implement it into practice before evaluation, the process of abstracting PTTT scores retrospectively from patient charts and medical records introduces a number of sources of potential bias or inaccuracy. For instance, several studies reported either high levels of missing data (ie, some of the observations required to populate the PTTT score being evaluated were not routinely collected or recorded and so were scored as ‘normal’)<sup>11 19 32 44 45</sup> or difficulty in abstracting certain descriptive or subjective PTTT components.<sup>19 28 41 49</sup> Assuming missing values are normal, or excluding some PTTT items for analysis are both likely to result in underscoring of the PTTT and skew the results. Finally, studies which evaluated a PTTT that had been implemented in practice are at risk of overestimating the ability of PTTT to predict proxy outcomes such as PICU transfer, inasmuch as high PTTT scores or triggers automatically direct staff towards escalation of care, or clinical actions which make escalation of care more likely.

The findings reported in several PTTT studies point towards two potential challenges for some centres in implementing and sustaining a PTTT in clinical practice. As noted above, a number of studies that retrospectively ‘bench-tested’ a PTTT reported that the observations that were required to score the tool were not always routinely collected or recorded in their centre. It may be that the introduction of a PTTT into practice would help create a framework to ensure that core vital signs and observations were collected more routinely (as demonstrated by Parshuram *et al*<sup>64</sup>), but this would obviously have resource implications that could be a potential barrier for some centres. Such considerations are important, as evidence from the adult literature points to the potential for tools to inadvertently mask deterioration when core observations are missing.<sup>76</sup> Second, PPV values reported in cohort studies, and case-control studies that adjusted for outcome prevalence, were uniformly low (between 2.3% and 5.9%).<sup>14 19 31–33 47</sup> They demonstrate that even PTTT which demonstrate good predictive performance are likely to generate a large amount of ‘false alarms’ because adverse outcomes are so rare. For some centres, these issues may be mitigated to some extent by dedicated response teams or other available resources, but other

hospitals may not be able to sustain the increased workload of responding to PTTT triggers.

### How effective are early warning systems for reducing mortality and morbidity?

We found limited evidence for early warning system interventions reducing mortality or arrest rates in hospitalised children. While some effectiveness papers did report significant reductions in rates of mortality (on the ward or in PICU) or cardiac arrests after implementation of different early warning system interventions,<sup>16–18 62 63</sup> they were all uncontrolled before-after studies which have inherent limitations in terms of establishing causality. They do not preclude the possibility that outcome rates would have improved over time regardless of the intervention<sup>77</sup> or changes were caused by other factors, and their inclusion is accordingly discouraged by some Cochrane review groups.<sup>78</sup> Three high-quality multicentre studies—two interrupted time series studies and a recent cluster randomised trial—found no changes in rates or trends of mortality or arrests postintervention.<sup>64 65 73</sup>

There was also limited evidence for early warning systems reducing PICU transfers or calls for urgent review. Again, a small number of uncontrolled before-after studies reported significant reductions postintervention,<sup>15 17 63</sup> but several other studies reported significant increases in transfers or calls for review<sup>66 72</sup> or no postintervention changes. We did find moderate evidence across four studies—including a controlled before-after study, a multicentre interrupted time series study and a multicentre cluster randomised trial—for early warning system interventions reducing rates of early critical interventions in children transferred to PICU.<sup>64 65 68 69</sup> Such results are promising, but corresponding reductions in hospital or PICU mortality rates have not yet been reported.

Implementing complex interventions in a health-care setting is challenging and evidence from the adult literature points to challenges and barriers to successfully implement TTT in practice.<sup>79–81</sup> However, given so few effectiveness studies reported on implementation outcomes, it is difficult to know whether negative findings reflect poor effectiveness or implementation of early warning systems. Again, effectiveness studies were predominantly carried out in specialist centres—and in all but three cases,<sup>67 68 70</sup> involved the use of a dedicated response team—which greatly limits the generalisability of findings outside of these contexts.

### Limitations of the review

There are several limitations of the current review. First, despite purposely widening the scope of the effectiveness review question to include paediatric ‘early warning systems’ with or without a PTTT, we identified very few studies that did not employ a PTTT as part of the intervention. In part, this likely reflects the fact that PTTT have become almost synonymous with early warning systems, but it is also possible that our search strategy may have missed some broader early warning system initiatives that



were not explicitly labelled as such. Second, our inclusion criteria for study selection were deliberately broad and so resulted in our including several validation and effectiveness studies that were subsequently excluded from analysis due to insufficient statistical detail or methodological issues. Third, the scope of the current review was limited to consideration of quantitative validation and effectiveness studies. We are mindful of research suggesting that implementing PTTT in practice may confer secondary benefits including, but not limited to improvements in communication, teamwork and empowerment of junior staff to call for assistance.<sup>82–84</sup> Finally, we opted not to conduct a meta-analysis of effectiveness findings due to the heterogeneity of outcome metrics, interventions and study designs, populations and settings. Given the large sample sizes required to detect changes in rare adverse events, we believe further work is needed to harmonise outcome measures used to evaluate early warning system interventions internationally, in order to facilitate pooling of findings across studies.

## CONCLUSION

The PTTT literature is currently characterised by an ‘absence of evidence’ rather than an ‘evidence of absence’. PTTT seem like a logical tool for helping staff detect and respond to deteriorating patients, but the existing evidence base is too limited to form clear judgements of their utility. We would argue that there has been too much confidence placed in the statistical findings of validation studies of PTTT, given methodological limitations in the study designs. There is evidence of consistently high false-alarm rates and bench-testing studies point to many PTTT parameters not being reliably recorded in practice: as such there is reason for caution in considering the viability of PTTT for all hospitals. Almost all of the early warning systems and PTTT reported in the literature have been developed and evaluated in specialist centres, typically in units with access to dedicated response teams—yet PTTT appear to be commonly adopted by non-specialist units with little modification. There is currently limited evidence that ‘early warning systems’ incorporating a PTTT reduce deterioration or death in practice. As such, we would urge caution among policymakers in calling for their use to become mandatory across all hospitals. We acknowledge the potential for PTTT to confer a range of secondary benefits in areas such as communication, teamwork and empowerment of junior staff. More work is required to understand the wider impact of PTTT implementation in different clinical settings before it is possible to evaluate their overall contribution to the wider safety mechanisms and systems aimed at identifying and responding to deteriorating in paediatric patients.

## Author affiliations

<sup>1</sup>Centre for Trials Research, Cardiff University, Cardiff, UK

<sup>2</sup>School of Healthcare Sciences, Cardiff University, Cardiff, UK

<sup>3</sup>Department of Paediatrics, Morriston Hospital, Swansea, UK

<sup>4</sup>Wirral University Teaching Hospital, Wirral, UK

<sup>5</sup>University Library Services, Cardiff University, Cardiff, UK

<sup>6</sup>Public Health Wales, Cardiff, UK

<sup>7</sup>Department of Paediatric Intensive Care, Noah's Ark Children's Hospital for Wales, Cardiff, UK

<sup>8</sup>Emergency Department, Paediatric Emergency Medicine Leicester Academic (PEMLA) Group, Leicester, UK

<sup>9</sup>Alder Hey Children's NHS Foundation Trust, Liverpool, UK

<sup>10</sup>Faculty of Health and Applied Sciences (HAS), University of the West of England Bristol, Bristol, UK

<sup>11</sup>Paediatrics, Children's Hospital for Wales, Cardiff, UK

<sup>12</sup>Department of Pediatric Emergency Medicine, Sidra Medical and Research Center, Doha, Qatar

**Acknowledgements** The authors would like to acknowledge the contribution of Dr James Bunn to the review.

**Contributors** RT: screening and review of papers, contribution to design of work, preparation of manuscript; CH: screening and review of papers, contribution to concept and design of work, review of manuscript; FVL-W: contribution to design of work, screening and review of papers, review of manuscript; KH: contribution to concept and design of work, screening and review of papers, review of manuscript; CP, DR, BM, AO, DE, RS, GS, DL, LT, DA, AL, ET-J: contribution to concept and design of work, screening and review of papers, review of manuscript; MM: information specialist, review of manuscript.

**Funding** This study is funded by the National Institute for Health Research (NIHR) Health Services and Delivery Research (HS&DR) programme (12/178/17).

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

## REFERENCES

1. Pearson GA, (ed). *Why Children Die: A Pilot Study 2006*. England (South West, North East and West Midlands), Wales and Northern Ireland, 2008.
2. Joint Commission. National Patient Safety Goals, Requirements and Rationales. *Jt Comm Perspect Patient Saf* 2009;2009:8–12.
3. Tume L. The deterioration of children in ward areas in a specialist children's hospital. *Nurs Crit Care* 2007;12:12–19.
4. NICE. Acutely ill patients in hospital: Recognition of and response to acute illness in adults in hospital. *NICE Clin Guidel* 2007;50:1–107.
5. Roland D, Oliver A, Edwards ED, et al. Use of paediatric early warning systems in Great Britain: has there been a change of practice in the last 7 years? *Arch Dis Child* 2014;99:26–9.
6. Chapman SM, Wray J, Oulton K, et al. Systematic review of paediatric track and trigger systems for hospitalised children. *Resuscitation* 2016;109:87–109.
7. Lambert V, Matthews A, MacDonell R, et al. Paediatric early warning systems for detecting and responding to clinical deterioration in children: a systematic review. *BMJ Open* 2017;7:e014497.
8. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
9. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;52:377–84.
10. Monaghan A. Detecting and managing deterioration in children. *Paediatr Nurs* 2005;17:32–5.
11. Parshuram CS, Hutchison J, Middaugh K. Development and initial validation of the Bedside Paediatric Early Warning System score. *Crit Care* 2009;13:R135.



12. Haines C, Perrott M, Weir P. Promoting care for acutely ill children-development and evaluation of a paediatric early warning tool. *Intensive Crit Care Nurs* 2006;22:73–81.
13. Tibballs J, Kinney S, Duke T, *et al.* Reduction of paediatric in-patient cardiac arrest and death with a medical emergency team: preliminary results. *Arch Dis Child* 2005;90:1148–52.
14. Mason BW, Edwards ED, Oliver A, *et al.* Cohort study to test the predictability of the NHS Institute for Innovation and Improvement Paediatric Early Warning System. *Arch Dis Child* 2016;101:552–5.
15. Brilli RJ, Gibson R, Luria JW, *et al.* Implementation of a medical emergency team in a large pediatric teaching hospital prevents respiratory and cardiopulmonary arrests outside the intensive care unit. *Pediatr Crit Care Med* 2007;8:236–46.
16. Hunt EA, Zimmer KP, Rinke ML, *et al.* Transition from a traditional code team to a medical emergency team and categorization of cardiopulmonary arrests in a children's center. *Arch Pediatr Adolesc Med* 2008;162:117–22.
17. Sharek PJ, Parast LM, Leong K, *et al.* Effect of a rapid response team on hospital-wide mortality and code rates outside the ICU in a Children's Hospital. *JAMA* 2007;298:2267–74.
18. Anwar-ul-Haque, Saleem AF, Zaidi S, *et al.* Experience of pediatric rapid response team in a tertiary care hospital in Pakistan. *Indian J Pediatr* 2010;77:273–6.
19. Zhai H, Brady P, Li Q, *et al.* Developing and evaluating a machine learning based algorithm to predict the need of pediatric intensive care unit transfer for newly hospitalized children. *Resuscitation* 2014;85:1065–71.
20. Garlick J, Furlong-Dillard J, Alexander M, *et al.* 679: Modified pediatric early warning scores, a predictor of appropriate transfer to higher level of care. *Crit Care Med* 2013;41:A166–A167.
21. Medar S, Weingarten-Arams J, Katyal C. 907: Pediatric early warning score (PEWS) is a reliable predictor of rapid response team (RRT) call. *Crit Care Med* 2015;43:228.
22. Bell D, Mac A, Ochoa Y, *et al.* The Texas Children's Hospital Pediatric Advanced Warning Score as a predictor of clinical deterioration in hospitalized infants and children: a modification of the PEWS tool. *J Pediatr Nurs* 2013;28:e2–e9.
23. McLellan MC, Connor JA. The Cardiac Children's Hospital Early Warning Score (C-CHEWS). *J Pediatr Nurs* 2013;28:171–8.
24. Rahman ZH, Leahy NE, Sessler K, *et al.* Application of the burn center Pediatric Early Warning Score (PEWS) tool to clinical practice: A pilot validation study. *J Burn Care Res* 2014;35:S187.
25. Hopkins M, Rowan C, Rigby M, *et al.* 684: PEWS score predicts the need for PICU transfer and critical care interventions. *Crit Care Med* 2013;41:A168.
26. Gawronski O, Bertaina A, Broccati F, *et al.* Bedside pews in an Italian pediatric bone marrow transplant unit: Preliminary results of a longitudinal retrospective chart review. *Intensive Care Med* 2013;39:S130.
27. Sefton G, Lane S, Carrol ED. Performance of individual predictors of deterioration used in a paediatric early warning score. *Pediatr Crit Care Med* 2014;15:29–30.
28. Robson MA, Cooper CL, Medicus LA, *et al.* Comparison of three acute care pediatric early warning scoring tools. *J Pediatr Nurs* 2013;28:e33–e41.
29. Chapman SM, Wray J, Oulton K, *et al.* 'The Score Matters': wide variations in predictive performance of 18 paediatric track and trigger systems. *Arch Dis Child* 2017;102:487–95.
30. Fuijkschot J, Vernhout B, Lemson J, *et al.* Validation of a paediatric early warning score: first results and implications of usage. *Eur J Pediatr* 2015;174:15–21.
31. Tucker KM, Brewer TL, Baker RB, *et al.* Prospective evaluation of a pediatric inpatient early warning scoring system. *J Spec Pediatr Nurs* 2009;14:79–85.
32. Edwards ED, Powell CV, Mason BW, *et al.* Prospective cohort study to test the predictability of the Cardiff and Vale paediatric early warning system. *Arch Dis Child* 2009;94:602–6.
33. Edwards ED, Mason BW, Oliver A, *et al.* Cohort study to test the predictability of the Melbourne criteria for activation of the medical emergency team. *Arch Dis Child* 2011;96:174–9.
34. O'Loughlin K, Ruparella K, Vince T, *et al.* The effectiveness of the paediatric early warning tool (PEWT) in identifying children requiring admission to a critical care unit. *Arch Dis Child* 2012;97:A154.
35. Wright D, Sefton G, Horan M. Charted observations and the use of a paediatric early warning tool did not predict the majority of cardio-respiratory arrest calls in a paediatric hospital. *Arch Dis Child* 2011;96:A59.
36. Ahmed M, Sobithadevi D, Lall R, *et al.* 1483 Burton paediatric early warning system score. *Arch Dis Child* 2012;97:A421.
37. McLellan MC, Gauvreau K, Connor JA. Validation of the cardiac children's hospital early warning score: an early warning scoring tool to prevent cardiopulmonary arrests in children with heart disease. *Congenit Heart Dis* 2014;9:194–202.
38. Clayton R, Selby A, Tume L. Is a paediatric early warning tool (PEWT) useful in cardiac patients? *Pediatr Crit Care Med* 2014;15:35–6.
39. Fenix JB, Gillespie CW, Levin A, *et al.* Comparison of Pediatric Early Warning Score to Physician Opinion for Deteriorating Patients. *Hosp Pediatr* 2015;5:474–9.
40. Mandell IM, Bynum F, Marshall L, *et al.* Pediatric Early Warning Score and unplanned readmission to the pediatric intensive care unit. *J Crit Care* 2015;30:1090–5.
41. Agulnik A, Forbes PW, Stenquist N, *et al.* Validation of a Pediatric Early Warning Score in Hospitalized Pediatric Oncology and Hematopoietic Stem Cell Transplant Patients. *Pediatr Crit Care Med* 2016;17:e146–e153.
42. Agulnik A, Méndez Aceituno A, Mora Robles LN, *et al.* Validation of a pediatric early warning system for hospitalized pediatric oncology patients in a resource-limited setting. *Cancer* 2017;123:4903–13.
43. Blackstone B, Sharmila A, Pienaar A, *et al.* G453(P) 'Between the flags' paediatric early warning system is a sensitive tool to detect deterioration in children from district general hospitals admitted to paediatric intensive care. *Arch Dis Child* 2017;102:A178 LP–A179.
44. Parshuram CS, Duncan HP, Joffe AR, *et al.* Multicentre validation of the bedside paediatric early warning system score: a severity of illness score to detect evolving critical illness in hospitalised children. *Crit Care* 2011;15:R184.
45. Akre M, Finkelstein M, Erickson M, *et al.* Sensitivity of the pediatric early warning score to identify patient deterioration. *Pediatrics* 2010;125:e763–e769.
46. Gawronski O, Ciofi Degli Atti ML, Di Ciommo V, *et al.* Accuracy of Bedside Paediatric Early Warning System (BedsidePEWS) in a Pediatric Stem Cell Transplant Unit. *J Pediatr Oncol Nurs* 2016;33:249–56.
47. Duncan H, Hutchison J, Parshuram CS. The pediatric early warning system score: a severity of illness score to predict urgent medical need in hospitalized children. *J Crit Care* 2006;21:271–8.
48. Skaletzky SM, Raszynski A, Totapally BR. Validation of a modified pediatric early warning system score: a retrospective case-control study. *Clin Pediatr* 2012;51:431–5.
49. Ross C, Harrysson I, Goel V, *et al.* 151: Sensitivity and specificity of a pediatric early warning system using data-driven vital signs. *Crit Care Med* 2015;43:39.
50. McLellan M, Gauvreau K, Connor JA. Validation of the children's hospital early warning scoring system for identifying hospitalized children at risk for arrest or ICU transfer. *Pediatr Crit Care Med* 2014;15:30.
51. Mistry KP, Turi J, Hueckel R, *et al.* Pediatric Rapid Response Teams in the Academic Medical Center. *Clin Pediatr Emerg Med* 2006;7:241–7.
52. Demmel KM, Williams L, Flesch L. Implementation of the pediatric early warning scoring system on a pediatric hematology/oncology unit. *J Pediatr Oncol Nurs* 2010;27:229–40.
53. Sandhu S, Ralph T, Bradbury K, *et al.* 726 The Paediatric Early Warning Tool in Combination with a Nurse-Led Outreach Team: Impact on Outcomes for Critically Ill Children. *Pediatr Res* 2010;68:368.
54. Randhawa S, Roberts-Turner R, Woronick K, *et al.* Implementing and sustaining evidence-based nursing practice to reduce pediatric cardiopulmonary arrest. *West J Nurs Res* 2011;33:443–56.
55. Camacho EA, Holman D, Badgett V, *et al.* Modification of a Pediatric Early Warning Tool for Use With Patients in a Specialized Cardiac/Renal Unit. *J Pediatr Nurs* 2011;26:e2–e3.
56. Heyden MV, Mudge B, Casella S. 593: Initiation of a dedicated pediatric rapid response team reduces cardiopulmonary arrests outside of the intensive care unit in a small children's hospital. *Crit Care Med* 2012;40:1–328.
57. Norville R, Wills-Bagnato P, Staton S, *et al.* Improvement in early recognition of deteriorating pediatric bone marrow transplant patients. *Biol Blood Marrow Transplant* 2013;19:S173.
58. Ambati SR, Sweberg T, Silver P, *et al.* 297: Effect of simulation based curriculum on the utilization of rapid response team activations (RRT). *Crit Care Med* 2014;42:A1432.
59. Ochoi T, Downing E, Rzeskiewicz A, *et al.* G204(P) Impact of the Bedside PEWS on early detection of deteriorating patients. *Arch Dis Child* 2014;99:A89–A90.
60. Zenker P, Schlesinger A, Hauck M, *et al.* Implementation and impact of a rapid response team in a children's hospital. *Jt Comm J Qual Patient Saf* 2007;33:418–25.
61. Hanson CC, Randolph GD, Erickson JA, *et al.* A reduction in cardiac arrests and duration of clinical instability after implementation of a paediatric rapid response system. *Postgrad Med J* 2010;86:314–8.

62. Tibballs J, Kinney S. Reduction of hospital mortality and of preventable cardiac arrest and death on introduction of a pediatric medical emergency team. *Pediatr Crit Care Med* 2009;10:306–12.
63. Kotsakis A, Lobos AT, Parshuram C, et al. Implementation of a multicenter rapid response system in pediatric academic hospitals is effective. *Pediatrics* 2011;128:72–8.
64. Parshuram CS, Dryden-Palmer K, Farrell C, et al. Effect of a pediatric early warning system on all-cause mortality in hospitalized pediatric patients: the EPOCH randomized clinical trial. *JAMA* 2018;319:1002–12.
65. Bonafide CP, Localio AR, Roberts KE, et al. Impact of rapid response system implementation on critical deterioration events in children. *JAMA Pediatr* 2014;168:25.
66. Duns N, Ceely B, Festa M, et al. Introduction of a track and trigger system is associated with increased paediatric intensive care outreach utilisation and a trend towards improved patient outcomes in a tertiary children's hospital. *Aust Crit Care* 2014;27:61.
67. Agulnik A, Mora Robles LN, Forbes PW, et al. Improved outcomes after successful implementation of a pediatric early warning system (PEWS) in a resource-limited pediatric oncology hospital. *Cancer* 2017;123:2965–74.
68. Sefton G, McGrath C, Tume L, et al. What impact did a Paediatric Early Warning system have on emergency admissions to the paediatric intensive care unit? An observational cohort study. *Intensive Crit Care Nurs* 2015;31:91–9.
69. Kolovos NS, Gill J, Michelson PH, et al. Reduction in Mortality Following Pediatric Rapid Response Team Implementation. *Pediatr Crit Care Med* 2018;19:477–82.
70. Parshuram CS, Bayliss A, Reimer J, et al. Implementing the Bedside Paediatric Early Warning System in a community hospital: A prospective observational study. *Paediatr Child Health* 2011;16:e18–e22.
71. Panesar R, Polikoff LA, Harris D, et al. Characteristics and outcomes of pediatric rapid response teams before and after mandatory triggering by an elevated Pediatric Early Warning System (PEWS) score. *Hosp Pediatr* 2014;4:135–40.
72. Douglas K, Collado JC, Keller S. Implementation of a pediatric early warning scoring system at an academic medical center. *Crit Care Nurs Q* 2016;39:363–70.
73. Kutty S, Jones PG, Karels Q, et al. Association of pediatric medical emergency teams with hospital mortality. *Circulation* 2018;137:38–46.
74. Guerra MW, Shults J, Amsterdam J, et al. The analysis of binary longitudinal data with time-dependent covariates. *Stat Med* 2012;31:931–48.
75. Selig JP, Preacher KJ, Little TD. Modeling time-dependent association in longitudinal data: a lag as moderator approach. *Multivariate Behav Res* 2012;47:697–716.
76. Clifton DA, Clifton L, Sandu DM, et al. 'Errors' and omissions in paper-based early warning scores: the association with changes in vital signs—a database analysis. *BMJ Open* 2015;5:e007376.
77. Joffe AR, Anton NR, Burkholder SC. Reduction in hospital mortality over time in a hospital without a pediatric medical emergency team: limitations of before-and-after study designs. *Arch Pediatr Adolesc Med* 2011;165:419–23.
78. Goodacre S. Uncontrolled before-after studies: discouraged by Cochrane and the EMJ. *Emerg Med J* 2015;32:507–8.
79. Niessch M, Fabritius ML, Anhøj J. Imperfect implementation of an early warning scoring system in a Danish teaching hospital: a cross-sectional study. *PLoS One* 2013;8:e70068.
80. Hands C, Reid E, Meredith P, et al. Patterns in the recording of vital signs and early warning scores: compliance with a clinical escalation protocol. *BMJ Qual Saf* 2013;22:719–26.
81. Lydon S, Byrne D, Offiah G, et al. A mixed-methods investigation of health professionals' perceptions of a physiological track and trigger system. *BMJ Qual* 2015:688–95.
82. Bonafide CP, Roberts KE, Weirich CM, et al. Beyond statistical prediction: qualitative evaluation of the mechanisms by which pediatric early warning scores impact patient safety. *J Hosp Med* 2013;8:248–53.
83. Ennis L. Paediatric early warning scores on a children's ward: a quality improvement initiative. *Nurs Child Young People* 2014;26:25–31.
84. Kaul M, Snethen J, Kelber ST, et al. Implementation of the Bedside Paediatric Early Warning System (BedsidePEWS) for nurse identification of deteriorating patients. *J Spec Pediatr Nurs* 2014;19:339–49.