

# The Domination Parameters of Cubic Graphs

Igor E. Zverovich<sup>\*†</sup>

RUTCOR – Rutgers Center for Operations Research

Rutgers, The State University of New Jersey

Piscataway, NJ 08854-8003

USA

Vadim E. Zverovich<sup>\*</sup>

Department of Mathematical Sciences

Brunel University, Uxbridge

Middlesex, UB8 3PH

UK

Vadim.Zverovich@brunel.ac.uk

January 27, 2005

## Abstract

Let  $ir(G)$ ,  $\gamma(G)$ ,  $i(G)$ ,  $\beta_0(G)$ ,  $\Gamma(G)$  and  $IR(G)$  be the irredundance number, the domination number, the independent domination number, the independence number, the upper domination number and the upper irredundance number of a graph  $G$ , respectively. In this paper we show that for any nonnegative integers  $k_1, k_2, k_3, k_4, k_5$  there exists a cubic graph  $G$  satisfying the following conditions:  $\gamma(G) - ir(G) \geq k_1$ ,  $i(G) - \gamma(G) \geq k_2$ ,  $\beta_0(G) - i(G) \geq k_3$ ,  $\Gamma(G) - \beta_0(G) \geq k_4$ , and  $IR(G) - \Gamma(G) \geq k_5$ . This result settles a problem posed in [9].

## 1 Introduction and Main Result

All graphs will be finite and undirected without multiple edges. If  $G$  is a graph,  $V(G)$  denotes the set, and  $|G|$  the number, of vertices in  $G$ . Let  $N(x)$  denote the neighborhood of a vertex  $x$ , and let  $\langle X \rangle$  denote the subgraph of  $G$  induced by  $X \subseteq V(G)$ . Also let  $N(X) = \cup_{x \in X} N(x)$  and  $N[X] = N(X) \cup X$ .

A set  $I \subseteq V(G)$  is called *independent* if no two vertices of  $I$  are adjacent. A set  $X$  is called a *dominating set* if  $N[X] = V(G)$ . An *independent dominating set* is a vertex subset that is both independent and dominating, or equivalently, is maximal independent.

---

<sup>\*</sup>Supported by the INTAS and the Belarus Government (Project INTAS-BELARUS 97-0093).

<sup>†</sup>Supported by RUTCOR.

The *independence number*  $\beta_0(G)$  is the maximum cardinality of a (maximal) independent set of  $G$ , and the *independent domination number*  $i(G)$  is the minimum cardinality taken over all maximal independent sets of  $G$ . The *domination number*  $\gamma(G)$  is the minimum cardinality of a (minimal) dominating set of  $G$ , and the *upper domination number*  $\Gamma(G)$  is the maximum cardinality taken over all minimal dominating sets of  $G$ . For  $x \in X$ , the set

$$PN(x, X) = PN(x) = N[x] - N[X - \{x\}]$$

is called the *private neighborhood* of  $x$ . If  $PN(x, X) = \emptyset$ , then  $x$  is said to be *redundant* in  $X$ . A set  $X$  containing no redundant vertex is called *irredundant*. The *irredundance number*  $ir(G)$  is the minimum cardinality taken over all maximal irredundant sets of  $G$ , and the *upper irredundance number*  $IR(G)$  is the maximum cardinality of a (maximal) irredundant set of  $G$ . An *ir-set*  $X$  of  $G$  is a maximal irredundant set of cardinality  $ir(G)$ . A  $\gamma$ -set, an  $i$ -set, a  $\beta_0$ -set, a  $\Gamma$ -set and an  $IR$ -set are defined analogously.

The following relationship among the parameters under consideration is well-known [2, 3]:

$$ir(G) \leq \gamma(G) \leq i(G) \leq \beta_0(G) \leq \Gamma(G) \leq IR(G).$$

The above and related parameters for regular graphs were investigated by many authors [1],[4]–[17]. For example, Cockayne and Mynhardt [4] and independently Rautenbach [15] disproved the Henning-Slater conjecture [12] that  $\Gamma(G) = IR(G)$  for any cubic graph  $G$ , while the Barefoot-Harary-Jones conjecture on the difference between the domination and independent domination numbers of cubic graphs was investigated in [5, 13, 14, 17].

In this paper, we deal with the next problem:

**Problem 1** ([9]) *Does there exist a cubic graph for which  $ir < \gamma < i < \beta_0 < \Gamma < IR$ ?*

We define the graph  $W_k$  ( $k \geq 0$ ) as follows. Take a disjoint union of the graphs

$$F_1 \cong F_2 \cong \dots \cong F_{2k+8}, G_1 \cong G_2 \cong \dots \cong G_{2k+6}, H_1 \cong H_2 \cong \dots \cong H_{3k+6},$$

where  $F_i, G_i$  and  $H_i$  are shown in Figure 1, and add the edges

$$\begin{aligned} &\{f'_i f_{i+1} : 1 \leq i \leq 2k + 7\}, f'_{2k+8} g_1, \\ &\{g'_i g_{i+1} : 1 \leq i \leq 2k + 5\}, g'_{2k+6} h_1, \\ &\{h'_i h_{i+1} : 1 \leq i \leq 3k + 5\}, h'_{3k+6} f_1. \end{aligned}$$

**Theorem 1** *For any nonnegative integers  $k_1, k_2, k_3, k_4, k_5$  there exists an integer  $k$  such that the cubic graph  $W_k$  satisfies the following conditions:  $\gamma(W_k) - ir(W_k) \geq k_1$ ,  $i(W_k) - \gamma(W_k) \geq k_2$ ,  $\beta_0(W_k) - i(W_k) \geq k_3$ ,  $\Gamma(W_k) - \beta_0(W_k) \geq k_4$ , and  $IR(W_k) - \Gamma(W_k) \geq k_5$ .*

It follows from Lemmas 1–5 of Section 2 that the graph  $W_0$  has the property

$$ir < \gamma < i < \beta_0 < \Gamma < IR,$$

thus solving Problem 1.

We conclude this section with the next conjecture.

**Conjecture 1** *For any integers  $k_1, k_2, k_3, k_4, k_5$  there exists a 3-connected cubic graph  $G$  satisfying the following conditions:  $\gamma(G) - ir(G) \geq k_1$ ,  $i(G) - \gamma(G) \geq k_2$ ,  $\beta_0(G) - i(G) \geq k_3$ ,  $\Gamma(G) - \beta_0(G) \geq k_4$ , and  $IR(G) - \Gamma(G) \geq k_5$ .*




Figure 1. Graphs  $F_i$ ,  $G_i$ , and  $H_i$ .

## 2 Proof of Theorem 1

The proof of Theorem 1 is based on five lemmas. Let us denote by  $F, G$  and  $H$  the graphs induced by the sets  $\cup_{i=1}^{2k+8} V(F_i)$ ,  $\cup_{i=1}^{2k+6} V(G_i)$ , and  $\cup_{i=1}^{3k+6} V(H_i)$ , respectively.

**Lemma 1**  $\gamma(W_k) - ir(W_k) \geq k + 1$ .

**Proof:** Let  $D$  denote a  $\gamma$ -set of  $W_k$ . It is straightforward to check that  $|D \cap V(G_i)| = 4$  whenever both  $g_i$  and  $g'_i$  are dominated by  $D - V(G_i)$ , and  $|D \cap V(G_i)| = 5$  otherwise. Moreover, if  $|D \cap V(G_i)| = 4$ , then  $g_i, g'_i \notin D$ . Thus, the number of components  $G_i$  satisfying  $|D \cap V(G_i)| = 4$  is at most  $k + 3$ . We obtain

$$|D \cap V(G)| \geq 4(k + 3) + 5(k + 3) = 9k + 27.$$

Consider the set  $J = (D - V(G)) \cup R$ , where

$$R = \{N(g_i) \cap V(G_i), N(g'_i) \cap V(G_i) : 1 \leq i \leq 2k + 6\}.$$

We have

$$|R| = 8k + 24.$$

Let us construct a maximal irredundant set of  $W_k$ . We first put  $J' = J$ . Further, if  $N[h_1] \cap J = \emptyset$ , then we put  $g'_{2k+6} \in J'$ . If  $N[f'_{2k+8}] \cap J = \emptyset$ , then we put  $g_1 \in J'$ . If  $h_1 \in D$  and  $PN(h_1, D) = g'_{2k+6}$ , then we put  $h_1 \notin J'$ . Finally, if  $f'_{2k+8} \in D$  and  $PN(f'_{2k+8}, D) = g_1$ , then we put  $f'_{2k+8} \notin J'$ . It is easy to see that the set  $J'$  is a maximal irredundant set, and  $|J'| \leq |J| + 2$ . We obtain

$$\gamma(W_k) - ir(W_k) \geq |D| - |J'| \geq |D| - |J| - 2 = |D \cap V(G)| - |R| - 2 \geq k + 1.$$

■

**Lemma 2**  $i(W_k) - \gamma(W_k) \geq k + 1$ .

**Proof:** We denote by  $I$  an  $i$ -set of  $W_k$ .

**Claim 1** *We have  $|I \cap V(H_i)| = 3$  or  $4$  for any  $i$ ,  $1 \leq i \leq 3k + 6$ . Moreover,  $|I \cap V(H_i)| = 3$  if and only if either  $h_i$  or  $h'_i$  is dominated by  $I - V(H_i)$ , and additionally  $h_i, h'_i \notin I$ .*

**Proof:** Assume that  $h_i, h'_i \in I$  for some  $i$ ,  $1 \leq i \leq 3k + 6$ . We obtain  $|I \cap V(H_i)| = 4$ . Suppose now that exactly one vertex from  $h_i, h'_i$  belongs to  $I$ , say  $h_i \in I$  and  $h'_i \notin I$ . If  $b_i, c_i \notin I$ , then these vertices cannot be dominated by an independent set, a contradiction. Therefore, without loss of generality,  $b_i \in I$  and  $c_i \notin I$ . Hence  $a'_i \in I$ , and either  $c'_i \in I$  or  $d'_i \in I$ . We have  $|I \cap V(H_i)| = 4$ . Consider the case  $h_i, h'_i \notin I$ . Since  $I \cap \{b_i, b'_i, c_i, c'_i\} \neq \emptyset$ , we may assume without loss of generality that  $b_i \in I$  and hence  $a'_i \in I$ . If  $c'_i \in I$ , then  $d_i \in I$  and  $|I \cap V(H_i)| = 4$ . If  $c'_i \notin I$ , then  $d'_i \in I$  and  $|I \cap V(H_i)| = 3$ . ■

By Claim 1, the number of components  $H_i$  satisfying  $|I \cap V(H_i)| = 3$  is at most  $2k + 4$ . Therefore,

$$|I \cap V(H)| \geq 10k + 20.$$

Let us consider the set  $D = \{h'_{3k+6}, h_i, b'_i, c'_i : i = 1, 2, \dots, 3k + 6\}$ . It is evident that the set  $J = (I - V(H)) \cup D$  is a dominating set of  $W_k$  and

$$i(W_k) - \gamma(W_k) \geq |I| - |J| = |I \cap V(H)| - |D| \geq 10k + 20 - 9k - 19 = k + 1.$$

■

Now we estimate the difference between the independence and independent domination numbers of  $W_k$ .

**Lemma 3**  $\beta_0(W_k) - i(W_k) \geq 2k + 4$ .

**Proof:** It is easy to construct a maximal independent set  $I$  of  $W_k$  such that  $|I \cap V(F_i)| = 6$ ,  $|I \cap V(G_i)| = 6$ , and  $|I \cap V(H_i)| = 4$ . We define the set  $R \subset V(H)$  as follows. For each  $i \in \{1, 2, \dots, 3k + 6\}$ , we put  $a_i, d_i, b'_i \in R$  if  $i \equiv 1 \pmod{3}$ ,  $h_i, h'_i, b'_i, c_i \in R$  if  $i \equiv 2 \pmod{3}$ , and  $a'_i, b_i, d'_i \in R$  if  $i \equiv 0 \pmod{3}$ . Now, the set  $J = (I - V(H)) \cup R$  is an independent dominating set and hence  $i(W_k) \leq |J|$ . We obtain

$$\beta_0(W_k) - i(W_k) \geq |I| - |J| = |I \cap V(H)| - |R| = 12k + 24 - 10k - 20 = 2k + 4.$$

■

**Lemma 4**  $\Gamma(W_k) - \beta_0(W_k) \geq 3k + 5$ .

**Proof:** We can split  $V(F_i)$  into three cycles  $C_3$  and one  $C_7$ ,  $V(G_i)$  into two cycles  $C_5$  and two cycles  $C_3$ , and  $V(H_i)$  into two cycles  $C_5$ . Therefore,

$$\beta_0(W_k) \leq 6(2k + 8) + 6(2k + 6) + 4(3k + 6) = 36k + 108.$$

It is easy to construct a maximal independent set  $I$  of  $W_k$  such that  $|I \cap V(F_i)| = 6$ ,  $|I \cap V(G_i)| = 6$  and  $g'_{2k+6} \in I$ , and  $|I \cap V(H_i)| = 4$ . Thus,  $|I| = 36k + 108$  and hence  $\beta_0(W_k) = |I|$ .

Consider the set  $S = \{h'_i, a'_i, b'_i, c'_i, d'_i : 1 \leq i \leq 3k + 6\} - \{h'_{3k+6}\}$ . It is evident that  $R = (I - V(H)) \cup S$  is a minimal dominating set. We have

$$\Gamma(W_k) - \beta_0(W_k) \geq |R| - |I| = |S| - |I \cap V(H)| = 15k + 29 - 12k - 24 = 3k + 5.$$

■

Denote by  $D$  a  $\Gamma$ -set of  $W_k$ .

**Proposition 1**  $|D \cap V(F)| \leq 13k + 53$ .

**Proof:** Let us label the vertices of  $F_i$  as shown in Figure 2, and put  $X = \{x, a, b, h, i, j\}$ .




Figure 2.

**Claim 2**  $|D \cap X| = 2$ . Moreover,  $f, e, m \notin D$  if  $c, d \in D$  and at least one of the vertices  $k, k', p$  belongs to  $D$ .

**Proof:** Since  $\{a, b, h, i\}$  is dominated by  $D \cap X$  and at least two vertices are required to dominate it,  $|D \cap X| \geq 2$ . Suppose  $|D \cap X| \geq 3$ . If  $|D \cap \{a, b, x\}| \geq 2$ , then without loss of generality  $a \in D$  and  $PN(a) = \{h\}$ . Thus  $h, i, j \notin D$ , so  $x \in D$  and  $PN(x) = \{d\}$ , whence  $c \notin D$ . Hence  $j$  is not dominated, a contradiction. A similar contradiction shows that  $|D \cap \{h, i, j\}| \geq 2$  is impossible. Therefore  $|D \cap X| = 2$ .

Suppose that  $c, d \in D$  and at least one of the vertices  $k, k', p$  belongs to  $D$ . We have  $\{x\} = PN(d)$  and hence  $a, b, x \notin D$ . Therefore  $h, i \in D$  and  $PN(c) = \{e\}$ . Hence  $m, e, f \notin D$ . ■

We define 16 types for the component  $F_i$  as follows:

- $F_i$  has type A1 if  $k', g' \in D$  and  $k \in PN(k'), g \in PN(g')$ ;
- $F_i$  has type A2 if  $k', g' \in D$  and  $k \notin PN(k'), g \in PN(g')$ ;
- $F_i$  has type A3 if  $k', g' \in D$  and  $k \in PN(k'), g \notin PN(g')$ ;
- $F_i$  has type A4 if  $k', g' \in D$  and  $k \notin PN(k'), g \notin PN(g')$ ;
- $F_i$  has type B1 if  $k' \in D, g' \notin D$  and  $k \in PN(k'), g' \in N(D - V(F_i))$ ;
- $F_i$  has type B2 if  $k' \in D, g' \notin D$  and  $k \notin PN(k'), g' \in N(D - V(F_i))$ ;
- $F_i$  has type B3 if  $k' \in D, g' \notin D$  and  $k \in PN(k'), g' \in PN(g)$ ;
- $F_i$  has type B4 if  $k' \in D, g' \notin D$  and  $k \notin PN(k'), g' \in PN(g)$ ;
- $F_i$  has type C1 if  $k' \notin D, g' \in D$  and  $k' \in N(D - V(F_i)), g \in PN(g')$ ;
- $F_i$  has type C2 if  $k' \notin D, g' \in D$  and  $k' \in N(D - V(F_i)), g \notin PN(g')$ ;
- $F_i$  has type C3 if  $k' \notin D, g' \in D$  and  $k' \in PN(k), g \notin PN(g')$ ;
- $F_i$  has type C4 if  $k' \notin D, g' \in D$  and  $k' \in PN(k), g \in PN(g')$ ;
- $F_i$  has type D1 if  $k', g' \notin D$  and  $k' \in N(D - V(F_i)), g' \in N(D - V(F_i))$ ;
- $F_i$  has type D2 if  $k', g' \notin D$  and  $k' \in PN(k), g' \in N(D - V(F_i))$ ;
- $F_i$  has type D3 if  $k', g' \notin D$  and  $k' \in N(D - V(F_i)), g' \in PN(g)$ ;
- $F_i$  has type D4 if  $k', g' \notin D$  and  $k' \in PN(k), g' \in PN(g)$ .

Let us denote  $D_i = D \cap V(F_i)$ .

**Claim 3** We have

- (a1)  $|D_i| = 5$  if  $F_i$  is of type A1;
- (a2)  $|D_i| = 6$  if  $F_i$  is of type A2;
- (a3)  $|D_i| = 5$  if  $F_i$  is of type A3;
- (a4)  $|D_i| = 6$  if  $F_i$  is of type A4;
- (b1)  $|D_i| = 5$  if  $F_i$  is of type B1;
- (b2)  $|D_i| = 6$  if  $F_i$  is of type B2;
- (b3)  $|D_i| = 5$  if  $F_i$  is of type B3;
- (b4)  $|D_i| = 7$  if  $F_i$  is of type B4;
- (c1)  $|D_i| = 6$  if  $F_i$  is of type C1;
- (c2)  $|D_i| = 6$  if  $F_i$  is of type C2;
- (c3)  $|D_i| = 7$  if  $F_i$  is of type C3;
- (c4)  $|D_i| = 6$  if  $F_i$  is of type C4;
- (d1)  $|D_i| = 6$  if  $F_i$  is of type D1;
- (d2)  $|D_i| = 7$  if  $F_i$  is of type D2;

- (d3)  $|D_i| = 7$  if  $F_i$  is of type  $D3$ ;  
(d4)  $|D_i| = 8$  if  $F_i$  is of type  $D4$ .

**Proof:** In what follows we will use the first part of Claim 2 without further reference.

(a1) Since  $k \in PN(k')$  and  $g \in PN(g')$ , we have  $d, k, p, g, f \notin D$ . Also,  $y \in D$ , for otherwise  $p$  is not dominated. Suppose that  $e \in D$ . We have  $m, n \notin D$ . Now we can use the vertex  $c$  and two vertices of  $X$  to construct  $D_i$  such that  $|D_i| = 5$ . Assume that  $e \notin D$ . We obtain  $n \in D$ . It is easy to see that exactly one of the vertices  $c, m$  belongs to  $D$  and hence  $|D_i| = 5$ .

(a2) We have  $p, g, f \notin D$ . If  $c \notin D$ , then  $|D_i - X| = 4$  and hence  $|D_i| = 6$ . Suppose that  $c \in D$ . If  $d \in D$ , then  $m, e \notin D$  by Claim 2. Hence  $n \in D$  and  $|D_i| = 6$ . If  $d \notin D$ , then again  $|D_i| = 6$ .

(a3) We have  $d, k, p \notin D$ . If  $c \notin D$ , then  $|D_i - X| = 3$  and hence  $|D_i| = 5$ . Consider the case  $c \in D$ . If  $y \in D$ , then  $|D_i| = 5$ . If  $y \notin D$ , then  $g \in D$ , for otherwise  $p$  is not dominated. To dominate  $y$  we must take either  $m$  or  $n$  and hence  $|D_i| = 5$ .

(a4) Assume that  $c, d \notin D$ . It is not difficult to see that  $|D_i - X| = 4$  and hence  $|D_i| = 6$ . Consider the case  $c \notin D$  and  $d \in D$ . If  $k \in D$ , then  $\{p\} = PN(k)$  and hence  $g, p, y \notin D$ . We have  $|D_i| = 6$ . If  $k \notin D$ , then one can easily check that again  $|D_i| = 6$ . The case  $c \in D$  and  $d \notin D$  is analogous. Finally, suppose that  $c, d \in D$ . By Claim 2,  $e, f, m \notin D$ . If  $k \in D$ , then  $\{p\} = PN(k)$ . Therefore,  $y, p, g \notin D$ ,  $n \in D$  and  $|D_i| = 6$ . If  $k \notin D$ , then exactly two vertices from  $\{n, y, p, g\}$  belong to  $D$  and  $|D_i| = 6$ .

(b1) We have  $d, k, p \notin D$ . Suppose that  $c \notin D$ . If  $y \in D$ , then  $|D_i - X| = 3$  and hence  $|D_i| = 5$ . If  $y \notin D$ , then  $g \in D$  to dominate  $p$ . Again,  $|D_i - X| = 3$  and  $|D_i| = 5$ . Consider the case  $c \in D$ . If  $y \in D$ , then  $f \in D$  or  $g \in D$ , for otherwise  $g$  is not dominated. We have  $|D_i| = 5$ . If  $y \notin D$ , then  $g \in D$ , for otherwise  $p$  is not dominated. Also, one of the vertices  $m, n$  belongs to  $D$  to dominate  $y$ . We obtain  $|D_i| = 5$ .

(b2) Suppose that  $c, d \notin D$ . It is not difficult to see that  $|D_i - X| = 4$  and hence  $|D_i| = 6$ . Consider the case  $|D \cap \{c, d\}| = 1$ . If  $k \in D$ , then  $\{p\} = PN(k)$  and hence  $g, p, y \notin D$ . We have  $|D_i| = 6$ . If  $k \notin D$ , then one can easily check that again  $|D_i| = 6$ . Finally, assume that  $c, d \in D$ . By Claim 2,  $f, e, m \notin D$ . If  $k \in D$ , then  $PN(k) = \{p\}$  and hence  $g, p, y \notin D$ . Now  $g$  is not dominated, a contradiction. If  $k \notin D$ , then  $|D_i| = 6$ .

(b3) We have  $d, k, p \notin D$  and  $g \in D$ . Suppose that  $c \notin D$ . If  $y \in D$ , then  $f, m \notin D$ ,  $e \in D$  and hence  $|D_i| = 5$ . If  $y \notin D$ , then again  $|D_i| = 5$ . Consider the case  $c \in D$ . To dominate  $y$ , exactly one of the vertices  $m, n, y$  belongs to  $D$ . Hence  $|D_i| = 5$ .

(b4) We have  $g \in D$ . Suppose that  $c, d \notin D$ . It is not difficult to see that  $|D_i - X| = 5$  and hence  $|D_i| = 7$ . Consider the case  $|D \cap \{c, d\}| = 1$ . If  $k \in D$ , then  $PN(k) = \emptyset$ , a contradiction. Therefore,  $k \notin D$ . It is easy to see that  $|D_i| = 7$ . Finally, assume that  $c, d \in D$ . By Claim 2,  $f, e, m \notin D$ . If  $k \in D$ , then  $PN(k) = \emptyset$ , a contradiction. Therefore,  $k \notin D$ . We obtain  $|D_i| = 6$ . Since  $D$  is a maximum minimal dominating set, we conclude that  $|D_i| = 7$ .

(c1) We have  $f, g, p \notin D$ . Suppose that  $k \notin D$ . We obtain  $y \in D$  to dominate  $p$ , and  $d \in D$  to dominate  $k$ . Therefore,  $|D_i| = 6$ . Consider the case  $k \in D$ . If  $c, d \notin D$ , then  $|D_i| = 5$ . If exactly one vertex from  $\{c, d\}$  is present in  $D$ , then it is checked directly that  $|D_i| = 6$ . Finally, suppose that  $c, d \in D$ . By Claim 2,  $e, m \notin D$ . We have  $|D_i| = 6$ . Since  $D$  is a maximum minimal dominating set, we conclude that  $|D_i| = 6$ .

(c2) Assume that  $c, d \notin D$ . It is not difficult to see that  $|D_i - X| = 4$  and hence  $|D_i| = 6$ . Consider the case  $c \notin D$  and  $d \in D$ . If  $k \in D$ , then  $\{p\} = PN(k)$  and hence

$g, p, y \notin D$ . We have  $|D_i| = 6$ . If  $k \notin D$ , then one can easily check that again  $|D_i| = 6$ . Consider the case  $c \in D$  and  $d \notin D$ . If  $k \notin D$ , then  $p \in D$  to dominate  $k$ . We obtain  $|D_i| = 6$ . If  $k \in D$ , then  $p \notin D$ , for otherwise  $PN(k) = \emptyset$ . It is easy to see that  $|D_i| = 6$ . Finally, suppose that  $c, d \in D$  and consider two cases.

*Case 1.*  $k \in D$ . By Claim 2,  $e, f, m \notin D$ . Further,  $\{p\} = PN(k)$ . Therefore,  $g, p, y \notin D, n \in D$  and  $|D_i| = 6$ .

*Case 2.*  $k \notin D$ . Suppose that  $p \in D$ . By Claim 2,  $e, f, m \notin D$ . Also,  $y \notin D$ , for otherwise  $PN(p) = \emptyset$ . We obtain  $n \in D$  and  $|D_i| = 6$ . Assume now that  $p \notin D$ . If  $y \in D$ , then  $|D_i| = 6$ . If  $y \notin D$ , then  $g \in D$  to dominate  $p$ . Moreover, exactly one vertex from  $\{m, n\}$  belongs to  $D$ . Thus,  $|D_i| = 6$ .

(c3) We have  $k \in D$ . Suppose that  $c, d \in D$ . By Claim 2,  $f, e, m \notin D$ . We see that  $|D_i| = 7$ . Consider the case  $|D \cap \{c, d\}| = 1$ . It is checked directly that  $|D_i| = 7$ . If  $c, d \notin D$ , then  $|D_i| = 6$ . Since  $D$  is a maximum minimal dominating set, we conclude that  $|D_i| = 7$ .

(c4) We have  $f, g, p \notin D$ . Suppose that  $c \notin D$ . It is checked directly that  $|D_i| = 6$ . Consider the case  $c \in D$ . If  $d \notin D$ , then  $|D_i| = 6$ . If  $d \in D$ , then  $e, m \notin D$  by Claim 2. Again,  $|D_i| = 6$ .

(d1) Assume that  $c, d \notin D$ . If  $k \notin D$ , then  $p \in D$  and  $|D_i| = 5$ . If  $k \in D$ , then it is not difficult to see that  $|D_i - X| = 4$  and hence  $|D_i| = 6$ . Consider the case  $c \notin D$  and  $d \in D$ . If  $k \in D$ , then  $\{p\} = PN(k)$  and hence  $g, p, y \notin D$ . We have  $|D_i| = 6$ . If  $k \notin D$ , then one can easily check that again  $|D_i| = 6$ . Consider the case  $c \in D$  and  $d \notin D$ . If  $k \notin D$ , then  $p \in D$  and  $|D_i| = 6$ . If  $k \in D$ , then  $p \notin D$ , for otherwise  $PN(k) = \emptyset$ . It is easy to see that  $|D_i| = 6$ . Finally, suppose that  $c, d \in D$ . By Claim 2,  $e, f, m \notin D$ . If  $k \in D$ , then  $\{p\} = PN(k)$ . Therefore,  $y, p, g \notin D, n \in D$  and  $|D_i| = 6$ . If  $k \notin D$ , then exactly two vertices from  $\{n, y, p, g\}$  belong to  $D$  and  $|D_i| = 6$ . Since  $D$  is a maximum minimal dominating set, we conclude that  $|D_i| = 6$ .

(d2) The proof is analogous to the case (c3).

(d3) We have  $g \in D$ . The only difference between this case and the case (b4) is that the vertex  $k$  is dominated by  $k'$  in the latter case. Hence, if  $d \in D$  or  $k \in D$ , then we use the corresponding reasoning of the case (b4) and obtain  $|D_i| = 7$ . Suppose now that  $d, k \notin D$ . We have  $p \in D$ , for otherwise  $k$  is not dominated. Obviously  $c, e, f \in D$  and  $|D_i| = 7$ .

(d4) We have  $k, g \in D$ . Suppose that  $c, d \notin D$ . It is not difficult to see that  $D_i - X = \{k, e, f, g, p\}$ . Hence  $|D_i| = 7$ . If  $|D \cap \{c, d\}| = 1$ , then  $|D_i| = 8$ . Finally, assume that  $c, d \in D$ . By Claim 2,  $f, e, m \notin D$  and hence  $|D_i| = 7$ . Since  $D$  is a maximum minimal dominating set, we conclude that  $|D_i| = 8$ . ■

**Claim 4** *If  $F_i$  ( $2 \leq i \leq 2k + 7$ ) has type  $D_4$ , then both (i) and (ii) hold; if  $F_i$  has type  $D_4$  and  $i = 2k + 8$ , then (i) holds. Furthermore, if  $F_i$  ( $2 \leq i \leq 2k + 7$ ) is of type  $B_4, C_3, D_2$  or  $D_3$ , then at least one of the properties (i) and (ii) holds.*

(i)  $F_{i-1}$  has type  $A_1, A_2, C_1$  or  $C_4$  and  $|D_{i-1}| \leq 6$ .

(ii)  $F_{i+1}$  has type  $A_1, A_3, B_1$  or  $B_3$  and  $|D_{i+1}| = 5$ .

**Proof:** This follows immediately from the definition and Claim 3. ■

Let  $F_i$  be a component of type  $D_4$  for some  $i \leq 2k + 7$ . By Claim 3,  $|D_i| = 8$ . By Claim 4,  $F_{i+1}$  has type  $A_1, A_3, B_1$  or  $B_3$  and  $|D_{i+1}| = 5$ . We denote by  $m$  the number of such



pairs. These components contain exactly  $13m$  vertices of  $D$ , and any other component  $F_j$  with  $j \leq 2k + 7$  has  $|D_j| \leq 7$ . Suppose that there exist three sequential components  $F_i, F_{i+1}, F_{i+2}$  such that  $|D_i| = |D_{i+1}| = |D_{i+2}| = 7$ , i.e., they are of type B4, C3, D2 or D3 by Claim 3. Applying Claim 4 to  $F_{i+1}$  we arrive at a contradiction. Consider two components  $F_i, F_{i+1}$  of type B4, C3, D2 or D3 such that  $i \leq 2k + 6$ . We have  $|D_i| = |D_{i+1}| = 7$ . Applying Claim 4 to  $F_{i+1}$ , we obtain  $|D_{i+2}| = 5$  for the component  $F_{i+2}$ . Denote by  $n$  the number of such triples. We see that these triples contain  $19n$  vertices of  $D$ .

Suppose that the component  $F_{2k+8}$  belongs to one of the above pairs or triples, and consider a maximal sequence

$$F_{i+1}, F_{i+2}, \dots, F_{i+r}$$

not containing the components from the above pairs and triples. It is obvious that either  $|D_{i+r+1}| = 8$  or  $|D_{i+r+1}| = |D_{i+r+2}| = 7$ . In the first case we know that  $F_{i+r+1}$  is of type D4 and  $|D_{i+r}| \leq 6$  by Claim 4. For the latter case we know that  $F_{i+r+1}$  must have type B4, C3, D2 or D3. Hence, by Claim 4,  $|D_{i+r}| \leq 6$ . Thus,

$$\sum_{j=1}^r |D_{i+j}| \leq 6.5r.$$

Taking into account all such maximal sequences, we obtain

$$|D \cap V(F)| \leq 13m + 19n + 6.5(2k + 8 - 2m - 3n) = 13k + 52 - 0.5n \leq 13k + 52.$$

Assume now that the component  $F_{2k+8}$  does not belong to any of the above pairs or triples, and denote by  $L$  a maximal sequence

$$F_{l+1}, F_{l+2}, \dots, F_{2k+8}$$

not containing the components from those pairs and triples. If  $|D_{2k+8}| = 8$ , then  $|D_{2k+7}| = 6$  by Claim 4. We have

$$\sum_{j=1}^{2k+8-l} |D_{l+j}| \leq 6.5(2k + 8 - l) + 1.5 = 6.5|L| + 1.5.$$

If  $|D_{2k+8}| = 7$ , then it is not difficult to see that

$$\sum_{j=1}^{2k+8-l} |D_{l+j}| \leq 6.5(2k + 8 - l) + 1 = 6.5|L| + 1.$$

We have already proved that if  $F_{i+1}, F_{i+2}, \dots, F_{i+r}$  ( $i + r < 2k + 8$ ) is a maximal sequence not containing the components of the pairs and triples, then

$$\sum_{j=1}^r |D_{i+j}| \leq 6.5r.$$

Taking into account all such maximal sequences and  $L$ , we obtain

$$|D \cap V(F)| \leq 13m + 19n + 6.5(2k + 8 - 2m - 3n - |L|) + 6.5|L| + 1.5 =$$

$$13k + 53.5 - 0.5n \leq 13k + 53.5.$$

Thus,

$$|D \cap V(F)| \leq 13k + 53,$$

as required. The proof of Proposition 1 is complete. ■

**Lemma 5**  $IR(W_k) - \Gamma(W_k) \geq k + 1$ .

**Proof:** Since  $D$  is a  $\Gamma$ -set, it follows that  $D$  is maximal irredundant. Adding to  $D - V(F)$  some new vertices, we will construct a set  $D'$  which is maximal irredundant and

$$|D' \cap V(F)| \geq 14k + 54.$$

We first put  $D' = D - V(F)$ . Taking into account the definition of the 16 types of the component  $F_1$ , we consider 4 cases. Suppose that  $k' \in D$  and  $k \in PN(k', D)$ . In this case, we put  $a, b, x, m, n, y \in D'$ . We do the same if  $k' \in D$  and  $k \notin PN(k', D)$ . Assume that  $k' \notin D$  and  $k' \in N(D - V(F_1))$ , say  $k'$  is adjacent to  $k''$ . Now, we put  $a, b, x, m, n, y \in D'$  if  $\{k'\} = PN(k'', D)$ , and we put  $h, i, j, k, m, n, p \in D'$  otherwise. Finally, suppose that  $k' \notin D$  and  $k' \in PN(k, D)$ . We put  $h, i, j, k, m, n, p \in D'$ .

Let us consider the component  $F_{2k+8}$ . Suppose that  $g' \in D$  and  $g \in PN(g', D)$ . We put  $a, b, x, m, n, y \in D'$ . Assume that  $g' \in D$  but  $g \notin PN(g', D)$ . We put  $a, b, x, m, n, y \in D'$ . Consider now the case  $g' \notin D$  and  $g' \in N(D - V(F))$ , say  $g'$  is adjacent to  $g''$ . We put  $a, b, x, m, n, y \in D'$  if  $\{g'\} = PN(g'', D)$ , and we put  $a, b, c, d, e, f, g \in D'$  otherwise. Finally, suppose that  $g' \notin D$  and  $g' \in PN(g, D)$ . We put  $a, b, c, d, e, f, g \in D'$ .

For  $2 \leq i \leq 2k + 7$ , we put  $a, b, c, d, e, f, g \in D'$  if  $i$  is even, and  $h, i, j, k, m, n, p \in D'$  if  $i$  is odd. It is easy to see that the resulting set  $D'$  is a maximal irredundant set and  $|D' \cap V(F)| \geq 14k + 54$ . Applying Proposition 1, we obtain

$$IR(W_k) - \Gamma(W_k) \geq |D'| - |D| = |D' \cap V(F)| - |D \cap V(F)| \geq 14k + 54 - 13k - 53 = k + 1.$$

■

Using Lemmas 1–5 we can easily choose the integer  $k$  such that the conditions of Theorem 1 are satisfied. The proof of Theorem 1 is complete.

**Acknowledgment** The authors thank the referee for valuable comments and suggestions.

## References

- [1] C. Barefoot, F. Harary and K.F. Jones, What is the difference between the domination and independent domination numbers of a cubic graph? *Graphs and Combin.* **7** (1991) 205–208.
- [2] E.J. Cockayne, O. Favaron, C. Payan and A.G. Thomason, Contributions to the theory of domination, independence and irredundance in graphs. *Discrete Math.* **33** (1981) 249–258.

- [3] E.J. Cockayne and C.M. Mynhardt, The sequence of upper and lower domination, independence and irredundance numbers of a graph. *Discrete Math.* **122** (1993) 89–102.
- [4] E.J. Cockayne and C.M. Mynhardt, Domination and irredundance in cubic graphs. *Discrete Math.* **167/168** (1997) 205–214.
- [5] E.J. Cockayne and C.M. Mynhardt, Independence and domination in 3-connected cubic graphs. *JCMCC* **10** (1991) 173–182.
- [6] J.E. Dunbar, S.T. Hedetniemi, M.A. Henning and A.A. McRae, Minus domination in regular graphs. *Discrete Math.* **49** (1996) 311–312.
- [7] J.E. Dunbar, S.T. Hedetniemi, M.A. Henning and P.J. Slater, Signed domination in graphs. *Graph Theory, Combinatorics, and Applications*, John Wiley & Sons, Inc. **1** (1995) 311–322.
- [8] O. Favaron, Signed domination in regular graphs. *Discrete Math.* **158** (1996) 287–293.
- [9] M.A. Henning, Domination functions in graphs. In *Domination in Graphs: Advanced Topics*, T.W. Haynes, S.T. Hedetniemi and P.J. Slater (eds), Marcel Dekker, Inc., New York, 1998, 31–57.
- [10] M.A. Henning, Domination in regular graphs. *Ars Combin.* **43** (1996) 263–271.
- [11] M.A. Henning and G. Kubicki, Real domination in graphs. *J. Combin. Math. Combin. Comput.* **26** (1998) 147–160.
- [12] M.A. Henning and P.J. Slater, Inequalities relating domination parameters in cubic graphs. *Discrete Math.* **158** (1996) 87–98.
- [13] A.V. Kostochka, The independent domination number of a cubic 3-connected graph can be much larger than its domination number. *Graphs Combin.* **9** (1993) 235–237.
- [14] C.M. Mynhardt, On the difference between the domination and independent domination numbers of cubic graphs. *Theory of Graphs and Applications*, New York: Wiley, 1991, 939–947.
- [15] D. Rautenbach, On the differences between the upper irredundance, upper domination and independence numbers of a graph. *Discrete Math.* **203** (1999) 239–252.
- [16] B. Zelinka, Some remarks on domination in cubic graphs. *Discrete Math.* **158** (1996) 249–255.
- [17] I.E. Zverovich and V.E. Zverovich, Disproof of a conjecture in the domination theory. *Graphs Combin.* **10** (1994) 389–396.