学校编码: 10384

学 号: 22620080150125

を の よう

博士学位论文

携带黑鲷 AS-prohepc2 及拟穴青蟹 Scy1 抗菌肽基因的减毒活菌载体系统作为 口服免疫途径的研究

Oral immunization studies relied on the live attenuated bacteria *in vivo* expressing antimicrobial peptides AS-prohepc2 and Scy1 in fish

陈贝

指导教师姓名:王克坚教授

专业名称:环境科学

论文提交日期: 2015年4月

论文答辩时间: 2015 年 5 月

2015 年 5 月

1

密级

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。 本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文 中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活 动规范(试行)》。

另外,该学位论文为(分子生物学与免疫毒理学)课题(组) 的研究成果,获得(国家海洋"863"重点项目《海洋动物抗菌类基 因工程产品作为饲料添加剂的研发及其应用研究》及福建省科技合作 计划项目《海水养殖鱼类抗菌肽口服免疫技术的研究》)课题(组) 经费或实验室的资助,在(王克坚教授)实验室完成。(请在以上 括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的, 可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》 等规定保留和使用此学位论文,并向主管部门或其指定机构送交学位 论文(包括纸质版和电子版),允许学位论文进入厦门大学图书馆及 其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、 硕士学位论文共建单位数据库进行检索,将学位论文的标题和摘要汇 编出版,采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于:

()1.经厦门大学保密委员会审查核定的保密学位论文,于年 月 日解密,解密后适用上述授权。

() 2.不保密,适用上述授权。

(请在以上相应括号内打"√"或填上相应内容。保密学位论文应 是已经厦门大学保密委员会审定过的学位论文,未经厦门大学保密委 员会审定的学位论文均为公开学位论文。此声明栏不填写的,默认为 公开学位论文,均适用上述授权。)

声明人(签名):

年 月 日

目 录

第一章	绪论	1
1.1 抗	ī菌肽研究概述	1
1.1.	1 抗菌肽来源、分类及理化特性	1
1.1.	2 抗菌肽的表达调控	1
1.1.	3 抗菌肽抗微生物特性	2
1.1.	4 抗菌肽免疫防御功能	3
1.1.	5 抗菌肽抗肿瘤功能	6
1.1.	6 Hepcidin 研究概述	6
1.1.	7 Scygonadin 研究概述	10
1.2 茲	迈物或疫苗运载体系	12
1.2.	1 生物工程及仿生载药系统	12
1.2.	2 基于细菌的载药策略	12
1.2.	3 减毒沙门氏菌在基因传递中研究进展	16
1.3 水	《产动物病害防治研究进展	29
1.3.	1 基于宿主的防治策略	30
1.3.	2 基于病原体的防治策略	35
1.4 本	际穷的技术路线、目的和意义	
1.4.	1 研究目的和意义	
1.4.	2 技术路线	41
参考了	文献	42
第二章	:黑鲷抗菌肽 HEPCIDIN 抗菌功能研究	62
2.1 实	会验材料	62
2.1.	1 原核表达载体、宿主菌及其它检测用菌株	62
2.1.	2 主要试剂和耗材	63

2.1.3 仪器	3
2.1.4 试剂配制64	4
2.2 实验方法65	5
2.2.1 AS-Hepc2 与 AS-Hepc6 成熟肽合成肽的制备及生物信息学分析65	5
2.2.2 前体肽 AS-prohepc2 的原核表达	5
2.2.3 最小抑菌浓度(MIC)和最小杀菌浓度(MBC)的测定	1
2.2.4 杀菌动力学测定71	1
2.3 实验结果	2
2.3.1 AS-hepc2 和 AS-hepc6 基因序列及成熟肽性质72	2
2.3.2 AS-hepc2 和 AS-hepc6 成熟肽抗菌谱74	4
2.3.3 AS-hepc2 和 AS-hepc6 成熟肽杀菌动力学曲线	5
2.3.4 前体肽 AS-prohepc2 的原核表达	5
2.3.5 AS-prohepc2 原核表达蛋白抗菌谱	2
2.4 讨论	3
2.4.1 黑鲷 hepcidin 成熟肽特性及抗菌活性	3
2.4.2 hepcidin 前体肽的原核表达	5
参考文献	9
第三章 海洋动物抗菌肽基因工程表达产物体内外抗菌活性研究92	2
3.1 实验材料	2
3.1.1 真核表达载体及宿主菌92	2
3.1.2 主要试剂和耗材92	2
3.1.3 仪器	3
3.1.4 试剂配制	3
3.2 实验方法	5
3.2.1 Scy1 在毕赤酵母中真核表达与纯化95	5
3.2.2 AS-prohepc2 在毕赤酵母中真核表达与纯化	5
3.2.3 黑鲷头肾巨噬细胞(BPMφ)体外培养条件优化96	5
3.2.4 抗菌肽真核表达产物共孵黑鲷巨噬细胞免疫相关基因表达检测97	7
3.2.5 抗菌肽真核表达产物共孵鲤鱼 EPC 细胞免疫相关基因表达检测 100)

3.2.6 Scyl 真核表达产物共孵黑鲷巨噬细胞吞噬活性检测	101
3.2.7 Scyl 真核表达产物注射斑马鱼的嗜水气单胞菌感染实验	102
3.3 实验结果	103
3.3.1 Scyl 在毕赤酵母中表达及纯化	103
3.3.2 AS-prohepc2 在毕赤酵母中表达及纯化	104
3.3.3 黑鲷巨噬细胞培养最适盐度	104
3.3.4 黑鲷巨噬细胞及鲤鱼 EPC 细胞免疫相关基因扩增效率标准曲线	106
3.3.5 Scy1 对黑鲷巨噬细胞和鲤鱼 EPC 细胞免疫相关基因表达的影响	107
3.3.6 AS-prohepc2 对黑鲷巨噬细胞和鲤鱼 EPC 细胞免疫相关基因表达	的影
响	109
3.3.7 Scy1 对黑鲷巨噬细胞吞噬效率的影响	110
3.3.8 Scy1 对嗜水气单胞菌感染的斑马鱼存活率的影响	
3.4 讨论	111
3.4.1 黑鲷巨噬细胞的体外培养	111
3.4.2 抗菌肽 Scyl 促进黑鲷巨噬细胞吞噬功能和免疫相关基因的上调	
	112
3.4.3 Scyl 真核表达产物提高斑马鱼嗜水气单胞菌感染的成活率	115
3.4.4 抗菌肽 AS-prohepc2 免疫功能	116
参考文献	117
第四章 体内外过表达海洋动物抗菌肽抗菌功能研究	.120
4.1 实验材料	120
4.1.1 表达载体、宿主菌株及其他检测用菌株	120
4.1.2 主要试剂和耗材	120
4.1.3 仪器	121
4.1.4 试剂配制	122
4.1.5 实验动物及细胞株	123
4.2 实验方法	123
4.2.1 抗菌肽目的基因的制备	123
4.2.2 真核表达载体的构建	125

4.2.3 AS-prohepc2 鼠多克隆抗体的制备	126
4.2.4 编码抗菌肽的真核表达载体在鲤鱼 EPC 细胞中的体外表达	128
4.2.5 过表达抗菌肽基因的鲤鱼 EPC 细胞抗菌功能检测	129
4.2.6 过表达抗菌肽基因的鲤鱼 EPC 细胞免疫相关基因表达检测	130
4.2.7 斑马鱼肌肉电转化真核质粒 pCMVHA/GFP 条件优化	130
4.2.8 斑马鱼肌肉电转化抗菌肽真核质粒蛋白表达检测	131
4.2.9 嗜水气单胞菌感染斑马鱼半数致死浓度 LD50 检测	131
4.2.10 斑马鱼肌肉过表达 Scy1 及 AS-prohepc2 后细菌感染死亡率检测	132
4.3 实验结果	132
4.3.1 重组真核表达质粒的鉴定及测序	
4.3.2 AS-prohepc2 鼠多克隆抗体效价	133
4.3.3 重组真核表达质粒在鲤鱼 EPC 细胞的表达	135
4.3.4 过表达抗菌肽基因的鲤鱼 EPC 细胞抗菌功能	136
4.3.5 过表达抗菌肽基因对鲤鱼 EPC 细胞免疫相关基因表达的影响	138
4.3.6 斑马鱼肌肉电转化真核质粒 pCMVHA/GFP 最适条件	140
4.3.7 肌肉电转化的斑马鱼在肌肉中检测到抗菌肽表达	141
4.3.8 斑马鱼肌肉注射嗜水气单胞菌半数致死浓度 LD50	142
4.3.9 肌肉过表达抗菌肽对嗜水气单胞菌感染的斑马鱼存活率的影响	143
4.4 讨论	143
参考文献	148
第五章 减毒活细菌载体携带抗菌肽基因运载系统的建立及其可	行性
与安全性研究	.151
5.1 实验材料	151
5.1.1 表达载体及宿主菌	151
5.1.2 主要试剂和耗材	152
5.1.3 仪器	152
5.1.4 试剂配制	153
5.1.5 实验动物及细胞株	154
5.2 实验方法	154

5.2.1 海洋动物抗菌肽原核表达载体的构建	154
5.2.2 Scy1 真核表达载体的构建	158
5.2.3 重组质粒电转化减毒志贺氏菌或减毒沙门氏菌	162
5.2.4 携带抗菌肽原核原核表达质粒的重组减毒细菌体外蛋白表达检测	163
5.2.5 编码 Scyl 的真核表达载体在鲤鱼 EPC 细胞的体外表达测定	164
5.2.6 转染 pCMV-ScySP-HA-Scy1-asd 的鲤鱼 EPC 体外抗菌功能检测	
5.2.7 重组菌体外稳定性试验	
5.2.8 重组菌生长曲线的测定	164
5.2.9 减毒志贺氏菌及减毒沙门氏菌安全性实验	165
5.2.10 鱼体口服减毒细菌条件优化	165
5.2.11 灌胃减毒沙门氏菌鱼体内细菌定植情况	166
5.2.12 灌胃 Scy1 原核表达重组沙门氏菌鱼体内各组织 Scy1 基因表达情	
测	166
5.2.13 灌胃 Scyl 真核表达重组沙门氏菌鱼体内各组织 Scyl 蛋白检测	169
5.2.14 灌胃 Scy1 重组沙门氏菌生理生化指标检测	169
5.2.15 灌胃 Scy1 重组沙门氏菌头肾免疫相关基因表达 qPCR 分析	171
5.2.16 灌胃 Scy1 重组沙门氏菌黑鲷肠道形态观察	
5.3 实验结果	172
5.3.1 抗菌肽原核表达载体构建	172
5.3.2 抗菌肽 Scy1 真核表达载体构建	175
5.3.3 重组减毒志贺氏菌 Fwl01 或减毒沙门氏菌 X3730/X4072 的建立	178
5.3.4 减毒志贺氏/沙门氏原核表达菌株蛋白体外表达	178
5.3.5 Scyl 重组真核表达载体在体外培养鲤鱼 EPC 细胞中的表达	179
5.3.6 转染 pCMV-ScySP-HA-Scy1-asd 的鲤鱼 EPC 体外抗菌功能	180
5.3.7 重组菌的体外稳定性	181
5.3.8 重组菌生长曲线	182
5.3.9 减毒志贺氏菌及减毒沙门氏菌安全性	183
5.3.10 减毒菌株和口服方式的确定	183
5.3.11 灌胃减毒沙门氏菌鱼体内细菌定植情况	185

5.3.12 灌胃原核重组减毒沙门氏菌后 Scyl 基因的组织转录水平	189
5.3.13 灌胃真核重组沙门氏菌后 Scyl 蛋白的组织表达	190
5.3.14 灌胃 Scy1 重组沙门氏菌对生理生化指标影响	191
5.3.15 灌胃 Scyl 重组沙门氏菌对免疫相关基因表达影响	193
5.3.16 灌胃 Scy1 重组沙门氏菌对黑鲷肠道形态的影响	194
5.4 讨论	
5.4.1 减毒沙门氏菌作为抗菌肽运载载体	197
5.4.2 平衡致死宿主载体表达系统	
5.4.3 沙门氏菌携带原核及真核表达载体蛋白递送策略	200
5.4.4 沙门氏菌肠道定植及组织分布	
5.4.5 抗菌肽重组表达载体体内体外表达	203
5.4.6 减毒沙门氏菌对生理生化指标和肠道结构影响	204
5.4.7 减毒沙门氏菌对鱼类细胞因子表达影响	207
参考文献	208
结 语	212
1 研究成果	
2 主要创新点	213
3 展望	214
在学期间参加的科研项目及成果	215
致 谢	

Contents

Abstract in Chinese]
Abstract in EnglishIV
Chapter 1 Introduction1
1.1 Development and progress in AMPs1
1.1.1 The source, classify and physicochemical properties of AMPs1
1.1.2 Expression and regulation of AMPs1
1.1.3 Antimicrobial activity of AMPs2
1.1.4 AMPs as a modulators of the immume system
1.1.5 Anticancer activity
1.1.6 Research progress of Hepcidin
1.1.7 Research progress of Scygonadin
1.2 The delivery system for drug and vaccine12
1.2.1 Bioengineered and biomimetic drug delivery carriers12
1.2.2 Bioengineered strategies based on bacteria12
1.2.3 Research progress of attenuated Salmonella delivery carriers16
1.3 Development and progress in disease management in aquaculture29
1.3.1 Disease control based on host
1.3.2 Disease control based on pathogen35
1.4 Technical proposal, aims and significance of present study
1.4.1 Aims and significance of present study
1.4.2 Technical proposal41
References42
Chapter 2 Antimicrobial activity of black pory hepcidin62
2.1 Materials62
2.1.1 Prokaryotic expression vectors and bacterial strains
2.1.2 Reagent and Consumables63

2.1.3 Instruments	63
2.1.4 Reagent preparation	······64
2.2 Methods	65
2.2.1 Analysis and synthesis of AS-Hepc2 and AS-Hepc6 mature peptide	65
2.2.2 Prokaryotic expression of propeptide AS-prohepc2	66
2.2.3 Minimum inhibitory concentration (MIC) and minimum bact	tericidal
concentration (MBC) tests	71
2.2.4 Bactericidal kinetics tests	71
2.3 Results	 72
2.3.1 Property of AS-hepc2 and AS-hepc6 mature peptide	72
2.3.2 Antimicrobial spectrum of AS-hepc2 and AS-hepc6 mature peptide	74
2.3.3 Bactericidal kinetics curves of AS-hepc2 and AS-hepc6 mature pept	ide ∙•75
2.3.4 Prokaryotic expression of propeptide AS-prohepc2	76
2.3.5 Antimicrobial spectrum of propeptide AS-prohepc2	82
2.4 Discussion ······	83
2.4.1 Property and antimicrobial activity of mature peptide of blac	k pogy
hepcidin	83
2.4.2 Prokaryotic expression of hepcidin propeptide	86
References	89
Chapter 3 In vivo and in vitro antibacterial activity of marine a	nimal
antimicrobial peptide expression product from genetic engin	oorina
system ·····	92
3.1 Materials	92
3.1.1 Eukaryotic expression vectors and bacterial strains	92
3.1.2 Reagent and Consumables	92
3.1.3 Instruments	93
3.1.4 Reagent preparation	93
3.2 Methods	95
3.2.1 Expression and purification of Scy1 in Pichia pastoris	95

3.2.2 Expression and purification of AS-prohepc2 in <i>P. pastoris</i> 96
3.2.3 Optimized condition for the in vitro cell culture of headkidney-derived
macrophages from black porgy96
3.2.4 Detection of immune-related gene expression in BPM ϕ incubated with
antimicrobial peptide97
3.2.5 Detection of immune-related gene expression in EPC incubated with
antimicrobial peptide100
3.2.6 The Phagocytosis of BPMφ after incubation with Scy1101
3.2.7 Bacterial challenge analysis in zebrafish pre-injected Scy1 protein102
3.3 Results103
3.3.1 Expression and purification of Scy1 in <i>P. pastoris</i> 103
3.3.2 Expression and purification of AS-prohepc2 in <i>P. pastoris</i> 104
3.3.3 Optimum salinity for BPMφ <i>in vitro</i> culture104
3.3.4 Standard curve for qPCR efficiency of immune-related gene in BPM ϕ and
EPC106
3.3.5 Effect of Scy1 on immune-related gene expression in BPM ϕ and EPC $\cdot 107$
3.3.6 Effect of AS-prohepc2 on immune-related gene expression in BPM ϕ and
EPC109
3.3.7 Effect of Scy1 on phagocytic index of BPMφ ······110
3.3.8 Effect of Scy1 on the survival rate of zebrafish infected with Aeromonas
hydrophila111
3.4 Discussion111
3.4.1 <i>In vitro</i> culture of BPM ϕ 111
3.4.2 Scy1increased phagocytosis function and the expression of immune-related
gene in BPM
3.4.3 Scy1 increased the survival rate of zebrafish infected with A. hydrophila
3.4.4 Immune function of AS-prohepc2 ······116
References117

Chapter 4 In vivo and in vitro overexpression and antibacterial
activity of antimicrobial peptide from marine animals 120
4.1 Materials120
4.1.1 Expression vector and bacterial strains120
4.1.2 Reagent and Consumables120
4.1.3 Instruments
4.1.4 Reagent preparation122
4.1.5 Animal and cell line123
4.2 Methods
4.2.1 Cloning of target gene
4.2.2 Construction of expression vector125
4.2.3 Preparation for the mouse polyclonal antibody of AS-prohepc2126
4.2.4 In vitro expression of the plasmid DNA encoded antimicrobial peptide in
EPC128
4.2.5 Evaluation of antimicrobial activity of EPC over-expressed antimicrobial
peptide129
4.2.6 Detection of immune-related gene expression of EPC over-expressed
antimicrobial peptide130
4.2.7 Identification of the optimal parameters for electrotransfer of plasmid DNA
pCMVHA/GFP into zebrafish muscle130
4.2.8 Detection the expression of antimicrobial peptide in the zebrafish after
intramuscular electroporation131
4.2.9 Detection LD50 of zebrafish infected by A. hydrophila131
4.2.10 Bacterial challenge analysis in zebrafish over-expressed Scy1 and
AS-prohepc2 in the muscle
4.3 Results
4.3.1Sequencing and analysis of recombinant plasmid132
4.3.2 Determine of AS-prohepc2 polyclonal antibody titer133
4.3.3 In vitro expression of the antimicrobial peptide in EPC135

4.3.4 Antimicrobial activity of EPC over-expressed antimicrobial peptide136
4.3.5 Effect of over-expressed antimicrobial peptide on immune-related gene
expression in EPC138
4.3.6 Optimum condition for electrotransfer of plasmid DNA pCMVHA/GFP
into zebrafish muscle
4.3.7 Antimicrobial peptide expressed in the muscle of zebrafish after
intramuscular electroporation141
4.3.8 LD50 of zebrafish infected by A. hydrophila142
4.3.9 Effect of intramuscular over-expressed antimicrobial peptide on the
survival rate of zebrafish infected with A. hydrophila143
4.4 Discussion143
References148
Chapter 5 Constuction, feasibility and security of the live attenuated
bacteria system delivered antimicrobial peptides151
5.1 Materials151
5.1.1 Expression vectors and bacterial strains151
5.1.2 Reagent and Consumables152
5.1.3 Instruments152
5.1.4 Reagent preparation153
5.1.4 Reagent preparation1535.1.5 Animal and cell lines154
5.1.5 Animal and cell lines154
5.1.5 Animal and cell lines
5.1.5 Animal and cell lines
5.1.5 Animal and cell lines
5.1.5 Animal and cell lines 154 5.2 Method 154 5.2.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 154 5.2.2 Construction of eucaryotic expression plasmid for Scy1 154
5.1.5 Animal and cell lines 154 5.2 Method 154 5.2.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 154 5.2.2 Construction of eucaryotic expression plasmid for Scy1 158 5.2.3 Attenuated Salmonella typhimurium and Shigella flexneri were
5.1.5 Animal and cell lines 154 5.2 Method 154 5.2.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 154 5.2.2 Construction of eucaryotic expression plasmid for Scy1 158 5.2.3 Attenuated Salmonella typhimurium and Shigella flexneri were electrotransfer with recombinant plasmid 162

in EPC164
5.2.6 Evaluation of antimicrobial activity of EPC transfected with
pCMV-ScySP-HA-Scy1-asd ·····164
5.2.7 Stability test of recombinant attenuated bacteria164
5.2.8 Detection of growth curve of recombinant attenuated bacteria164
5.2.9 Safety test of attenuated S. typhimurium and S. flexneri to black porgy 165
5.2.10 Optimized condition for oral administration of attenuated bacteria165
5.2.11 Observation of tissue colonization of attenuated S. typhimurium166
5.2.12 In vivo transcription analysis of recombinant attenuated S. typhimurium
carried Scy1 prokaryotic plasmid166
5.2.13 In vivo Scy1 protein expression analysis of recombinant attenuated S.
typhimurium carried eukaryotic plasmid169
5.2.14 Physiological parameters analysis in intestine169
5.2.15 qPCR analysis of the expression of immune-related genes in headkidney
5.2.16 Investigation of intestinal histomorphology171
5.2.16 Investigation of intestinal histomorphology171
5.2.16 Investigation of intestinal histomorphology ······171 5.3 Results ·····172
 5.2.16 Investigation of intestinal histomorphology171 5.3 Results172 5.3.1 Construction of procaryotic expression plasmid for antimicrobial peptide
5.2.16 Investigation of intestinal histomorphology
5.2.16 Investigation of intestinal histomorphology
5.2.16 Investigation of intestinal histomorphology 171 5.3 Results 172 5.3.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 172 5.3.2 Construction of eucaryotic expression plasmid for Scy1 175 5.3.3 Construction of recombinant atteuated S. flexneri Fwl01 and S.
5.2.16 Investigation of intestinal histomorphology
5.2.16 Investigation of intestinal histomorphology 171 5.3 Results 172 5.3.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 172 5.3.2 Construction of eucaryotic expression plasmid for Scy1 175 5.3.3 Construction of recombinant atteuated S. flexneri Fwl01 and S. typhimurium X3730/X4072 178 5.3.4 In vitro expression of recombinant attenuated S. flexneri and S.
5.2.16 Investigation of intestinal histomorphology 171 5.3 Results 172 5.3.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 172 5.3.2 Construction of eucaryotic expression plasmid for Scy1 175 5.3.3 Construction of recombinant atteuated S. flexneri Fwl01 and S. typhimurium X3730/X4072 5.3.4 In vitro expression of recombinant attenuated S. flexneri and S. typhimurium carried prokaryotic expression plasmid 178
5.2.16 Investigation of intestinal histomorphology 171 5.3 Results 172 5.3.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 172 5.3.2 Construction of eucaryotic expression plasmid for Scy1 175 5.3.3 Construction of recombinant atteuated S. flexneri Fwl01 and S. typhimurium X3730/X4072 5.3.4 In vitro expression of recombinant attenuated S. flexneri and S. typhimurium carried prokaryotic expression plasmid 178 5.3.5 In vitro expression of the reconstructed eukaryotic plasmid encoded Scy1 178
5.2.16 Investigation of intestinal histomorphology 171 5.3 Results 172 5.3.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 172 5.3.2 Construction of eucaryotic expression plasmid for Scy1 175 5.3.3 Construction of recombinant attenuated S. flexneri Fwl01 and S. flexneri typhimurium X3730/X4072 178 5.3.4 In vitro expression of recombinant attenuated S. flexneri and S. flexneri 178 5.3.5 In vitro expression of the reconstructed eukaryotic plasmid encoded Scy1 in EPC 179
5.2.16 Investigation of intestinal histomorphology 171 5.3 Results 172 5.3.1 Construction of procaryotic expression plasmid for antimicrobial peptide from marine animal 172 5.3.2 Construction of eucaryotic expression plasmid for Scy1 175 5.3.3 Construction of recombinant atteuated S. flexneri Fwl01 and S. typhimurium X3730/X4072 5.3.4 In vitro expression of recombinant attenuated S. flexneri and S. typhimurium carried prokaryotic expression plasmid 178 5.3.5 In vitro expression of the reconstructed eukaryotic plasmid encoded Scy1 in EPC 179 5.3.6 Antimicrobial activity of EPC transfected with pCMV-ScySP-HA-Scy1-asd

5.3.9 Safety of attenuated S. typhimurium and S. flexneri to black porgy	183
5.3.10 Optimum condition for oral administration of attenuated bacteria	183
5.3.11 The ability of attenuated S. typhimurium to colonize different tissues	185
5.3.12 In vivo transcription of recombinant attenuated S. typhimurium c	arried
Scy1 prokaryotic plasmid	189
5.3.13 In vivo Scy1 protein expression analysis of recombinant attenua	ted S.
typhimurium carried eukaryotic plasmid	190
5.3.14 Effect of Scy1 recombinant attenuated S. typhimurium on physiol	ogical
parameters	191
5.3.15 The immune-related genes responses after oral treatment with	Scy1
recombinant attenuated S. typhimurium	193
5.3.16 Effect of Scy1 recombinant attenuated S. typhimurium on inte	estinal
histomorphology	194
4 Discussion	••197
5.4.1 Attenuated S. typhimurium acted as a delivery vector for antimic	robial
peptide	197
5.4.2 Balanced lethal host plasmid expression system	199
5.4.3 Strategies for heterologous antigen delivery by S. typhimurium	200
5.4.4 Intestine colonization and tissue distribution of attenuated S. typhim	urium
	201
5.4.5 In vivo and in vitro expression of target protein by attenuat	ed S.
typhimurium ·····	203
5.4.6 Effect of Scy1 recombinant attenuated S. typhimurium on inte	estinal
histomorphology and physiological parameters	204
5.4.7 Effect of Scy1 recombinant attenuated S. typhimurium on the express	ion of
immune-related genes	207
Reference	

Degree papers are in the "Xiamen University Electronic Theses and Dissertations Database". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on http://etd.calis.edu.cn/ and submit requests online, or consult the interlibrary loan department in your library.

2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.