



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The Impact of Educational Attainment on Black Women's Obesity Rate in the United States

Citation for published version:

Curry, GD 2019, 'The Impact of Educational Attainment on Black Women's Obesity Rate in the United States', *Journal of racial and ethnic health disparities*, vol. 7, no. 2, pp. 345-354.
<https://doi.org/10.1007/s40615-019-00663-z>

Digital Object Identifier (DOI):

[10.1007/s40615-019-00663-z](https://doi.org/10.1007/s40615-019-00663-z)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of racial and ethnic health disparities

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Author: Dr. Gwenetta Denise Curry

Title: The Impact of Educational Attainment on Black Women's Obesity rate in the United States

Affiliation: Research Fellow at the University of Edinburgh and Assistant Professor at The University of Alabama

Address: 101 Manly Hall

Tuscaloosa, Alabama 35487

Email: dr.gdc Curry@gmail.com

Phone: 618-203-6399

ORCID: [0000-0003-3427-6630](https://orcid.org/0000-0003-3427-6630)

Abstract: Over the last two decades, Black women have been disproportionately impacted by the obesity epidemic in the United States. According to the Centers for Disease Control and Prevention (CDC), 56.6% of Black women are overweight or obese compared to 44.4% Hispanic and 32.8% of white women. Social scientists and public health researchers have argued that increasing educational attainment would lead to overall improvements in health outcomes. Using the National Health and Nutrition Examination Survey (NHANES), a nationally representative cross-sectional survey, Cycles 1999-2010, I examined how educational attainment impacts Black women's rate of obesity and c-reactive protein levels (N= 2,685). Multiple linear regression was used to analyze the association between body mass index (BMI) and educational attainment. C-reactive protein, inflammation response, was used to measure the body's reaction to being exposed to stress. The results demonstrated that educational attainment among Black women does not decrease their risk of being obese or levels of c-reactive protein. This article provides evidence to support a need to increase awareness of health disparities that disproportionately impact Black women.

Keywords: Health Disparities, Race, Obesity, Gender, Education

Acknowledgments: I would like to thank my husband, Dr. Tommy J. Curry, for his patience and extraordinary support while conducting this research. Special thank you to Dr. Alex McIntosh for your constant support and encouragement with my many projects. My mother, father aunts, and uncles for encouraging me to use my dedication to research in improving the lives of Black people.

Bio: Dr. Gwenetta Denise Curry, a research fellow at the University of Edinburgh and Assistant Professor in Gender and Race Studies at The University of Alabama. Her current areas of study include Critical Race Theory, Africana Womanism, Black Family Studies, Black Male Studies, Medical Sociology, and Food Insecurity. Her previous research investigated the relationship between educational attainment and body mass index among Black women.

Introduction

Over the last decade, obesity has become a major pandemic on a global scale driven by local markets and changes in food distribution [26]. According to the Center for Disease Control and Prevention (CDC), 39.8% of American adults are considered obese and affects 93.3 million U.S. adults in 2015-2016 [9]. Healthy People 2020 has set a target to reduce the number of adults who are considered obese to 30% by the year 2020 [12]. Ogden and others found that among men, 27.4% with a college degree are obese compared with 32.1% of those with less than a high school education, although these differences were not statistically significant [31]. However, there is a trend showing that women with college degrees were less likely to be obese than those without a college degree [31]. National data indicate that the obesity epidemic disproportionately impacts black women compared to their male and white male and female counterparts [24, 29]. The research on the contributing factors of obesity among Black women has been inconsistent [4]. Black women's economic disadvantages impact their access to resources, and stress levels have a negative impact on their health outcomes [9]. While there are many contributing factors to the condition of Black women, studies no longer deny the impact that one's environment has on health outcomes. Black women's biological response to stress plays a major role in obesity rates and obesity-related illnesses in this group [16]. Elevated concentrations of c-reactive protein have been seen in adults with exposure to stress associated with socioeconomic disadvantages [7, 28, 32], and there is a confirmed link between CRP and obesity [28, 5, 15]. More research is needed to evaluate the impact education has on obesity rates among Black women and the influence of stress on this relationship.

This study aims to examine the association of educational attainment, c-reactive protein, and obesity among Black women in the United States. Understanding the factors that affect obesity risk among Black women is important because four out of every five Black women are overweight or obese [7], and there are serious health-related consequences for individuals and their families, including the development of cardiometabolic disease(s) and lower quality of life. Black women have higher levels of degree attainment than their Black male counterparts [19]; it is critical to understand the relationship between education and obesity.

Education and Health

The idea that years of higher levels of education might protect against obesity is generally consistent with a larger body of research on socioeconomic status and health. For example, several studies have uncovered an inverse relationship between socioeconomic status and various indicators of physical health [1, 2, 17, 29]. Although education is sometimes included in composite measures of socioeconomic status, researchers have not begun to understand the unique ways that higher levels of education

influence obesity among Black women. Previous research has shown that there is a relationship between psychosocial stressors, such as racism, to health outcomes among Blacks [5]. It remains clear that prolonged exposure to stress has an impact on health outcomes, and Geronimus' [16] research supports this claim by showing that Black women aged more rapidly than whites due to stress. Black women's subjective experience of racism- the stigmas others have of them- can pose serious health threats and appear to be related to obesity and weight gain. Despite these well-established facts, medical sociologists continue to focus on race as only a demographic measure without considering the multiple effects impact that racial inequality has on the overall health outcomes of Blacks.

Previous research by Holmes and others [19, 22-23], focused on educational differences in obesity among blacks, as a part of their larger study of the links between educational attainment and obesity across racial/ethnic groups. These investigators argued that educational attainment reduces the rate of obesity through several contributing factors. These factors include differences in income, occupation, and access to resources that are necessary to live a healthy lifestyle. Holmes and Zajacova used the National Health Interview Survey 1997-2012 to determine if whites experience greater benefits from educational attainment than other minorities [19]. They found that higher educational attainment affects the health of whites more than minorities, even when potential sociodemographic, behavioral, and economic mediators are considered.

Supporting research by Zhang and Wang demonstrated that while obesity increased in all race-sex groups from 1971 to 2000, white women had a clear inverse association between obesity and educational attainment over time, and white men in the low socioeconomic status group experienced a decrease in obesity from 1999 to 2002 [42]. In the same study, among Black women, the association between obesity and education switched from an inverse relationship to the medium SES group having the highest prevalence of obesity by 1999. Black women at every educational level are disproportionately affected by the obesity epidemic compared to their white counterparts [20]. Jackson et al. found that Black women with greater than a high school education had substantially higher BMIs than white women with less than a high school education [20].

The National Health and Nutrition Examination Survey (NHANES) data from 1976 to 2004, African American women age 20 to 74 obesity rates increased from 15.6 percent to 20 percent when compared to white women overtime [14]. The longitudinal data on the development of obesity indicates that the rise in obesity rates in women has been the greatest in African-American women, followed by Hispanic women, then white women [10]. Obesity was significantly more prevalent among black men as of NHANES II but by a much smaller margin than women in the survey [14].

With over half the Black women in America being considered overweight or obese [10, 23], scholars must consider the possibility that some psycho-social factors are impacting Black women differently than other populations. Among men, there was little difference in the prevalence of obesity by race/ethnicity [18]. Among women, however,

the overall 1999-2010 prevalence among non-Hispanic Black (51%) was 10% higher than that among Mexican-Americans and 20 percentage points higher than that among non-Hispanic whites [18]. According to the CDC, “Black women with a bachelor’s degree had a median income of \$33, 877, which was 111 percent of the \$30, 413 median income figures for non-Hispanic white women who held a college degree. It is clear then that the strong income performance of black colleges is largely due to the earnings performance of black women, while higher education has failed to produce similar income gains for black men in comparison to white men [19].” Although these figures show Black women with bachelor’s degrees making slightly more money than white women with the same education, it does not tell the complete story. Compared with white women, Black women possessed less access to education, high paying occupations, and were less able to maximize those resources given the societal structure [26]. Fewer resources and larger amounts of debt contribute to the difficulty Black women have with upward mobility [5]. The lack of social mobility among Black women increases their proximity to poverty and decreases their access to healthy foods and safe places to exercise [5]. In combination, these conditions make Black women more vulnerable to negative health outcomes.

Previous studies have shown that individuals with lower socioeconomic status have high levels of stress and were at greater risk of obesity compared to individuals with high socioeconomic status [9, 27]. Recently, Black women have become the most educated population in the United States, but their health outcomes have not followed the same trajectory. There are mediating factors that need to be investigated to gain a clear understanding of what is preventing Black women from having better health outcomes with higher socioeconomic status.

Methods

Dataset

The National Health and Nutrition Examination Survey (NHANES) is a program of studies designed to assess the health and nutritional status of adults and children in the United States beginning in 1960 [15]. NHANES is a major program of the National Center for Health Statistics as a part of the Centers for Disease Control and Prevention and has the responsibility for producing vital and health statistics for the Nation. “The survey combines an in-home interview and a standardized physical examination at a mobile examination center (MEC). In each survey, a nationally representative sample of the US civilian noninstitutionalized population was selected using a complex, stratified, multistage probability cluster sampling design [7].”

Beginning in 1999, NHANES became a continuous survey without a break between cycles to focus on a variety of health and nutrition measurements to meet emerging needs [7]. The procedures followed in selecting the sample and conducting the

interview and examination were similar to those for previous surveys. Household interviews and physical examinations were conducted for each survey participant. During the physical examination, conducted in mobile examination centers, height and weight were measured as part of a more comprehensive set of body measurements [11]. These measures were taken by trained health technicians, using standardized measuring procedures and equipment. The NHANES examines a nationally representative sample of roughly 5,000 persons each year [11]. The participants were in counties across the country, 15 of which are visited every year.

Participants

This study will focus on Black women age 20-84 who participated in the Continuous NHANES from the years 1999 -2010. Observations for pregnant women and women missing a valid height or weight measurements were excluded from the data analysis. For the current research, the final NHANES analytical sample consisted of 3,488 Black women over the age of 20 years.

Measurements

Dependent Measures

Body Mass Index and Obesity

Participants' weight and height were measured in a mobile examination center using standardized techniques and equipment. Body mass index (BMI) was expressed as weight in kilograms divided by the square of height in meters. Adults are commonly classified as overweight (BMI 25.0-29.9), obese (BMI greater than or equal to 30.0), and extreme obesity (BMI greater than or equal to 40) among adults (age 20 years and over). Body Mass Index will be used to measure obesity in three different ways: Continuous, dichotomous (0=Not Obese, 1= Obese), and Categorical (1=underweight, 2= Normal, 3= Overweight, 4= Obese).

Independent Measures

Educational attainment

Education will be used to measure socioeconomic status (SES) and has been suggested by many scholars to be the most stable and robust indicator of SES. Other SES variables, such as income and occupation, vary over time and have a lower response rate than education items. Educational attainment will be measured using the question "What is the highest grade or level of school completed or the highest degree received?" Education was coded as a categorical variable where 1= Less than High School, 2=High School, 3 Some College or Associate, 4= College Graduate or Greater. The reference group will be those with "high school" education. Education will be treated as an independent variable rather than a moderator when testing the hypotheses.

Serum C-reactive protein

Participants' serum C – reactive protein measures were taken in the mobile examination center with values ranging from 0.01 to 18.01 mg/dL. C-reactive protein measures inflammation response, and participants with levels greater than or equal to 0.3 indicate high risk. Analysis will be measured two ways: continuous and dichotomous (0 = <0.3; 1 = ≥0.3).

Control Measures

Food Insecurity

Household food security status was based on the NHANES household food security category designation, which was determined using the 18-item U.S. Food Security Survey Module known as the “Core food security module.” Only one respondent per household answered the Module, even if more than one family is living in a household. Screening into the module depended on responses to previous questions, and those households screened out were assumed to have negative responses to all module questions. Households that give uniformly negative responses to the early questions will have a very low likelihood of having experienced any conditions of food insecurity and can safely be deemed to be food secure [10]. The two levels of screening are as follows: The First-level screen, including optional Q1. “We always have enough to eat and the kinds of food we want” and respond “never true” (or “don’t know” or “refuse” to all five of the questions Q2 to Q6 [10]. Second level screening for households not previously screened out responded “never true” (or “don’t know” or “refuse”) to “The children were not eating enough because we just out couldn’t afford enough food” were omitted. Food security question referred to circumstances over the 12 months preceding the survey. The NHANES categories are as follows: “fully food secure,” “marginally food secure,” “food insecure without hunger,” and “food insecure with hunger.” Participants were considered “fully food secure” if they had no affirmative response in any items from the Food Module. Participants who were considered “marginally food secure” answered affirmatively to 1 or 2 items in the module. The participants considered having “low food security” answered affirmatively to 3-5 items in the module. The participants who were “very low food security” answered 6 to 10 items in the module in the affirmative.

Hours Worked

Hours worked was measured through a series of questions related to the number of hours worked at all their jobs or businesses in the last week. (Continuous numbers)

Physical Activity

Physical activity was measured with items concerning daily activity and activity over the last 30 days. If they did not participate in vigorous or moderate activity, their response

was coded as 0=No and 1=Yes if they exercised. Responses of “Unable to do activity,” “Refused,” “Don’t Know,” coded as missing (.).

Food Measures

Twenty-four-hour dietary recall interviews were conducted where foods and beverages consumed the previous 24 hours ending at midnight were solicited and recorded using the standardized Automated Multiple Pass Method (AMPM) [23]. “With the exception of a few entry screens, all of the Dietary Recall section of the interview was collected using the AMPM program and all of the post-recall session is collected using the wrapper program. The 24hr dietary recall collects a list of all the foods and beverages consumed within a 24hr period; the time of consumption and the name of the eating occasion; detailed food descriptions and amounts of the reported foods; where it was obtained; and whether it was eaten at home or not. The recall data is coded and linked to the database of foods and their nutrient composition. Calculations of total daily nutrient intakes were derived from these data” [10].

Additionally, Food measure guidelines were given to participants for assistance in estimating portion sizes during the interview. Fruit intake will be measured using a dummy variable (0= No fruit, 1=Fruit). Vegetable intake will be measured using a dummy variable (0=No Vegetable 1=Vegetable).

Income

Participants were asked to disclose information regarding their total annual family income. “National Center for Health Statistics used the U.S. Bureau of Census Current Population Survey definition of “family” to household group members into one or more families. The CPS defines a family as “a group of two people or more (one of whom is the householder) related by birth, marriage or adoption and residing together”; all such people (including related subfamily members) are considered to be members of one family” [7]. Total family income reported as a range of values in dollars was measured on a 15-point scale (1= <4,999 and 15= >100,000). The income groups were combined to generate three income categories (1=<20,000, 2= 21,000-54,999 and 3=55,000 to >100,000).

Age

Participants were asked, “Best age in years of the sample person at the time of HH screening. Individuals 80 and over were top coded at 80 years of age [11].” Age will be broken down into three different groups 1= 20-39, 2= 40-59, 3= 60-80. Age was also measured as a continuous variable.

Marital Status

Marital status measured as a (0= Not Married and 1= Married/Living with a partner). The not married category will include divorced, widowed, separated, and never married. Participants who responded “refused” or answered “don’t know” were coded as missing. The “Married” category will be the reference group.

Statistical Analysis

Data will be analyzed using Stata version 13.1, College Station, Texas, for Windows software programs. All analyses included sample weights that account for the unequal probabilities of selection, oversampling and nonresponse. To account for the complex survey design survey set (SVYSET), primary sampling unit (PSU), and stratified sampling (STRATA) codes were used for unequal probabilities of sample selection in NHANES.

Descriptive statistics were calculated for sociodemographic characteristics, food insecurity and C-reactive protein, waist measurements, body mass index, and weighted with the sample weights calculated for the 12-year sample in NHANES. Multiple linear regressions will be used to analyze the association between BMI, waist, and education. Also, some of the regression models will contain interaction terms to test whether education moderates the effect of c-reactive protein on obesity.

To test for multicollinearity, I ran the variance inflation factor analysis and found all the results to be below 10. Indicating there was not a problem with variance inflations.

Results and Discussion

Descriptive Analysis

Table 1 is a descriptive table that shows the makeup of the participants in the dataset. The mean/percentages and standard deviations for the overall sample of Black Women by education group--less than high school, high school, some college & associate, and college. Most of the Black women in the study had Some College or Associate Degree education 32.4 percent compared to 31.1, 22.6, and 13.9 percent for Less Than High School, High School, and College, respectively. The overall average age is 47.1 years old. The average age across education groups is 36.4, 41.2, 45.1, and 44.9 for the Less Than High School, High School, Some College and Associate, and College-educated, respectively.

The average overall body mass index was 31.1 kg/m², and across education groups, the numbers exceed 30.0; 30.2, 31.3, 34.0, and 30.2 kg/m² for Less than High School, High School, Some College, and College, respectively. These results demonstrate that for all levels of education, the average BMI is in the obese range. The Some College and Associate Education group had the highest average BMI, which is very different from what other studies have found with white populations. According to the NHANES Survey from 2005 to 2008, white women with less than high school education had the highest rate of obesity (42.1%) compared to 21.8% with a college degree [30].

The results show that overall, 50.8 percent of Black women in this study are considered to be obese, and 27.4 percent are overweight. Black women with Some College and Associate degrees had the highest percentage of obese women, with 56.5

percent being categorized as obese compared to 46.9, 50.0, and 46.3 percent for Less than High School, High School, and College, respectively. College-educated Black women had the highest percentage of women who were considered “normal” weight, 26.9% compared to 21.8, 23.1, and 21.8 percent for Less than High School, High School, and Some College & Associate, respectively.

The overall waist circumference for Black women was 97.0cm. Across education groups, the average waist circumference was 94.3, 96.9, 102.0, and 95.3cm for Less than High School, High School, Some College & Associate, and College, respectively. Similar to what was seen with the average BMI scores, waist circumference was the highest for the Some College and Associate education groups. Forty percent of Black women in this study participated in some moderate or vigorous physical activity in the past thirty days. Black women the Some College & Associate degrees had the highest participation in physical activity with 51.2% participation compared to 28.3%, 40.9%, and 47.7% for Less than High School, High School, and College, respectively. C-reactive protein levels decreased as the level of education increased but remained in the high-risk range after obtaining a College Degree. Black women with less than a High School education had the highest C-reactive protein level 0.70 compared to .66, .57, and 0.53 for the High School, Some College or Associate, and College degree, respectively. The majority of the participants had a high-risk c-reactive protein level, 53.7 percent, compared to 46.2 percent. Black women with a college degree had the lowest percentage with high-risk C-reactive protein levels, 49.3 compared to 56.6, 55.9, and 51.3 for less than high school, high school, and some college or associate, respectively.

Majority of the population worked less than 40 hours a week (65%), and those with less than a High School education had the highest percentage working under 40 hours, 80.9 percent compared to 64.2, 58.4, 47.3 for High School, Some College or Associate, and College-educated, respectively. Black women with a college degree had the highest percentage of working more than 55 hours (15%) and 40 hours a week (52%). Interestingly, work hours increased as the level of education increased, demonstrating that education doesn’t protect Black women from working long hours.

The majority of the Black women in this study made between 20,000 to 54,000 dollars in the last year. As expected, Black women with less than a High School education had the highest percentage of women making less than 20,000 dollars in the last year. Also, Black women with college degrees had the highest percentage of women making more than 50,000. The percentage of women making between 20,000 to 54,000 dollars was practically the same for the High School, Some College & Associate, and College education women, 46.7%, 46.2%, and 46.6%, respectively.

Overall, 36.3 percent of the Black women were either married or cohabitating with a partner. Across education groups, the majority of the women were not married or cohabitating, and the women with Less than High School education had the highest percentage not married, 68.9% compared to 63.8%, 62.6%, and 55.5% for High School, Some College or Associate and College educated respectively. The College-educated women had the highest percentage of marriage 44.5% compared to 31.1%, 36.2%, and

37.4% for the Less than High School, High School, and Some College and Associate, respectively.

Fruit and Vegetable consumption was rather low across all groups. The lowest fruit consumption was found among the population with Less than High School population, 91.1 percent did not consume any fruit during the 24 hr dietary recall. The High School educated women had the highest percentage of fruit consumption 10.3 percent compared to 8.9%, 6.9% 6.5% for the Less than High School, Some College and Associate and College-educated, respectively. Vegetable consumption was much higher than fruit consumption. Overall 21.9% consumed vegetables during the 24-Hour dietary recall. Black women with Some College or Associate degrees had the highest consumption of vegetables 24.2% compared to 20.4%, 19.5%, and 23.7% for the Less than High School, High School, and College educated respectively.

The majority of the population was considered to be Fully Food Secure with College-educated Black women having the highest percentage of participants who were Fully Food Secure, 83.5% compared to 63.8%, 64.5% and 69.5% for the Less than High school, High School, and Some College & Associate, respectively. Overall only 7.3% of the total group was considered to be Food Insecure with hunger.

Table 1: Descriptive Statistics

	Education Category				
	Overall	Less Than HS	High School/GED	Some College or Associate	College Graduate
Less than High School	31.1%				
High School/GED	22.6%				
Some College or Associate	32.4%				
College Grade+	13.9%				
BMI (kg/m**2) (ave)	31.1	30.2	31.3	34.0	30.2
Waist Circumference (cm)	97.0	94.3	96.9	102.0	95.3
BMI Category					
Underweight	0.8%	1.2%	0.8%	0.6%	0.7%
Normal Weight	21.0%	21.8%	23.1%	21.8%	26.9%
Overweight	27.4%	30.1%	26.2%	21.2%	26.1%
Obese	50.85	46.9%	50.0%	56.5%	46.3%

CRP (%)					
Low Risk	46.2%	43.4%	44.1%	48.8%	50.7%
High Risk	53.7%	56.6%	55.9%	51.3%	49.3%
Age (ave yrs)	47.1	36.4	41.2	45.1	44.9
Work Hours (ave)	22.8	13.9	22.9	25.5	32.8
Work Hours (%)					
<40hrs	65.4%	80.9%	64.2%	58.4%	47.3%
≥40hrs	34.6%	19.1%	35.8%	41.6%	52.7%
<55hrs	92.5%	97.2%	92.8%	91.4%	84.0%
≥55hrs	7.47%	2.8%	7.2%	8.7%	16.0%
Marital Status (%)					
Not Married	63.8%	68.9%	63.8%	62.6%	55.5%
Married/cohabitating	36.3%	31.1%	36.2%	37.4%	44.5%
Income (%)					
<\$20,000	39.9%	54.4%	38.6%	35.9%	14.3%
\$20,000 to \$54,000	42.8%	33.4%	46.7%	46.2%	46.6%
>\$55,000	17.3%	12.2%	14.7%	17.9%	39.2%
Physical Activity (%)					
No	58.6%	71.8%	59.2%	48.8%	52.3%
Yes	41.4%	28.2%	40.9%	51.2%	47.7%

Food Security (%)					
Fully Food Secure	68.5%	63.8%	64.5%	69.5%	83.5%
Marginally Food Secure	13.5%	13.9%	15.8%	13.7%	8.3%
Food Insecure w/o Hunger	10.6%	13.1%	11.2%	10.2%	5.1%
Food Insecure w/hunger	7.3%	9.2%	8.5%	6.6%	2.9%
Fruit Intake					
No	91.8%	91.1%	89.7%	93.1%	93.5%
Yes	8.2%	8.9%	10.3%	6.9%	6.5%
Vegetable Intake					
No	78.1%	79.6%	80.5%	75.8%	76.3%
Yes	21.9%	20.4%	19.5%	24.2%	23.7%
Number of Observations	N=2,658	N=826	N=600	N=862	N=370

Table 2 contains six different models measuring variables association with body mass index of Black women. Model one indicates that relative to Black women with a college degree, women with some college education were more likely to have a higher body mass index. Model 2 shows that c-reactive protein is positively associated with body mass index with a p-value of 0.00. Model 3 includes education and c-reactive protein. The results show when education and c-reactive protein are in the model together, the women with Some College and Associate Education remained statistically different from those with College Education. Also, high-risk c-reactive protein continued to be significantly different from low-risk c-reactive protein in a positive direction. In Model 4, the control variables married, age and cycle intake were all added to the equation along with c-reactive protein (a measure of stress). Model 5 included Food Security, Fruit, and Vegetable, and Physical activity. Model 6 added the education, c-reactive protein variable.

Body mass index for Black women with some college or associate education continued to differ significantly from those having a college degree. The results show that age was significantly associated with Body Mass Index. Also, serum C-reactive protein level ≥ 0.3 mg/dl (High Risk) was significantly associated with body mass index with a p-value ≤ 0.01 . Stated another way, exposure to stress has a significant positive impact on body mass index compared to the reference group of C-reactive protein level < 0.3 (Low Risk).

Model 5 included all of the measures for the previous four models with the addition of Food Security, Fruit and Vegetable Intake, and Physical activity. Participants with some college or associate degree were not significantly associated with body mass index. Compared to the reference category of Fully Food Secure, being Marginally Food Secure was not associated with body mass index (p-value < 0.01). There was a statistical difference between the reference groups and “Food Insecure w/ Hunger” and “Food Insecure w/o hunger.” Participants being physically active and eating fruits and vegetables was not statically different from the reference categories. The findings indicate that these positive health behaviors did not have an association with body mass index. These results do not imply that healthy behaviors (physical activity and fruit and vegetable consumption) could not potentially have a positive impact on maintaining healthy body weight. Instead, these results, confirm the finding of previous work by Cozier and other [13] demonstrating that more efforts should be made to understand how psychosocial factors (e.g., stress, experiences of racism, sexism, and discrimination) impact health outcomes in Black women. Said differently, while exercise and healthy lifestyle habits should be recommended for Black women, these findings suggest that stress must also be considered as a factor increasing BMI and decreasing the benefits of exercise and diet.

Model 6 included all the measures for the previous five models with the addition of interaction variables. Compared to the reference category, Black women with some college or associate degree was significantly associated with body mass index

(p-value<0.01). There was no statistical difference between education and c-reactive protein interaction. Food insecurity with hunger and with hunger were both statistically significant. Compared to the reference group working long hours was significantly associated with BMI. These results mirror that of other research findings showing long work hours increase not only stress levels but also reduce the time one has for physical activity and healthy eating [26-27].

Table 2: Linear Regression Body Mass Index of Black Women in the National Health and Nutrition Examination Survey Cycles 1999-2010

	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
	Education	CRP	Education and CRP	Control	Food	Interaction
	b/se	b/se	b/se	b/se	b/se	b/se
Less than High School	0.998		0.402	3.168**	0.341	0.875
	-0.58		-0.55	-1.08	-0.58	-0.6
High School	0.95		0.446	2.843**	0.358	0.419
	-0.57		-0.55	-1.05	-0.59	-0.59
Some College or Associate	1.483**		1.171*	3.124**	1.039	1.257*
	-0.56		-0.54	-1.05	-0.56	-0.56
High-Risk CRP		7.103***	7.100***	14.694***	7.094***	7.591***
		-0.34	-0.34	-0.68	-0.35	-0.74
Married/cohabitating				0.366	-0.06	-0.072
				-0.62	-0.32	-0.32
Age				0.103***	0.012	0.012
				-0.02	-0.01	-0.01
NHANES Cycle				0.674***	0.198	0.201
				-0.2	-0.1	-0.1

Vegetable Intake	-0.24	-0.246
	-0.43	-0.43
Fruit Intake	0.536	0.543
	-0.51	-0.51
Physical Activity	-0.285	-0.275
	-0.31	-0.3
Food Security		
Marginally Food Secure	0.302	0.293
	-0.36	-0.36
Food Insecure w/o Hunger	1.641**	1.648**
	-0.58	-0.58

Table 2 Continued

	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
Food Insecure with Hunger					1.987**	1.970**
					-0.6	-0.61
Work Hours					0.018*	0.018*
					-0.01	-0.01
Less than High School X CRP						-1.058
						-0.95
High School/ GED X CRP						-0.186
						-0.84
Some College or Associate X CRP						-0.48
						-0.92
constant	30.649***	27.957***	27.342***	82.732***	25.722***	25.500***
	-0.46	-0.18	-0.42	-1.35	-0.72	-0.76
Observation	2,597	2,597	2,597	2,564	2,564	2,564
R2	0.003	0.177	0.18	0.206	0.189	0.19
Degrees of Freedom	90	90	90	89	89	89

Standard Errors in Parenthesis

***P<0.00, **P<0.01, *p<0.05

In summary, other studies found there to be a gradient decrease in body mass index and the risk of obesity as education increases [20, 23, 38]. Educational attainment has been linked directly to health outcomes among white populations. The current data challenges this view by demonstrating there is not a gradient decrease in obesity and body mass index with educational attainment among Black women. Black women with some college and associate education were more likely to be overweight and had a higher risk of obesity compared to women with a college degree.

Also, these results support the previous literature by Brooks and others [4,7-8, 38], demonstrating that C-reactive protein has a strong association with body mass index and increases the odds of being obese when levels are $\geq 0.3\text{mg/dL}$ for Black women.

Conclusion: While other populations' obesity rates have remained steady over the years, Black women's rate has increased and continues to outnumber their white female and male counterparts. Educational attainment has long been touted as a solution to decreasing negative health outcomes. This research challenges the current understanding of the relationship between educational attainment and obesity. My research sought to 1) analyze the obesity trends among Black women from 1999-2010 by education and age, 2) gain a better understanding of the psycho-social factors contributing to Black women's risk of obesity across levels of education, and 3) research the impact of stress on Black women's risk of obesity.

In the current study, Black women with some college or associate degree were more likely to be overweight or obese compared to other education groups. While the causes of this increased risk are unclear, one possible explanation is women with some college education are working multiple jobs, which can result in limited time for food preparation and exercise. Cozier's [13] study found a strong association between experiences of discrimination and obesity rates. While the current study did not measure discrimination and racism, stress was measured using serum C-reactive protein levels. In line with other research, body mass index and being obese was strongly associated with C-reactive protein levels equal to or greater than 0.03 mg/dl . Waist circumference also had a strong positive association with C-reactive protein levels. Numerous possible factors could contribute to Black women's stress levels, but that data was not available for the study. However, the research performed by others demonstrates that Black women experience racism and discrimination. Perceived discrimination based on race and gender are key factors in chronic stress-related health disparities among ethnic/racial and other minority groups [19, 30]. Cozier's research suggests workplace and community-based programs to combat racism and interventions to reduce racism-related stress could be important components of strategies for the prevention of obesity [13]. Her findings also suggest that experiences of racism possibly explain why the obesity epidemic disproportionately impacts US Black women. Other studies have linked maternal stress as a potential explanation for excess preterm delivery among Black women because of exposure to racism associated stress [13].

Broader Implications: Being overweight or obese can lead to several life-threatening diseases and a lower quality of life. According to the CDC, obesity has been linked to increased risk of

hypertension, diabetes, high cholesterol, stroke, and some cancers. High rates of obesity will have an overall increase in the cost of medical care and disability services provided by federal funds.

Conflicts of Interest: This is an original article with no known conflicts of interest.

References

1. Adler, Nancy and David Rehkopf. U.S. Disparities in Health: Descriptions, Causes, and Mechanisms. *Annual Review Public Health*. 2008; 29:235-252.
2. Adler, Nancy and Alana Snibbe. The Role of Psychosocial Processes in Explaining the Gradient Between Socioeconomic Status and Health. *Journal of the Association for Psychological Science*. 2003;12(4):119-123.
3. Agyemang, Priscilla and Tiffany M. Powell-Wiley. Obesity and Black Women: Special Considerations Related to Genesis and Therapeutic Approaches. *Current Cardiovasc Risk Rep*. 2013; 7(5):378-386.
4. Alley, Dawn; Seeman, T.E.; Kim, J.K.; Karlamangla, Arun; Hu, Peifeng; Crimmins, E.M. Socioeconomic status and C-reactive protein levels in the US Population: NHANES IV.” *Brian, Behavior, and Immunity*. 2006;20(5): 498-504.
5. Assari, Shervin, Race. Intergenerational Social Mobility and Stressful Life Events. *Behavioral Science (Basel)*. 2018 Oct; 8(10):86.
6. Brondolo, E, LC Gallo, and HF Myers. Race, racism and health: disparities, mechanisms, and interventions. *Journal of Behavioral Medicine*. 2009 Vol. 32(1) 1-8.
7. Brooks, GC, Blaha, MJ, Blumenthal, RS. Relation of C-reactive protein to abdominal adiposity. *American Journal of Cardiology*. 2010. Vol. 106(1). 56-61.
8. Carpenter, LL, Gawuga, CE, Tyrka, AR, and LH Price. “C-reactive protein, early life stress, and wellbeing in healthy adults.” *Acta Psychiatr Scand*. 2012. Vol. 126(6): 402-410.
9. Center for Disease Control. Adult Obesity Facts. 2015-2016. (<https://www.cdc.gov/obesity/data/adult.html>). Accessed October 21, 2019
10. Center for Disease Control and Prevention. National Health and Nutrition Examination Survey: MEC In-Person Dietary Interviewers Procedures Manuel. 2006. (http://www.cdc.gov/nchs/data/nhanes/nhanes_05_06/DIETARY_MEC.pdf). Accessed April 22, 2016.
11. Centers for Disease Control and Prevention. Health data interactive. Atlanta: Center for Disease Control and Prevention, National Center for Health Statistics. 2014. (<http://www.cdc.gov/nchs/hdi.htm>). Accessed July 10, 2015.

12. Cossrow, N. a. Race/Ethnic Issues in Obesity and Obesity-Related Comorbidities. 2004. *The Journal of Clinical Endocrinology and Metabolism*. 2004; 2590-2593.
13. Cozier, Yvette, Jeffrey Yu, Patricia Coogan, Traci Bethea, Lynn Rosenberg, and Julie Palmer. Racism, Segregation, and Risk of Obesity in the Black Women's Health Study. *American Journal of Epidemiology*. 2014;179(7):875-883.
14. Department of Health and Human Services Healthy People 2020. Nutrition and Weight Status. Office of Disease Prevention and Health Promotion. 2016. Washington (DC): HHS; Available from: <https://www.healthypeople.gov/2020/topics-objectives/topic/nutrition-and-weight-status>. Accessed July 15, 2018.
15. Flegal, Katherine, Margaret Carroll, Cynthia Ogden, and Lester Curtin. Prevalence and Trends in Obesity Among US Adults 1999-2008. *The Journal of the American Medical Association*. 2010;303(3):235-241.
16. Geronimus, Arline, Margaret Hicken, Jay Pearson, Sarah Seashols, Kelly Brown, and Tracey Cruz. Do US Black Women Experience Stress-Related Accelerated Biological Aging? *Human Nat* 2010; 21:19-38.
17. Hamer, M, Stamatakis, E. 2008. Inflammation as an intermediate pathway in the association between psychosocial stress and obesity. *Physiol Behavior*. Vol 94. Pp. 536-539.
18. Heiland, M. A. Race, Obesity and the Puzzle of Gender Specificity. Working Paper Series (Federal Reserve Bank of Boston), 2008;1-103.
19. Hogue, Carol J Rowland, and j. Douglas Bremner. Stress Model for Research into Preterm Delivery Among Black Women. *American Journal of Obstetrics and Gynecology*. 2005;192(5): S47-S55.
20. Holmes, Christopher and Anna Zajacova. Education is the Great Equalizer: Health Benefits for Black and White adults. *Social Science Quarterly*. 2014;95(4):1064-1085.
21. Jackson, Chandra, Moyes Szklo, Hsin-Chieh Yeh, Nae-Yuh Wang, Rosemary Dray-Spira, Roland Thorpe, and Frederick and Brancati. Black-White Disparities in Overweight and Obesity Trends by Educational Attainment in the United States, 1997-2008. *Journal of Obesity*. 2013:1-9.
22. Journal of Blacks in Higher Education. *Journal of Blacks in Higher Education. News & Views*. 2007 http://www.jbhe.com/news_views/55_closing_the_income_gap.html Accessed March 9, 2015.
23. Marmot, Sir Michael. *The Status Syndrome: How Social Standing Affects Our Health and Longevity*. Bloomsbury Publishing Plc. London; 2005.

24. May, Ashleigh, David Freedman, Betty Lou Sherry, and Heidi Blanck. Center for Disease Control and Prevention. Morbidity and Mortality Weekly Report. 2013 <http://www.cdc.gov/mmwr/preview/mmwrhtml/su6203a20.htm> Accessed 10 July 2016
25. Meyer, I. H., Schwartz, S., & Frost, D. M. Social patterning of stress and coping: Does disadvantaged social statuses confer more stress and fewer coping resources? *Social Science and Medicine*, 2008; 67(3), 368–379.
26. Mirowsky, J., Ross, C. Education, personal control, lifestyle, and health: A human capital hypothesis. *Research on Aging*, 1998; 20:415-449.
27. Mirowsky, J., Ross, C. Education, Social Status, and Education. New York: Aldine de Gruyter, Inc. 2003
28. Moore, Carla, and Solveig Cunningham. Social Position, Psychological Stress, and Obesity: A Systemic Review. *Journal of the Academy of Nutrition and Dietetics*, 2012; 112:4:518-526
29. Moshfegh, AJ; Rhodes DG; Baer, DJ; Murayi, T; Clemens, JC; Rumpler, WV; Paul, DR; Sebastian, RS; Kuczynski, KJ; Ingwersen, LA; Staples, RC; and LE Cleveland. The US Department of Agriculture Automated Multiple-Pass Method reduces bias in the collection of energy intakes. *American Journal of Clinical Nutrition*. 2008;88(2):324-332.
30. Ogden, Cynthia. Disparities in Obesity Prevalence in the United States: Black Women at Risk. *The American Journal of Clinical Nutrition*. 2009;89(4):1001-1002.
31. Ogden, Cynthia et al. Obesity and Socioeconomic Status in Adults: United States, 2005-2008. National Center of Health Statistics. 2010; 50:1-8.
32. Park, HS, Park JY, Yu R. 2005 Relationship of obesity and visceral adiposity with serum concentrations of CRP, TNF-alpha and IL-6. *Diabetes Res Clin Pract*. Vol 69 pp. 29-35.
33. Pollitt, RA, Kaufman JS, Rose KM, Diez-Roux, AV, Zeng D, Heiss, G. Early-Life and adult socioeconomic status and inflammatory risk markers in adulthood. *Eur J Epidemiol*. 2007; 22(1):55-66.
34. Swinburn, Boyd A; G Sacks; KD Hall; K McPherson; DT Finegood; M Moodie; SL Gortmaker. The global obesity pandemic: shaped by global drivers and local environments. *The Lancet*. 2011;378(9793): 804-814.
35. U.S. Census Bureau. Current Population Survey (CPS) Definitions and Explanations. Revised 1/20/04. <http://www.census.gov/population/www/cps/cpsdef.html>. Accessed 10 June 2019.

36. U.S. Department of Labor. Women in the labor force: A Databook (Report No.1011). Washington, DC 2008.
37. Van Dyke, ME., Vaccarino, V., Dunbar, S.B., Pemu, P., Gibbons, G.H., Quyyumi, A.A., and Lewis, T.T. 2017 "Socioeconomic status discrimination and C-reactive protein in African-American and White adults." *Psychoneuroendocrinology*. Vol 82. Pp 9-16
38. Wang, Youfa and May Baydoun. The Obesity Epidemic in the United States--Gender, age, socioeconomic, Race/Ethnic and Geographic Characteristic: A Systematic Review and Meta-Regression Analysis. *Epidemiol Review*, 2007; May 17, pp. 6-28
39. Williams, David, and Selina Mohammed. Discrimination and racial disparities in health: evidence and needed research. *Journal Behavior Med.* 2009;32(1):1-38.
40. Winkleby, M.A., Jatulis, D.E., Frank, E. and Fortmann, S.P. Socioeconomic Status and Health How Education, Income, and Occupation Contribute to Risk Factors for Cardiovascular Disease. *American Journal of Public Health*, 1992; 82, 816-820.
41. Wisenivesky, Juan, Jessica Lorenzo, Jonathan Feldman, Howard Leventhal, and Ethan Halm. The Relationship Between Perceived Stress and Morbidity among Adult Inner-City Asthmatics. *Journal of Asthma*.2010; 47(1):100-104.
42. Zhang, Qi and Youfa Wang. Socioeconomic inequality of obesity in the United States: do gender, age, and ethnicity matter? *Social Science & Medicine*. 2004; 58(6):1171-1180.
43. Zhang, Q, and Y Wang. Trends in the association between obesity and socioeconomic status in US adults: 1971 to 2000. *Obesity Research* 2004:1622-1632

Supplemental Data

Matrix of correlations

Variables	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
(1) BMI	1.000								
(2) Age	-0.044	1.000							
(3) Educ	0.066	-0.251	1.000						
(4) CRP	0.409	0.015	0.025	1.000					
(5) Physical Act	-0.022	-0.092	0.269	-0.052	1.000				
(6) Fruit and Veg	0.009	0.012	0.044	-0.014	0.030	1.000			
(7) Food Security	-0.010	-0.132	-0.021	-0.004	0.049	-0.032	1.000		
(8) Married	-0.067	-0.047	0.063	-0.135	0.033	0.014	-0.099	1.000	
(9) Smoker	-0.057	0.061	-0.030	0.008	0.007	-0.011	0.111	-0.139	1.000

Variance inflation factor

	VIF	1/VIF
education	1.881	.532
C-Reactive Pro	5.812	.172
Age	1.177	.849
Married	1.13	.885
smoker	1.107	.903
Family Inc	1.248	.801
Alcohol	1.12	.893
Physical Act	1.08	.926
Fruit and Veg	1.008	.992
Food Security	1.106	.904
2.Educ#c.CRP	2.977	.336
3.Educ#c.CRP	5.144	.194
4.Educ#c.CRP	2.435	.411
Mean VIF	2.094	.