

1 **Microbial ‘gardening’ by a seaweed holobiont: surface metabolites attract**  
2 **protective and deter pathogenic epibacterial settlement**

3

4 Mahasweta Saha<sup>1,2,3</sup> and Florian Weinberger<sup>1</sup>

5 <sup>1</sup>Benthic Ecology, GEOMAR Helmholtz Centre for Ocean Research, Düsternbrookerweg 20,  
6 24105 Kiel, Germany.

7 <sup>2</sup>School of Biological Sciences, University of Essex, Wivenhoe Park, CO4 3SQ, Colchester,  
8 United Kingdom.

9 <sup>3</sup>Current address: Marine Ecology and Biodiversity, Plymouth Marine Laboratory, Prospect  
10 Place, PL1 3DH Plymouth, United Kingdom.

11 Author for correspondence:

12 Mahasweta Saha

13 Email: [sahamahasweta@gmail.com](mailto:sahamahasweta@gmail.com)

14 Phone: +44 1752633415

15 **Running Head:** Chemical ‘gardening’ of beneficial epibacteria by an invasive seaweed

16 **Summary**

17 1. Epimicrobial communities on seaweed surfaces usually contain not only potentially  
18 pathogenic, but also potentially beneficial microorganisms. Capacity of terrestrial  
19 plants for chemically mediated recruitment i.e. ‘gardening’ of bacterial communities  
20 in the rhizosphere was recently demonstrated. Empirical evidence directly linking  
21 such chemical ‘gardening’ with the beneficial role of gardened microbes in terrestrial  
22 plants is rare and largely missing for aquatic macrophytes.

23 2. Here we demonstrate that our model invasive seaweed holobiont *Agarophyton*  
24 *vermiculophyllum* possesses beneficial microbiota on its surface that provide

25 protection from bacterial pathogens. Metabolites from the algal holobiont's surface  
26 reduced settlement of opportunistic pathogens but attracted protective epibacterial  
27 settlement.

28 3. We tested 58 different bacterial species (isolated from the surface of *A.*  
29 *vermiculophyllum*) individually in tip bleaching assays. *Kordia algicida* was identified  
30 as a 'significant pathogen' inducing a bleaching disease. In addition, 9 other species  
31 significantly reduced the risk of algal bleaching and were thus 'significantly  
32 protective'. Additionally, 2 'potential pathogens' and 10 'potential protectors' were  
33 identified. When 19 significant and potential protectors and 3 significant and  
34 potential pathogens were tested together, the protective strains fully prevented  
35 bleaching, suggesting that a component of *A. vermiculophyllum*'s epimicrobiome  
36 provides an associational defence against pathogens. Chemically mediated selective  
37 recruitment of microbes was demonstrated in bioassays, where *A. vermiculophyllum*  
38 surface metabolites attracted the settlement of protective strains, but reduced  
39 settlement of pathogens.

40 4. Synthesis: The capacity of an aquatic macrophyte to chemically 'garden' protective  
41 microorganisms to the benefit of strengthened disease resistance is demonstrated  
42 for the first time. Such a role of surface chemistry in 'gardening' of microbes as  
43 found in the current study could also be applicable to other host plant – microbe  
44 interactions. Our results may open new avenues towards manipulation of the  
45 surface microbiome of seaweeds via chemical 'gardening', enhancing sustainable  
46 production of healthy seaweeds.

47 **Key words:** *Agarophyton vermiculophyllum*, Macrophyte, Chemical Defence, Plant-microbe  
48 interactions, *Gracilaria vermiculophylla*, Bleaching, Gardening, Invasive, Seaweed,  
49 Holobiont.

50

## 51 **Introduction**

52 All eukaryotes including terrestrial plants and aquatic macrophytes are influenced by  
53 complex interactions with microbial communities. The animal gut microbiome is very well  
54 known to influence the health and nutritional status of its host (Hooper *et al.*, 2002; Flint *et*  
55 *al.*, 2012), ultimately forming a metaorganism or holobiont that consists of the host and  
56 associated microbiomes (Bordenstein & Theis, 2015). These microbes form an integral part  
57 of a plant or animal phenotype, influencing the fitness and ecological traits of their hosts.

58

59 The outer body surface is the primary physiological and ecological interface of multicellular  
60 aquatic organisms like water plants or seaweeds with the environment (Wahl, 2008). Apart  
61 from exchange and uptake of nutrients, this interface is involved in the exchange of  
62 chemical cues and signals that mediate the recognition of an organism by a partner, a  
63 parasite, an epibiont or a predator. This surface is often colonized by complex microbial  
64 communities, a biofilm-like epimicrobiome that has also been denoted as 'second skin'  
65 (Wahl *et al.*, 2012). Marine macroalgae i.e. seaweeds have an additional diffusive boundary  
66 layer (Hurd, 2000) along with their 'second skin' that serves as the micro-niche of chemically  
67 mediated ecological interactions. This micro-niche is analogous to the rhizosphere of plant  
68 roots (Hartmann *et al.*, 2009) or the phycosphere of phytoplankton (Bell & Mitchell, 1972).

69 This niche is an ecological interface of seaweed-microbe relationships, modulates most of  
70 the interactions between the seaweed host and the environment and is typically  
71 characterized by a specific chemical fingerprint.

72 Seaweeds are omnipresent organisms in photic coastal zones, play key roles in carbon  
73 fixation, biogeochemical cycling and food web formation. They can drive the biogeochemical  
74 pump and release climate cooling gases like dimethyl sulphide (Van Alstyne & Houser,  
75 2003). They act as nursery ground and protective shelters for many animals (Schiel & Foster,  
76 2006; Pereira *et al.*, 2017). Seaweeds also provide substrate for numerous sessile organisms,  
77 ranging from bacteria to macro-invertebrates (Wahl, 1989). Epibacteria that colonize the  
78 surfaces of seaweeds vary taxonomically with host, space and time (Cundell *et al.*, 1977;  
79 Lachnit *et al.*, 2011) and can affect the well-being of their host in multiple ways.

80 The epimicrobial communities on seaweeds consist not only of pathogenic species but also  
81 of potentially beneficial ones. Interactions with the surface epimicrobiome have the  
82 potential to influence seaweed health and development in two different ways: they can be  
83 detrimental, as seaweeds can be plagued by bacterial and eukaryotic pathogens (see  
84 Gachon *et al.*, 2010; Egan *et al.*, 2013 and references therein). The epimicrobiome also often  
85 provides inductive settlement cues to algal spores and invertebrate larvae, causing heavy  
86 detrimental fouling (see Wahl *et al.*, 2012 and references therein). Alternatively, seaweed  
87 epimicrobiomes can also be beneficial, supplying essential nutrients (see Hollants *et al.*,  
88 2013 and references therein) and chemical cues for morphogenesis (see Wichard *et al.*,  
89 2015 and references therein).

90 A suspected yet relatively undemonstrated beneficial role of the epimicrobiome is the  
91 protection from pathogens and other detrimental microorganisms (but see Longford *et al.*,

92 2019). A certain component of these epimicrobial communities on seaweed surfaces is quite  
93 host-specific (Lachnit *et al.*, 2009; Bengtsson & Ovreas, 2010) and the same is true for the  
94 rhizosphere of terrestrial plant roots (Raaijmakers *et al.*, 2009). However, the principles  
95 governing the assemblages of microbes on surfaces of seaweeds or any other aquatic  
96 macrophytes are unclear. Based upon recent independent studies with terrestrial plants and  
97 aquatic macrophytes the following models for the association of microbial communities  
98 have been proposed: 1. The 'neutral' hypothesis assumes that species are ecologically  
99 equivalent, and the community structure is determined randomly (Hubbel 2001, 2006;  
100 Woodcock, 2007). 2. The 'niche' model stresses that only microorganisms which are  
101 adapted to the specific conditions on a host surface will be able to settle on it (Dumbrell *et*  
102 *al.*, 2010). 3. The 'lottery' hypothesis combines both neutral and functional aspects and  
103 predicts that multiple microorganisms could make use of the same niche, but those that  
104 reach it first have a larger chance of settlement success (Burke *et al.*, 2011). 4. Untargeted  
105 recruitment of microorganisms by the host via the release of exuded nutrients has also been  
106 proposed, as well as targeted deterrence by processes like induced defence (Weinberger,  
107 2007).

108 By comparing bacterial root microbiomes between wildtype *Arabidopsis thaliana* and  
109 mutants that could not produce the defence phytohormone salicylic acid, Lebeis *et al.*,  
110 2015 recently demonstrated that salicylic acid signalling can modulate root microbial  
111 communities. While such studies on the role of chemical manipulation of root microbiota  
112 have started to appear for land plants, no parallel study exists for aquatic macrophytes that  
113 demonstrates an active 'deliberate' recruitment or 'gardening' of beneficial microbes.  
114 Surface associated metabolites may shape the microbial communities on seaweed surfaces.  
115 For example, halogenated furanones excreted by the host *Delisea pulchra* were

116 demonstrated to shape the microbiome of the seaweed (Longford *et al.*, 2019). Also, in the  
117 brown alga *Fucus vesiculosus* surface metabolites were found to have an effect on the  
118 biofilm composition both under field and lab conditions (Lachnit *et al.*, 2010). The authors  
119 used an experimental system that simulated the delivery of *Fucus* surface associated  
120 metabolites on artificial substrates and tested the effect of algal surface chemistry on  
121 bacterial community composition. Bacterial communities that developed on test surfaces  
122 loaded with *Fucus* surface metabolites were found to be quite similar to communities on the  
123 surfaces of *Fucus*, but different from communities on solvent controls, which hinted at the  
124 strong selective force of these surface metabolites of *Fucus*. However, for the investigation  
125 with *Fucus vesiculosus* no evidence could be demonstrated for the beneficial role of such  
126 microbes and thus the purpose of such chemically mediated recruitment of microbes. Also,  
127 studies of the rhizosphere of terrestrial plants already reported selective ‘gardening’ of  
128 microbes (Currier & Strobel, 1976; Bacilio-Jiménez *et al.*, 2003). For example, root exudates  
129 of different developmental stages of *Arabidopsis* promoted the formation of microbial  
130 communities with different compositions when the influence of environmental and soil  
131 edaphic factors was experimentally excluded (Yuan *et al.*, 2015). Although there have been  
132 demonstrations of the possible beneficial roles of such active microbial gardening for plant  
133 growth and development in terrestrial environments (Lebeis *et al.*, 2015) and marine  
134 environments (Kessler *et al.*, 2018), none of the studies in the aquatic realm have yet been  
135 able to empirically link chemically mediated microbial ‘gardening’ with resistance to disease.

136

137 Thallus whitening, bleaching or ‘ice-ice disease’ is a common problem in certain farmed and  
138 wild red seaweeds, such as *Gracilaria ‘conferta’* (Weinberger *et al.*, 1994; Weinberger *et al.*,

139 1997), *Kappaphycus* and *Eucheuma* or *Delisea pulchra* (Case *et al.*, 2011; Campbell *et al.*,  
140 2011). It was repeatedly shown that this depigmentation symptom can be induced by  
141 multiple opportunistic bacterial pathogens and in the case of *G. 'conferta'* a component of  
142 the microbiome was shown to prevent the disease. Also, in *Delisea pulchra* early  
143 successional epibacterial strains protected the host from a later successional strain that was  
144 pathogenic when the host microbiome was experimentally disturbed (Longford *et al.*, 2019).  
145 In the context of the 'gardening' hypothesis the present study investigated whether (a)  
146 epibacteria originating from healthy specimens of the invasive red seaweed *Agarophyton*  
147 *vermiculophyllum* can also induce thallus bleaching in *A. vermiculophyllum*, whether (b) a  
148 subset of epibacterial strains of the algal microbiome offers protection towards pathogenic  
149 strains and whether (c) *A. vermiculophyllum* has a capacity for chemically mediated  
150 recruitment of such protective microbes while deterring the settlement by pathogens.

## 151 **Materials and Methods**

### 152 **Isolation and identification of epibacterial strains**

153 Five invasive and five native populations of *Agarophyton vermiculophyllum* (Gurgel *et al.*,  
154 2018) (Synonym: *Gracilaria vermiculophyllum* (Ohmi) Papenfuss, hereafter: *Agarophyton*)  
155 were sampled along the Danish-German Peninsula of Jutland and Schleswig-Holstein and  
156 the South Korean peninsula, respectively (see Table S1 in Supporting Information). Using  
157 standard protocols, bacterial strains were isolated from the surface of *Agarophyton*. Thus,  
158 the tested bacterial strains were ecologically relevant. 5 g of pooled algal individuals arising  
159 from each population were rinsed three times in 35 ml of Bacto Marine Broth (MB; Difco  
160 2216, Becton Dickinson and Company, Heidelberg, Germany) to remove loosely attached  
161 bacteria. Then, the samples were immediately transferred to 10 ml of MB and vortexed  
162 vigorously for 20 s to detach the epibacteria. The suspension was subsequently diluted in

163 MB using the log dilution method and plated out directly on MB agar (37.3 g<sup>-1</sup> MB, 15.0 g<sup>-1</sup>  
164 agar; pH 7.6) in standard Petri dishes. Incubation was performed in the dark at 28°C for 7  
165 days. Pure cultures were obtained through several subsequent picking and culturing steps  
166 for individual colonies on MB agar plates. The isolates were cryopreserved at -80°C using the  
167 Cryobank System (Mast Diagnostica GmbH, Reinfeld, Germany) according to the  
168 manufacturer's instructions, until processed further. Strains were identified by 16S rRNA  
169 sequencing as described in Saha *et al.*, (2016) and tested in the bioassays described below.  
170 For methodological details see Appendix S1 in Supporting Information.

### 171 ***Agarophyton* tip bleaching assay with epibacterial strains**

#### 172 **(A) Bleaching assay with single isolates**

173 To test the potential capacity of epibacterial strains for induction of thallus bleaching in  
174 *Agarophyton*, 58 of the cryopreserved bacterial strains were reanimated in November 2015.  
175 They were then maintained on MB agar medium in darkness. All cultures were incubated at  
176 25°C, except *Psychroserpens mesophilus* and *Pseudoalteromonas lipolytica* as they exhibited  
177 no growth at this temperature and were incubated at 15°C.

178 In November 2015, *Agarophyton* individuals were sampled from Nordstrand  
179 (53°29'10.25"N, 8°38'35.33"E) and brought to the laboratory in a cooler box. They were  
180 maintained in 20 L aquaria at a salinity of 33 psu (approximate salinity value at the collection  
181 site) and a temperature of 16°C under constant aeration and a photon flux density of 75  
182  $\mu\text{mol m}^2 \text{s}^{-1}$  (12 h of light per d).

183 For the experiment, *Agarophyton* thallus tips (n = 6 in total for each bacterial strain, each tip  
184 was ca. 2-3 cm long) were individually placed into separate wells of 24 well plates (Sarstedt,  
185 GmbH) containing 2 ml of sterile sea water (SSW, 33 psu). To eliminate epibacteria from the  
186 algal surface, two antibiotics, Vancomycin and Cefotaxim (each at concentration of 0.1 mg



187 ml<sup>-1</sup>) were added to each well (Weinberger et al., 1997). The wells were then incubated for 2  
188 days at 16 °C and a photon flux density of 75 μmol m<sup>-2</sup> s<sup>-1</sup>. Following this pre-treatment, the  
189 wells were carefully emptied of SSW and antibiotics. Remaining antibiotics were removed  
190 from *Agarophyton* tips and the wells by washing with 1 ml of SSW. Finally, 2 ml of SSW were  
191 again added into each well and bacteria cultures were immediately inoculated.

192 Prior to inoculation all bacterial cultures were grown in sterile MB medium for 3-7 d at the  
193 same temperature that was used for their maintenance (25°C or 15°C, see above) in  
194 darkness until they had reached an OD<sub>610</sub> of 0.2 to 0.3. A volume of 20 μl bacterial cells  
195 along with the medium was then added into the wells containing *Agarophyton* tips (n = 6).  
196 Controls consisted of the same volume of sterile bacterial culture medium added into the  
197 wells containing *Agarophyton* tips (n = 6) and treated in a similar manner as above.  
198 Following five days of incubation (16 °C and a photon flux density of 75 μmol m<sup>-2</sup> s<sup>-1</sup>) all wells  
199 were checked under the binocular microscope (magnification factor: 20, see Supporting  
200 Information Fig. S1) and numbers of bleached and non-bleached tips in each well were  
201 counted, using a black background (Weinberger *et al.*, 1997).

202 Relative risk of thallus tip bleaching in treatments with addition of bacteria relative to  
203 control treatments were calculated as odds ratios of numbers of bleached and non-  
204 bleached tips.

205

$$206 \text{ Relative risk of bleaching} = \frac{\text{(Bleached tips in treatments} \div \text{healthy tips in treatments)}}{\text{(Bleached tips in controls} \div \text{healthy tips in controls)}}$$

208 95% Confidence intervals of these ratios were constructed following Fisher and Van Belle  
209 (1993). The divergence of these ratios from 1 (= no effect of the treatment on the risk) was  
210 tested for significance, using Fisher's exact test (Fisher & Van Belle, 1993). Isolates that

211 significantly induced thallus tip bleaching were retested in one or two independent  
212 repetitions of the experiment to confirm the result. The Mantel-Haenssel-extension of  
213 Fisher's exact test for replicated test designs was used for the statistical analysis in these  
214 cases (Fisher & Van Belle, 1993; Weinberger *et al.*, 1997). To reduce the risk of type I error a  
215 Bonferroni correction was applied if multiple tests had to be done (Fisher & Van Belle,  
216 1993). Isolates that turned out to be significantly pathogenic after applying Bonferroni  
217 correction (i.e.  $p < 0.00086$ ) were called 'significant pathogens' while the ones which were  
218 non-significant after Bonferroni correction (i.e.  $p < 0.05$ ) were called 'potential pathogens'.  
219 Isolates that reduced the risk of thallus tip bleaching were all designated as 'protectors'.  
220 However, isolates that significantly reduced the risk of thallus tip bleaching after applying  
221 Bonferroni correction (i.e.  $p < 0.00086$ ) were called 'significant protectors', while the other  
222 isolates that also reduced tip bleaching ( $p < 0.05$ ) but were not significant after applying  
223 Bonferroni correction were designated as 'potential protectors'.

#### 224 **(B) Bleaching assay with combined 'protectors' and 'pathogens'**

225 The combined effect of all 'protectors' ('significant' and 'potential' protectors) on the  
226 virulence of confirmed pathogens ('significant' and 'potential' pathogens) identified in the  
227 above experiment was tested in an additional experiment. In order to observe any  
228 community effect of these epibacteria, we included all the significant and potential strains  
229 because bonferroni correction is known to be relatively conservative (Moran 2003). The  
230 general design was as described above, but the method of inoculation differed: bacterial  
231 cultures were incubated until their  $OD_{610}$  was between 0.1 and 0.5. Different aliquots of all  
232 identified 'protectors' were then pooled so that each culture contributed the same  $OD_{610}$  to  
233 the mixture, which had a final  $OD_{610}$  of 0.25. Cells in the mixture were separated from the  
234 medium by centrifugation (10 000 g, 20 min) and resuspended in SSW. A mixture of three

235 pathogens was prepared in an analogous manner. *Agarophyton* was then inoculated with 10  
236  $\mu\text{l}$  of either (i) all of the ‘protectors’ (19 bacterial strains, thereof 10 ‘significant protectors’  
237 and 9 ‘potential protectors’) or (ii) all of the pathogens (3 bacterial strains, thereof 1  
238 ‘significant pathogen’ and 2 ‘potential pathogens’) or (iii) pathogens and protectors together  
239 (22 bacterial strains). Final volumes of either protectors or pathogens were brought up to 20  
240  $\mu\text{l}$  with SSW, while controls received just 20  $\mu\text{l}$  of SSW. This experiment was repeated in one  
241 fully independent repetition ( $n = 2 \times 6$ ). Numbers of bleached thallus tips relative to all tips  
242 were counted and significant differences were identified using Kruskal-Wallis-ANOVA and  
243 Dunn’s post hoc test.

#### 244 **Extraction of surface associated metabolites of *Agarophyton***

245 To generate surface associated metabolites, *Agarophyton* individuals ( $n=5$ ) were collected  
246 from the same location as above, i.e. Nordstrand. Surface-associated metabolites  
247 originating from single *Agarophyton* specimens were extracted immediately upon collection  
248 according to Saha *et al.*, (2016). Briefly, *Agarophyton* branches were dipped into a solvent  
249 mixture of dichloromethane and hexane 1:4 (v/v) for 5 s. This process is benign and does not  
250 harvest intracellular metabolites (see Saha *et al.*, (2016) for details). The prepared extract  
251 ( $n=5$ ) containing the surface associated metabolites was filtered through GF/A filter paper  
252 (Whatman  $\varnothing = 15$  mm) to remove particles, and the solvent was evaporated under a  
253 vacuum at 20°C, using a rotary evaporator (Laborota 3000, Heidolph, Germany). The extract  
254 was then taken up in acetonitrile in such a way that 1.5  $\mu\text{l}$  contained metabolites extracted  
255 from an algal surface of 99.64  $\text{mm}^2$ . The extract was used to coat each replicate well with a  
256 surface area 99.64  $\text{mm}^2$ . Acetonitrile was then evaporated and metabolites originating from  
257 the surface of the alga remained on the surface of the well, allowing us to test at an  
258 ecologically realistic 1-fold concentration.

259

260 **Defence capacity test of *Agarophyton* surface metabolites against pathogens and**  
261 **protectors**

262 Inhibition or reduction of bacterial settlement and attachment represents the first line of  
263 defence against microbial challenge (Lane & Kubanek, 2009). Thus, an antisettlement assay  
264 was employed as the most relevant criterion for determining antimicrofouling defence  
265 because it quantifies both repellent and toxic effects (Wahl, Jensen & Fenical, 1993). The  
266 assay was performed according to Saha *et al.* (2016). Briefly, the assay was conducted in 96-  
267 well plates that were surface-impregnated with *Agarophyton* surface extract metabolites at  
268 a 1-fold natural concentration (Saha *et al.*, 2016) and with solvent residue as a control. In  
269 total five extracts – originating from five *Agarophyton* individuals - were tested and  
270 regarded as replicates. Each *Agarophyton* extract was then subdivided and tested in three  
271 pseudo replicates against each bacterial isolate to account for the variability in the bacterial  
272 settlement rates. Results obtained for pseudo replicates were averaged before statistical  
273 analyses were conducted. The tested target strains were chosen based on results from the  
274 tip bleaching assay described above. All three pathogens (both the ‘significant pathogen’  
275 *Kordia algicida* and the ‘potential pathogens’ *Croceitalea eckloniae*, *Pseudoalteromonas*  
276 *arctica*) were tested in the anti-settlement assays, but due to shortage of surface extracts  
277 was it not possible to test all nineteen ‘protective’ strains. Thus, only five ‘significant  
278 protectors’ i.e. *Ralstonia sp.*, *Shewanella aquimarina*, *Tenacibaculum skagerrakense*,  
279 *Alteromonas stellipolaris*, *Tenacibaculum aestuarii* and two ‘potential protectors’ i.e.  
280 *Cobetia marina* and *Nonlabens dokdonensis* were tested. 106 µL suspensions of these  
281 bacterial strains (O.D. 0.6-0.8) precultured in MB liquid medium (as described above) were  
282 pipetted into the wells. The bacteria were allowed to settle for 3 h, and the settled cells that

283 could not be removed by rinsing two times with 110µL sterile seawater were stained with  
284 the fluorescent DNA-binding dye Syto 9 (Invitrogen, GmbH). Fluorescence was subsequently  
285 measured (excitation, 477–491 nm; emission, 540 nm) with a plate reader as a proxy for  
286 bacterial settlement in terms of the attached cell density. All tested strains were allowed to  
287 settle on all extracts.

288 The defence strength of *Agarophyton* surface metabolites is expressed as the ‘log effect  
289 ratio,’ i.e., the logarithm of the fluorescence attributable to the settled bacteria of strain Y in  
290 the presence of surface metabolites, divided by the fluorescence attributable to the settled  
291 bacteria of strain Y in the absence of surface metabolites. A log effect ratio value of 0 (i.e.,  
292 an equal number of settled bacteria in wells with and without surface metabolites)  
293 indicated that the tested surface metabolites had no effect on settlement, whereas a  
294 negative log effect ratio value indicated a deterrent effect, and a positive log effect ratio  
295 value indicated an attractant effect. Thus, a log effect ratio of -1 represents a 10-fold  
296 reduction, whereas a value of +1 represents a 10-fold enhancement of bacterial settlement  
297 caused by surface metabolites.

298

299 Defence strength =  $\log \left( \frac{\text{bacterial settlement in presence of } \textit{Agarophyton} \text{ surface metabolites}}{\text{bacterial settlement in absence of } \textit{Agarophyton} \text{ surface metabolites}} \right)$

300

301

## 302 **Results**

### 303 ***Agarophyton* tip bleaching assay with epibacterial strains**

#### 304 **(A) Bleaching assay with single isolates**

305 Out of 58 tested bacterial isolates *Kordia algicida* was found to significantly increase the risk  
306 of tip bleaching (Table 1; Fig. 1,  $p < 0.00086$ ), compared to control treatments without  
307 inoculation of bacteria and was a 'significant pathogen' after Bonferroni correction. Two  
308 additional isolates (*Pseudoalteromonas arctica* and *Croceitalea eckloniae*) had the same  
309 effect but were not significantly pathogenic after Bonferroni correction (Table 1; Fig. 1,  
310  $p < 0.05$ ) and were thus 'potential pathogens'. Out of the remaining 55 isolates, 9 were found  
311 to significantly reduce the risk of tip bleaching (Table 1; Fig. 1,  $p < 0.00086$ ) and were grouped  
312 under 'significant protectors'. 10 others had the same effect, although they were not  
313 significantly protective after Bonferroni correction (Table 1; Fig. 1,  $p < 0.05$ ) and were called  
314 'potential protectors'. The remaining 36 isolates were found to be neutral, neither inducing  
315 nor preventing bleaching (see Table S2; Fig. S2 in Supporting Information).

316 Similar numbers of microbiota that originated from native and non-native populations (30  
317 and 28, respectively, see Table 1 and Table S2) of *Agarophyton* were tested in our bleaching  
318 assay and double numbers of protective microbiota were detected from the non-native  
319 range (3 from the native range and 6 from the non-native range, respectively, Table 1).

#### 320 **(B) Bleaching assay with combined 'protectors' and 'pathogens'**

321 When all three isolates (one 'significant pathogen' and two 'potential pathogen') that  
322 induced bleaching individually at least with  $p < 0.05$  were combined, a significant increase in  
323 bleaching relative to the control was again observed (Fig. 2,  $p < 0.05$ ). However, combined  
324 application of these three 'pathogens' and the nineteen 'protective' isolates that prevented  
325 bleaching individually at least with  $p < 0.05$  resulted in no such increase (Fig. 2,  $p < 0.05$ ). No  
326 bleaching was observed when all 19 'protectors' and no 'pathogens' were inoculated.

327

#### 328 **Defence capacity test of *Agarophyton* against pathogens and protectors**

329 The effect of *Agarophyton* surface associated metabolites on bacterial settlement differed  
330 significantly between the two groups of bacteria, i.e. 'protectors' and 'inducers' (Fig. 3,  
331 Welch-corrected t-test,  $p < 0.0001$ ). While the surface associated metabolites significantly  
332 increased the settlement of 'protectors', the settlement of the bleaching 'inducers' was  
333 significantly reduced by the surface associated metabolites.

334

### 335 **Discussion**

336 The data presented here demonstrate for the first time that aquatic macrophytes can use  
337 surface associated chemicals not only to directly reduce settlement of pathogenic bacteria,  
338 but also to recruit bacterial strains that provide protection from such pathogens. The  
339 epimicrobiome of *Agarophyton* contains a component that protects the alga from  
340 pathogens in a similar way as earlier demonstrated for two other seaweeds (*G. conferta*  
341 (Weinberger *et al.*, 1997) and *D. pulchra* (Longford *et al.*, 2019)), and the settlement of such  
342 protective bacteria on the surface of *Agarophyton* is not random. Surface associated  
343 metabolites from the *Agarophyton* holobiont significantly deterred three strains that were  
344 significantly and potentially pathogenic, while the metabolites had a probiotic effect  
345 towards seven significantly and potentially protective strains that were tested. This confirms  
346 the surface chemistry of *Agarophyton* has a similarly strong selective effect on bacterial  
347 colonization as in *Fucus vesiculosus* (Lachnit *et al.*, 2010) or *Delisea pulchra* (Longford *et al.*,  
348 2019). Moreover, it demonstrates for the first time that this selection is not only targeted to  
349 exclude pathogens, but also targeted to attract protectors. Together with Lachnit *et al.*,  
350 (2010) and Kessler *et al.*, (2018) our data strongly support the concept of chemically  
351 mediated recruitment of microbes and not the 'neutral' hypothesis, according to which the  
352 microbial community structure is determined randomly. Our data clearly support the

353 targeted deterrence hypothesis, as settlement of detrimental bacteria was chemically  
354 suppressed. On the other hand, we cannot reject the 'niche' model, as multiple microbiota  
355 were attracted by *Agarophyton* and possibly able to make use of resources provided by it.  
356 Also, the 'lottery' hypothesis cannot be currently rejected, since the capacity of attracted  
357 microbiota to coexist and share host resources is unknown.

358 Only 5% of the marine bacterial strains are cultivable (Haglund et al., 2002) and to date no  
359 alternative technique has been developed to separate selected microbial components from  
360 natural microbial communities and to test them in infection assays. Thus, only a small  
361 fraction of all bacteria that are associated with the surface of *Agarophyton* could be isolated  
362 and tested in our study. One representative out of 58 tested bacterial species, *Kordia*  
363 *algicida*, was significantly capable to induce the tip bleaching symptom in *Agarophyton*. *K.*  
364 *algicida* is already known to be detrimental to other organisms. It can kill diatom blooms in  
365 a protease mediated molecular interaction (Paul & Pohnert, 2011) and a similar mechanism  
366 cannot be excluded in the present case. Bleaching is often correlated with microbial cell wall  
367 matrix degrading activity (Weinberger et al., 1994, 1997), but this was not the case in the  
368 present study, as *Kordia* is incapable of agar degradation.

369 Two other isolates – which were also not agar degraders - also exhibited the potential to  
370 induce bleaching symptoms in *Agarophyton*, which strongly suggests that this capacity is not  
371 unique. Interestingly, all three detrimental isolates originated from virtually healthy host  
372 specimens. Thus, a relevant fraction of *Agarophyton's* surface microbiome is obviously  
373 composed of opportunistic pathogens that can induce bleaching symptoms under certain  
374 conditions, similar as in several red seaweeds belonging to other species (Case et al., 2011;  
375 Weinberger et al., 1994; Weinberger et al., 1997). Given that three out of 58 culturable



376 strains were significant or potential pathogens this fraction can be estimated to include  
377 approximately 5 % of the microbiome. However, this percentage calculation is based on the  
378 culturable proportion which is just a representative sample of the whole microbiome.

379 Of the remaining strains, 19 (9 'significant protectors' and 10 'potential protectors') could  
380 reduce the risk of 'spontaneous' bleaching in thalli that were not intentionally inoculated  
381 with pathogenic bacteria (Fig. 1). All the specimens of *Agarophyton* tested in our bleaching  
382 induction assays were subjected to a pretreatment with two antibiotics that inhibited  
383 bacterial cell wall synthesis, with the dual goal to remove opportunistic pathogens and to  
384 disturb and weaken any protective components of the algal microbiome. The circumstance  
385 that bleaching occurred 'spontaneously' at a low rate but could be prevented by an  
386 important percentage of all tested isolates suggests that some opportunistic pathogens  
387 survived the treatment with antibiotics but could then not become virulent when protective  
388 bacteria were inoculated - similar as previously reported for bacteria that had been isolated  
389 from *Gracilaria 'conferta'* and prevented thallus tip bleaching in this alga (Weinberger *et al.*,  
390 1997). The protective effect of various isolates on *Agarophyton* was further confirmed when  
391 all the 19 protective strains (nine 'significantly protective' and ten 'potentially protective')  
392 were tested together in combination with the 3 pathogenic strains (one 'significant  
393 pathogen' and two 'potential pathogens') and a bleaching reduction was still documented.  
394 Altogether, our observations strongly hint at the presence of protective epibacteria on the  
395 surface of *Agarophyton*. They could (again estimated from the number of identified isolates  
396 in our tested strain collection) comprise at least 15% of all taxa present in this microbiome.  
397 The presence of such beneficial bacteria has been previously demonstrated not only for  
398 other Gracilarioids (Weinberger *et al.*, 1997), but also for *Delisea pulchra* (Longford *et al.*,  
399 2019), corals (Rosenberg *et al.*, 2007) and other seaweeds like the brown alga *Fucus*

400 *vesiculosus*, in which surface associated bacteria were found to inhibit the settlement of  
401 macrofoulers (Nasrolahi *et al.*, 2012).

402 The biofilms on seaweed surfaces represent highly competitive environments, with  
403 microbes competing for refuge, nutrients and substratum, and interspecific antagonistic  
404 effects of bacterial strains are not rare. For example, such effects have been previously  
405 demonstrated for the brown alga *Saccharina latissima* (Wiese *et al.*, 2009), the red alga  
406 *Delisea pulchra* and the green alga *Ulva australis* (Penesyan *et al.*, 2009). Release of  
407 inhibitory components like antibiotics (Wiese *et al.*, 2009) and/or quorum sensing inhibitors  
408 (Romero *et al.*, 2010) has been observed and could also explain the 'protective' effect  
409 observed by us. Interestingly, one of the significant protective strains, *Pseudoalteromonas*  
410 *piscicida*, belongs to a genus which comprises several species that are known to produce  
411 antibacterial products to outcompete other bacteria for space and nutrients (Holmström &  
412 Kjelleberg, 1999). Also *Pseudoalteromonas piscicida* has been recently demonstrated to  
413 inhibit and/or kill competing bacteria - including several marine pathogens, such as *Vibrio*  
414 *vulnificus*, *Vibrio parahaemolyticus*, *Vibrio cholerae*, *Photobacterium damsela*, and  
415 *Shewanella algae* - through secretion of antimicrobial substances and the direct transfer of  
416 digestive vesicles to competing bacteria (Richards *et al.*, 2017). On the other hand,  
417 *Shewanella aquimarina* exhibited a strong protective effect against *Agarophyton* tip  
418 bleaching in the current study and the same was observed with two other potentially  
419 protective species of the genus *Shewanella*, *i.e.* *S. marisflavi* and *S. loihica*. These  
420 observations contrast with the findings that *S. marisflavi* is a pathogen of sea cucumbers (Li  
421 *et al.*, 2010) and other bacteria of the genus are pathogenic towards humans. The  
422 mechanisms behind the protective effects on *Agarophyton* deserve further investigation.  
423 The exact (additive or synergistic) contributions of the active epibacterial players in the

424 cross-infection experiment with all 19 ‘protectors’ combined with 3 ‘pathogens’ are not  
425 known yet.

426 Beneficial roles of certain components in the bacterial communities around the rhizosphere  
427 of terrestrial plants are well known. They can not only facilitate nutrient acquisition, but also  
428 support plant growth under biotic and abiotic plant stress (Lareen *et al.*, 2016; Mendes *et al.*,  
429 *al.*, 2013). Seaweed-associated bacteria may also facilitate nutrient acquisition and provide  
430 essential vitamins and growth factors (Wahl *et al.*, 2012), and – as confirmed in the present  
431 study – mediate biotic stress. However, while we have started to understand these benefits  
432 and to gather evidence of a selective recruitment of bacteria both in terrestrial (Lebeis *et al.*,  
433 2015) and aquatic environments, empirical links between this selective recruitment of  
434 communities and a health benefit for the host are still very rare in the aquatic realm. Kessler  
435 *et al.*, 2018 recently demonstrated for the marine macroalga *Ulva mutabilis* (Chlorophyta) a  
436 mediating role of the algal osmolyte DMSP (dimethylsulfoniopropionate) in the attraction of  
437 the beneficial bacterium *Roseovarius* sp. MS2, responsible for release of morphogenetic  
438 compounds that ensure proper algal morphogenesis. In absence of these morphogenetic  
439 compounds under axenic conditions, *Ulva mutabilis* develops into callus-like colonies  
440 consisting of undifferentiated cells and abnormal cell walls. While microbial ‘gardening’ via  
441 use of chemicals has been documented in terms of growth and development of seaweeds  
442 (Kessler *et al.*, 2018), our study demonstrates for the first time such a link between the  
443 disease resistance capacity of a seaweed and beneficial selective gardening of ‘protective’  
444 bacteria based upon surface chemistry.

445 Metabolites present on the surface of seaweeds or in the rhizosphere are a cocktail of  
446 metabolites originating both from the algal or plant host and from associated surface

447 microbiota. Such surface associated metabolites from the algal holobiont are also known to  
448 function as a defence against fouling by microfoulers (e.g. bacteria, diatoms) and  
449 macrofoulers (e.g. barnacle larvae, mussels) (reviewed by Da Gama, 2014; Saha *et al.*, 2017).  
450 Also epibacteria from seaweeds are well known to have inhibitory activities against other  
451 fouling organisms (Singh & Reddy, 2014). Thus, it is possible that beneficial bacteria  
452 recruited by *Agarophyton* will not only act as a defence against pathogens but also against  
453 other foulers, like filamentous algae. Using a transcriptomic approach, de Oliveira *et al.*,  
454 2012 demonstrated that the red seaweed host *Laurencia dendriodea* (rather the surface  
455 associated bacteria) is involved in the biosynthesis of terpenoids (chemical defence  
456 compounds against bacterial colonization and infection) through the mevalonate  
457 independent pathway. For the *Agarophyton* holobiont, we do not know yet the identity of  
458 surface associated bioactive compounds. Thus, it was not possible for us to distinguish the  
459 relative contributions of surface metabolites originating from the algal host *Agarophyton*  
460 and from surface associated microbiota. Mutants of *Arabidopsis thaliana* with suppressed  
461 salicylic acid signalling pathways formed abnormal root microbiomes when compared to the  
462 wild plants (Lebeis *et al.*, 2015), which could suggest that the role of the host was more  
463 important in this specific case. The contribution of seaweed microbiome metabolites  
464 depends on the community composition, abundance and metabolic activity (Wahl *et al.*,  
465 2010) and may be expected to be more variable than that of the host. Selective effects  
466 observed with surface associated metabolites coming from different algal individuals varied  
467 relatively little in our study, which could suggest that metabolites generated by the host  
468 have more importance. However, our knowledge of the species-species interactions of  
469 cultivatable and non-cultivatible taxa associated with *Agarophyton* or other plants is  
470 rudimentary at best. The involvement of multiple protective microorganisms in our and

471 several other cases (see above) strongly supports the view that the traditional conceptual  
472 model emphasizing direct interactions of hosts and single microbes needs to be expanded to  
473 a holobiont concept if seaweed- or plant-microbe interactions are to be understood (Egan *et*  
474 *al.*, 2013).

475 In conclusion, our study demonstrates for the first-time selective chemical ‘gardening’ of  
476 protective epibacterial strains by a seaweed holobiont with regard to disease resistance  
477 capacity. The combined effect of metabolites generated by the host alga and the protection  
478 offered by associated microbial partners determines the virulence of harmful opportunistic  
479 bacterial pathogens. A major component of the epibacterial community appears capable of  
480 contributing to this protection against co-occurring pathogens, which suggests that  
481 microbiota of very different taxonomic groups may provide the holobiont with the same  
482 ecological function, which could be pivotal for the establishment of *Agarophyton* in new  
483 environments. Thus, absence of protective microbiota in new environments might not be a  
484 factor limiting the invasion success of *Agarophyton*.

485 As known for other seaweeds like the brown alga *Fucus vesiculosus*, bioactive surface  
486 metabolites often act in synergism or additively and/or antagonistically, producing an  
487 overall defensive or prebiotic effect on bacterial recruitment (Saha *et al.*, 2011; Saha *et al.*,  
488 2012). The identification of metabolites responsible for such chemical ‘gardening’ effects via  
489 classical bioassay guided fractionation techniques in the near future may allow us to  
490 manipulate algal thallus microbiomes to enhance seaweed health, prevent bleaching  
491 diseases and ensure production and sustainability in *Agarophyton* aquaculture.

## 492 **Acknowledgements**

493 This research was supported by a grant (CP1215) from the DFG Cluster of Excellence “Future  
494 Ocean” to M. Saha. We thank the Institute of Clinical Molecular Biology of the Christian-  
495 Albrechts-University Kiel (Germany) for performing Sanger sequencing, supported in part by  
496 the DFG Clusters of Excellence “Inflammation at Interfaces” and “Future Ocean”.

#### 497 **Author contributions**

498 M.S. isolated and identified the bacterial isolates, designed and performed the anti-  
499 settlement experiments. F.W. designed and performed the tip bleaching assays. M.S. and  
500 F.W. analysed the data. M.S. wrote the paper and F.W. contributed to the editing.

#### 501 **Author Declaration**

502 The authors declare no conflict of interest. Data underlying this publication are freely  
503 accessible and can be downloaded from the DRYAD data repository (Provisional DOI:  
504 doi:10.5061/dryad.52j8p1r).

#### 505 **References**

506

507 Bacilio-Jiménez, M., Aguilar-flores, S., Ventura-zapata, E., & Eduardo, P. (2003). Chemical  
508 characterization of root exudates from rice (*Oryza sativa*) and their effects on the  
509 chemotactic response of endophytic bacteria, *Plant and Soil* **249**: 271–277.

510 Bell, W., & Mitchell, R. (1972). Chemotactic and growth responses of marine bacteria to algal  
511 extracellular products. *The Biological Bulletin*, **143(2)**, 265–277.

512 Bengtsson, M. M., & Ovreas, L. (2010). Planctomycetes dominate biofilms on surfaces of the  
513 kelp *Laminaria hyperborea*. *BMC Microbiology*, **10(1)**, 261.

514 Bordenstein, S. R., & Theis, K. R. (2015). Host biology in light of the microbiome: ten

515 principles of holobionts and hologenomes. *PLoS Biology*, **13(8)**, e1002226.

516 Burke, C., Steinberg, P., Rusch, D. B., Kjelleberg, S., & Thomas, T. (2011). Bacterial community  
517 assembly based on functional genes rather than species. *Proceedings of the National  
518 Academy of Sciences of the USA*, **108(34)**, 14288–14293.

519 Campbell, A. H., Harder, T., Nielsen, S., Kjelleberg, S., & Steinberg, P. D. (2011). Climate  
520 change and disease: bleaching of a chemically defended seaweed. *Global Change  
521 Biology*, **17(9)**, 2958–2970.

522 Case, R. J., Longford, S. R., Campbell, A. H., Low, A., Tujula, N., Steinberg, P. D., & Kjelleberg,  
523 S. (2011). Temperature induced bacterial virulence and bleaching disease in a  
524 chemically defended marine macroalga. *Environmental Microbiology*, **13(2)**, 529–537.

525 Cundell, A. M., Sleeter, T. D., & Mitchell, R. (1977). Microbial populations associated with the  
526 surface of the brown alga *Ascophyllum nodosum*. *Microbial Ecology*, **4(1)**, 81–91.

527 Currier, W. W., & Strobel, G. A. (1976). Chemotaxis of *Rhizobium* spp. to plant root exudates.  
528 *Plant Physiology*, **57(5)**, 820–823.

529 Egan, S., Harder, T., Burke, C., Steinberg, P., Kjelleberg, S., & Thomas, T. (2013). The seaweed  
530 holobiont: Understanding seaweed-bacteria interactions. *FEMS Microbiology Reviews*,  
531 **37(3)**, 462–476.

532 Flint, H. J., Scott, K. P., Louis, P., & Duncan, S. H. (2012). The role of the gut microbiota in  
533 nutrition and health. *Nature Reviews Gastroenterology and Hepatology*, **9(10)**, 577.

534 Gachon, C. M. M., Sime-Ngando, T., Strittmatter, M., Chambouvet, A., & Kim, G. H. (2010).  
535 Algal diseases: spotlight on a black box. *Trends in Plant Science*, **15(11)**, 633–40.

536 Gurgel, C. F. D., Norris, J. N., Schmidt, W. E., Le, H. A. U. N. H. U., & Fredericq, S. (2018).  
537 Systematics of the *Gracilariales* (Rhodophyta) including new subfamilies, tribes,  
538 subgenera, and two new genera, *Agarophyton* gen. nov. and *Crassa* gen. nov., **374(1)**,  
539 1–23.

540 Haglund, A.-L., Törnblom, E., Boström, B., & Tranvik, L. (2002). Large differences in the  
541 fraction of active bacteria in plankton, sediments, and biofilm. *Microbial Ecology*, **43(2)**,  
542 232–241.

543 Hartmann, A., Schmid, M., Van Tuinen, D., & Berg, G. (2009). Plant-driven selection of  
544 microbes. *Plant and Soil*, **321(1–2)**, 235–257.

545 Hollants, J., Leliaert, F., De Clerck, O., & Willems, A. (2013). What we can learn from sushi: A  
546 review on seaweed-bacterial associations. *FEMS Microbiology Ecology*, **83(1)**, 1–16.

547 Holmström, C., & Kjelleberg, S. (1999). Marine *Speudoalteromonas* species are associated  
548 with higher organisms and produce biologically active extracellular agents. *FEMS*  
549 *Microbiology Ecology*, **30**, 285–293.

550 Hooper, L. V., Midtvedt, T., & Gordon, J. I. (2002). How host-microbial interactions shape the  
551 nutrient environment of the mammalian intestine. *Annual Review of Nutrition*, **22(1)**,  
552 283–307.

553 Hurd, C. L. (2000). Review water motion, marine macroalgal physiology, and production.  
554 *Journal of Phycology*, **36 (3)**, 453–472.

555 Kessler, R. W., Weiss, A., Kuegler, S., Hermes, C., & Wichard, T. (2018). Macroalgal–bacterial  
556 interactions: Role of dimethylsulfoniopropionate in microbial gardening by *Ulva*  
557 (Chlorophyta). *Molecular Ecology*, **27(8)**, 1808–1819.



558 Lachnit, T., Blümel, M., Imhoff, J., & Wahl, M. (2009). Specific epibacterial communities on  
559 macroalgae: phylogeny matters more than habitat. *Aquatic Biology*, **5**, 181–186.

560 Lachnit, T., Wahl, M., & Harder, T. (2010). Isolated thallus-associated compounds from the  
561 macroalga *Fucus vesiculosus* mediate bacterial surface colonization in the field similar to  
562 that on the natural alga. *Biofouling*, **26(3)**, 247–55.

563

564 Lachnit, T., Meske, D., Wahl, M., Harder, T., & Schmitz, R. (2011). Epibacterial community  
565 patterns on marine macroalgae are host-specific but temporally variable. *Environmental*  
566 *Microbiology*, **13(3)**, 655–65.

567 Lane, A. L., Nyadong, L., Galhena, A. S., Shearer, T. L., Stout, E. P., Parry, R. M., ... Kubanek, J.  
568 (2009). Desorption electrospray ionization mass spectrometry reveals surface-mediated  
569 antifungal chemical defense of a tropical seaweed. *Proceedings of the National*  
570 *Academy of Sciences of the USA*, **106 (18)**, 7314–7319.

571 Lebeis, S. I., Paredes, S. H., Lundberg, D. S., Breakfield, N., Gehring, J., McDonald, M., ...  
572 Dangl, J. L. (2015). Research reports 27. *Science*, **349(6250)**, 860–864.

573 Li, H., Qiao, G., Li, Q., Zhou, W., Won, K. M., Xu, D., & Park, S. (2010). Biological  
574 characteristics and pathogenicity of a highly pathogenic *Shewanella marisflavi* infecting  
575 sea cucumber, *Apostichopus japonicus*. *Journal of Fish Diseases*, **33(11)**, 865–877.

576 Longford, S. R., Campbell, A. H., Nielsen, S., & Case, R. J. (2019). Interactions within the  
577 microbiome alter microbial interactions with host chemical defences and affect disease  
578 in a marine holobiont, **9**, 1–13.

579 Mendes, R., Garbeva, P., & Raaijmakers, J. M. (2013). The rhizosphere microbiome:

580 significance of plant beneficial, plant pathogenic, and human pathogenic  
581 microorganisms. *FEMS Microbiology Reviews*, **37(5)**, 634–663.

582 Moran, M. D., Colledge, H., & Ae, W. (2003). Arguments for rejecting the sequential  
583 Bonferroni in ecological studies, *Oikos*, **2**, 1–3.

584 Nasrolahi, A., Stratil, S. B., Jacob, K. J., & Wahl, M. (2012). A protective coat of  
585 microorganisms on macroalgae: inhibitory effects of bacterial biofilms and epibiotic  
586 microbial assemblages on barnacle attachment. *FEMS Microbiology Ecology*, **81(3)**,  
587 583–95.

588 Paul, C., & Pohnert, G. (2011). Interactions of the algicidal bacterium *Kordia algicida* with  
589 diatoms: Regulated protease excretion for specific algal lysis. *PLoS ONE*, **6(6)**, e21032.

590 Penesyan, A., Marshall-Jones, Z., Holmstrom, C., Kjelleberg, S., & Egan, S. (2009).  
591 Antimicrobial activity observed among cultured marine epiphytic bacteria reflects their  
592 potential as a source of new drugs. *FEMS Microbiology Ecology*, **69(1)**, 113–124.

593 Raaijmakers, J. M., Paulitz, T. C., Steinberg, C., Alabouvette, C., & Moënne-Loccoz, Y. (2009).  
594 The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial  
595 microorganisms. *Plant and Soil*, **321(1–2)**, 341–361.

596 Richards, G. P., Watson, M. A., Needleman, D. S., Uknalis, J., Boyd, E. F., & Fay, P. (2017).  
597 Mechanisms for *Pseudoalteromonas piscicida*-Induced Killing of Vibrios and Other  
598 Bacterial Pathogens, *Applied Environmental Ecology*, **83(11)**, 1–17.

599 Romero, M., Martin-Cuadrado, A.-B., Roca-Rivada, A., Cabello, A. M., & Otero, A. (2010).  
600 Quorum quenching in cultivable bacteria from dense marine coastal microbial  
601 communities. *FEMS Microbiology Ecology*, **75(2)**, 205–217.

- 602 Rosenberg, E., Koren, O., Reshef, L., Efrony, R., & Zilber-Rosenberg, I. (2007). The role of  
603 microorganisms in coral health, disease and evolution. *Nature Reviews Microbiology*,  
604 **5(5)**, 355.
- 605 Saha, M., Rempt, M., Gebser, B., Grueneberg, J., Pohnert, G., & Weinberger, F. (2012).  
606 Dimethylsulphopropionate (DMSP) and proline from the surface of the brown alga  
607 *Fucus vesiculosus* inhibit bacterial attachment. *Biofouling*, **28(6)**, 593–604.
- 608 Saha, M., Rempt, M., Grosser, K., Pohnert, G., & Weinberger, F. (2011). Surface-associated  
609 fucoxanthin mediates settlement of bacterial epiphytes on the rockweed *Fucus*  
610 *vesiculosus*. *Biofouling*, **27(4)**, 423–433.
- 611 Schiel, D. R., & Foster, M. S. (2006). The population biology of large brown seaweeds:  
612 ecological consequences of multiphase life histories in dynamic coastal environments.  
613 *Annual Review of Ecology, Evolution, and Systematics*, **37**, 343–372.
- 614 Singh, R. P., & Reddy, C. R. K. (2014). Seaweed-microbial interactions: Key functions of  
615 seaweed-associated bacteria. *FEMS Microbiology Ecology*, **88(2)**, 213–230.
- 616 Van Alstyne, K., & Houser, L. (2003). Dimethylsulfide release during macroinvertebrate  
617 grazing and its role as an activated chemical defense. *Marine Ecology Progress Series*,  
618 **250**, 175–181.
- 619 Wahl, M. (1989). Marine epibiosis. I. Fouling and antifouling: some basic aspects. *Marine*  
620 *Ecology Progress Series*, **58**, 175–189.
- 621 Wahl, M. (2008). Ecological lever and interface ecology: epibiosis modulates the interactions  
622 between host and environment. *Biofouling*, **24(6)**, 427–38.
- 623 Wahl, M., Goecke, F., Labes, A., Dobretsov, S., & Weinberger, F. (2012). The second skin:

624 Ecological role of epibiotic biofilms on marine organisms. *Frontiers in Microbiology*, **3**,  
625 1–21.

626 Wahl, M., ShahnazL., Dobretsov, S., Saha, M., Symanowski, F., DavidK., ... Weinberger, F.  
627 (2010). Ecology of antifouling resistance in the bladder wrack *Fucus vesiculosus*:  
628 Patterns of microfouling and antimicrobial protection. *Marine Ecology Progress Series*,  
629 **411**, 33–48.

630 Weinberger, F. (2007). Pathogen induced defense and innate immunity in macroalgae.  
631 *Biological Bulletin*, **35(1)**, 29–54.

632 Weinberger, F., Friedlander, M., & Gunkel, W. (1994). A bacterial facultative parasite of  
633 *Gracilaria conferta*. *Diseases of Aquatic Organisms*, **18(2)**, 135–141.

634 Weinberger, F., Hoppe, H. G., & Friedlander, M. (1997). Bacterial induction and inhibition of  
635 a fast necrotic response in *Gracilaria conferta* (Rhodophyta). *Journal of Applied*  
636 *Phycology*, **9(3)**, 277–285.

637 Wiese, J., Thiel, V., Nagel, K., Staufenberg, T., & Imhoff, J. F. (2009). Diversity of antibiotic-  
638 active bacteria associated with the brown alga *Laminaria saccharina* from the Baltic  
639 Sea. *Marine Biotechnology*, **11(2)**, 287–300.

640 Yuan, J., Chaparro, J. M., Manter, D. K., Zhang, R., Vivanco, J. M., & Shen, Q. (2015). Roots  
641 from distinct plant developmental stages are capable of rapidly selecting their own  
642 microbiome without the influence of environmental and soil edaphic factors. *Soil*  
643 *Biology and Biochemistry*, **89**, 206–209.

644



646 **Tables**


647 Table 1. Epibacterial strains [isolated from *Agarophyton vermiculophyllum* (GV) populations]  
 648 that were tested in tip bleaching assays. IR=bacteria isolated from the invasive populations  
 649 of GV; NR=bacteria isolated from the native populations of GV. Significantly protective  
 650 strains are bacteria that were still protective after Bonferroni correction. Asterisks indicate  
 651 strains that were tested in anti-settlement assays with surface associated compounds of GV.

Isolate code, Range of isolation	Closest match on RDF	Designation
G-NY6, IR	<i>Ralstonia sp.*</i>	Significantly Protective
G-JI1, NR	<i>Shewanella aquimarina*</i>	Significantly Protective
G-NORD3, IR	<i>Gaetbulibacter lutimaris</i>	Significantly Protective
G-G2, NR	<i>Vibrio marisflavi</i>	Significantly Protective
G-MAN7, IR	<i>Tenacibaculum skagerrakense*</i>	Significantly Protective
G-HO9, IR	<i>Alteromonas stellipolaris*</i>	Significantly Protective
G-ODO3, NR	<i>Maribacter polysiphoniae</i>	Significantly Protective
G-FALK1, IR	<i>Nonlabens dokdonensis*</i>	Potentially Protective
G-NORD11, IR	<i>Pseudoalteromonas piscicida</i>	Significantly Protective
G-G4, NR	<i>Shewanella marisflavi</i>	Potentially Protective
G-DA3, NR	<i>Shewanella loihica</i>	Potentially Protective
G-NORD6, IR	<i>Tenacibaculum aestuarii*</i>	Significantly Protective
G-JI4, NR	<i>Cobetia marina*</i>	Potentially Protective
G-NY1, IR	<i>Alteromonas simiduii</i>	Potentially Protective

G-DA5, NR	<i>Pseudoalteromonas atlantica</i>	Potentially Protective
G-FALK2, IR	<i>Pseudoalteromonas ulvae</i>	Potentially Protective
G-MAN5, IR	<i>Bacillus amyloliquefaciens</i>	Potentially Protective
G-JI5, NR	<i>Marinomonas communis</i>	Potentially Protective
G-HO8, IR	<i>Rhodobacter sp.</i>	Potentially Protective
G-MAN6, IR	<i>Pseudoalteromonas arctica*</i>	Potentially Pathogenic
G-NORD9, IR	<i>Croceitalea eckloniae*</i>	Potentially Pathogenic
G-MAN4, IR	<i>Kordia algicida*</i>	Significantly Pathogenic

652


653



655

656 **Fig. 1:** Risk of thallus tip bleaching in *A. vermiculophyllum* after inoculation of 22 bacterial  
 657 strains relative to control thalli without such inoculation. Numbers of independent infection  
 658 experiments (each with n = 6) were three in the case of *K. algicida*, two in the cases of *C.*  
 659 *eckloniae* and *P. arctica* and one in all other cases. Only isolates that affected the risk at p <  
 660 0.05 are shown. Asterisks indicate isolates which were significantly pathogenic or protective  
 661 after Bonferroni-correction (p<0.00086). Error bars±95% CI.






662

663 **Fig. 2:** Relative amounts of bleached thallus tips in *A. vermiculophyllum* after inoculation  
 664 with 3 pathogenic bacterial isolates with 19 protective bacterial isolates, with all 22 isolates  
 665 together and in controls without any inoculation. Different letters indicate treatments that  
 666 are significantly different (n = 12; Kruskal-Wallis-ANOVA (p < 0.0001) and Dunn's post hoc  
 667 test (p < 0.05)). Pathogenic strains include both 'significant pathogens' and 'potential  
 668 pathogens'. Protective strains include both 'significant protectors' and 'potential  
 669 protectors'. Median ± quartiles.

670



671

672 **Fig. 3:** Mean anti-settlement activity of *Agarophyton* surface metabolites against three  
 673 pathogenic (one 'significant pathogen' and two 'potential pathogen') and seven protective  
 674 (five 'significant protector' and two 'potential protector') strains. Error bars  $\pm$  SD (n=5);  
 675 Welch-corrected t-test,  $p < 0.0001$ .