

1 **Understanding women's help-seeking with intimate partner violence in Tanzania**

2

3 Bathsheba Mahenge ^{a,b,*} & Heidi Stöckl ^b

4 ^a Department of Psychiatry and Mental Health, University of Dodoma, P.O Box 395,

5 Dodoma, Tanzania

6 ^b Gender Violence & Health Centre, Department of Global Health and Development, London

7 School of Hygiene and Tropical Medicine, London, UK

8

9

10 *Corresponding author

11 Email: bmahenge@gmail.com

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32 **Abstract**

33 Intimate partner violence (IPV) is a serious global health problem affecting millions of women
34 worldwide. Despite increased investments into its reduction, little research has been conducted
35 into how women in low and middle-income countries deal with IPV. This study seeks to
36 explore this by looking in-depth into help seeking strategies abused women utilize in Tanzania,
37 using the 2015-2016 Tanzania Demographic and Health Survey. The prevalence of lifetime
38 physical and/or sexual IPV was 41.6% in this study, but only half of all affected women sought
39 help from anyone. The only clear association found with help-seeking was severity of IPV.

40

41 **Keywords:** intimate partner violence, help seeking, disclosure, Tanzania

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 **Introduction**

60 Over the years intimate partner violence (IPV) has been recognised as a serious global health
61 problem affecting millions of women, irrespective of their socio-economic status, educational
62 attainment or marital status (García-Moreno & Stöckl, 2009). It is estimated that almost 30%
63 of women worldwide have experienced some form of physical and/or sexual IPV in their
64 lifetime while one out of four women in Tanzania have experienced physical and/or sexual IPV
65 in their lifetime (Garcia-Moreno, HA, Ellsberg, L, & Watts, 2005; Ministry of Health, 2016).
66 IPV has been linked with a wide array of both physical and mental health effects such as
67 depression, suicidality, PTSD, miscarriages, injuries and in severe cases death (Devries et al.,
68 2013; Mahenge, Likindikoki, Stockl, & Mbwambo, 2013; Mapayi et al., 2013; Stockl et al.,
69 2013). Not only women get affected by IPV, their children and family's economic wellbeing
70 suffers as well (Adams, Tolman, Bybee, Sullivan, & Kennedy, 2012; Jouriles et al., 2018;
71 Neamah et al., 2018).

72 Despite the fact that IPV has tragic consequences, most instances of IPV go unreported and
73 only few women seek help (Okenwa, Lawoko, & Jansson, 2009; Roelens, Verstraelen, Van
74 Egmond, & Temmerman, 2008). Women who have experienced IPV develop different patterns
75 to cope with IPV. Types of help-seeking that exists for those who seek help are either termed
76 "informal", women seek help from their families or their partner's family, friends or
77 neighbours and they are also termed "formal," when women seek help from social services,
78 doctors, lawyers or the police (Ragusa, 2013; Sylaska & Edwards, 2014). According to a study
79 of Demographic and Health Surveys (DHS) from 24 developing countries between 2004 and
80 2011, only 40% of women who experienced physical and/or sexual IPV sought any help, out
81 of which 36.8% sought informal help and 7% sought formal help (Palermo, Bleck, & Peterman,
82 2014).

83 Literature from several studies from Western countries highlight the conditions under which
84 women try to seek both formal and informal help to end IPV and the factors they are associated
85 with are women's socio-economic status, ethnicity, culture and religion (Taket, O'Doherty,
86 Valpied, & Hegarty, 2014). Although social norms have also been said to play part, as IPV
87 might be viewed as a private matter and seeking help comes with loss of privacy (Liang,
88 Goodman, Tummala-Narra, & Weintraub, 2005), research has moved over the years to
89 understand individual factors that may hinder help-seeking. One of the factors highlighted is
90 stigma. Stigma can range from anticipated stigma from the formal or informal support or
91 service providers or internalized stigma associated with feelings of shame and embarrassment
92 that one has experienced IPV or lastly cultural stigma which is fear of judgemental attitudes
93 and victim blaming from service providers (Kennedy & Prock, 2016; Overstreet & Quinn,
94 2013) .

95 The phenomena of stigma and shame was a prominent finding in a qualitative study on help
96 seeking in Tanzania among 96 male and female community members. Both men and women
97 understood what constituted violence against women, still most women normalized IPV as part
98 of a normal relationship and were reluctant to seek help due to stigma, shame, fear and lack of
99 trust in existing response systems (McCleary-Sills et al., 2016). The Palermo 2010 et al. multi-
100 country study which included. data from the 2010 Tanzanian DHS found in its multivariate
101 analysis that seeking help from formal sources among Tanzanian women who experienced IPV
102 was associated with being previously married and being in the bottom 40% wealth quartile
103 while women who had a secondary or higher level of education were less likely to seek help
104 (Palermo et al., 2014).

105 While these existing studies underlined that help-seeking is an important issue to investigate to
106 address IPV, there is a scarcity of quantitative studies from low and middle-income countries
107 like Tanzania with high prevalence rates of IPV on the different forms of help seeking for IPV

108 and from whom help is sought beyond formal services. Studies investigating factors associated
109 with help-seeking have mainly looked at the associations with women's characteristics,
110 ignoring the importance of their partner's characteristics, the type of IPV experienced and
111 relationship characteristics. The aim of this study is to bridge the current research gap and
112 establish help seeking strategies women use and its associated factors in Tanzania, using the
113 national representative 2015-2016 Tanzanian DHS.

114 .

115 A

116 **Methodology**

117 This study employed secondary data analysis of the 2015-2016 Tanzanian DHS. that covered
118 all 30 regions in Tanzania mainland and Zanzibar. The the National Bureau of Statistics (NBS)
119 and Office of the Chief Government Statistician (OCGS), Zanzibar, the Ministry of Health,
120 Community Development, Gender, Elderly and Children, Mainland, and the Ministry of
121 Health, Zanzibar implemented the DHS and ICF provided technical assistance. The survey was
122 funded by the Government of Tanzania; United States Agency for International Development
123 (USAID); Global Affairs Canada; Irish Aid; United Nations Children's Fund (UNICEF); and
124 United Nations Population Fund (UNFPA).

125 The DHS used a multi-stage cluster sampling from the 2012 Tanzanian census, whereby 608
126 clusters were selected, ending up with a representative probability sample of 13,376 households
127 and a total of 13,266 women aged 15 to 49 were interviewed in this survey. In this study we
128 excluded all women who did not take part in the domestic violence module and thus ending
129 with 7,597 women.. The DHS domestic violence module adheres to strict protocols on safety
130 and confidentiality of the study participants and fieldworkers (Ministry of Health, 2016).

131 Permission to conduct data analysis was sought through DHS program website
132 (<http://dhsprogram.com/data/available-datasets.cfm>)

133 **Measures**

134 The help seeking outcome was determined by the first question which all women who reported
135 any form of physical or sexual violence were asked “Thinking about what you yourself have
136 experienced among the different things we have been talking about, have you ever tried to seek
137 help?” If she responded positively, she was asked who she sought help from, with the answer
138 categories a) own family, b) husband’s / partner’s family, c) friend, d) neighbour, e) religious
139 leader, f) doctor/medical personnel g) police, h) lawyer, i) social services. In the analysis,
140 seeking help was categorised into four groups, seeking help from anyone included seeking help
141 from anyone in a)-i), seeking help from family, either their own family or partner’s family,
142 seeking help from friends and neighbours and lastly seeking help from official sources
143 including the police, lawyers, social services or religious leaders. Given the high prevalence
144 rates for informal help-seeking suggested in other studies (REF the other African studies) in
145 sub-Saharan Africa, informal help-seeking was broken down into own family, his family and
146 friends and neighbours.

147 The domestic violence module in the DHS 2015-2016 is based on the conflict tactics scale that
148 asks respondents if they have experienced physical or sexual IPV, psychological abuse and
149 controlling behaviours. Women were asked if they had experienced the above forms of IPV in
150 their lifetime and the past 12 months. For physical IPV women were asked seven questions if
151 they had ever been pushed, shook or something thrown at them, ever been slapped, ever been
152 kicked or dragged or beat up, ever been strangled or burnt, arm got twisted or hair pulled, ever
153 been punched with a fist or hit by something harmful and lastly if they had ever been threatened
154 by a knife/gun or any other weapon. For sexual IPV, women were asked three questions, if

155 they have ever been physically forced into unwanted sexual intercourse, physically forced to
156 perform any other sexual acts and if they have ever been threatened to perform any other sexual
157 acts. Psychological IPV had a total of four questions, if the partner had insulted them, being
158 belittled in front of other people, intimidate on purpose and if the partner threatened to hurt the
159 respondent. For controlling behaviours five questions were asked, if partner had tried to restrict
160 seeing friend, restrict contact with family of birth, insisted on knowing where the respondent
161 was most of the times, acted jealous or angry if the respondent spoke to another man and often
162 suspicious if is unfaithful. A woman was considered to have experienced IPV if she answered
163 yes to any question within any of forms of IPV.

164 Factors considered important influencers that are perceived to influence women's help-seeking
165 that were considered in the analysis include women's age, marital status, educational
166 attainment, partner age, partner's education, partner's occupation, duration of relationship, if
167 the woman is working, number of living children, decision on earnings, partner's alcohol intake
168 and other outcomes of IPV such as being afraid of the partner most of the time and having eyes
169 injuries, sprains, dislocations, or burns as a result of IPV.

170 **Analysis**

171 This study made use of the Tanzania DHS's individual women's data of 2015-16, who
172 participated in the domestic violence module and were either in a relationship at the time of the
173 interview or had been previously in a relationship meaning during the time of the interview
174 they were either separated, divorced or widowed. Data was analysed by STATA 15, weights
175 recommended by the DHS were used to adjust for sampling design and domestic violence
176 survey participation.

177 Frequencies were run to estimate the prevalence of different forms of IPV among all women
178 who participated in the domestic violence module and the prevalence of different forms of help

179 seeking among all women who experienced IPV. In the first step, we ran separate frequencies
180 for all women on women's and their partner's socio-demographic and relationship
181 characteristics and for all women who reported they had experienced IPV and those who sought
182 help. The sample was later reduced to women who reported any lifetime physical and/or sexual
183 IPV, to determine factors associated with help seeking among women who had experienced
184 physical and /or sexual IPV. We then screened for all potential factors that could influence our
185 dependent variable any help-seeking, with separate analyses conducted for the different types
186 of help-seeking: help seeking from anyone, help seeking from his or her family, friends and
187 neighbours and official sources. Cross tabulations and chi-square statistics were carried out to
188 determine associations between IPV and all the different forms of helping seeking and other
189 socio-demographic characteristics. Variables that were significant in each of the different
190 categories of help seeking were then used in the multivariate logistic regression model.. We
191 used a probability value of $p \leq 0.05$ to define the level of statistical significance, and an odds
192 ratio < 1 represents a protective factor, where as an odds ratio > 1 was considered a risk factor.

193 **Results**

194 A total of 7,597 women were included in the analysis, aged 15- 49 with a mean age of 32. Half
195 of the women (52%, $n=3899$) were between 31 and 49 years old, and a small proportion (6.6%
196 $n=466$), were between 15 to 19 years at the time of the interview. The majority of women had
197 primary education (66.8%, $n=4,833$), were working at the time of the interview (80.9%, $n=$
198 6,048), married (82.7%, $n=6,479$) and had 3 to 4 living children (55.4%, $n=4,329$) Women's
199 partner's age ranged from 17 to 91 with a mean age of 38. Of them, 39.1 % ($n=2,679$) were
200 between 30 and 40 years old and 22.5% ($n=1,438$) were 17 to 29 years old. Primary education
201 was the most frequent among the partners (69.0%, $n = 4,290$) and more than half of them were
202 working in the agricultural sector (55.8%, $n=3,722$) (see ref. Table 1for details).

203 The lifetime prevalence of physical and/or sexual IPV was 41.6% (n =2,913) and that of past
204 12 months was 29.3% (n=2,037). Among the 2,913 women who experienced lifetime physical
205 and/or sexual IPV, 51% (n=1,472) reported help seeking from anyone, with 43.6% (n=1,233)
206 of women reporting they either sought help from his/her family, 32.1% (n=890) reporting
207 seeking help from her own family, 28.4% (n=796) reporting seeking help from his family,
208 10.8% (n=336) seeking help from either their friends or neighbours, 4.5% (n=110) seeking help
209 from the police, 2.6% (n=45) from religious leaders, 1.1% (n= 27) from social services, 0.4%
210 (n=10) from medical doctors and lastly 1.4% (n= 44) from lawyers (see ref. Table 2 for details).

211 Results displayed in Table X show that help seeking from anyone was associated with being
212 afraid of the partner most of the times (AOR 1.8, 95% CI: 1.4, 2.3), ever having had eye
213 injuries, sprains, dislocations, or burns because of husband/partner (AOR 1.6, 95% CI: 1.1,
214 2.2), emotional IPV (AOR 2.3 , 95% CI: 1.9, 2.8), partner's alcohol intake (AOR 1.3, 95% CI
215 1.1,1.6) and severe physical IPV (AOR 2.5, 95% CI: 1.7, 3.7) .

216 Help seeking from his or her family was associated with being previously married (AOR 1.4,
217 95% CI: 1.1, 1.7), having 1 to 2 children (AOR 1.4, 95% CI: 1.1, 1.7), being afraid of the
218 partner most of the times (AOR 1.7, 95% CI: 1.3, 2.2), ever having had eye injuries, sprains,
219 dislocations or burns because of husband/partner (AOR 1.4, 95% CI: 1.0, 1.9), emotional IPV
220 (AOR 1.9, 95% CI: 1.5, 2.3) and severe physical IPV(AOR 2.4, 95% CI: 1.5, 3.7).

221 Seeking help from official sources (police, social services ,lawyers, doctors & religious leaders)
222 was associated with being previously married (AOR 2.3, 95% CI: 1.6, 3.4), being afraid of the
223 husband/partner most of the time (AOR 1.8, 95 % CI: 1.1, 2.9), ever having had eye injuries,
224 sprains, dislocations or burns because of husband/partner (AOR 2.0, 95% CI:1.4 ,3.0) and
225 emotional IPV (AOR 1.9, 95% CI:1.2, 3.1).

226 Seeking help from neighbours and friends was associated with partner's alcohol intake (AOR
227 1.7, 95% CI: 1.2, 2.3) and emotional IPV (AOR 2.3, 95% CI: 1.6, 3.5).

228 **Discussion**

229 This study found that four out of 10 women experienced lifetime physical and/or sexual IPV
230 and three out of 10 women experienced physical and/or sexual IPV in the past 12 months in
231 Tanzania. Of those women who experienced lifetime physical and/or sexual IPV, half sought
232 help and help seeking from the respondents own family ranked the highest followed by seeking
233 help from the perpetrators family.

234 The findings are similar to a study of can you give an n? or some more infom on the study?
235 pregnant women in Tanzania, of which only a quarter of women who experienced IPV during
236 pregnancy sought help and reported similar low levels of formal help seeking (Katiti, Sigalla,
237 Rogathi, Manongi, & Mushi, 2016). A number of reasons have been suggested that prevent
238 women from seeking help in Tanzania. For example, IPV is normalized in the society and
239 therefore seen as insignificant, it is associated with shame and stigma and women do not trust
240 the available structures, corruption and the feeling they may not be able to attain the justice
241 they deserve (McCleary-Sills et al., 2016). Other factors that have been mentioned include
242 being threatened by the partner, being afraid that the family would find out about the violence
243 or the woman not wanting the family to know (Frias, 2013).

244 Another finding in our study is that many women who sought help, sought it mainly from the
245 their own family and the perpetrators family which is in contrast to the above mentioned study
246 from northern Tanzania among pregnant women (Katiti et al., 2016). That study found out that
247 pregnant women disclosed IPV more often to their own family then followed by friends. The
248 trend of women seeking help from the partner's family can be explained as a cultural aspect
249 that women in patrilineal societies become part of the men's family and all problems should be

250 reported to the family. Similar suggestions have been reported in Kenya where women sought
251 help from partner's family as they are the ones to settle marital disputes (Odero et al., 2013).

252 Another contributory factor across all forms of help seeking among women who have
253 experienced physical and/or sexual IPV is the severity of IPV and its overlap with other forms
254 of intimate partner abuse. Women who sought help also reported controlling behaviours and
255 emotional IPV. Not only were these women afraid of their partner most of the time, they also
256 suffered eyes injuries, sprains, dislocations, or burns because of the IPV. Across studies we see
257 that women wait for IPV to become serious before seeking help which can become lethal
258 (Evans & Feder, 2016; Stockl et al., 2013). Our results are also in line with a study conducted
259 in Ghana that showed that women who had a perceived risk of injury from physical violence
260 were more likely to seek help (Tenkorang et al.,2018). (

261 The findings also illustrate that family is an important cultural aspect in Tanzania. Currently,
262 interventions addressing IPV in sub-Saharan Africa are focused on the community level,
263 individual women and men or the couple as a whole. While those interventions are important
264 to empower women and men, challenges existing gender and social norms at both the
265 individual and community level, up to now they do not focus on the natal and in law family,
266 who may play a significant role in reducing IPV. Any intervention designed either to challenge
267 cultural norms or provide education on the effects of IPV to society at large should start with
268 the family unit. Our study, as several other studies indicate the importance of the family unit
269 in sub-Saharan Africa on reporting violence among couples (Odero et al., 2013; Okenwa et
270 al., 2009).

271 The DHS are rigorous done surveys with national representation, but it is important to note the
272 limitations of this study as it a quantitative cross-sectional study and thus failing to understand
273 in detail why women chose to report to their own family and partner's family instead of others.

274 In addition, due its cross-sectional nature, it is impossible to understand when women sought
275 help for the violence they experienced in their relationship and whether associated factors were
276 causes or consequences of help-seeking. Another limitation is that IPV is a sensitive topic
277 coupled with social stigma so IPV might be under-reported. Unfortunately, important
278 measures related to help-seeking such as social norms, shame stigma, trust in system were not
279 available to be included in model. Lastly, due to the study's cross-sectional nature we cannot
280 establish causality between IPV and help seeking, as in other cases help-seeking has been said
281 to cause more violence (McCleary-Sills et al., 2016).

282 **Conclusion**

283 IPV is a serious problem in Tanzania, with four out 10 women having experienced physical
284 and/or sexual IPV and only half of the women of the women who experienced lifetime IPV
285 sought help from anyone. The findings of the study highlight the need for interventions to
286 incorporate the wider family unit, including parents of married couples instead of only focusing
287 on empowering individual women. Another important aspect is the need for further research to
288 understand the dynamics of seeking help, especially on how to improve formal help seeking
289 by women in Tanzania, so that tailored recommendations can be made to the government to
290 approve existing services and their accessibility. The Tanzanian government has made an
291 important pledge to reduce the prevalence of IPV through its National Action Plan and further
292 research is needed on how to effectively support Tanzanian women who experience IPV.

293

294 **Declaration of Conflicting Interest**

295 The authors declare they have no conflicting interest with respect to the data analysis or
296 authorship and publication of this study.

297

298 **Funding**

299 Bathsheba Mahenge is currently funded by the Commonwealth Rutherford Fellowship. Heidi
300 Stöckl is funded by an ERC Starting Grant. The funders had no influence on this study.

301

302

303 **References**

304 Adams, A. E., Tolman, R. M., Bybee, D., Sullivan, C. M., & Kennedy, A. C. (2012). The impact of
305 intimate partner violence on low-income women's economic well-being: the mediating role
306 of job stability. *Violence Against Women, 18*(12), 1345-1367.

307

308 Ahmad, F., Driver, N., McNally, M. J., & Stewart, D. E. (2009). "Why doesn't she seek help for partner
309 abuse?" An exploratory study with South Asian immigrant women. *Social Science &*
310 *Medicine, 69*(4), 613-622.

311

312 Devries, K. M., Mak, J. Y., Bacchus, L. J., Child, J. C., Falder, G., Petzold, M., et al. (2013). Intimate
313 partner violence and incident depressive symptoms and suicide attempts: a systematic
314 review of longitudinal studies. *PLoS Med, 10*(5), e1001439.

315

316 Ellsberg, M., & Heise, L. (2002). Bearing witness: ethics in domestic violence research. *The Lancet,*
317 *359*(9317), 1599-1604.

318

319 Evans, M. A., & Feder, G. S. (2016). Help-seeking amongst women survivors of domestic violence: a
320 qualitative study of pathways towards formal and informal support. *Health Expect, 19*(1), 62-
321 73.

322

323 Fanslow, J. L., & Robinson, E. M. (2010). Help-seeking behaviors and reasons for help seeking
324 reported by a representative sample of women victims of intimate partner violence in New
325 Zealand. *J Interpers Violence, 25*(5), 929-951.

326

327 Frias, S. M. (2013). Strategies and help-seeking behavior among Mexican women experiencing
328 partner violence. *Violence Against Women, 19*(1), 24-49.

329

330 Garcia-Moreno, C., HA, J., Ellsberg, M., L, H., & Watts, C. (2005). WHO multi-country study on
331 womens health and domestic violence against women: initial results on prevalence health
332 outcomes and women responses. *Geneva. World health Organization.*

333

334 García-Moreno, C., & Stöckl, H. (2009). Protection of sexual and reproductive health rights:
335 addressing violence against women. *International Journal of Gynecology & Obstetrics,*
336 *106*(2), 144-147.

337

338 Jansen, H. A., Watts, C., Ellsberg, M., Heise, L., & Garcia-Moreno, C. (2004). Interviewer training in
339 the WHO multi-country study on women's health and domestic violence. *Violence against*
340 *women, 10*(7), 831-849.

341

342 Jouriles, E. N., McFarlane, J., Vu, N. L., Maddoux, J., Rosenfield, D., Symes, L., et al. (2018). Mothers'
343 posttraumatic stress and child adjustment problems in families seeking services for intimate
344 partner violence. *J Consult Clin Psychol, 86*(7), 604-614.

345

346 Katiti, V., Sigalla, G. N., Rogathi, J., Manongi, R., & Mushi, D. (2016). Factors influencing disclosure
347 among women experiencing intimate partner violence during pregnancy in Moshi
348 Municipality, Tanzania. *BMC Public Health, 16*, 715.

349

350 Kennedy, A. C., & Prock, K. A. (2016). "I Still Feel Like I Am Not Normal": A Review of the Role of
351 Stigma and Stigmatization Among Female Survivors of Child Sexual Abuse, Sexual Assault,
352 and Intimate Partner Violence. *Trauma, Violence, & Abuse, 19*(5), 512-527.

353

354 Liang, B., Goodman, L., Tummala-Narra, P., & Weintraub, S. (2005). A theoretical framework for
355 understanding help-seeking processes among survivors of intimate partner violence. *Am J*
356 *Community Psychol, 36*(1-2), 71-84.

357

358 Mahenge, B., Likindikoki, S., Stockl, H., & Mbwambo, J. (2013). Intimate partner violence during
359 pregnancy and associated mental health symptoms among pregnant women in Tanzania: a
360 cross-sectional study. *Bjog, 120*(8), 940-946.

361

362 Mapayi, B., Makanjuola, R. O., Mosaku, S. K., Adewuya, O. A., Afolabi, O., Aloba, O. O., et al. (2013).
363 Impact of intimate partner violence on anxiety and depression amongst women in Ile-Ife,
364 Nigeria. *Arch Womens Ment Health, 16*(1), 11-18.

365

366 McCleary-Sills, J., Namy, S., Nyoni, J., Rweyemamu, D., Salvatory, A., & Steven, E. (2016). Stigma,
367 shame and women's limited agency in help-seeking for intimate partner violence. *Global*
368 *Public Health, 11*(1-2), 224-235.

369

370 Ministry of Health, C. D., Gender, Elderly and Children (MoHCDGEC) (Tanzania Mainland), Ministry of
371 Health (MoH) (Zanzibar), National Bureau of Statistics (NBS), Office of Chief Statistician (OCGS)
372 and ICF International (2016). Tanzania Demographic survey and malaria indicator survey
373 (TDHS_MIS) 2015-2016. *Dar es Salaam, Tanzania and Rockville, Maryland, (MoHCDGEC) USA:*
374 *MoHSW, MoH, NBS, OCGS and ICF International.*

375

376 Neamah, H. H., Sudfeld, C., McCoy, D. C., Fink, G., Fawzi, W. W., Masanja, H., et al. (2018). Intimate
377 Partner Violence, Depression, and Child Growth and Development. *Pediatrics, 142*(1).

378

379 Odero, M., Hatcher, A. M., Bryant, C., Onono, M., Romito, P., Bukusi, E. A., et al. (2013). Responses
380 to and Resources for Intimate Partner Violence: Qualitative Findings From Women, Men,
381 and Service Providers in Rural Kenya. *Journal of Interpersonal Violence, 29*(5), 783-805.

382

383 Okenwa, L., Lawoko, S., & Jansson, B. (2009). Factors Associated with Disclosure of Intimate Partner
384 Violence among Women in Lagos, Nigeria. *Journal of Injury and Violence Research*, 1(1), 37-
385 47.

386

387 Overstreet, N. M., & Quinn, D. M. (2013). The Intimate Partner Violence Stigmatization Model and
388 Barriers to Help-Seeking. *Basic and applied social psychology*, 35(1), 109-122.

389

390 Palermo, T., Bleck, J., & Peterman, A. (2014). Tip of the iceberg: reporting and gender-based violence
391 in developing countries. *Am J Epidemiol*, 179(5), 602-612.

392

393 Ragusa, A. T. (2013). Rural Australian women's legal help seeking for intimate partner violence:
394 women intimate partner violence victim survivors' perceptions of criminal justice support
395 services. *J Interpers Violence*, 28(4), 685-717.

396

397 Roelens, K., Verstraelen, H., Van Egmond, K., & Temmerman, M. (2008). Disclosure and health-
398 seeking behaviour following intimate partner violence before and during pregnancy in
399 Flanders, Belgium: A survey surveillance study. *European Journal of Obstetrics & Gynecology*
400 *and Reproductive Biology*, 137(1), 37-42.

401

402 Stockl, H., Devries, K., Rotstein, A., Abrahams, N., Campbell, J., Watts, C., et al. (2013). The global
403 prevalence of intimate partner homicide: a systematic review. *Lancet*, 382(9895), 859-865.

404

405 Sylaska, K. M., & Edwards, K. M. (2014). Disclosure of intimate partner violence to informal social
406 support network members: a review of the literature. *Trauma Violence Abuse*, 15(1), 3-21.

407

408 Taket, A., O'Doherty, L., Valpied, J., & Hegarty, K. (2014). What do Australian Women Experiencing
409 Intimate Partner Abuse Want From Family and Friends? *Qual Health Res*, 24(7), 983-996.

410

411

412

413

414

415

416

417

418

419

420 Table 1: The association of women life time experiences of physical and/or sexual IPV and
 421 seeking help for IPV

422

Variables	Respondent's characteristics (n=7,597)	Ever experienced Physical and/or sexual IPV(n=2913)	Sought Help for IPV(n = 1,472)
Woman's age			
31+	3,899(52.5%)	1517(53.0%)*	800(54.9%)
20-30	3,232(40.9%)	1242(41.5%)	612(40.6%)
15-19	466(6.6%)	154(5.5%)	60(4.5%)
Woman's education			
none/incomplete primary	1,439(17.8%)	533(18.1%)*	258(17.4%)
Primary	4,833(66.8%)	2059(69.9%)	1048(70.6%)
secondary/higher	1,325(15.4%)	321(12.0%)	166(12.0%)
Woman is working			
No	1,549(19.1%)	464(16.5%)*	218(16.1%)
Yes	6,048(80.9%)	2449(83.5%)	1254(83.9%)
Marital status			
Was previously married	1,118(17.3%)	634(25.2%)*	401(31.1%)*
Currently married	6,479(82.7%)	2279(74.9%)	1071(68.9%)
Type of partnership			
polygamy	1250(16.0%)	488(17.1%)	227(15.9%)
Monogamy	6347(84.0%)	2425(82.9%)	1245(84.1%)
Duration of relationship			
10 years +	4333(57.1%)	1769(61.2%)*	929(62.9%)
5-9 years	1508(19.6%)	599(20.0%)	299(19.0%)
0-4 years	1756(23.2%)	545(18.8%)	244(18.0%)
Number of living children			
None	569(8.1%)	153(5.3%)*	62(4.3%)
3-4	4329(55.4%)	1762(59.8%)	919(59.1%)
1-2	2699(36.5%)	998(34.9%)	491(35.6%)
Partner's age			
41 +	2361(38.4%)	766(37.1%)	776(37.1%)
30 – 40	2679(39.1%)	988(40.9%)	988(40.9%)
17 -29	1428(22.5%)	514(22.0%)	514(22.0%)
Partners education level			
None /incomplete primary	835(11.9%)	292(13.9%)*	292(13.9%)
Primary	4290(69.0%)	1661(71.1%)	1661(71.1%)
Secondary/higher	1354(19.1%)	326(15.1%)	326(15.1%)
Partner's occupation			
Agriculture	3722(55.7%)	1380(59.8%)*	1380(59.8%)
Unskilled labour	852(31.0%)	296(13.0%)	296(13.0%)
Professional, clerical & sale	190(13.4%)	603(27.2%)	603(27.2%)

Decision making on how earnings are spent			
Woman only	1206(36.5%)	349(36.5%)	349(36.5%)
Woman & partner	1602(54.9%)	560(53.2%)	560(53.2%)
Partner alone	267(8.7%)	115(10.2%)	115(10.3%)
Who earns more			
Woman earns more than him	276(11.2%)	92(13.8%)	92(13.8%)
Husband/partner earns more	2198(68.7%)	743(68.2%)	743(68.2%)
About the same	605(20.1%)	189(18.0%)	189(18.0%)
Urban / Rural			
Rural	5560(67.6%)	2132(70.3%)*	1081(71.0%)
Urban	2067(32.4%)	781(29.8%)	391(29.0%)
Wealth index			
Poorest	1,379(18.2%)	617(45.6%)*	196(34.0%)
Poorer	1,414(18.6%)	586(45.5%)	165(29.7%)
Middle	1,559(20.5%)	623(43.2%)	185(29.9%)
Richer	1,753(23.1%)	614(40.3%)	201(33.9%)
Richest	1,492(20.1%)	473(35.1%)	143(32.8%)
Number of people living in the household			
1-5	4212(48.3%)	1591(46.0%)**	804(45.9%)
6-48	3385(51.7%)	1322(54.0%)	668(53.8%)
Partner's alcohol intake			
No	5208(66.5%)	1518(52.0%)*	688(46.2%)*
Yes	2389(33.5%)	1395(48.0%)	784(53.8%)
Eye injuries, sprains, dislocations or burns			
No	2502(85.4%)	2497(85.4%)	1224(85.0%)*
Yes	416(14.6%)	416(14.6%)	248(15.0%)
Afraid of the partner			
Never	4313(56.0%)	1028(35.2%)*	396(27.6%)*
Most	1123(16.5%)	833(29.3%)	538(36.5%)
Sometimes	2163(27.5%)	1052(35.5%)	538(36.0%)
Past year physical IPV			
No	5726(73.1%)		467(31.9%)*
Yes	1871(27.0%)		1005(68.1%)
Past year sexual IPV			
No	6898(90.1%)		1107((76.7%)
Yes	699(9.9%)		365(23.3%)
Past year physical and/or sexual IPV			
No	5560(70.7%)		430(29.2%)
Yes	2037(29.3%)		1042(70.8%)
Moderate physical IPV			
No	6240 (80.7%)		614(41.2%)*
Yes	1357(19.3%)		858(58.8%)

Severe Physical IPV			
No	4952(62.2%)		614(41.2%)* **
Yes	2645(37.8%)		858(58.8%)
Psychological /emotional IPV			
No	5145(64.1%)	1003(32.4%)* **	348(22.1%)* **
Yes	2452(36.0%)	1910(67.4%)	1124(71.9%)
Controlling behaviours			
No	2180(25.8%)	401(13.1%)* **	159(10.2%)* **
Yes	5417(74.2%)	2512(86.9%)	1313(89.8%)

423 * \leq 05, ** \leq 005, *** \leq 0005

424

425

426

427 Table 2: From who the women sought help for lifetime physical and/or sexual IPV

	n(%)
	N= 2913
Sought help from her family	
No	2023(67.9%)
Yes	890(32.1%)
Sought help from his family	
No	2117(71.6%)
Yes	796(28.4%)
Sought help from her friends	
No	2811(96.7%)
Yes	102(3.3%)
Sought help from the neighbour	
No	2637(91.6%)
Yes	276(8.4%)
Sought help from official sources	
No	2686(91.5%)
Yes	227(8.5%)

428

429

430

431

432

433

434 Table 3: Factors associated with women help seeking for lifetime physical and/or sexual IPV
 435 from anyone

Variable	AOR	95% CI	P value
Marital Status			
Previously married	REF		
Currently married	.63	(.49 ; .82)	< 0.0001
Partners alcohol intake			
No	REF		
Yes	1.3	(1.1 ; 1.6)	0.013
Afraid of the partner			
Never	REF		
Most of the times	1.8	(1.4 ; 2.3)	< 0.0001
Sometimes	1.4	(1.1 ; 1.8)	0.008
Eye injuries, sprains, dislocations or burns			
No	REF		
Yes	1.6	(1.2 ; 2.2)	0.003
Emotional IPV			
No	REF		
Yes	2.3	(1.9 ; 2.8)	< 0.0001
Controlling behaviours			
No	REF		
Yes	1.2	(.93 ; 1.6)	0.138
Severe physical IPV			
No	REF		
Yes	2.5	(1.7 ; 3.7)	< 0.0001