

1 **What is the Risk Posed to the Lateral Femoral Cutaneous Nerve During**
2 **the Use of the Anterior Portal of Supine Hip Arthroscopy and the**
3 **Minimally Invasive Anterior Approach for Total Hip Arthroplasty?**

4
5 **Abstract**

6
7 *Purpose*

8 The purpose of the study was to determine 1) What is the proximity of the lateral femoral
9 cutaneous nerve (LFCN) to the anterior portal (AP) used in supine hip arthroscopy (SHA)? 2)
10 What is the proximity of the LFCN to the incision in the minimally invasive anterior
11 approach (MIAA) for total hip arthroplasty? 3) What effect does lateralizing the AP have on
12 the likelihood of nerve injury? 4) What branching patterns are observable in the LFCN?

13
14 *Methods*

15 Forty-five hemipelves were dissected. The LFCN was identified and its path dissected. The
16 positions of the nerve in relation to the AP and the MIAA incision were measured.

17
18 *Results*

19 The AP intersected with 38% of nerves. In the remainder, the LFCN was located $5.7 \pm$
20 4.5mm from the portal's edge. Additionally, 44% of nerves crossed the incision of the MIAA.
21 Of those that did not, the average minimum distance from the incision was $14.4 \pm 7.0\text{mm}$. We
22 found a significant reduction in risk if the AP is moved medially by 5mm or laterally by
23 15mm ($P = .0054$ & $P = 0.0002$). The LFCN showed considerable variation with four
24 branching variants.

25

26 *Conclusions*

27 These results show the LFCN is at high risk during SHA and the MIAA, emphasizing the
28 need for meticulous dissection. We suggest that relocation of the AP 5mm medially or 15mm
29 laterally will reduce the risk to the LFCN.

30

31 *Clinical Relevance*

- 32 - These results provide insight into the location of the LFCN in relation to the AP and
33 the incision of the MIAA.
- 34 - Relocation of the AP 5mm medially or 15mm laterally may significantly reduce the
35 risk of LFCN injury.

36

37 **Introduction**

38

39 The growing use of minimally invasive surgical techniques combined with the widespread
40 implementation of enhanced recovery programs has led to dramatic reductions in the length
41 of stay for patients undergoing elective hip procedures¹⁻⁴. These advances are exemplified by
42 two techniques: arthroscopic hip surgery and the minimally invasive anterior approach to the
43 hip (MIAA). Improvements in technical skills and instrumentation have advanced our ability
44 to accurately diagnose and arthroscopically treat an increasing number of hip pathologies,
45 with patients experiencing greatly reduced recovery time compared to traditional open
46 procedures^{2,5-8}. Hip arthroscopy can be performed with the patient in the lateral or supine
47 positions, with the surgeon utilizing a variety of portals in order to gain access to the joint.
48 For example, when operating with the patient in the supine position, the anterior portal (AP)
49 penetrates the muscle bellies of Sartorius and Rectus Femoris before entering the joint
50 capsule^{9,10}. This portal is one of the three main arthroscopy portals, and is placed at the

51 intersection of a sagittal line drawn inferiorly from the ASIS and a line drawn medially along
52 the superior margin of the greater trochanter ¹¹.

53

54 The MIAA involves significantly shortening and lateralizing the Smith-Petersen incision
55 whilst continuing to utilize the intermuscular plane between Sartorius and Tensor Fascia Lata
56 to access the hip joint ¹². The skin incision used in the MIAA begins proximally at a point
57 20mm lateral and 20mm distal to the Anterior-Superior Iliac Spine (ASIS), and runs distally
58 for 100mm towards a point 20mm lateral to the head of the fibula ¹². Although similar to the
59 Hueter approach's incision, this approach utilizes a slightly more distal incision, and the
60 angle of incision is less oblique. This technique's small incision and soft tissue preservation
61 means it is now considered highly effective for minimally invasive total hip arthroplasty, and
62 its lateral position is thought to reduce the risk of nerve injury ^{1,12,13}.

63

64 During both supine hip arthroscopy (SHA) using the AP and the MIAA, the structure often
65 quoted as being most at risk is the lateral femoral cutaneous nerve (LFCN), with a reported
66 incidence of injury ranging from 0.1-16.5% for SHA, and 0.2-25% for the MIAA¹⁴⁻²⁰. Such
67 injury can have serious consequences for a patient's recovery and quality of life, with
68 symptoms ranging from transient neurapraxia to long term sensorineural dysfunction ^{20,21}.
69 Although previous cadaveric studies into the risk during AP placement have emphasized the
70 close proximity of the LFCN to the AP, these studies are few in number and small in size
71 ^{19,22-24}. Previous cadaveric studies into this risk during the MIAA have focused on the path of
72 this nerve in relation to bony landmarks, not the nerve's proximity to the incision at its
73 lateralized site ^{25,26}. Furthermore, despite repeated documentation into the variability of the
74 LFCN anatomy, only one study has attempted to categorize the anatomical variants and
75 assess the impact these variants have on surgical procedures ²⁷.

76

77 The purpose of this cadaveric study was to determine 1) What is the proximity of the LFCN
78 to the AP used in SHA? 2) What is the proximity of the LFCN to the incision in the MIAA?
79 3) What effect does lateralizing the AP have on the likelihood of nerve injury? 4) What
80 branching patterns are observable in the LFCN? We hypothesize that the LFCN will be in
81 close proximity to the AP of SHA and will intersect with the incision line of the MIAA in the
82 majority of cadavers, despite its more lateral site.

83 **Methods**

84

85 A total of 45 hemipelves from 39 formalin preserved cadavers (20 males and 19 females,
86 with no lower limb or spinal abnormalities and no history of spinal or hip surgery) with a
87 mean age of 83.9 years (range 62-100) were dissected in this study. 45 cadavers were used as
88 this was the maximum number available for use. Though an a priori power calculation was
89 not performed, a post-hoc power calculation showed our study to have a power of 93% (with
90 $\alpha = 0.05$). Use of cadavers in this study was approved by the Human Tissue Authority
91 and all specimens were assessed prior to their inclusion in the study by JL, with no specimens
92 excluded from this study. Midline abdominal and bilateral subcostal incisions were made and
93 the abdominal viscera removed, allowing visualization of the psoas muscle. The LFCN was
94 identified in the abdomen based on its position lateral to the psoas muscle and traced distally
95 to the inguinal ligament (IL). The nerve was dissected out of the thigh to an inferior limit at
96 the level of the patella.

97

98 To locate the site of AP placement, one length of thread was traced inferiorly from the ASIS
99 and a second was traced medially along the line of the superior margin of the greater
100 trochanter. A pin was placed at the intersection of these two lines to indicate the position of
101 the AP. Two observers (JB and JL) were used for all measurements. The distance between all
102 the LFCN's branches and the pin were measured in the transverse plane using Vernier
103 Callipers. Additionally, the closest point the nerve travelled to the pin was marked and its
104 distance from the pin measured. This point's distance from the pin in the sagittal (y) and
105 transverse (x) planes was also noted, thus giving an x,y coordinate of the nerve's location
106 relative to the portal's center. For all measurements greater than 2.5mm from the pin, 2.5mm

107 was subtracted, thus allowing calculation of the nerve's distance from a 5mm arthroscopy
108 portal's edge.

109

110 In order to measure the LFCN in relation to the MIAA, a length of thread was used to mark a
111 line from a point 20mm lateral and 20mm distal to the ASIS to a point 20mm lateral to the
112 head of the fibula (lateralized Smith-Petersen Line). A pin was placed 100mm distal to the
113 proximal end of the thread, thus demarcating the incision used in the Minimally Invasive
114 Anterior Approach. The following were measured using Vernier Callipers:

- 115 - If the LFCN crossed the incision line, the distance from the proximal end of the
116 incision at which it crossed.
- 117 - If the LFCN did not cross the incision line, the minimum distance between the nerve
118 and the incision line, and the distance along the incision at which this occurred.
- 119 - If the LFCN crossed the lateralized Smith-Petersen (LSP) line distal to the incision,
120 the distance from the distal end of the incision line to the LFCN along the LSP line
121 (this is the distance the incision could safely be extend without risk of nerve injury).
- 122 - The length of the inguinal ligament (IL) and the distance from the ASIS to the point at
123 which the LFCN crosses it.
- 124 - The distance proximal or distal to the IL that the nerve first divides.

125

126 *Statistical Analysis*

127 All analyses were conducted using R version 3.2.1 (Vienna, Austria) (25). Using the position
128 of all LFCN branches in the transverse plane, the number of nerve-AP intersections and
129 misses were calculated for the current AP location and for AP location following
130 lateralization and medialization by 5mm, 10mm, 15mm, 20mm, 25mm or 30mm. 2 x 2 tables
131 were produced comparing the frequency of AP intersection and misses for the portal's current

132 location with each of the new locations. These tables were then analyzed using Pearson's
133 Chi-square test to assess significant differences between the current location and the new
134 location's AP intersection frequency. The relative risk reductions of nerve-portal
135 intersections at the new locations were also calculated. All other data are presented as mean \pm
136 standard deviation.

137 **Results**

138

139 *What is the proximity of the LFCN to the AP used in SHA?*

140 Out of the 45 nerves dissected, 17 (38%) were found to be within a 2.5mm radius of the pin
141 placed at the site of AP, with the remaining 28 (62%) nerves passing 5.7 ± 4.5 mm from the
142 portal's edge (**figure 1**). If the nerve passed laterally to the portal, the average closest
143 distance to the nerve in the transverse plane was found to be 5.4 ± 4.3 mm. If the nerve passed
144 medially, it was found to do so at a distance of 6.6 ± 5.1 mm.

145

146 *What is the proximity of the LFCN to the incision used in the MIAA?*

147 Of the 45 nerves dissected in this study, 20 (44%) crossed the incision line used in the MIAA
148 at a mean distance of 47.0 ± 28.0 mm distal along the incision (**figure 2a**). Of the 25 nerves
149 that did not cross the incision line, the mean minimum distance between the nerve and
150 incision was 14.4 ± 7.0 mm. This point occurred, on average, $74. \pm 37.3$ mm along and medial
151 to the incision. For 15 of these 25 nerves, this closest point occurred at the distal limit of the
152 incision (100mm). 20 of these 25 nerves went on to cross the LSP line at a point distal to the
153 incision. In these instances, the incision could have been safely extended a mean distance of
154 56.7 ± 46.1 mm along the LSP line before coming into contact with the LFCN (**figure 2b**).

155

156 *What effect does lateralizing the AP have on the likelihood of nerve injury?*

157 Based on all LFCN branches, we found a significant reduction in risk if the portal is moved to
158 a point 5mm medial or 15mm lateral to its current location, as only 5 and 2 nerves
159 respectively would have been immediately below the 2.5mm radius of the insertion point (P
160 = .0054 & $P = 0.0002$, relative risk reduction: 0.71 & 0.88) (**figure 3**).

161

162 *What branching patterns are observable in the LFCN?*

163 In our specimens, we observed four distinct branching patterns. Of the 45 nerves dissected,
164 29 (15 male and 14 female) displayed the classical branching pattern of the LFCN, giving rise
165 to distinct femoral and gluteal branches around the level of the IL (**figure 4**). In the remaining
166 16 cadavers, three novel branching patterns were seen: late, primary femoral and trifurcate. In
167 the 8 nerves (5 male and 3 female) that were described as ‘late branching’, the nerve was seen
168 to offer two branches after passing beyond the upper thigh (**figure 5**). 6 nerves (2 male and 4
169 female) offered no evident gluteal branch of the LFCN and instead continued down the
170 antero-lateral thigh. This was described as primary femoral branching (**figure 6**). In two
171 instances (one male and one female), the LFCN gave rise to three equally sized trunks that
172 travelled across the thigh. This was described as trifurcate branching (**figure 7**). In all the
173 dissected hips, the LFCN was observed to be located medial to the ASIS. In 5 cases (11%)
174 the LFCN divided before passing under the inguinal ligament (IL), in 28 (62%) cases the
175 LFCN divided after passing below the IL and in the remaining 12 cases (27%), at the level of
176 the IL. The average distance beyond the IL the nerve travelled before branching was $15.1 \pm$
177 23.7mm (**table 1**). When it was observed that the nerve divided before the IL, the branches
178 travelled together under the IL in all except two cases where the nerves passed under the IL
179 9mm and 19mm apart respectively. In one of these instances, the LFCN appeared to be
180 composed of two completely separate nerve trunks that originated in the abdomen and
181 travelled separately under the IL (**figure 8**).

182 **Discussion**

183

184 We found that in over a third of cases the LFCN intersected the AP, and that the LFCN
185 passed across the incision used in the MIAA in nearly half of cases, even with its 2cm of
186 lateralization, thus confirming our hypotheses. Additionally, we were able to show a
187 reduction in risk to the LFCN during SHA if the AP were moved 5mm medially or 15mm
188 laterally, and that there were four main LFCN branching patterns identifiable in our study
189 population.

190

191 Due to its close proximity to the AP's insertion point, the LFCN is considered to be the
192 structure most at risk during SHA ¹⁹. Previous smaller cadaveric studies have found the nerve
193 to lie 3-15mm from the AP insertion point on average, with considerable variation in the path
194 of the nerve relative to the AP ²²⁻²⁴. Similarly, a study by Wastson et al examined the path of
195 the LFCN in one hundred MRI scans and found the mean distance of the LFCN from the AP
196 insertion point to be 6.37mm (though this study did not account for portal diameter in its
197 measurements) ²⁸. These results are similar to those from our study, which found that in one
198 third of cases the nerve was directly deep to the AP's insertion point, and, when not deep to
199 the insertion point, was 5.7 ± 4.5 mm from the portal edge. This further highlights the
200 considerable risk to the LFCN during AP insertion and the need for meticulous dissection
201 before portal placement. In order to aid identification of the LFCN before portal insertion, we
202 assessed the proximity of the nerve's closest branch in the transverse plane, finding it to be
203 on average 5.4 ± 4.3 mm lateral or 6.6 ± 5.1 mm medial to the portal.

204

205 In our study, we found that the LFCN would lie in the path of the MIAA incision in nearly
206 half of patients, even with its 2cm of lateralization. This is important as the lateralized

207 incision site has been suggested to reduce the risk of LFCN injury ¹². Though higher than
208 Rudin et al's suggestion that in 33% of instances injury to the LFCN is unavoidable, this
209 difference is explained by differing definitions used in our studies ²⁷. In Rudin et al's work,
210 branching pattern classification was used an indicator of 'definite' injury and not each
211 branch's proximity to the incision used. Thus, branching patterns defined as not at definite
212 risk may still have offered branches that intersected the MIAA's incision line ²⁷.
213 Additionally, though Ropars and colleagues were able to suggest a region 27mm to 92mm
214 distal to the ASIS where the nerve was most at risk, we found considerable variance in the
215 distance along the incision at which the nerve crossed, meaning that a 'probable' location for
216 intraoperative use cannot be determined ²⁶. Furthermore, this study did not thoroughly assess
217 the nerve's location lateral to the ASIS. However, in cases where the nerve and incision line
218 do not cross, the closest point between the two most commonly occurred at the distal end of
219 the incision. The distal end of the incision therefore represents the primary area where the
220 LFCN is at risk of blind injury through soft tissue handling and muscle retraction.
221 Additionally, this study characterized the risk posed to the LFCN by extending the MIAA
222 incision. Twenty of the twenty-five nerves that did not cross the incision line went on to cross
223 the LSP line distally. However, there was large variation in the point at which the nerve
224 crossed the LSP line and therefore a recommendation of a safe extension distance for the
225 incision cannot be made.

226

227 Owing to the high number of AP-nerve intersects, we found a significant reduction in risk to
228 the LFCN if the portal were moved 5mm medial or 15mm lateral to its current location. At
229 these new locations only 5 and 2 nerves respectively were found below the portal, compared
230 with 17 at its current position ($P = .0054$ & $P = 0.0002$; relative risk reduction: 0.71 & 0.88).
231 These new locations offer preferable sites for portal placement as there is both a reduction in

232 the risk to the LFCN during the initial incision, and reduced need to dissect out the LFCN
233 trunk. These new locations are distinct from the mid-anterior portal, a portal devised to
234 reduce the risk of LFCN injury, with the mid-anterior portal placed approximately 7cm distal
235 and lateral to the AP. As a capsulotomy is performed immediately following insertion of the
236 AP, these alterations to the portal location are unlikely to significantly impact the
237 arthroscopic field of view. However, the effect this alteration may have on the work space is
238 unclear and may result in hand-to-hand abutting.

239

240 In this study, we have found the anatomy of LFCN to be highly variable, exhibiting four
241 distinct branching patterns - classical, late, primary femoral and trifurcate. As the late branch
242 variants give rise to the gluteal branch distal to the level of the greater trochanter, it is highly
243 likely to be at risk of injury during the MIAA, as the gluteal branch traverses the thigh
244 perpendicular to the incision line. Conversely, classical branch variants are more likely to
245 give rise to their gluteal branches proximal to the level of the incision and are thus at lower
246 risk of damage from the incision. Trifurcate branch variants are also at high risk of injury
247 owing to the close proximity of their 'middle' branch to the incision as it traverses the thigh.
248 Primary femoral branch variants lack a gluteal branch and thus carry a lower risk of injury,
249 with the nerve travelling down the anterolateral thigh, medial to the incision (**figure 7**). These
250 branch variants are similar to those described by Rudin et al, however we found a lower
251 frequency of 'fan-type' branching pattern, described here as trifurcate ²⁷. Additionally, the
252 trifurcate branching pattern we noted consisted of fewer branches and travelled more
253 medially than Rudin et al's 'fan-type'. They also describe the classical and late branching
254 patterns as a 'posterior-type' of branching and do not distinguish between the distance
255 travelled before the gluteal branch arises, meaning it is unclear how many of these branches

256 would have been at risk. However, the ‘sartorial-type’ of branching pattern they describe
257 follows the same definition as our primary femoral branching type.

258

259 *Limitations*

260

261 Although this study gives detailed information about the location of the LFCN and its risk of
262 injury, the primary limitation of this study is that our data gives no indicator of injury
263 severity, nor the impact on patients’ quality of life. Though we were able to accurately
264 quantify the risk of LFCN injury, the size of the nerve trunk intersecting with the AP, or
265 crossing the incision line, varied greatly. Thus, some of the nerves we deemed to have been at
266 risk may have only provided a minor contribution to the sensory innervation of the thigh and
267 therefore injury would not lead to significant sensory impairment. Redundancy within the
268 sensory innervation of the thigh could also reduce the impact LFCN injury has on a patient’s
269 quality of life. These facts are likely to explain the discrepancy between the high nerve-portal
270 intersection and nerve-incision rate found in this study, and the prevalence of LFCN injuries
271 in the literature. Additionally, during our study, the cadaver’s hips were not in traction as they
272 may be during SHA, nor were they extended as in the MIAA.

273

274 Furthermore, several factors may have altered the proximity of the LFCN to the AP and
275 MIAA when compared to real-life. The formalin used in the embalming process may have
276 altered the position of the LFCN by affecting the dimensions of the tissues surrounding it,
277 thus influencing its proximity to the MIAA incision site and the AP. The age of the
278 specimen’s dissected in this study may have had an impact on the anatomy. Age-related
279 quadriceps muscle atrophy may also have influenced the structures surrounding the LFCN,
280 particularly when compared to the younger population undergoing THA and SHA.

281 Measurements taken were also undertaken following removal of the soft-tissues overlying the
282 LFCN and may therefore have altered the relative position of the portal and incision to the
283 nerve.

284

285 Additionally, this study only offers insight into the risk posed to the LFCN when using the
286 AP. Another commonly used portal, the mid-anterior portal (MAP), though more distal and
287 lateral to the AP, may also place the LFCN at risk. In this study, we chose not to assess the
288 proximity of the LFCN to this portal as its location is based relative to the AP and the
289 anterolateral portals, not bony landmarks, thus limiting its reproducibility.

290

291 **Conclusions**

292

293 These results show the LFCN is at high risk during SHA and the MIAA, emphasizing the
294 need for meticulous dissection. We suggest that relocation of the AP 5mm medially or 15mm
295 laterally will reduce the risk to the LFCN.

296

297 **References**

- 298 1. Chimento GF, Pavone V, Sharrock N, Kahn B, Cahill J, Sculco TP. Minimally invasive
299 total hip arthroplasty: a prospective randomized study. *J Arthroplasty*. 2005;20(2):139-
300 144.
- 301 2. Rath E, Tsvieli O, Levy O. Hip arthroscopy: an emerging technique and indications. *Isr*
302 *Med Assoc J IMAJ*. 2012;14(3):170-174.
- 303 3. Wainwright T, Middleton R. An orthopaedic enhanced recovery pathway. *Curr Anaesth*
304 *Crit Care*. 2010;21(3):114-120. doi:10.1016/j.cacc.2010.01.003.
- 305 4. Wojciechowski P, Kusz D, Kopeć K, Borowski M. Minimally invasive approaches in
306 total hip arthroplasty. *Ortop Traumatol Rehabil*. 2007;9(1):1-7.
- 307 5. Byrd JWT. *Operative Hip Arthroscopy*. Springer Science & Business Media; 2012.
- 308 6. Colvin AC, Harrast J, Harner C. Trends in hip arthroscopy. *J Bone Joint Surg Am*.
309 2012;94(4):e23. doi:10.2106/JBJS.J.01886.
- 310 7. Glick JM, Valone F, Safran MR. Hip arthroscopy: from the beginning to the future--an
311 innovator's perspective. *Knee Surg Sports Traumatol Arthrosc Off J ESSKA*.
312 2014;22(4):714-721. doi:10.1007/s00167-014-2859-y.
- 313 8. Griffiths EJ, Khanduja V. Hip arthroscopy: evolution, current practice and future
314 developments. *Int Orthop*. 2012;36(6):1115-1121. doi:10.1007/s00264-011-1459-4.
- 315 9. Bedi A, Kelly BT, Khanduja V. Arthroscopic hip preservation surgery: current concepts
316 and perspective. *Bone Jt J*. 2013;95-B(1):10-19. doi:10.1302/0301-620X.95B1.29608.

- 317 10. Thomas Byrd JW. Modified Anterior Portal for Hip Arthroscopy. *Arthrosc Tech.*
318 2013;2(4):e337-e339. doi:10.1016/j.eats.2013.05.006.
- 319 11. Aprato A, Giachino M, Masse A. Arthroscopic approach and anatomy of the hip.
320 *Muscles Ligaments Tendons J.* 2016;6(3):309-316. doi:10.11138/mltj/2016.6.3.309.
- 321 12. Bal BS, Vallurupalli S. Minimally invasive total hip arthroplasty with the anterior
322 approach. *Indian J Orthop.* 2008;42(3):301-308. doi:10.4103/0019-5413.41853.
- 323 13. Siguier T, Siguier M, Brumpt B. Mini-incision anterior approach does not increase
324 dislocation rate: a study of 1037 total hip replacements. *Clin Orthop.* 2004;(426):164-
325 173.
- 326 14. Bal BS, Haltom D, Aleto T, Barrett M. Early complications of primary total hip
327 replacement performed with a two-incision minimally invasive technique. *J Bone Joint*
328 *Surg Am.* 2005;87(11):2432-2438. doi:10.2106/JBJS.D.02847.
- 329 15. Byrd JWT. Complications Associated with Hip Arthroscopy. In: MD JWTB, ed.
330 *Operative Hip Arthroscopy.* Springer New York; 2005:229-235. doi:10.1007/0-387-
331 27047-7_16.
- 332 16. Chan K, Farrokhyar F, Burrow S, Kowalczyk M, Bhandari M, Ayeni OR.
333 Complications following hip arthroscopy: a retrospective review of the McMaster
334 experience (2009–2012). *Can J Surg.* 2013;56(6):422-426. doi:10.1503/cjs.021712.
- 335 17. Macheras GA, Christofilopoulos P, Lepetsos P, Leonidou AO, Anastasopoulos PP,
336 Galanakos SP. Nerve injuries in total hip arthroplasty with a mini invasive anterior
337 approach. *Hip Int.* April 2016:0-0. doi:10.5301/hipint.5000352.

- 338 18. Simpson J, Sadri H, Villar R. Hip arthroscopy technique and complications. *Orthop*
339 *Traumatol Surg Res.* 2010;96(8, Supplement):S68-S76. doi:10.1016/j.otsr.2010.09.010.
- 340 19. Thorey F, Ezechieli M, Ettinger M, Albrecht U-V, Budde S. Access to the hip joint from
341 standard arthroscopic portals: a cadaveric study. *Arthrosc J Arthrosc Relat Surg Off*
342 *Publ Arthrosc Assoc N Am Int Arthrosc Assoc.* 2013;29(8):1297-1307.
343 doi:10.1016/j.arthro.2013.05.017.
- 344 20. Larson CM, Clohisy JC, Beaulé PE, et al. Intraoperative and Early Postoperative
345 Complications After Hip Arthroscopic Surgery: A Prospective Multicenter Trial
346 Utilizing a Validated Grading Scheme. *Am J Sports Med.* 2016;44(9):2292-2298.
347 doi:10.1177/0363546516650885.
- 348 21. Goulding K, Beaulé PE, Kim PR, Fazekas A. Incidence of Lateral Femoral Cutaneous
349 Nerve Neuropraxia After Anterior Approach Hip Arthroplasty. *Clin Orthop.*
350 2010;468(9):2397-2404. doi:10.1007/s11999-010-1406-5.
- 351 22. Byrd JW, Pappas JN, Pedley MJ. Hip arthroscopy: an anatomic study of portal
352 placement and relationship to the extra-articular structures. *Arthrosc J Arthrosc Relat*
353 *Surg Off Publ Arthrosc Assoc N Am Int Arthrosc Assoc.* 1995;11(4):418-423.
- 354 23. Elsaidi GA, Ruch DS, Schaefer WD, Kuzma K, Smith BP. Complications associated
355 with traction on the hip during arthroscopy. *Bone Jt J.* 2004;86-B(6):793-796.
356 doi:10.1302/0301-620X.86B6.14426.
- 357 24. Robertson WJ, Kelly BT. The safe zone for hip arthroscopy: a cadaveric assessment of
358 central, peripheral, and lateral compartment portal placement. *Arthrosc J Arthrosc Relat*
359 *Surg Off Publ Arthrosc Assoc N Am Int Arthrosc Assoc.* 2008;24(9):1019-1026.
360 doi:10.1016/j.arthro.2008.05.008.

- 361 25. Chen LH, Huang QW, Wang WJ, He ZR, Ding WL. The applied anatomy of anterior
362 approach for minimally invasive hip joint surgery. *Clin Anat N Y N*. 2009;22(2):250-
363 255. doi:10.1002/ca.20750.
- 364 26. Ropars M, Morandi X, Hutten D, Thomazeau H, Berton E, Darnault P. Anatomical study
365 of the lateral femoral cutaneous nerve with special reference to minimally invasive
366 anterior approach for total hip replacement. *Surg Radiol Anat SRA*. 2009;31(3):199-204.
367 doi:10.1007/s00276-008-0433-3.
- 368 27. Rudin D, Manestar M, Ullrich O, Erhardt J, Grob K. The Anatomical Course of the
369 Lateral Femoral Cutaneous Nerve with Special Attention to the Anterior Approach to
370 the Hip Joint. *J Bone Joint Surg Am*. 2016;98(7):561-567. doi:10.2106/JBJS.15.01022.
- 371 28. Watson JN, Bohnenkamp F, El-Bitar Y, Moretti V, Domb BG. Variability in locations
372 of hip neurovascular structures and their proximity to hip arthroscopic portals. *Arthrosc*
373 *J Arthrosc Relat Surg Off Publ Arthrosc Assoc N Am Int Arthrosc Assoc*.
374 2014;30(4):462-467. doi:10.1016/j.arthro.2013.12.012.

375

376 **Tables**

377

378 ***Table 1 – The Lateral Femoral Cutaneous Nerve’s Branching and Location with Regards***
379 ***to the Inguinal Ligament***

380

Distance between the anterior superior iliac spine and nerve along the inguinal ligament in mm n = 47	Nerve location medial to the anterior superior iliac spine as % distance along inguinal ligament n = 47	Distance beyond inguinal ligament nerve travels before branching in mm n = 44
19.5 ± 15.07	15.1 ± 11.91	15.1 ± 23.7

381

382

383

384 **Figure Legends**

385

386 **Figure 1.** Position of LFCN's closest point in relation to an anterior portal on the right hip
387 and diagram of AP position on a hip for orientation. A: The x-axis represents the transverse
388 plane, negative values indicate distances laterally and positive values medially. The y-axis
389 represents the sagittal plane, positive values indicate distances superiorly and negative values
390 inferiorly. The x,y intercept represents the center of the anterior portal and the red ring
391 indicates the portal's position. Blue dots indicate the position of the LFCN at its closest point.
392 It should also be noted that 15 data points are at the origin. B: The anterior portal location
393 depicted as the intersection of two lines – a sagittal line drawn inferiorly from the anterior
394 superior iliac spine and a transverse line drawn medially from the greater trochanter.

395

396 **Figure 2.** Position of LFCN in relation to incision line of the MIAA on a right hip with
397 extension and diagram of the MIAA on a hip for orientation. A: X-axis represents the incision
398 line. Red dots indicate points where the nerve crossed the incision line, green dots indicate
399 position of the nerve at its closest point if it did not cross the incision line and orange dots
400 represent points where the nerve crossed upon extension of the incision. B: The MIAA
401 incision beginning 2cm inferior and 2cm lateral to the anterior superior iliac spine, and
402 continue along the lateralized Smith-Petersen line for 10cm.

403

404 **Figure 3.** Position of all LFCN branches in the transverse plane and effect of moving portal.
405 Red bar indicates nerve branches beneath current portal location. Blue bars indicate nerve
406 branches beneath portal location following medial and lateral relocation. Differences between
407 original portal location and relocalised position are all significant ($p<0.05$) unless otherwise
408 indicated in the graph (n/s).

409

410 **Figure 4.** Classical LFCN Branching Pattern on left thigh. The A flag indicates the position
411 of the ASIS. The B flag indicates the position of the nerve at the level of the inguinal
412 ligament. The C and D flags indicate the femoral and gluteal branches of the LFCN
413 respectively. The path of the incision used in the MIAA is represented by the red string
414 between the E and F flags.

415

416 **Figure 5.** LFCN Late Branching Pattern on right thigh. The A flag indicates the position of
417 the ASIS. The B flag indicates the position of the nerve at the level of the inguinal ligament.
418 The C and D flags indicate the femoral and gluteal branches of the LFCN respectively. The
419 path of the incision used in the MIAA is represented by the red string between the E and F
420 flags.

421

422 **Figure 6.** LFCN Primary Femoral Branching Pattern on left thigh. The A flag indicates the
423 position of the ASIS. The B flag indicates the position of the nerve at the level of the inguinal
424 ligament. The C flags indicate the various femoral branches of the LFCN. The path of the
425 incision used in the MIAA is represented by the red string between the D and E flags.

426

427 **Figure 7.** LFCN Trifurcate Branching Pattern on right thigh. The A flag indicates the
428 position of the ASIS. The B flag indicates the position of the nerve at the level of the inguinal
429 ligament. The C flags indicate the three main nerve trunks of the LFCN. The path of the
430 incision used in the MIAA is represented by the red string between the D and E flags.

431

432 **Figure 8.** Novel two-trunk LFCN anatomy on left side. The LFCN was composed of two
433 completely separate nerve trunks that originated in the abdomen and travelled separately

434 under the inguinal ligament. A flag indicates the position of the ASIS. The B & C flags
435 highlight the two separate trunks as they pass beneath the inguinal ligament.