

**COMPARISON OF DNA EXTRACTION METHODS IN THE DETECTION
OF GENETICALLY MODIFIED FOODS USING POLYMERASE
CHAIN REACTION**

NGUYEN CHAU THANH TUNG

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2004

**COMPARISON OF DNA EXTRACTION METHODS IN THE DETECTION
OF GENETICALLY MODIFIED FOODS USING POLYMERASE
CHAIN REACTION**

By

NGUYEN CHAU THANH TUNG

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

December 2004

DEDICATION

I wish to dedicate this work to my beloved parents, Mr. Nguyen Chau Boi and Mrs. Nguyen Thi Thanh Lan, who sacrificed their lives for my study

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Science

**COMPARISON OF DNA EXTRACTION METHODS IN THE DETECTION
OF GENETICALLY MODIFIED FOODS USING POLYMERASE
CHAIN REACTION**

By

NGUYEN CHAU THANH TUNG

December 2004

Chairman: Professor Son Radu, PhD

Faculty: Food Science and Technology

Genetically modified organisms (GMOs) can be defined as organisms in which genetic materials have been altered in the way that does not occur naturally by mating or natural combination. According to Novel Food Regulation (EC/258/97, EC/1139/98, EC/49/2000, EC/50/2000 and EC/1829/2003), foods and food ingredients derived from GMOs are strictly regulated and are labeled mandatorily. Polymerase chain reaction (PCR) method is used to detect GM events in foods. The specific objectives of this study are to establish a sensitive, robust and rapid operation method for detection of GM events by using PCR and to conduct a preliminary survey for distribution of animal feeds and foods derived from GM events in both Malaysia and Vietnam.

The two critical factors taken into account to achieve these objectives are applicability of different DNA extraction methods for each kind of samples and PCR amplification conditions.

Five DNA extraction methods (the Wizard method from Switzerland, the modified Wizard method with addition of beta-mercaptoethanol, the combination of Wizard and CTAB method, the CTAB method from Germany and the modified CTAB method with addition of beta-mercaptoethanol method) were optimized. The yield and quality of DNA obtained from raw soyabean, raw maize, animal feed, smooth tofu and soya milk samples were examined to determine the optimum method for each kind of sample.

The results of comparative analysis showed that the CTAB method was the most optimal protocol for extracting total DNA from raw soyabean, raw maize and animal feed samples with 32.7, 28.4, 33.4 ng DNA/mg sample in yield, respectively. In addition, the DNA quality (the ratio of A_{260}/A_{280} ranged from 1.9 to 2) was good enough not only for PCR amplification but also for DNA sequencing. However, the Wizard method was the best candidate for DNA isolation from smooth tofu (13.2 ng DNA/mg sample) and soya milk (3.4 ng DNA/mg sample) with relatively high quality of DNA (A_{260}/A_{280} ratio was 1.7 and 1.9, respectively).

The results of this survey showed that 20 out of 24 animal feed samples contaminated by at least one of three introduced genetic elements as promoter (P35S), terminator (NOS) and structural gene (EPSPS). In particular, all of the 16 animal feed samples from Malaysia and four out of eight animal feeds from Vietnam were GM-contaminant products. In contrast, neither soybean samples (12 samples) nor maize samples (24 samples) were positive with these assays. Therefore they were categorized as non-GM products.

These results revealed that PCR amplification method provides the key advantages of high sensitivity, and robust and rapid operation whilst providing the requisites of careful experimental design that avoids both false-negative and/or false-positive results. Five primer pairs of LEC1/LEC2; ZE03/ZE04; P35S 1-5'/P35S 2-3'; HA-NOS118-F/HA-NOS118-R and EPSPS 1-5'/EPSPS 3-3' chosen in this study produced the amplicons of 164, 277, 101, 118 and 118 base pair, respectively that fulfilled the product-size requirement and completed the whole detection procedure of GM events for raw soybean and raw maize as well animal feed samples. In addition, PCR amplification and DNA sequencing protocols presented in this study should provide a very useful tool for routine GM event detection in foods and feeds with regards to false-negative and/or false-positive results.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia dalam memenuhi keperluan untuk ijazah Master Sains

**PERBANDINGAN BAGI KAEDAH PENGEKSTRAKAN DNA DI DALAM
PENGESANAN MAKANAN TERUBAHSUAI GENETIK MENGGUNAKAN
TINDAKBALAS RANTAIAN POLIMERASE**

Oleh

NGUYEN CHAU THANH TUNG

December 2004

Pengerusi: Professor Son Radu, PhD

Fakulti: Sains Makanan dan Teknologi

Organisma terubahsuai genetik (GMO) boleh di definisikan sebagai organisma di mana pengubahsuaian kandungan genetikanya tidak berlaku secara kombinasi semulajadi. Berdasarkan “Novel Food Regulation” (EC/258/97, EC/1139/98, EC/49/2000, EC/50/2000 dan EC/1829/2003), makanan dan kandungan makanan terbitan dari GMOs adalah dikawal rapi dan penglabelan diwajibkan. Kaedah tindakbalas rantaian polimerase (PCR) digunakan untuk mengecam GMOs dalam makanan. Objektif-objektif spesifik dalam kajian ini adalah untuk mengukuhkan kaedah operasi dalam pengecaman GMO dengan PCR yang sensitif, tegap and pantas serta mengendalikan pemeriksaan saringan terhadap pengagihan makanan haiwan dan makanan terbitan dari GMOs dalam Malaysia dan Vietnam.

Dua faktor kritikal yang diambil kira dalam mengecapi objektif-objektif tersebut adalah aplikasi kaedah ekstraksi DNA yang berlainan untuk setiap sampel dan keadaan amplifikasi PCR.

Lima kaedah ekstraksi DNA (kaedah Wizard dari Switzerland, kaedah Wizard yang diubahsuai dengan penambahan beta-mercaptoethanol, kombinasi kaedah Wizard dan CTAB, kaedah CTAB dari Germany dan kaedah CTAB yang diubahsuai dengan penambahan beta-mercaptoethanol) telah diperbaikpulihan. Hasil and ketulenan DNA yang didapati dari sampel seperti kacang soya, jagung mentah, makanan haiwan, tauhu lembut dan susu soya diselidik untuk menentukan kaedah yang paling optimum untuk setiap sampel berkenaan.

Keputusan dari analisis perbandingan memaparkan bahawa kaedah CTAB merupakan protokol yang paling optimam untuk mengekstrak jumlah DNA dari sampel kacang soya mentah, jagung mentah dan makanan dengan nilai yang diperolehi ialah 32.7, 28.4, 33.4 ng DNA/mg masing-masing. Kualiti DNA (nisbah A_{260}/A_{280} antara 1.9 to 2) adalah mencukupi untuk amplifikasi PCR dan jujukan DNA. Akan tetapi, kaedah Wizard merupakan pilihan untuk isolasi DNA dari tauhu lembut (13.2 ng DNA/mg sampel) dan susu soya (3.4 ng DNA/mg sampel) dengan kualiti DNA (nisbah A_{260}/A_{280} adalah 1.7 dan 1.9 masing-masing) yang tinggi perbandingannya.

Keputusan tinjauan menunjukkan bahawa 20 daripada 24 sampel makanan haiwan mengandungi kandungan GM sekurang-kurang satu daripada tiga unsur-unsur genetik iaitu “promoter” (P35S), “terminator” (NOS) dan “structural gene” (EPSPS). Kesemua 16 sampel makanan haiwan dari Malaysia dan 4 daripada 8 makanan

haiwan dari Vietnam merupakan produk kontaminasi-GM. Sebaliknya, tiada sampel kacang soya (12 sampel) atau sample jagung (24 sampel) adalah positif dengan analisis tersebut. Oleh yang demikian, sampel tersebut boleh dikategorikan sebagai produk bukan GM.

Keputusan kajian menunjukkan bahawa kaedah amplifikasi PCR membekalkan kunci kelebihan dari segi sensitiviti, ketegapan dan operasi yang pantas sejurus membekalkan keperluan dalam rekabentuk eksperimen yang teliti untuk mengelakkan keputusan negatif-tiruan dan positif-tiruan. Lima pasang primer khususnya LEC1/LEC2; ZE03/ZE04; P35S 1-5'/P35S 2-3'; HA-NOS118-F/HA-NOS118-R dan EPSPS 1-5'/EPSPS 3-3' yang dipilih dalam kajian ini menghasilkan amplikons dengan pasangan bes sebanyak 164, 277, 101, 118 dan 118 masing-masing telah memenuhi syarat saiz-produk dan menyelesaikan keseluruhan kaedah pengesanan GMO untuk kacang soya mentah dan jagung mentah serta sampel haiwan makanan. Tambahan pula, protokol amplifikasi PCR dan jujukan DNA yang disampaikan dalam kajian ini boleh dikatakan kaedah yang bersesuaian dalam pengesanan GMO harian khasnya makanan manusia dan haiwan tanpa syak atas keputusan negatif-tiruan dan positif-tiruan.

ACKNOWLEDGEMENTS

I would like to express my sincere gratitude and deep appreciation to my supervisor, Professor Dr. Son Radu, and my former supervisor, Dr. Michael Wong Vui Ling for their dedicated supervision, advice and encouragement during my study in Universiti Putra Malaysia. My appreciation and gratitude are also extended to Associate Professor Dr. Raha Abdul Rahim and Associate Professor Dr. Lai Oi Ming for their advice and suggestions in the preparation of the thesis. In addition, I take this opportunity to thank Associate Professor Dr. Norihan Mohd Saleh for her valuable contributions to my presentation in the faculty. Finally, I would like to express my deep thanks to Professor Dr. Gulam Rusul Rahmat and Dr. Ho Chai Ling for their valuable help to arrange my viva earlier.

My sincere gratitude also goes to all my Vietnamese housemates as Mr. Dang, Mr. Thach and Mr. Ky and my colleagues especially Mr. Ong, Kqueen, Rozila, Daniel and Tuan in my workplace for their contributions and encouragements that enable me to complete this project. I am grateful to Cantho University of Vietnam for the financial sponsor which helps me to pursue my study in Universiti Putra Malaysia.

Finally, I would like to dedicate my work as a moral gift to my parents and my “tiny” friend from Vietnam for their support and spiritual encouragements.

I certify that an Examination Committee met on 20th December 2004 to conduct the final examination of Nguyen Chau Thanh Tung on his degree thesis entitled “Comparison of DNA Extraction Methods in the Detection of Genetically Modified (GM) Foods using Polymerase Chain Reaction (PCR)” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Ho Chai Ling, PhD

Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Suhaimi Napis, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Janna Ong Abdullah, PhD

Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Kasing Apun, PhD

Associate Professor
Faculty of Resource Science and Technology
Universiti Malaysia Sarawak
(Independent Examiner)

GULAM RUSUL RAHMAT ALI, PhD

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

SON RADU, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

RAHA ABDUL RAHIM, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

AINI IDERIS, PhD

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

NGUYEN CHAU THANH TUNG

Date: 24 December 2004

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xviii
LIST OF SYMBOLS AND UNITS	xix
CHAPTER	
1. INTRODUCTION	1
2. LITERATURE REVIEW	4
2.1 Definition of Genetically Modified Organisms (GMOs) and Typical Structure of a Genetically Modified Organism (GMO)	4
2.2 The Novel Food Regulation	9
2.3 Methods for Detection of a GMO or Its Molecular Derivatives	10
2.3.1 What is the Benefit of Detecting DNA Instead of Proteins or RNA?	10
2.3.2 Principles of the PCR Method as Applied to GMO Testing	14
2.3.3 PCR Reaction Kinetics	17
2.3.4 Different Methods of DNA Extraction Used for GMO Analyses	20
2.3.5 Choice of Different Target DNA Sequences	25
2.3.6 Methodology for the Analysis of a Sample	27
2.4 The Process of Detection of a GMO or Its Molecular Derivatives	30
2.5 The Current Possibility and Limitations of the Detection Methods	33
2.6 Multiplex PCR-based Detection Methods	45
3. MATERIALS AND METHODS	47
3.1 Certified Reference Materials	47
3.2 Sample Collection and Preparation	49
3.3 Methods for DNA Extraction	55
3.3.1 DNA Extraction Methods for the Experiment of Comparative Analysis	55
3.3.2 DNA Extraction Method Used for All Samples of the GMO Detection Experiment	58
3.4 Methods for DNA Quantification	59
3.5 Polymerase Chain Reaction (PCR)	59
3.6 Agarose Gel Electrophoresis	59
3.7 Statistical Analysis	62

3.8 Gel Purification for DNA Sequencing Using QIAquick® Gel Extraction Kit	62
3.9 Sequencing Analysis	63
4. RESULTS AND DISCUSSION	64
4.1 Comparative Analysis of Five Plant DNA Extraction Methods on Five Categories of Samples	64
4.1.1 Evaluation on DNA Yield (ng DNA/mg sample) of Five Plant DNA Extraction Methods on Five Categories of Samples	65
4.1.2 Evaluation on DNA Quality (A_{260}/A_{280} Ratio) of Five Plant DNA Extraction Methods on Five Categories of Samples	71
4.1.3 Confirmatory Assay of DNA Quality by Using PCR Amplification	80
4.2 Detection of Genetically Modified (GM) Events using Polymerase Chain Reaction (PCR)	84
4.2.1 Methodology for GM Event Analysis of A Sample	84
4.2.2 Total DNA Extraction	86
4.2.3 A Flow Model for PCR Detection of GM Event in Foods	90
4.2.4 Application of PCR Method on GM Events Screening	96
4.2.5 Application of PCR Method on Detection of Specific Traits in GM Events	102
4.2.6 Confirmatory Assay of PCR Products by Using DNA Sequencing	108
5. CONCLUSIONS AND RECOMMENDATIONS	115
5.1 Conclusions	115
5.2 Recommendations	116
REFERENCES	117
APPENDIX	126
BIODATA OF THE AUTHOR	137

LIST OF TABLES

Table	Page
2.1 The common traits of interest	7
2.2 Procedures for DNA extraction and purification from food matrixes	22
2.3 PCR methods grouped according to categories of specificity in GMO detection for maize	36
2.4 PCR methods grouped according to categories of specificity in GMO detection for soybean	41
3.1 Gene constructs in RR soybean	48
3.2 Description of the analyzed soybean samples in this study	50
3.3 Description of the analyzed maize samples in this study	51
3.4 Description of the analyzed animal feed samples in this study	53
3.5 Sequences of oligonucleotide primers used in this study	60
3.6 Amplification condition for PCR	61
4.1 Comparison of DNA yield (ng DNA/mg sample) obtained by five DNA extraction methods on five categories of samples	67
4.2 Single analysis on DNA yield (ng DNA/sample) for each kind of samples on five extraction methods	70
4.3 Comparison of DNA quality (A_{260}/A_{280}) obtained by five DNA extraction methods on five categories of samples	72
4.4 Quality and purity of total DNA from soybean, maize and animal feed samples extracted by the original CTAB protocol	87
4.5 Summary of sample status with three basic assays in GMO detection	106
4.6 Alternative promoters, terminators and structural genes in other GM plants	107

LIST OF FIGURES

Figure		Page
2.1	Transformation process	5
2.2	The integration site of a gene construct and the various molecules derived from reading and translating the DNA	8
2.3	Polymerase chain reaction (PCR) technique	12
4.1	Agarose gel electrophoresis (1%) of total DNA from five categories of samples from the original Wizard method	75
4.2	Agarose gel electrophoresis (1%) of total DNA from five categories of samples from the modified Wizard method	76
4.3	Agarose gel electrophoresis (1%) of total DNA from five categories of samples from the Wizard-CTAB combination method	77
4.4	Agarose gel electrophoresis (1%) of total DNA from five categories of samples from the original CTAB method	78
4.5	Agarose gel electrophoresis (1%) of total DNA from five categories of samples from the modified CTAB method	79
4.6	Agarose gel electrophoresis of PCR products produced by the original Wizard (lanes 1 to 5), the modified Wizard (lanes 7 to 11) and the Wizard-CTAB combination method (lane 13 to 17)	81
4.7	Agarose gel electrophoresis of PCR products produced by the original CTAB (lanes 1 to 5), the modified CTAB (lanes 7 to 11)	82
4.8	Agarose gel electrophoresis (1%) of total DNA from soybean samples extracted by the original CTAB protocol	88
4.9	Agarose gel electrophoresis (1%) of total DNA from maize samples extracted by the original CTAB protocol	89
4.10	Agarose gel electrophoresis (1%) of total DNA from AF samples extracted by the original CTAB protocol	89
4.11	Flow chart of PCR method for detection GMO in foods	91
4.12	Agarose gel electrophoresis (1.8%) of lectin PCR product from soybean samples	93
4.13	Agarose gel electrophoresis (1.8%) of lectin PCR product from AF	93

	samples	
4.14	Agarose gel electrophoresis (1.8%) of zein PCR product from maize samples	94
4.15	Agarose gel electrophoresis (1.8%) of zein PCR product from AF samples	94
4.16	Agarose gel electrophoresis (1.8%) of amplification products of 35S promoter from soybean samples	99
4.17	Agarose gel electrophoresis (1.8%) of amplification products of 35S promoter from maize samples	99
4.18	Agarose gel electrophoresis (1.8%) of amplification products of 35S promoter from AF samples	100
4.19	Agarose gel electrophoresis (1.8%) of amplification products of NOS terminator from soybean samples	100
4.20	Agarose gel electrophoresis (1.8%) of amplification products of NOS terminator from maize samples	101
4.21	Agarose gel electrophoresis (1.8%) of amplification products of NOS terminator from AF samples	101
4.22	Agarose gel electrophoresis (1.8%) of amplification product of EPSPS gene from AF samples	104
4.23	The comparison of partial sequence of P35S promoter from Cauliflower mosaic virus with the most five-respective sequences in database through BLASTN program	110
4.24	The comparison of partial sequence of EPSPS gene referring to glyphosate tolerance with the most two-respective sequences in database through BLASTN program	112
4.25	The comparison of partial sequence of NOS terminator from <i>A. tumefaciens</i> with the most five-respective sequences in database through BLASTN program	114

LIST OF ABBREVIATIONS

BLAST	Basic local alignment search tool
CaMV	Cauliflower mosaic virus
CTAB	Cetyl-trimethyl ammonium bromide
DNA	Deoxyribonucleic acid
dNTP	Deoxynucleoside triphosphate
EC	European Council
EDTA	Ethylene-diamine-tetra acetic acid
EPSPS	5-enolpyruvylshikimate-3-phosphate synthase
EtBr	Ethidium bromide
EU	European Union
GM	Genetically modified
GMO	Genetically modified organism
GMOs	Genetically modified organisms
NaOAc	Sodium acetate
NCBI	National Center for Biotechnology Information
P35S	35S promoter
PCR	Polymerase chain reaction
RNA	Ribonucleic acid
RNase	Ribonuclease
RR soybean	Roundup Ready soybean
SDS	Sodium dodecyl sulfate
TBE	Tris-Boric acid-EDTA buffer
UV	Ultraviolet

LIST OF SYMBOLS AND UNITS

bp	Base pair
β	Beta
$^{\circ}\text{C}$	Degree Celsius
U	Enzyme unit
kb	Kilo-base pair
μg	Microgram
μL	Microliter
mg	Milligram
mL	Milliliter
mM	Millimolar
M	Molar
ng	Nanogram
OD	Optical density
%	Percent
pmol	Picomole
rpm	Revolutions per minute
vol	Volume
v/v	Volume per volume
w/v	Weight per volume

CHAPTER 1

INTRODUCTION

Agricultural biotechnology has brought tremendous triumphs in the development of plants for the production of human food, animal feed and other products. In the few years since the commercial introduction of the first genetically modified product (FlavrSavr tomato), the cultivation of several transgenic crop species has increased rapidly by more than 25-fold, from 1.7 million hectares in 1996 (Roush, 2001) to 44.2 hectares worldwide in 2000 (Roush, 2001; Anklam *et al.*, 2002).

Genetically modified organisms (GMOs) contain specific traits which have been added to the organisms to improve their properties that have not occurred by mating or natural combination (Taverniers *et al.*, 2004; Anklam *et al.*, 2002). The addition of foreign genes has often been used in plants to produce novel protein that confer pest and disease tolerance and, more recently, to improve the chemical profile of process product, for example vegetable oils (Hemmer, 1997). Generally, the introduced gene works under the control of a promoter and a terminator. All of them (promoter, introduced gene and terminator) forms a gene construct and is inserted into the plants by means of biological, physical or chemical transformation methods. A genetically modified (GM) plant or product can thus be distinguished from its wild type counterpart by testing for the presence of the introduced DNA (Ahmed, 2002).

In European Union (EU) and other regions, the use of this technology, the consequent release of GMOs in the environment and the marketing of GMO-derived food products are strictly regulated (Huggett and Conzelmann, 1997; Anklam *et al.*, 2002).

The European Parliament and the Council of the EU have demanded in the Novel Food Regulation (EC/258/97, EC/1139/98, EC/49/2000, EC/50/2000 and EC/1829/2003), a clear and mandatory labeling of GMOs in food (Regulation EC, 1997; 1998; 2000; 2003). Because the Roundup-Ready soy bean and the Bt-176 maize were put on the market before Regulation EC/258/97 came into force, labeling of products derived from these GMOs is demanded by the separate Council Regulation EC/1139/98. On April 18th 2004, Regulation EC/1829/2003 on GM-food and feed came into force within the EU countries, regulating the placing on the market of such products and provided mandatory labeling thresholds of 0.9% for authorized GMOs and 0.5% for un-authorized GM events within the EU (Regulation EC, 2003). In addition, methods for specific detection of the most economically important GMOs are already available.

In contrast, Malaysian legislation is finalizing the enactment of food regulation for the presence of GMOs in animal feeds, human foods and ingredients. For this reason, there is a fundamental necessity to carry out a preliminary survey for the distribution of foods and feeds derived from GMOs and to establish an effective, accurate and reliable method for GM-events detection.

The objectives of this study are:

- i. to conduct a preliminary survey on distribution of animal feeds and foods derived from GM events in both Malaysia and Vietnam;
- ii. to optimize the DNA extraction methods for raw materials, animal feeds and processed foods;
- iii. to optimize the Polymerase Chain Reaction (PCR) method for the detection of GM foods.

CHAPTER 2

LITERATURE REVIEW

2.1 Definition of Genetically Modified Organisms (GMO) and Typical Structure of a GMO

Genetically Modified Organisms (GMOs) can be defined as organisms in which the genetic materials (DNA) have been altered in a way that does not occur naturally by mating or natural combination, that is by being genetically modified (GM) or by recombination DNA technology (Anklam *et al.*, 2002). In other words, GM products contain at least an additional trait encoded by an introduced gene(s), which generally produce an additional protein(s) that confers the trait of interest (Ahmed, 2002).

A GMO is a living organism, for example a plant, yeast or an organism whose genetic composition has been altered by means of gene technology. The genetic modification usually involves insertion of a piece of DNA (the insert), a synthetic combination of several smaller pieces of DNA, into the genome of the organism to be modified (Gachet *et al.*, 1999; Holst-Jensen, 2001). This process is called transformation (Figure. 2.1).