

**EMBRYOGENIC CALLUS FORMATION IN LOCAL COCOA  
(*Theobroma cacao* L.) CLONES**

**By**

**ASFALIZA RAMLI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the requirement for the Degree of Master of Science**

**Mac 2006**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the degree of Master of Science

**EMBRYOGENIC CALLUS FORMATION IN LOCAL COCOA  
(*Theobroma cacao* L.) CLONES**

By

**ASFALIZA RAMLI**

**March 2006**

**Chairman: Associate Professor Radzali Muse, PhD**

**Faculty: Biotechnology and Biomolecular Sciences**

A study on the potential of embryogenic callus formation in KKM 1, KKM 15, KKM 17, KKM 22, KKM 27, KKM 28, PBC 123, PBC 159, MHP 78, MHP 79, MHP 296, AMAZ, GS 29, EET 339 and MJS 47 local cocoa clones which had been released by MARDI was carried out. A sigmoid curve for callus growth, viability and total protein content of stem, leaf, matured and immature cotyledon derived callus were obtained. Based on this curve, the subculture interval was determined and it shows that the appropriate timing or period for subculture was between 20 to 24 days after the incubation. Obviously in this interval, the growth of callus was enormous and the total soluble protein was the highest. However, the viability was slowly decreased due to browning of the callus. Therefore only the white and friable callus was selected for subcultures.

Embryogenic callus was successfully induced for PBC 123 and MHP 296 cocoa clones by using immature cotyledon derived callus. Negative results were obtained

for all clones and explants tested using MS as the embryogenic induction medium but positive results were obtained for PBC 123 and MHP 296 clones using DKW and WPM basic media. The combinations of 2 mg/L 2,4-D and 0.25 mg/L Kin was the most successful auxin and cytokinin combination for embryogenic callus induction.. The nodular embryogenic callus of PBC 123 and MHP 296 emerged after the second subculture. Browning of callus occurred during incubation especially at the end of the culture period and was obvious for both clones. Enhancement of the embryogenic induction media by adding nitrogen rich compounds and amino acid such as casein hydrolysate, malt extract and proline did not improve the initiation of embryogenic callus. Instead, more friable callus were obtained from these enhanced media.

In direct somatic embryogenesis, 15 cocoa clones had been tested and four clones showed positive results. These clones (MJS 47, GS 29, PBC 123 and PBC 159) were able to produce embryogenic callus from staminodes explants using the protocol previously developed by Li et al (1998). The nodular embryogenic callus of MJS 47, PBC 123, PBC 159 and GS 29 emerged at the cut ends and were transferred onto a secondary callus growth medium and continued to proliferate on this medium.

Maturation of the embryogenic callus was carried out in a medium described by Li et al. (1998). The globular callus were pale brown in colour at the early maturation period but turned dark brown after 3 weeks of incubation period. None of the globular callus turned to heart shaped somatic embryo which is the second stage of somatic embryogenesis. The maturation of the globular callus was not achieved and the

embryogenic callus became brown and turned non viable at the end of the incubation period. All embryogenic clones showed similar response to the maturation medium.

Total soluble polyphenol content of embryogenic callus of PBC 123 after 16 days of incubation was  $259.94 \pm 15.53$   $\mu\text{g/g}$  fresh weight which was relatively lower than the non embryogenic callus ( $451.19 \pm 5.42$   $\mu\text{g/g}$  fresh weight). The increment or accumulation of the TSP was almost similar in MHP 296 immatured cotyledon derived callus especially towards the end of the incubation period. Furthermore, the embryogenic callus which were induced from immature cotyledon of MHP 296 and PBC 123 showed high polyphenol accumulation as compared to embryogenic callus induced from staminode of GS 29, PBC 123, PBC 159 and MJS 47. The browning of non embryogenic callus was greater than embryogenic callus. The specific peroxidase activity was slowly increased as the callus became embryogenic. The POD activity was  $164.27 \pm 9.42$  unit/mg protein for GS 29 embryogenic callus while for non embryogenic callus, the activity was around  $63.31 \pm 9.24$  unit/mg protein at 16 days of incubation. Even though a similar pattern of increment could be observed between the embryogenic and non embryogenic callus, the POD activities was higher in embryogenic callus as compared to non embryogenic callus.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGARUHAN KALUS EMBRIOGENIK BAGI KLON KOKO  
(*Theobroma cacao* L.)TEMPATAN**

Oleh

**ASFALIZA RAMLI**

**Mac 2006**

**Pengerusi: Profesor Madya Radzali Muse, PhD**

**Fakulti: Bioteknologi dan Sains Biomolekul**

Kajian mengenai potensi klon koko tempatan yang telah diisytiharkan oleh MARDI untuk menghasilkan embrio somatic telah dilakukan. Keluk sigmoid bagi pertumbuhan, kebolehidupan dan kandungan protein kalus dari batang, daun, kotiledon tidak matang dan kotiledon diperolehi. Berdasarkan keluk ini, sela tempoh untuk pengkulturan semula ialah diantara 20 hingga 24 hari selepas inkubasi. Sela tempoh ini jelas menunjukkan pertumbuhan kalus adalah tertinggi dan kandungan protein adalah tertinggi. Walau bagaimanapun, kebolehidupan kalus semakin berkurang disebabkan proses pencoklatan. Dengan itu, hanya kalus yang peroi dan berwarna putih sahaja dipilih untuk pengkulturan semula.

Pengaruh kalus embriogenik telah berjaya dilakukan untuk klon koko PBC 123 dan MHP 296 dengan menggunakan kalus yang terbentuk dari kotiledon tidak matang. Keputusan negatif bagi pengaruh kalus embriogenik ditunjukkan oleh semua klon koko dan eksplan yang dirawat dengan media MS. Keputusan yang positif didapati bagi

klon koko PBC 123 dan MHP 296 dengan menggunakan media DKW dan WPM sebagai media asas. Kombinasi 2 mg/L (w/v) 2,4-D dan 0.25 mg/L(w/v) Kin merupakan kombinasi auksin dan sitokinin terbaik bagi pengaruh kalus embriogenik. Kalus embryogenik nodular bagi klon PBC 123 dan MHP 296 mula kelihatan semasa pengkulturan semula yang ke dua. Kalus menjadi keperangan semasa dalam tempoh pengeraman terutama sekali pada akhir tempoh pengeraman bagi kedua dua klon tersebut. Penambahbaikan komposisi media dengan penambahan beberapa jenis kompaun kaya nitrogen dan asid amino seperti kasein hidrolisat, ekstrak malta dan prolin didapati tidak membantu di dalam pengaruh kalus embriogenik. Walaubagaimana pun penambahan kompaun ini dapat menjadikan kalus lebih peroi.

Bagi pengaruh embryo secara langsung, 15 jenis klon telah diuji tetapi hanya 4 jenis sahaja yang menunjukkan hasil positif. Klon-klon MJS 47, GS 29, PBC 123 dan PBC 159 berupaya menghasilkan kalus embriogenik daripada eksplan staminod dengan menggunakan protokol yang telah dibangunkan oleh Li et al., (1998). Kalus embriogenik nodular bagi klon – klon MJS 47, GS 29, PBC 123 dan PBC 159 kemudiannya dipindahkan ke media pertumbuhan sekunder untuk perkembangan kalus selanjutnya.

Pematangan kalus embriogenik dilakukan dengan menggunakan media yang dibangunkan oleh Li et al., (1998). Kalus embriogenik nodular berwarna coklat pucat semasa awal pematangan dan bertukar ke coklat gelap selepas tiga minggu tempoh pengeraman. Kesemua kalus embriogenik gagal melepasi tahap pembentukan embrio soma berbentuk hati yang merupakan tahap ke dua proses pengaruh embrio somatik.

Proses pematangan kalus embriogenik gagal dimana ianya menjadi keperangan dan hilang kebolehidupan pada akhir tempoh pengeraman. Kesemua klon menunjukkan tindakbalas yang sama terhadap media pematangan.

Kandungan keseluruhan fenol bagi kalus embriogenik bagi klon PBC 123 selepas 16 hari pengeraman adalah  $259.94 \pm 15.53 \mu\text{g/g}$  berat basah dimana ianya lebih rendah berbanding kalus tidak embriogenik ( $451.19 \pm 5.42 \mu\text{g/g}$  berat basah). Kenaikan atau pengumpulan kandungan polifenol ini didapati lebih kurang sama dengan klon MHP 296 yang menggunakan kalus dari kotiledon tidak matang sebagai eksplan terutamanya pada akhir tempoh pengeraman. Disamping itu, kalus embriogenik yang teraruh daripada eksplan kotiledon tidak matang menunjukkan kandungan polifenol yang tinggi berbanding kalus embriogenik dari eksplan staminod. Proses pencoklatan kalus yang tidak embriogenik adalah lebih teruk berbanding kalus embriogenik. Aktiviti enzim peroksida di dapati bertambah secara perlahan-lahan apabila kalus menjadi embriogenik. Aktiviti enzim peroksida bagi kalus embriogenik klon GS 29 ialah  $164.27 \pm 9.42$  unit/mg protein berbanding  $63.31 \pm 9.24$  unit/mg protein bagi kalus tidak embriogenik pada 16 hari selepas pengeraman. Walaupun corak peningkatan aktiviti enzim peroksida yang seakan sama bagi kalus embriogenik dan kalus yang tidak embriogenik, didapati aktiviti enzim lebih tinggi bagi kalus embriogenik berbanding kalus tidak embriogenik.

## **ACKNOWLEDGEMENTS**

First and foremost, I would like to express my deepest appreciation to my supervisors, Associate Professor Dr. Radzali Muse, Prof. Dr. Marziah Mahmood and Dr. Misri Kusnan for their guidance, knowledge, encouragement and critical review towards the completion of my thesis. I would like to thank Dr. Mohd Yunos Shukor (UPM), Dr Othman Omar (Deputy Director of RIC), En. Mansor Puteh (Deputy Director of RIC), Dr Md Jusoh Mamat (Director of RIC) and Dr Habibbudin Hashim (MARDI) and the manager of FELCRA Bukit Sedanan for their encouragement, support and suggestions through out the completion of this thesis. I also would like to thank En. Ibrahim, En. Onn, En Khalid, En Zamros, En. Hussin and Pn. Siti Rodiah (Biochemistry Department), En Ismail Sharif and Pn. Rashidah Jasman (MARDI) for their technical assistance. Not forgetting the Government of Malaysia and Universiti Putra Malaysia, I am truly grateful for the financial support from which I was able to complete my research successfully. Last but not least, I would like to thank my family and friends for their love, support and encouragement throughout my study.



I certify that an Examination Committee has met on 3<sup>rd</sup> March 2006 to conduct the final examination of Asfaliza Ramli on her Master of Science thesis entitled “Embryogenic Callus Formation in Local Cocoa (*Theobroma cacao* L.) Clones” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Norhani Abdullah, PhD**

Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Chairman)

**Maheran Abd. Aziz, PhD**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

**Nor’Aini Mohd Fadzillah, PhD**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Douglas Bruce Furtek, PhD**

Researcher  
Malaysia Cocoa Board  
(External Examiner)

---

**HASANAH MOHD. GHAZALI, PhD**

Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as partial fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**Radzali Muse, PhD**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences  
(Chairman)

**Marziah Mahmood, PhD**

Professor

Faculty of Biotechnology and Biomolecular Sciences  
(Member)

**Misri Kusnan, PhD**

Lecturer

Faculty of Science  
(Member)

---

**AINI IDERIS, PhD**

Professor/Dean

School of Graduate Studies  
Universiti Putra Malaysia

Date:

## **DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any degree at UPM or other institutions

---

**ASFALIZA RAMLI**

Date:

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	<b>ii</b>
<b>ABSTRAK</b>	<b>v</b>
<b>ACKNOWLEDGEMENTS</b>	<b>viii</b>
<b>APPROVAL</b>	<b>ix</b>
<b>DECLARATION</b>	<b>xi</b>
<b>LIST OF TABLES</b>	<b>xv</b>
<b>LIST OF FIGURES</b>	<b>xvii</b>
<b>LIST OF ABBREVIATIONS</b>	<b>xx</b>
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>6</b>
2.1 <i>Theobroma cacao</i> Tissue Culture Studies	6
2.2 Somatic Embryogenesis	8
2.3 Indirect Somatic Embryogenesis	10
2.4 Direct Somatic Embryogenesis	12
2.5 Induction of Somatic Embryogenesis in <i>T. cacao</i>	14
2.5.1 Induction via Zygotic embryos	14
2.5.2 Induction via Leaves and Nucellar Tissue	16
2.5.3 Induction via Cotyledon and Flower Buds	17
2.6 Factors Influencing Embryogenic Callus Induction	20
2.6.1 Culture Medium	21
2.6.2 Plant Growth Regulators	21
2.6.3 Genotype	24
2.6.4 Casein Hydrolysate and Malt Extract	25
2.7 Biochemicals Parameters in Embryogenic and Non Embryogenic Callus	26
2.7.1 Total Soluble Polyphenol Content	26
2.7.2 Peroxidase Activity	27
<b>3 MATERIALS AND METHODS</b>	<b>29</b>
3.1 Indirect Somatic Embryogenesis	29
3.1.1 Initiation of Callus from Aseptically Seedling of <i>T.cacao</i>	30
3.1.1.1 In vitro Seedling Initiation Medium	30
3.1.1.2 Pods Surface Sterilization Procedure	31
3.1.1.3 Seedling Initiation	31
3.1.1.4 Callus Initiation Medium	31
3.1.1.5 Callus Initiation and Maintenance	32
3.1.2 The Callus Growth, Viability and Total Protein Determination of Stem, Leaf, Cotyledon and Immatured Cotyledon Derived Callus	33
3.1.2.1 Callus Growth Determination	33
3.1.2.2 Callus Viability Determination	33

3.1.2.3	Callus Total Protein Determination	34
3.1.3	Initiation of Somatic Embryogenesis from Stem, Leaf, Cotyledon and Immatured Cotyledon Derived Callus	35
3.1.3.1	Medium for Embryogenic Callus Induction	35
3.1.3.2	Induction of Embryogenic Callus	37
3.1.3.3	Manipulation of Medium for Embryogenic Callus Induction	37
3.1.3.4	Medium for Embryogenic Callus Maturation	38
3.1.4	Embryogenic Callus Maturation	39
3.2	Direct Somatic Embryogenesis	40
3.2.1	Medium for Direct Embryogenic Initiation	41
3.2.1.1	Primary Callus Growth Medium	41
3.2.1.2	Secondary Callus Growth Medium	41
3.2.1.3	Embryogenic Callus Maturation Medium	42
3.2.2	Explants Sterilization Procedure	42
3.2.3	Direct Embryogenic Callus Induction	43
3.2.4	Embryogenic Callus Maturation	45
3.3	Biochemical Parameters in Embryogenic and Non Embryogenic Callus	46
3.3.1	Total Soluble Polyphenol Content	46
3.3.2	Specific Peroxidase Activity	47
3.4	Statistical Analysis	48
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	<b>49</b>
4.1	Indirect Somatic Embryogenesis	49
4.1.1	Initiation of Callus from Various Explants of <i>T.cacao</i>	49
4.1.2	The Callus Growth, Viability and Total Protein of Stem, Leaf, Cotyledon and Immatured Cotyledon Derived Callus	57
4.1.2.1	Callus Growth	57
4.1.2.2	Callus Viability	61
4.1.2.3	Callus Total Soluble Protein	65
4.1.3	Indirect Embryogenic Callus Induction	69
4.1.4	Medium Manipulation for Embryogenic Callus Induction	73
4.1.5	Maturation of Embryogenic Callus	81
4.2	Direct Somatic Embryogenesis	82
4.2.1	Initiation of Embryogenic Callus from Staminodes	82
4.2.2	Embryogenic Callus Maturation	90
4.3	Biochemical Parameters in Embryogenic and Non Embryogenic Callus	93
4.3.1	Total Soluble Polyphenol	93
4.3.2	Specific Peroxidase Activity	96
<b>5</b>	<b>CONCLUSIONS</b>	<b>102</b>
	<b>REFERENCES</b>	<b>105</b>

<b>APPENDICES</b>	117
<b>BIODATA OF THE AUTHOR</b>	121