

Fruit consumption and the risk of bladder cancer

Jochems, Sylvia; Reulen, Raoul; Van Osch, Frits; Witlox, Willem J.A; Goossens, Maria E; Brinkman, Maree; Giles, Graham G; Milne, Roger L.; van den Brandt, Piet A. ; White, Emily; Weiderpass, Elisabete; Huybrechts, Inge ; Hémon, Bertrand ; Agudo, Antonio ; Bueno-de-Mesquita, Bas; Cheng, KK; van Schooten, Frederik J; Bryan, Rik; Wesselius, Anke; Zeegers, Maurice

DOI:
[10.1002/ijc.33008](https://doi.org/10.1002/ijc.33008)

License:
Other (please specify with Rights Statement)

Document Version
Peer reviewed version

Citation for published version (Harvard):
Jochems, S, Reulen, R, Van Osch, F, Witlox, WJA, Goossens, ME, Brinkman, M, Giles, GG, Milne, RL, van den Brandt, PA, White, E, Weiderpass, E, Huybrechts, I, Hémon, B, Agudo, A, Bueno-de-Mesquita, B, Cheng, KK, van Schooten, FJ, Bryan, R, Wesselius, A & Zeegers, M 2020, 'Fruit consumption and the risk of bladder cancer: a pooled analysis by the BLadder cancer Epidemiology and Nutritional Determinants study', *International Journal of Cancer*. <https://doi.org/10.1002/ijc.33008>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Jochems, SHJ, Reulen, RC, van Osch, FHM, et al. Fruit consumption and the risk of bladder cancer: A pooled analysis by the Bladder Cancer Epidemiology and Nutritional Determinants Study. *Int. J. Cancer*. 2020; 1– 10., which has been published in final form at: <https://doi.org/10.1002/ijc.33008>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **Fruit consumption and the risk of bladder cancer: a pooled analysis**
2 **by the BLadder cancer Epidemiology and Nutritional Determinants**
3 **study**

4
5 Sylvia H.J. Jochems^{1,2}, Raoul C. Reulen³, Frits H.M. van Osch^{1,2}, Willem J.A.
6 Witlox^{1,4}, Maria E. Goossens⁵, Maree Brinkman^{1,6,7}, Graham G. Giles^{7,8,9}, Roger L.
7 Milne^{7,8,9}, Piet A. van den Brandt¹⁰, Emily White¹¹, Elisabete Weiderpass¹², Inge
8 Huybrechts¹², Bertrand Hémon¹², Antonio Agudo¹³, Bas Bueno-de-Mesquita^{14,15,16,17},
9 K.K. Cheng³, Frederik J. van Schooten¹⁸, Richard T. Bryan², Anke Wesselius¹,
10 Maurice P. Zeegers^{1,2,19}

- 11
12 1. Department of Complex Genetics and Epidemiology, School of Nutrition and
13 Translational Research in Metabolism, Maastricht University, Maastricht, the
14 Netherlands
15 2. Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham,
16 United Kingdom
17 3. Institute of Applied Health Research, University of Birmingham, Birmingham, United
18 Kingdom
19 4. Maastricht University Medical Centre+, Department of Clinical Epidemiology and
20 Medical Technology Assessment, Maastricht, The Netherlands
21 5. Cancer Centre of Sciensano, OD Public Health and Surveillance, Belgium
22 6. Department of Clinical Studies and Nutritional Epidemiology, Nutrition Biomed
23 Research Institute, Melbourne, Australia
24 7. Cancer Epidemiology Division, Cancer Council Victoria, Melbourne, VIC, Australia
25 8. Centre for Epidemiology and Biostatistics, School of Population and Global Health,
26 The University of Melbourne, Melbourne, VIC, Australia
27 9. Precision Medicine, School of Clinical Sciences at Monash Health, Monash University,
28 Clayton, Victoria Australia
29 10. Department of Epidemiology, Schools for Oncology and Developmental Biology and
30 Public Health and Primary Care, Maastricht University Medical Centre, Maastricht, the
31 Netherlands
32 11. Fred Hutchinson Cancer Research Center, Seattle, Washington, United States
33 12. International Agency for Research on Cancer (IARC), World Health Organization,
34 Lyon, France
35 13. Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Institut Català
36 d'Oncologia, L'Hospitalet de Llobregat, Spain
37 14. Former senior scientist, Dept. for Determinants of Chronic Diseases (DCD), National
38 Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands
39 15. Former associate professor, Department of Gastroenterology and Hepatology,
40 University Medical Centre, Utrecht, The Netherlands
41 16. Former visiting professor, Dept. of Epidemiology and Biostatistics, The School of
42 Public Health, Imperial College London, St Mary's Campus, Norfolk Place, London,
43 United Kingdom
44 17. Former Academic Icon / visiting professor, Dept. of Social & Preventive Medicine,
45 Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia
46 18. Department of Pharmacology and Toxicology, NUTRIM School for Nutrition and
47 Translational Research in Metabolism, Maastricht University, Maastricht, the
48 Netherlands
49 19. CAPHRI School for Public Health and Primary Care, University of Maastricht,
50 Maastricht, The Netherlands
51

52

53 **Abbreviations**

54 95% confidence interval = 95% CI

55 BLadder Cancer Epidemiology and Nutritional Determinants study = BLEND

56 Carcinoma In Situ = CIS

57 European Prospective Investigation into Cancer and Nutrition study = EPIC

58 Food Frequency Questionnaire = FFQ

59 Hazard Ratio = HR

60 Melbourne Collaborative Cohort Study = MCCS

61 Muscle invasive bladder cancer = MIBC

62 Netherlands Cohort Study = NLCS

63 Non-muscle invasive bladder cancer = NMIBC

64 VITamins And Lifestyle cohort study = VITAL

65 World Cancer Research Fund = WCRF

66

67

68 **Novelty and impact statement**

69 Previous studies often lacked adequate numbers with bladder cancer to detect

70 associations between fruit consumption and bladder cancer risk, especially for

71 specific types of fruit and for women. In this large prospective study, we pooled data

72 from 13 cohort studies and found that increasing total fruit consumption may reduce

73 the risk of bladder cancer in women.

74

75 Abstract

76 While the association between fruit consumption and bladder cancer risk has been
77 extensively reported, studies have had inadequate statistical power to investigate
78 associations between types of fruit and bladder cancer risk satisfactorily.

79 Fruit consumption in relation to bladder cancer risk was investigated by pooling
80 individual data from 13 cohort studies. Cox regression models with attained age as
81 time scale were used to estimate hazard ratios (HRs) for intakes of total fruit and
82 each of citrus fruits, soft fruits, stone fruits, tropical fruits, pome fruits, and fruit
83 products. Analyses were stratified by sex, smoking status, and bladder cancer
84 subtype.

85 During on average 11.2 years of follow-up, 2836 individuals developed incident
86 bladder cancer. Increasing fruit consumption (by 100 gram/day) was inversely
87 associated with the risk of bladder cancer in women (HR=0.92; 95% CI 0.85-0.99).
88 Although in women the association with fruit consumption was most evident for
89 higher-risk non-muscle invasive bladder cancer (NMIBC) (HR=0.72; 95% CI 0.56-
90 0.92), the test for heterogeneity by bladder cancer subtype was non-significant (p-
91 heterogeneity=0.14). Increasing fruit consumption (by 100 gram/day) was not
92 associated with bladder cancer risk in men (HR=0.99; 95% CI 0.94-1.03), never
93 smokers (HR=0.96; 95% CI 0.88-1.05), former smokers (HR=0.98; 95% CI 0.92-
94 1.05), or current smokers (HR=0.95; 95% CI 0.89-1.01). The consumption of any
95 type of fruit was not found to be associated with bladder cancer risk (p-values>0.05).
96 This study supports no evidence that the consumption of specific types of fruit
97 reduces the risk of bladder cancer. However, increasing fruit consumption may
98 reduce bladder cancer risk in women.

99

100 **Introduction**

101 Bladder cancer is the ninth most common cancer worldwide, with almost 550,000
102 newly diagnosed cases in 2018 (1). Although cigarette smoking is the primary risk
103 factor for bladder cancer, the sex-based difference in bladder cancer incidence is
104 independent of differences in smoking status (2). Dietary factors may contribute to
105 bladder cancer risk considering that many dietary compounds are excreted in urine
106 (3). Fruits contain high levels of phytochemicals, minerals, and antioxidant nutrients,
107 that may hold anti-carcinogenic properties (4). According to a panel of experts of the
108 World Cancer Research Fund (WCRF) Continuous Update Project report 'Diet,
109 nutrition, physical activity and bladder cancer', there is limited evidence from cohort
110 studies that greater consumption of fruits and vegetables may decrease bladder
111 cancer risk (4). Moreover, adherence to the Mediterranean diet, rich in fruits, may
112 decrease the risk of bladder cancer (5). Results from case-control studies have
113 mainly shown inverse associations with fruit consumption, especially for the intake of
114 citrus fruits (6–8). However, recall bias and selection bias may have influenced the
115 reporting of fruit intake in case-control studies and most studies often lacked an
116 adequate number of individuals to detect associations between fruit intake and
117 bladder cancer risk, especially for types of fruits and bladder cancer subtypes. By
118 pooling individual data from multiple cohort studies, the number of bladder cancer
119 cases can be substantially increased with the advantage that the association
120 between fruit intake and bladder cancer risk can be investigated with greater power
121 for different types of fruits and by sex, smoking status, and bladder cancer subtype.
122 In addition, fruit intake categories and covariates can be standardized across studies
123 (unlike in systematic reviews or meta-analyses). The aim of this large-scale study
124 was to build on previous results of the European Prospective Investigation into
125 Cancer and Nutrition (EPIC) studies (9,10) and investigates the association between
126 fruit consumption and bladder cancer risk by pooling data for 535,713 individuals in
127 13 cohort studies included in the BLadder Cancer Epidemiology and Nutritional
128 Determinants (BLEND) study.

129

130 **Methods**

131 *Study population*

132 The BLEND study is an international consortium that pools individual participant data
133 from international cohort studies and case-control studies. Details of the

134 methodology of the BLEND study have been described elsewhere (11). Briefly, a
135 total of 13 cohort studies had sufficient data to be eligible for inclusion in this study
136 (i.e. method of dietary assessment, geographical region, disease status). About 75%
137 of the study populations originated from centers in Europe including the EPIC studies
138 (12,13) and the Netherlands Cohort Study (NLCS) (14). Other populations originated
139 from Australia (Melbourne Collaborative Cohort Study (MCCS)) (15) and North
140 America (VITamins And Lifestyle cohort study (VITAL)) (16). All studies have been
141 ethically approved and all study participants provided informed consent.

142

143 *Bladder cancer ascertainment*

144 Each study ascertained incident bladder cancers with International Classification of
145 Diseases (ICD-O-three code C67) using population-based cancer registries, health
146 insurance records, cancer registries, or medical records. Linkages to mortality
147 registries were conducted during the follow-up period of each study. The term
148 bladder cancer is used for all urinary bladder neoplasms. Bladder cancers were
149 classified into non-muscle invasive bladder cancer (NMIBC) and muscle invasive
150 bladder cancer (MIBC). NMIBCs included noninvasive carcinomas confined to the
151 urothelium (stage Ta), carcinomas that invaded the lamina propria of the bladder wall
152 (stage T1), and high grade flat noninvasive carcinomas confined to the urothelium
153 (carcinoma in situ; CIS). MIBCs included carcinomas that invaded into the detrusor
154 muscle (stage T2), carcinomas that invaded into the peri vesical tissue (stage T3),
155 and carcinomas that invaded adjacent tissues and organs (most often the prostate or
156 uterus) (stage T4). With bladder cancer representing a heterogeneous group of
157 tumours, that possibly develop through different but interrelated pathways (17) and
158 could have implications for treatments and outcomes, NMIBCs were further divided
159 into “lower” risk (stage Ta with a low grade (grade 1 or grade 2)) and “higher” risk
160 (stage Ta with grade 3, stage T1, and CIS). Whilst lower-risk NMIBC often occurs
161 from papillary tumours, higher-risk NMIBC and MIBC are more likely to develop from
162 non-papillary tumours (18).

163

164 *Dietary assessment*

165 For each study, participants were asked to report on their usual fruit consumption
166 during the preceding year before study enrolment. All the studies assessed usual
167 dietary intake with a validated food frequency questionnaire (FFQ). To harmonise

168 data collected from the study specific FFQs and to consider the varying portions
169 sizes between different populations, frequency intakes were converted to grams
170 using the portion sizes described in the FFQ of each study. Where applicable, fruit
171 intakes were converted from weekly, monthly, or yearly intakes, to daily intakes. The
172 consumption of fruits in grams per day was then standardised across studies by
173 making use of the Eurocode 2 food coding system (19). Total fruit intake was
174 computed as the sum of grams of all fruit items or fruit groups (excluding fruit juices)
175 provided by each study. The following types of fruits were defined: citrus fruits, pome
176 fruits, soft fruits, stone fruits, tropical fruits, and fruit products (Table 1).

177

178 *Statistical analysis*

179 Person-years of follow-up for each participant were calculated from date of study
180 enrolment until the date of a first bladder cancer diagnosis, death, emigration, last
181 known contact, or end of study follow-up, whichever came first. For the NLCS, a
182 nested case-cohort approach was applied, in which the number of person-years at
183 risk was estimated based on a subcohort that was randomly sampled after baseline
184 (14). Total fruit consumption was analysed both as a continuous variable (expressing
185 results per 100 grams per day in usual total fruit consumption), and a categorical
186 variable. For the categorical variable, total fruit consumption was divided into four
187 intake categories: <100 grams of fruit per day (less than approximately one piece of
188 fruit), 100-200 grams of fruit per day (approximately one to two pieces of fruit), 200-
189 300 grams of fruit per day (approximately two to three pieces of fruit), and >300
190 grams of fruit per day (more than approximately three pieces of fruit), using the
191 lowest intake category as a reference. Fruit types (citrus fruits, soft fruits, stone fruits,
192 tropical fruits, pome fruits, and fruit products) were each analysed as a continuous
193 variable per 25 grams of fruits per day increase and were modelled into quartiles,
194 using the lowest quartile as a reference. Cox proportional hazard models with
195 attained age as time scale were used to calculate hazard ratios (HR) and 95%
196 confidence intervals (95% CI) for bladder cancer. The assumption of proportional
197 hazards was examined for the relationship of scaled Schoenfeld residuals with time
198 and appeared to be violated when considering all participants together (20). Based
199 on *a priori* reasons and the violation of the proportional hazard assumption for all
200 participants, analyses were performed for sex, smoking status (never smokers,
201 former smokers, current smokers), and bladder cancer subtype (NMIBC and MIBC,

202 and further classification into lower-risk NMIBC and higher-risk NMIBC); the
203 assumption of proportional hazards was now found to be satisfied in all models.
204 Heterogeneity was calculated by the duplication method for Cox regression as
205 described by Lunn *et al.* (21), using a likelihood ratio test to compare the model with
206 and without interaction terms between total fruit consumption and sex, smoking
207 status, and bladder cancer subtype. Within the regression models, all analyses were
208 stratified by cohort, sex, and age at study enrolment. Adjustment was made for the
209 potential confounders smoking status (current smoker/former smoker/never smoker),
210 pack-years of cigarette smoking (continuous in years), ethnicity
211 (Asian/Black/Caucasian), total vegetable consumption (continuous in gram/day),
212 alcohol intake (continuous in gram/day), and total energy intake (continuous in
213 kcal/day). A sensitivity analysis was performed on pre-defined sex-specific energy
214 intake cut offs (800-4000 kcal/day for women and 1500-6000 kcal/day for men). All
215 statistical analyses were performed using Stata software version 14 and a two-sided
216 p-value of <0.05 was considered statistically significant.

217

218 *Data Availability*

219 The data that support the findings of this study are not publicly available, but data will
220 be made available upon reasonable request.

221

222

223 **Results**

224 Baseline characteristics of the study samples included are presented in Table 2. Of
225 597,231 potentially eligible participants, 61,327 individuals were excluded from the
226 statistical analyses for having missing data on total fruit consumption (n=28,929),
227 total vegetable consumption (n=173), ethnicity (n=472), pack-years of smoking
228 (n=27,476), or for missing and extreme values (<800 kcal/day and >6000 kcal/day)
229 of total energy intake (n=46,906). In addition, individuals with incident bladder
230 cancers diagnosed within the first two years of study follow-up were excluded
231 (n=191) (Figure 1). During an average of 11.2 years of follow-up, 2836 of the
232 remaining 535,713 participants were diagnosed with an incident bladder cancer. A
233 total of 1135 cases were classified as NMIBC and 706 as MIBC; 995 bladder
234 cancers could not be classified due to missing data on tumour characteristics.

235

236 *Total fruit*

237 In men, increasing fruit consumption by 100 grams per day was not associated with
238 overall bladder cancer risk (HR=0.99; 95% CI 0.94-1.03) (Table 3), or with any
239 bladder cancer subtype (p-heterogeneity=0.33) (Table 4). The sensitivity analysis on
240 sex-specific cut offs for total energy intake in men per 100 grams of fruit per day
241 increase showed a comparable result (HR=0.99; 95% CI 0.94-1.04). Compared with
242 the lowest category of fruit intake (<100 grams of fruit per day), the highest total fruit
243 intake category (>300 grams of fruit per day) was associated with a lower risk of
244 overall bladder cancer in women (HR=0.75; 95% CI 0.59-0.97) (Table 3). In the
245 continuous analysis for women, increasing total fruit consumption by 100 grams per
246 day was inversely associated with the risk of overall bladder cancer (HR=0.92; 95%
247 CI 0.85-0.99) (Table 3). A similar result for increasing total fruit consumption and
248 bladder cancer risk in women was obtained from the sensitivity analysis when sex-
249 specific cut offs for total energy intake were used (HR=0.92; 95% 0.85-0.99).
250 Although in women the association was stronger for higher-risk NMIBC (HR=0.72;
251 95% CI 0.56-0.92) than for all NMIBCs combined (HR=0.79; 95% CI 0.67-0.94), the
252 test for heterogeneity by bladder cancer subtype did not reach significance (p-
253 heterogeneity=0.14) (Table 4). In the subgroup analysis on smoking status, the
254 consumption of fruit was not associated with the risk of bladder cancer in never
255 smokers (HR=0.96; 95% CI 0.88-1.05), former smokers (HR=0.98; 95% CI 0.92-
256 1.05), current smokers (HR=0.95; 95% CI 0.89-1.01) (Table 3), or ever smokers
257 (current and former smokers combined) (HR=0.96; 95% CI 0.92-1.01).

258

259 *Subtypes of fruit*

260 In women, no associations were found between increasing consumption by 25
261 grams per day of citrus fruits (HR=0.97; 95% CI 0.92-1.03), soft fruits (HR=0.95;
262 95% CI 0.84-1.09), stone fruits (HR=0.94; 95% CI 0.85-1.03), pome fruits (HR=0.95;
263 95% CI 0.87-1.03), or fruit products (HR=1.00; 95% CI 0.76-1.32), and overall
264 bladder cancer risk (Table 3). Although for tropical fruit intake an association was
265 found with the risk of overall bladder cancer in women in the categorical analysis
266 (highest quintile vs. lowest quintile HR=0.78; 95% CI 0.62-0.99, p-trend=0.05), the
267 continuous analysis for increasing tropical fruit consumption by 25 grams per day
268 showed no association (HR=0.97; 95% CI 0.91-1.04). In the analysis for men and on

269 smoking status, no associations were found between any specific type of fruit and
270 the risk of overall bladder cancer ($p>0.05$) (Table 3).

271

272 **Discussion**

273 In this analysis of pooled data from 13 prospective cohort studies, comprising 2836
274 individuals with incident bladder cancer, an association was found between
275 increasing total fruit consumption and a decreased risk of bladder cancer in women.
276 No associations were found between fruit consumption and the risk of bladder
277 cancer for men, current smokers, former smokers, or never smokers (9,10). With
278 bladder cancer being a heterogeneous disease, attention has increasingly focused
279 on investigating subtypes of bladder cancer. While in the EPIC study, tumours were
280 defined as non-aggressive urothelial cell carcinomas or aggressive urothelial cell
281 carcinomas (10), the classification of bladder tumours for this BLEND study included
282 lower-risk NMIBC, higher-risk NMIBC and MIBC. However, there were no significant
283 differences between the risk associations for the bladder cancer subtypes in relation
284 to the consumption of fruit using the duplication method for Cox regression as
285 described by Lunn *et al.* (21). The addition of incident bladder cancers from three
286 additional large cohorts (NLCS, MCCS, and VITAL) could explain the novel finding
287 for the inverse association between total fruit consumption and bladder cancer risk
288 for women. Although most prospective studies on bladder cancer risk found no
289 associations with fruit consumption (22–25), the findings for women are in partial
290 agreement with results of the Multiethnic Cohort study (26). Park *et al.* (26) found
291 that only for women, total fruit and citrus fruit consumptions were inversely
292 associated with the risk of bladder cancer (HR=0.54; 95% CI 0.34-0.85 and
293 HR=0.56; 95% CI 0.34-0.90, respectively). Interestingly, the authors showed that
294 there was only a significant association with fruit consumption for women when
295 considering invasive bladder cancer as an endpoint, not non-invasive bladder cancer
296 (26). Results from the Nurses' Health study on lung cancer (a smoking-related
297 cancer as bladder cancer) also showed that especially women with greater intakes of
298 fruit had a reduced risk of cancer (27). It cannot be excluded that the inverse
299 association found for women but not men may be partially explained by differences
300 in hormonal factors (e.g. estrogen) and urination habits between men and women, or
301 by residual confounding by smoking habits, though the inverse association for
302 women in the Multiethnic Cohort study was found after rigorous adjustment for

303 cigarette smoking and reproductive factors (26). Although statistical power was more
304 limited for women compared with men (683 incident bladder cancers in women and
305 2153 incident bladder cancers in men), especially in the categorical analysis of fruit
306 intake, the number of incident bladder cancers in the continuous analyses for
307 increasing total fruit consumption by 100 grams per day in women had adequate
308 power. All types of fruit showed non-significant associations with the risk of bladder
309 cancer (all $p > 0.05$) and therefore the inverse association between fruit consumption
310 and bladder cancer risk in women cannot be attributed to increased consumption of
311 a specific type of fruit.

312 This study has several strengths, including the large sample size providing statistical
313 power to examine different types of fruits, the possibility to classify risks by sex,
314 smoking status, and bladder cancer subtype, and the inclusion of studies from 12
315 different countries. Although the use of a calibration method might have reduced
316 between-country heterogeneity in dietary intake, results of both the EPIC study (9)
317 and the Multiethnic Cohort study (26) on fruit consumption and bladder cancer risk
318 indicated that after applying a calibration method (28), there were no substantial
319 differences between their observed findings and their calibrated estimates. Although
320 by making use of the Eurocode 2 Food Coding System (19) the potential for
321 misclassification for the types of fruit is limited, measurement error in the dietary
322 assessment by limitations of the FFQs, including over- and under-reporting of usual
323 fruit consumption, and the inability to investigate dietary changes over time with only
324 one single measurement of fruit at time of study entry, cannot be excluded. However,
325 if changes in dietary intake were made by individuals during follow-up, it would still
326 be questionable whether these changes would have influenced bladder cancer risk
327 in this relatively short period of time. Other limitations of this study were the limited
328 information on covariates that may be associated with the risk of bladder cancer (and
329 that are highly correlated with fruit consumption), such as body mass index, physical
330 activity, and socioeconomic status. However, it has been indicated that these factors
331 may probably account for only a small percentage of bladder cancer cases overall
332 (29,30).

333 In conclusion, there was no evidence that the consumption of specific types of fruit
334 reduces the risk of bladder cancer. However, increasing consumption of the total
335 amount of fruits may reduce bladder cancer risk in women.

336

337

338 Funding

339 This study was partly funded by the World Cancer Research Fund.

340

341 Conflict of interest

342 On behalf of all authors, the corresponding author states that there is no conflict of
343 interest.

344

345 DISCLAIMER

346 Where authors are identified as personnel of the International Agency for Research
347 on Cancer / World Health Organization, the authors alone are responsible for the
348 views expressed in this article and they do not necessarily represent the decisions,
349 policy or views of the International Agency for Research on Cancer / World Health
350 Organization.

351

352 **References**

- 353 1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global
354 cancer statistics 2018: GLOBOCAN estimates of incidence and mortality
355 worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018
356 Nov;68(6):394–424.
- 357 2. Dobruch J, Daneshmand S, Fisch M, Lotan Y, Noon AP, Resnick MJ, et al.
358 Gender and Bladder Cancer: A Collaborative Review of Etiology, Biology, and
359 Outcomes. *Eur Urol [Internet]*. 2016 Feb [cited 2018 Sep 8];69(2):300–10.
360 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26346676>
- 361 3. Piyathilake C. Dietary factors associated with bladder cancer. *Investig Clin*
362 *Urol [Internet]*. 2016 Jun [cited 2017 Feb 22];57(Suppl 1):S14. Available from:
363 <http://www.ncbi.nlm.nih.gov/pubmed/27326403>
- 364 4. World Cancer Research Fund International/American Institute for Cancer
365 Research. Continuous Update Project Report: Diet, Nutrition, Physical Activity
366 and Bladder Cancer. 2018 [Internet]. Available from: [wcrf.org/bladder-cancer-](http://wcrf.org/bladder-cancer-2018)
367 [2018](http://wcrf.org/bladder-cancer-2018)
- 368 5. Witlox WJA, van Osch FHM, Brinkman M, Jochems S, Goossens ME,
369 Weiderpass E, et al. An inverse association between the Mediterranean diet
370 and bladder cancer risk: a pooled analysis of 13 cohort studies. *Eur J Nutr*.
371 2019 Feb;
- 372 6. Al-Zalabani AH, Stewart KFJ, Wesselius A, Schols AMWJ, Zeegers MP.
373 Modifiable risk factors for the prevention of bladder cancer: a systematic
374 review of meta-analyses. *Eur J Epidemiol [Internet]*. 2016 Sep 21 [cited 2017
375 Feb 21];31(9):811–51. Available from:
376 <http://www.ncbi.nlm.nih.gov/pubmed/27000312>
- 377 7. Yao B, Yan Y, Ye X, Fang H, Xu H, Liu Y, et al. Intake of fruit and vegetables
378 and risk of bladder cancer: a dose–response meta-analysis of observational
379 studies. *Cancer Causes Control [Internet]*. 2014 Dec 24 [cited 2017 Feb
380 22];25(12):1645–58. Available from: [http://link.springer.com/10.1007/s10552-](http://link.springer.com/10.1007/s10552-014-0469-0)
381 [014-0469-0](http://link.springer.com/10.1007/s10552-014-0469-0)
- 382 8. Liu H, Wang X-C, Hu G-H, Guo Z-F, Lai P, Xu L, et al. Fruit and vegetable
383 consumption and risk of bladder cancer. *Eur J Cancer Prev [Internet]*. 2015
384 Nov [cited 2017 Feb 22];24(6):508–16. Available from:
385 <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=0>

- 386 0008469-201511000-00008
- 387 9. Büchner FL, Bueno-de-Mesquita HB, Ros MM, Kampman E, Egevad L,
388 Overvad K, et al. Consumption of vegetables and fruit and the risk of bladder
389 cancer in the European Prospective Investigation into Cancer and Nutrition. *Int*
390 *J cancer* [Internet]. 2009 Dec 1 [cited 2018 Aug 2];125(11):2643–51. Available
391 from: <http://doi.wiley.com/10.1002/ijc.24582>
- 392 10. Ros MM, Bueno-de-Mesquita HB, Kampman E, Büchner FL, Aben KKH,
393 Egevad L, et al. Fruit and vegetable consumption and risk of aggressive and
394 non-aggressive urothelial cell carcinomas in the European Prospective
395 Investigation into Cancer and Nutrition. *Eur J Cancer* [Internet]. 2012 Nov
396 [cited 2018 Sep 3];48(17):3267–77. Available from:
397 <http://linkinghub.elsevier.com/retrieve/pii/S0959804912004753>
- 398 11. Goossens ME, Isa F, Brinkman M, Mak D, Reulen R, Wesselius A, et al.
399 International pooled study on diet and bladder cancer: the bladder cancer,
400 epidemiology and nutritional determinants (BLEND) study: design and baseline
401 characteristics. *Arch Public Heal* [Internet]. 2016;74(1):30. Available from:
402 <http://archpublichealth.biomedcentral.com/articles/10.1186/s13690-016-0140-1>
- 403 12. Riboli E, Kaaks R. The EPIC Project: rationale and study design. *European*
404 *Prospective Investigation into Cancer and Nutrition. Int J Epidemiol* [Internet].
405 1997 [cited 2018 Jul 4];26 Suppl 1:S6-14. Available from:
406 <http://www.ncbi.nlm.nih.gov/pubmed/9126529>
- 407 13. Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, et al. *European*
408 *Prospective Investigation into Cancer and Nutrition (EPIC): study populations*
409 *and data collection. Public Health Nutr.* 2002 Dec;5(6B):1113–24.
- 410 14. van den Brandt PA, Goldbohm RA, Van 't Veer P, Volovics A, Hermus RJJ,
411 Sturmans F. A large-scale prospective cohort study on diet and cancer in the
412 Netherlands. *J Clin Epidemiol* [Internet]. 1990 Jan [cited 2018 Jul
413 4];43(3):285–95. Available from:
414 <http://linkinghub.elsevier.com/retrieve/pii/089543569090009E>
- 415 15. Milne RL, Fletcher AS, MacInnis RJ, Hodge AM, Hopkins AH, Bassett JK, et
416 al. Cohort Profile: The Melbourne Collaborative Cohort Study (Health 2020).
417 *Int J Epidemiol* [Internet]. 2017 Dec 1 [cited 2018 Jul 4];46(6):1757-1757i.
418 Available from: <http://academic.oup.com/ije/article/46/6/1757/3882696>
- 419 16. White E, Patterson RE, Kristal AR, Thornquist M, King I, Shattuck AL, et al.

- 420 VITamins And Lifestyle cohort study: study design and characteristics of
421 supplement users. *Am J Epidemiol* [Internet]. 2004 Jan 1 [cited 2018 Jul
422 4];159(1):83–93. Available from:
423 <http://www.ncbi.nlm.nih.gov/pubmed/14693663>
- 424 17. Droller MJ. Biological considerations in the assessment of urothelial cancer: a
425 retrospective. *Urology* [Internet]. 2005 Nov [cited 2018 Sep 1];66(5 Suppl):66–
426 75. Available from:
427 <http://linkinghub.elsevier.com/retrieve/pii/S0090429505010174>
- 428 18. Dinney CPN, McConkey DJ, Millikan RE, Wu X, Bar-Eli M, Adam L, et al.
429 Focus on bladder cancer. *Cancer Cell* [Internet]. 2004 Aug [cited 2018 Aug
430 31];6(2):111–6. Available from:
431 <http://linkinghub.elsevier.com/retrieve/pii/S1535610804002120>
- 432 19. Kohlmeier L. The Eurocode 2 food coding system. *Eur J Clin Nutr* [Internet].
433 1992 Dec [cited 2018 Jul 4];46 Suppl 5:S25-34. Available from:
434 <http://www.ncbi.nlm.nih.gov/pubmed/1486871>
- 435 20. D. Schoenfeld. Partial Residuals for The Proportionnal Hazards Regression
436 Model. *Biometrika* [Internet]. 1982;69(1):239–41. Available from:
437 <http://www.jstor.org/stable/2335876>
- 438 21. Lunn M, McNeil D. Applying Cox regression to competing risks. *Biometrics*.
439 1995 Jun;51(2):524–32.
- 440 22. Holick CN, De Vivo I, Feskanich D, Giovannucci E, Stampfer M, Michaud DS.
441 Intake of Fruits and Vegetables, Carotenoids, Folate, and Vitamins A, C, E and
442 Risk of Bladder Cancer Among Women (United States). *Cancer Causes
443 Control* [Internet]. 2005 Dec [cited 2018 Jul 4];16(10):1135–45. Available from:
444 <http://www.ncbi.nlm.nih.gov/pubmed/16215863>
- 445 23. Michaud DS, Spiegelman D, Clinton SK, Rimm EB, Willett WC, Giovannucci
446 EL. Fruit and vegetable intake and incidence of bladder cancer in a male
447 prospective cohort. *J Natl Cancer Inst* [Internet]. 1999 Apr 7 [cited 2018 Jul
448 4];91(7):605–13. Available from:
449 <http://www.ncbi.nlm.nih.gov/pubmed/10203279>
- 450 24. Zeegers MP, Goldbohm RA, van den Brandt PA. Consumption of vegetables
451 and fruits and urothelial cancer incidence: a prospective study. *Cancer
452 Epidemiol Biomarkers Prev* [Internet]. 2001 Nov [cited 2018 Jul
453 4];10(11):1121–8. Available from:

- 454 <http://www.ncbi.nlm.nih.gov/pubmed/11700259>
- 455 25. Larsson SC, Andersson S-O, Johansson J-E, Wolk a. Fruit and Vegetable
456 Consumption and Risk of Bladder Cancer: A Prospective Cohort Study.
457 *Cancer Epidemiol Biomarkers Prev* [Internet]. 2008;17(9):2519–22. Available
458 from: <http://cebp.aacrjournals.org/cgi/doi/10.1158/1055-9965.EPI-08-0407>
- 459 26. Park S-Y, Ollberding NJ, Woolcott CG, Wilkens LR, Henderson BE, Kolonel
460 LN. Fruit and vegetable intakes are associated with lower risk of bladder
461 cancer among women in the Multiethnic Cohort Study. *J Nutr* [Internet]. 2013
462 Aug 1 [cited 2018 Jul 4];143(8):1283–92. Available from:
463 <https://academic.oup.com/jn/article/143/8/1283/4637671>
- 464 27. Feskanich D, Ziegler RG, Michaud DS, Giovannucci EL, Speizer FE, Willett
465 WC, et al. Prospective study of fruit and vegetable consumption and risk of
466 lung cancer among men and women. *J Natl Cancer Inst* [Internet]. 2000 Nov
467 15 [cited 2018 Jul 4];92(22):1812–23. Available from:
468 <http://www.ncbi.nlm.nih.gov/pubmed/11078758>
- 469 28. Ferrari P, Slimani N, Ciampi A, Trichopoulou A, Naska A, Lauria C, et al.
470 Evaluation of under- and overreporting of energy intake in the 24-hour diet
471 recalls in the European Prospective Investigation into Cancer and Nutrition
472 (EPIC). *Public Health Nutr*. 2002 Dec;5(6B):1329–45.
- 473 29. Burger M, Catto JWF, Dalbagni G, Grossman HB, Herr H, Karakiewicz P, et al.
474 Epidemiology and risk factors of urothelial bladder cancer. *Eur Urol* [Internet].
475 2013 Feb [cited 2018 Aug 30];63(2):234–41. Available from:
476 <http://linkinghub.elsevier.com/retrieve/pii/S0302283812008780>
- 477 30. Koenig C, Michaud D, Moore SC, Park Y, Hollenbeck A, Ballard-Barbash R,
478 et al. Body mass index, physical activity, and bladder cancer in a large
479 prospective study. *Cancer Epidemiol Biomarkers Prev*. 2008 May;17(5):1214–
480 21.
- 481

Table 1. Classification of types of fruit based on composition

Types of fruit	Composition
Citrus fruits	lemons, oranges, tangerines, grapefruits, pomelos, limes, kumquats
Soft fruits	strawberries, raspberries, white grapes, black grapes, loganberries, blackberries, dewberries, cloudberrries, gooseberries, black currants, red currants, white currants, cranberries, bilberries, cowberries, blueberries, elderberries, rowanberries, physalis, mulberries, bearberries, sea buckthorns
Stone fruits	apricots, peaches, nectarines, plums, damsons, mirabelles, greengages, sweet cherries, sour cherries, chickasaws, susinas, sloes, dates, lychees, persimmons, barbados cherries
Pome fruits	apples, pears, quinces, medlars, and loquats
Tropical fruits	bananas, pineapples, kiwi fruits, (water)melons, figs, mangos, pomegranates, passion fruits, cashew fruits, guavas, papayas, rose hips, sapodillas, carambolas, durians, jack fruit, chayotes, rambutans, tamarinds
Fruit products	dried mixed fruits, mixed peels, glace cherries, crystallized pineapple, apple sauce, cranberry sauce
Fruit mixtures	fruit salads, fruit cocktails

Table 2. Baseline characteristics of individuals from the 13 cohort studies included in the pooled analysis

	EPIC Denmark	EPIC France	EPIC Germany	EPIC Greece	EPIC Italy	EPIC The Netherlands	EPIC Norway	EPIC Spain	EPIC Sweden	EPIC United Kingdom	NLCS The Netherlands (case-cohort design)	MCCS Australia	VITAL USA	Total in BLEND study
	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))	No. ^a (%) / (mean (SD))
Total participants	56,005 (9)	64,866 (11)	49,457 (8)	25,268 (4)	45,204 (8)	37,102 (6)	33,856 (6)	40,782 (7)	49,328 (8)	75,035 (13)	5632 (1)	38,263 (6)	76,433 (13)	597,231 (100)
Men	26,764 (13)	0	21,551 (11)	10,438 (6)	14,084 (7)	9801 (5)	0	15,439 (8)	22,546 (11)	22,476 (11)	3052 (2)	15,798 (8)	36,453 (18)	198,402 (100)
Women	29,241 (7)	64,866 (16)	27,906 (7)	14,830 (4)	31,120 (8)	27,301 (7)	33,856 (8)	25,343 (6)	26,782 (7)	52,559 (13)	2580 (1)	22,465 (6)	39,980 (10)	398,829 (100)
All incident bladder cancers ^b	391 (11)	31 (<1)	207 (6)	50 (1)	187 (5)	107 (3)	24 (<1)	152 (4)	303 (9)	248 (7)	940 (27)	520 (15)	378 (11)	3538 (100)
Lower-risk NMIBC	87 (17)	17 (3)	79 (16)	-	46 (9)	71 (14)	-	21 (4)	-	0 (<1)	-	188 (37)	-	509 (100)
Higher-risk NMIBC	51 (8)	5 (<1)	35 (5)	-	58 (9)	22 (3)	-	29 (4)	-	1 (<1)	409 (61)	47 (7)	15 (2)	672 (100)
MIBC	44 (5)	5 (<1)	40 (4)	-	20 (2)	23 (2)	-	7 (<1)	-	6 (<1)	443 (47)	232 (24)	121 (13)	941 (100)
Mean age at study entry (yrs)	56.7 (4.4)	52.8 (6.6)	50.6 (8.6)	53.3 (12.6)	50.5 (7.9)	48.9 (12.0)	48.1 (4.3)	49.2 (8.0)	52.0 (10.9)	49.1 (14.4)	62.1 (4.2)	55.0 (8.7)	61.4 (7.5)	52.9 (10.2)
Never smoker	19,624 (7)	45,797 (15)	22,658 (7)	14,060 (4)	20,540 (7)	14,171 (6)	12,057 (4)	22,599 (8)	24,205 (8)	41,948 (14)	1848 (1)	22,057 (7)	35,818 (12)	297,324 (100)
Former smoker	17,070 (10)	13,121 (7)	16,386 (9)	4232 (2)	12,096 (7)	11,572 (7)	10,438 (6)	7207 (4)	13,410 (8)	23,924 (14)	2018 (1)	11,848 (7)	33,648 (18)	176,970 (100)
Current smoker	19,624 (16)	5948 (5)	10,413 (9)	6976 (6)	12,568 (10)	11,359 (9)	11,361 (9)	10,976 (9)	11,713 (10)	9163 (7)	1766 (1)	4358 (4)	6412 (5)	122,324 (100)
Mean total fruit intake (g/day)	179.3 (149)	263.2 (168)	138.8 (100)	358.8 (201)	340.5 (213)	196.4 (137)	156.9 (121)	335.2 (223)	175.9 (130)	250.2 (201)	173.3 (119)	241.0 (150)	93.9 (90)	211.4 (183)

BLEND= BLadder cancer Epidemiology and Nutritional Determinants study, EPIC=The European Prospective Investigation into Cancer and Nutrition study, NLCS=The Netherlands Cohort Study, MCCS=The Melbourne Collaborative Cohort Study, VITAL=The VITamins And Lifestyle cohort, NMIBC=non-muscle invasive bladder cancer, MIBC=muscle invasive bladder cancer

^aAs a result of the exclusion criteria, cohort study size and number of cases included in BLEND may differ from original study-specific publications

^bFor a total of 1416 bladder cancers the histological bladder cancer subtype was not specified

Table 3. Adjusted hazard ratios for all bladder cancers by total fruit consumption and the consumption of specific types of fruit

	Full cohort		Males		Females		Never smokers		Former smokers		Current smokers	
	Cases	HR ^{a,b} (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)
Total fruit												
<100 grams of fruit per day	1044	1.00 (ref)	866	1.00 (ref)	178	1.00 (ref)	169	1.00 (ref)	424	1.00 (ref)	451	1.00 (ref)
100–200 grams of fruit per day	824	0.93 (0.84-1.02)	620	0.95 (0.85-1.06)	204	0.83 (0.68-1.03)	187	1.10 (0.87-1.40)	318	0.85 (0.73-1.00)	319	0.94 (0.81-1.09)
200–300 grams of fruit per day	492	0.92 (0.82-1.04)	341	0.95 (0.83-1.10)	151	0.83 (0.65-1.04)	123	0.96 (0.73-1.26)	215	0.96 (0.80-1.16)	154	0.86 (0.71-1.05)
>300 grams of fruit per day	476	0.90 (0.79-1.02)	326	0.96 (0.83-1.12)	150	0.75 (0.59-0.97)	134	0.93 (0.70-1.23)	197	0.91 (0.74-1.11)	145	0.87 (0.70-1.08)
p for trend		>0.05		>0.05		0.04		>0.05		>0.05		>0.05
Per 100 grams a day	2836	0.97 (0.93-1.01)	2153	0.99 (0.94-1.03)	683	0.92 (0.85-0.99)	613	0.96 (0.88-1.05)	1154	0.98 (0.92-1.05)	1069	0.95 (0.89-1.01)
Citrus fruit												
Q1	773	1.00 (ref)	612	1.00 (ref)	161	1.00 (ref)	129	1.00 (ref)	292	1.00 (ref)	352	1.00 (ref)
Q2	626	0.96 (0.86-1.07)	470	0.97 (0.85-1.09)	156	0.93 (0.74-1.17)	118	0.87 (0.68-1.12)	259	1.03 (0.87-1.23)	249	0.94 (0.80-1.11)
Q3	558	0.97 (0.87-1.08)	393	0.98 (0.86-1.12)	165	0.92 (0.74-1.15)	144	1.10 (0.86-1.41)	198	0.88 (0.73-1.06)	216	0.98 (0.82-1.17)
Q4	608	0.97 (0.87-1.09)	430	1.01 (0.88-1.15)	178	0.88 (0.70-1.11)	142	0.95 (0.76-1.28)	247	1.04 (0.86-1.25)	219	0.91 (0.75-1.09)
p for trend		>0.05		>0.05		>0.05		>0.05		>0.05		>0.05
Per 25 grams a day	2565	1.00 (0.97-1.03)	1905	1.01 (0.97-1.04)	660	0.97 (0.92-1.03)	533	1.02 (0.96-1.09)	996	1.00 (0.96-1.05)	1036	0.98 (0.93-1.02)
Soft fruit												
Q1	523	1.00 (ref)	400	1.00 (ref)	123	1.00 (ref)	100	1.00 (ref)	216	1.00 (ref)	207	1.00 (ref)
Q2	952	1.02 (0.90-1.15)	754	1.04 (0.91-1.20)	198	0.94 (0.72-1.23)	199	1.13 (0.86-1.48)	393	1.06 (0.88-1.27)	360	0.95 (0.77-1.16)
Q3	857	0.94 (0.83-1.07)	653	1.00 (0.87-1.15)	204	0.78 (0.60-1.01)	198	1.00 (0.76-1.32)	333	0.90 (0.75-1.10)	326	0.98 (0.80-1.20)
Q4	504	1.00 (0.87-1.14)	346	1.08 (0.92-1.26)	158	0.79 (0.60-1.04)	116	0.86 (0.64-1.16)	212	1.11 (0.90-1.37)	176	1.00 (0.79-1.25)
p for trend		>0.05		>0.05		0.05		>0.05		>0.05		>0.05
Per 25 grams a day	2836	1.00 (0.93-1.08)	2153	1.03 (0.94-1.13)	683	0.95 (0.84-1.09)	613	0.91 (0.78-1.07)	1154	1.09 (0.97-1.22)	1069	0.98 (0.86-1.12)
Stone fruit												
Q1	469	1.00 (ref)	379	1.00 (ref)	90	1.00 (ref)	117	1.00 (ref)	208	1.00 (ref)	144	1.00 (ref)
Q2	620	1.11 (0.97-1.28)	475	1.08 (0.93-1.26)	145	1.26 (0.91-1.75)	136	0.93 (0.70-1.24)	233	1.06 (0.86-1.31)	251	1.39 (1.08-1.79)
Q3	526	1.00 (0.86-1.16)	357	1.00 (0.84-1.19)	169	1.04 (0.75-1.43)	130	0.79 (0.58-1.06)	210	1.03 (0.81-1.30)	186	1.19 (0.92-1.54)
Q4	422	1.01 (0.84-1.20)	280	1.04 (0.85-1.28)	142	0.98 (0.69-1.39)	119	0.82 (0.58-1.15)	165	1.05 (0.79-1.39)	138	1.14 (0.84-1.53)
p for trend		>0.05		>0.05		>0.05		>0.05		>0.05		>0.05
Per 25 grams a day	2037	0.99 (0.94-1.04)	1491	1.02 (0.95-1.08)	546	0.94 (0.85-1.03)	502	0.97 (0.88-1.08)	816	1.00 (0.92-1.09)	719	1.00 (0.91-1.09)

^a Model stratified by cohort, age at study entry, and sex (where applicable), and adjusted for smoking status and pack-years of cigarette smoking (where applicable), ethnicity, total vegetable consumption, alcohol intake, and total energy intake

^b The assumption of proportional hazards was violated

Table 3. continued

	Full cohort		Males		Females		Never smokers		Former smokers		Current smokers	
	Cases	HR ^{a,b} (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)
Tropical fruit												
Q1	1053	1.00 (ref)	824	1.00 (ref)	229	1.00 (ref)	189	1.00 (ref)	387	1.00 (ref)	477	1.00 (ref)
Q2	539	0.84 (0.75-0.93)	417	0.87 (0.77-0.99)	122	0.71 (0.56-0.90)	90	0.63 (0.48-0.82)	233	1.01 (0.85-1.20)	213	0.81 (0.68-0.96)
Q3	599	0.90 (0.81-1.01)	453	0.96 (0.84-1.08)	146	0.74 (0.59-0.94)	152	0.83 (0.65-1.06)	245	0.95 (0.80-1.13)	202	0.93 (0.78-1.11)
Q4	645	0.94 (0.83-1.05)	459	0.98 (0.86-1.12)	186	0.78 (0.62-0.99)	179	0.82 (0.64-1.05)	289	1.07 (0.89-1.28)	177	0.87 (0.72-1.06)
p for trend		>0.05		>0.05		0.05		>0.05		>0.05		>0.05
Per 25 grams a day	2836	0.98 (0.95-1.02)	2153	0.99 (0.95-1.03)	683	0.97 (0.91-1.04)	613	0.99 (0.93-1.06)	1154	1.01 (0.96-1.07)	1069	0.93 (0.87-1.00)
Pome fruit												
Q1	393	1.00 (ref)	331	1.00 (ref)	62	1.00 (ref)	52	1.00 (ref)	156	1.00 (ref)	185	1.00 (ref)
Q2	213	0.86 (0.73-1.02)	172	0.87 (0.72-1.05)	41	0.83 (0.56-1.24)	39	0.80 (0.53-1.22)	85	0.74 (0.56-0.96)	89	1.03 (0.80-1.32)
Q3	179	0.83 (0.69-0.99)	139	0.83 (0.68-1.02)	40	0.79 (0.52-1.18)	44	0.82 (0.54-1.23)	86	0.81 (0.62-1.06)	49	0.84 (0.61-1.14)
Q4	286	0.90 (0.77-1.05)	226	0.91 (0.77-1.09)	60	0.83 (0.58-1.20)	61	0.81 (0.55-1.18)	153	0.93 (0.74-1.17)	72	0.78 (0.60-1.03)
p for trend		>0.05		>0.05		>0.05		>0.05		>0.05		>0.05
Per 25 grams a day	1071	0.98 (0.94-1.01)	868	0.98 (0.94-1.02)	203	0.95 (0.87-1.03)	196	0.96 (0.88-1.04)	480	1.00 (0.94-1.05)	395	0.95 (0.89-1.01)
Fruit products												
Q1	345	1.00 (ref)	278	1.00 (ref)	67	1.00 (ref)	56	1.00 (ref)	161	1.00 (ref)	128	1.00 (ref)
Q2	69	0.98 (0.74-1.30)	52	1.05 (0.76-1.45)	17	0.77 (0.43-1.38)	20	0.75 (0.44-1.28)	36	1.05 (0.71-1.55)	13	1.25 (0.65-2.39)
Q3	216	0.95 (0.80-1.12)	174	0.98 (0.81-1.19)	42	0.80 (0.54-1.18)	38	0.82 (0.54-1.25)	107	1.05 (0.82-1.34)	71	0.84 (0.63-1.13)
Q4	441	0.86 (0.74-1.00)	364	0.90 (0.76-1.06)	77	0.71 (0.50-1.00)	82	0.86 (0.60-1.24)	176	0.82 (0.66-1.03)	183	0.88 (0.69-1.12)
p for trend		>0.05		>0.05		>0.05		>0.05		>0.05		>0.05
Per 25 grams a day	1071	0.98 (0.87-1.11)	868	0.97 (0.85-1.12)	203	1.00 (0.76-1.32)	196	1.12 (0.86-1.45)	480	0.85 (0.68-1.06)	395	1.03 (0.85-1.24)

^a Model stratified by cohort, age at study entry, and sex (where applicable), and adjusted for smoking status and pack-years of cigarette smoking (where applicable), ethnicity, total vegetable consumption, alcohol intake, and total energy intake

^b The assumption of proportional hazards was violated

Table 4. Adjusted hazard ratios for subtypes of bladder cancer by total fruit consumption

	All NMIBC		Lower-risk NMIBC		Higher-risk NMIBC		MIBC	
	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)	Cases	HR ^a (95% CI)
Total fruit intake in men								
<100 grams of fruit per day	386	1.00 (ref)	80	1.00 (ref)	166	1.00 (ref)	239	1.00 (ref)
100–200 grams of fruit per day	242	0.96 (0.81-1.15)	74	1.00 (0.72-1.39)	152	0.93 (0.74-1.16)	164	0.92 (0.74-1.14)
>200 grams of fruit per day	239	0.92 (0.75-1.12)	92	0.87 (0.61-1.24)	139	0.96 (0.75-1.24)	168	1.03 (0.82-1.30)
p for trend		>0.05		>0.05		>0.05		>0.05
Per 100 grams a day	867	0.96 (0.87-1.06)	246	0.93 (0.78-1.11)	457	0.98 (0.86-1.11)	571	1.01 (0.90-1.14)
Total fruit intake in women								
<100 grams of fruit per day	83	1.00 (ref)	31	1.00 (ref)	33	1.00 (ref)	35	1.00 (ref)
100–200 grams of fruit per day	77	0.75 (0.54-1.04)	39	0.85 (0.53-1.39)	38	0.74 (0.46-1.19)	44	0.91 (0.57-1.45)
>200 grams of fruit per day	108	0.63 (0.45-0.88)	59	0.76 (0.47-1.23)	46	0.52 (0.31-0.85)	56	0.94 (0.59-1.49)
p for trend		0.01		>0.05		0.01		>0.05
Per 100 grams a day	268	0.79 (0.67-0.94)	129	0.87 (0.69-1.11)	117	0.72 (0.56-0.92)	135	0.97 (0.77-1.23)
Total fruit intake in never smokers								
<100 grams of fruit per day	83	1.00 (ref)	16	1.00 (ref)	21	1.00 (ref)	43	1.00 (ref)
100–200 grams of fruit per day	75	1.11 (0.77-1.61)	33	1.22 (0.67-2.25)	36	1.07 (0.62-1.87)	33	0.87 (0.51-1.49)
>200 grams of fruit per day	90	0.80 (0.54-1.19)	50	0.97 (0.53-1.79)	34	0.58 (0.31-1.06)	50	0.99 (0.58-1.69)
p for trend		>0.05		>0.05		>0.05		>0.05
Per 100 grams a day	248	0.88 (0.72-1.06)	99	0.95 (0.71-1.26)	91	0.74 (0.56-1.00)	126	1.01 (0.78-1.32)
Total fruit intake in former smokers								
<100 grams of fruit per day	203	1.00 (ref)	39	1.00 (ref)	72	1.00 (ref)	118	1.00 (ref)
100–200 grams of fruit per day	125	0.85 (0.66-1.10)	37	0.80 (0.51-1.28)	80	0.91 (0.65-1.27)	85	0.84 (0.61-1.15)
>200 grams of fruit per day	152	0.89 (0.68-1.16)	58	0.77 (0.48-1.24)	89	1.03 (0.73-1.45)	107	0.98 (0.72-1.35)
p for trend		>0.05		>0.05		>0.05		>0.05
Per 100 grams a day	480	0.94 (0.82-1.08)	134	0.88 (0.69-1.12)	241	1.02 (0.85-1.21)	310	1.00 (0.85-1.17)
Total fruit intake in current smokers								
<100 grams of fruit per day	183	1.00 (ref)	56	1.00 (ref)	106	1.00 (ref)	113	1.00 (ref)
100–200 grams of fruit per day	119	0.92 (0.72-1.17)	43	1.01 (0.67-1.52)	74	0.88 (0.65-1.19)	90	1.00 (0.75-1.33)
>200 grams of fruit per day	105	0.81 (0.61-1.07)	43	0.86 (0.54-1.36)	62	0.82 (0.58-1.17)	67	0.96 (0.69-1.33)
p for trend		>0.05		>0.05		>0.05		>0.05
Per 100 grams a day	407	0.90 (0.79-1.03)	142	0.93 (0.74-1.17)	242	0.90 (0.76-1.07)	270	0.98 (0.84-1.15)

^aModel stratified by cohort, age at study entry, and sex (where applicable), and adjusted for smoking status and pack-years of cigarette smoking (where applicable), ethnicity, total vegetable consumption, alcohol intake, and total energy intake

