

Two bifurcation sets arising from the beta transformation with a hole at 0

Baker, Simon; Kong, Derong

DOI:

[10.1016/j.indag.2020.03.001](https://doi.org/10.1016/j.indag.2020.03.001)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Baker, S & Kong, D 2020, 'Two bifurcation sets arising from the beta transformation with a hole at 0', *Indagationes Mathematicae*, vol. 31, no. 3, pp. 436-449. <https://doi.org/10.1016/j.indag.2020.03.001>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

TWO BIFURCATION SETS ARISING FROM THE BETA TRANSFORMATION WITH A HOLE AT 0

SIMON BAKER AND DERONG KONG

ABSTRACT. Given $\beta \in (1, 2]$, the β -transformation $T_\beta : x \mapsto \beta x \pmod{1}$ on the circle $[0, 1)$ with a hole $[0, t)$ was investigated by Kalle et al. (2019). They described the set-valued bifurcation set

$$\mathcal{E}_\beta := \{t \in [0, 1) : K_\beta(t') \neq K_\beta(t) \forall t' > t\},$$

where $K_\beta(t) := \{x \in [0, 1) : T_\beta^n(x) \geq t \forall n \geq 0\}$ is the survivor set. In this paper we investigate the dimension bifurcation set

$$\mathcal{B}_\beta := \{t \in [0, 1) : \dim_H K_\beta(t') \neq \dim_H K_\beta(t) \forall t' > t\},$$

where \dim_H denotes the Hausdorff dimension. We show that if $\beta \in (1, 2]$ is a multinacci number then the two bifurcation sets \mathcal{B}_β and \mathcal{E}_β coincide. Moreover we give a complete characterization of these two sets. As a corollary of our main result we prove that for β a multinacci number we have $\dim_H(\mathcal{E}_\beta \cap [t, 1]) = \dim_H K_\beta(t)$ for any $t \in [0, 1)$. This confirms a conjecture of Kalle et al. for β a multinacci number.

1. INTRODUCTION

Given $\beta \in (1, 2]$, the β -transformation T_β on the circle $\mathbb{R}/\mathbb{Z} \sim [0, 1)$ is defined by

$$T_\beta : [0, 1) \rightarrow [0, 1); \quad x \mapsto \beta x \pmod{1}.$$

Following the pioneering work of Rényi [11] and Parry [9] there has been a great interest in the study of T_β . In general, the system $\Phi_\beta = ([0, 1), T_\beta)$ does not admit a Markov partition (cf. [12]), this makes describing the dynamics of Φ_β more challenging.

When $\beta = 2$, Urbański considered in [14, 15] the open dynamical system under the doubling map T_2 with a hole at zero. More precisely, for $t \in [0, 1)$ let

$$K_2(t) := \{x \in [0, 1) : T_2^n(x) \geq t \forall n \geq 0\}.$$

Here we use a slightly different definition of $K_2(t)$ from that by Urbański. By [14, Theorem 1 and Corollary 1] it follows that the dimension function $t \mapsto \eta_2(t) := \dim_H K_2(t)$ is a Devil's staircase on $[0, 1)$, that is (i) η_2 is decreasing and continuous on $[0, 1)$; (ii) η_2 is locally constant almost everywhere on $[0, 1)$; and (iii) η_2 is not constant on $[0, 1)$. Here and throughout the paper \dim_H denotes the Hausdorff dimension. Moreover, Urbański investigated the bifurcation sets

$$\mathcal{E}_2 := \{t \in [0, 1) : K_2(t') \neq K_2(t) \forall t' > t\} \quad \text{and} \quad \mathcal{B}_2 := \{t \in [0, 1) : \eta_2(t') \neq \eta_2(t) \forall t' > t\}.$$

Clearly, $\mathcal{B}_2 \subseteq \mathcal{E}_2$. It can be easily deduced from the proof of Theorem 1 in [14] that $\mathcal{B}_2 = \mathcal{E}_2$, and its topological closure $\overline{\mathcal{B}_2}$ is a *Cantor set*, i.e., a non-empty compact set that has neither

Date: 20th January 2020.

2010 Mathematics Subject Classification. Primary: 37B10, Secondary: 28A78, 11A63.

Key words and phrases. Bifurcation sets; beta transformation; local dimension; survivor set.

isolated nor interior points. Furthermore, the following local dimension property was shown to hold: $\lim_{r \rightarrow 0} \dim_H(\mathcal{E}_2 \cap (t-r, t+r)) = \eta_2(t)$ for all $t \in \mathcal{E}_2$. Recently, Carminati and Tiozzo in [1] showed that the local Hölder exponent of the dimension function η_2 at any $t \in \mathcal{E}_2$ equals $\eta_2(t)$.

Inspired by the work of Urbański [14, 15], Kalle et al. in [6] considered the analogous problem for the β -transformation with a hole $[0, t)$. More precisely, for $t \in [0, 1)$ they investigated the survivor set

$$K_\beta(t) := \{x \in [0, 1) : T_\beta^n(x) \geq t \forall n \geq 0\},$$

and showed that the dimension function $t \mapsto \dim_H K_\beta(t)$ is also a Devil's staircase on $[0, 1)$. Furthermore, they characterized the *set-valued bifurcation set*

$$\mathcal{E}_\beta := \{t \in [0, 1) : K_\beta(t') \neq K_\beta(t) \forall t' > t\},$$

and proved that \mathcal{E}_β is a Lebesgue null set of full Hausdorff dimension for any $\beta \in (1, 2)$. Note that the bifurcation set \mathcal{E}_β defined here coincides with the set

$$E_\beta^+ := \{t \in [0, 1) : T_\beta^n(t) \geq t \forall n \geq 0\}$$

in [6]. Interestingly, they showed that \mathcal{E}_β contains infinitely many isolated points for Lebesgue almost every $\beta \in (1, 2)$. This is in contrast to the case where $\beta = 2$ and \mathcal{E}_2 has no isolated points. For β -transformation with an arbitrary hole we refer to the work of Clark [2]. We also mention that the study of bifurcation sets plays an important role in one-dimensional dynamics (cf. [5]).

Since for each $\beta \in (1, 2)$ the dimension function $\eta_\beta : t \mapsto \dim_H K_\beta(t)$ is a Devil's staircase, it is natural to consider the *dimension bifurcation set*

$$\mathcal{B}_\beta := \{t \in [0, 1) : \eta_\beta(t') \neq \eta_\beta(t) \forall t' > t\}.$$

This set records those t for which the dimension function η_β has a ‘change’ within any right neighborhood. Since η_β is continuous, \mathcal{B}_β cannot have isolated points. On the other hand, the set-valued bifurcation set \mathcal{E}_β contains (infinitely many) isolated points for Lebesgue almost every $\beta \in (1, 2)$. So in general we cannot expect the coincidence of the two bifurcation sets \mathcal{B}_β and \mathcal{E}_β . That being said, in this paper we show that if β is a multinacci number, i.e., the unique root in $(1, 2)$ of the equation

$$x^{m+1} = x^m + x^{m-1} + \dots + x + 1$$

for some $m \in \mathbb{N}$, then the two bifurcation sets indeed coincide. Importantly, if β is a multinacci number then its quasi-greedy expansion of 1 is of the form $((1^m 0)^\infty)$. This property will be useful in our analysis. Here for $\beta \in (1, 2]$ the *quasi-greedy* β -expansion $\delta(\beta) = \delta_1(\beta)\delta_2(\beta)\dots$ of 1 is the lexicographically largest zero-one sequence not ending with an infinite string of zeros and satisfying $1 = \sum_{i=1}^{\infty} \delta_i(\beta)/\beta^i$ (see Section 2 for more details). Furthermore, throughout the paper we will use lexicographical order ‘ \prec, \preceq, \succ ’ and ‘ \succcurlyeq ’ between sequences and words.

When $\beta \in (1, 2)$ is a multinacci number, the following result for the set-valued bifurcation set \mathcal{E}_β was established in [6, Theorems C and D]. We record it here for later use.

Theorem 1.1 ([6]). *Let $\beta \in (1, 2]$ be a multinacci number. Then the topological closure $\overline{\mathcal{E}_\beta}$ is a Cantor set. Furthermore, $\max \overline{\mathcal{E}_\beta} = 1 - 1/\beta$.*

In order to give a complete description of the dimension bifurcation set \mathcal{B}_β we introduce a class of basic intervals.

Definition 1.2. Let $\beta \in (1, 2]$. A word $s_1 \dots s_m$ is called β -Lyndon if

$$s_{i+1} \dots s_m \succ s_1 \dots s_{m-i} \quad \forall 1 \leq i < m, \quad \text{and} \quad \sigma^n((s_1 \dots s_m)^\infty) \prec \delta(\beta) \quad \forall n \geq 0.$$

Accordingly, an interval $[t_L, t_R) \subset [0, 1)$ is called a β -Lyndon interval if there exists a β -Lyndon word $s_1 \dots s_m$ such that

$$t_L = \sum_{i=1}^m \frac{s_i}{\beta^i} \quad \text{and} \quad t_R = \frac{\beta^m}{\beta^m - 1} \cdot t_L.$$

Here we mention that in Definition 1.2 the left endpoint $t_L = (s_1 \dots s_m 0^\infty)_\beta$ has a finite β -expansion and the right endpoint $t_R = ((s_1 \dots s_m)^\infty)_\beta$ has a periodic β -expansion, see Section 2 for more explanations.

We will show that the β -Lyndon intervals are pairwise disjoint for all $\beta \in (1, 2]$, and when β is multinacci they cover the interval $[0, 1 - 1/\beta)$ up to a Lebesgue null set. The latter statement can be seen as a consequence of our main result for the coincidence of the two bifurcation sets, which we state below.

Theorem 1. *Let $\beta \in (1, 2]$ be a multinacci number. Then*

$$\begin{aligned} \mathcal{B}_\beta &= \mathcal{E}_\beta = \left[0, 1 - \frac{1}{\beta}\right) \setminus \bigcup [t_L, t_R) \\ &= \left\{ t \in [0, 1) : \lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t, t+r)) = \dim_H K_\beta(t) > 0 \right\}, \end{aligned}$$

where the union is taken over all pairwise disjoint β -Lyndon intervals.

By Theorem 1 it follows that the topological closure $[t_L, t_R]$ of each β -Lyndon interval is indeed a maximal interval where the dimension function η_β is constant. As a corollary of Theorem 1 we confirm a conjecture of [6] for β a multinacci number.

Corollary 2. *If $\beta \in (1, 2]$ is a multinacci number, then*

$$\dim_H(\mathcal{E}_\beta \cap [t, 1]) = \dim_H K_\beta(t) \quad \forall t \in [0, 1).$$

The rest of the paper is organized as follows. In Section 2 we recall some properties from symbolic dynamics and the dimension formula for the survivor set $K_\beta(t)$. The proof of Theorem 1 and Corollary 2 will be given in Section 3. In Section 4 we make some remarks and point out that the method of proof for Theorem 1 can be applied to some other special values of $\beta \in (1, 2]$.

2. PRELIMINARIES AND β -LYNDON INTERVALS

Given $\beta \in (1, 2]$, for each $x \in I_\beta := [0, 1/(\beta - 1)]$ there exists a sequence $(d_i) = d_1 d_2 \dots \in \{0, 1\}^\mathbb{N}$ such that

$$x = \sum_{i=1}^{\infty} \frac{d_i}{\beta^i} =: ((d_i))_\beta.$$

The sequence (d_i) is called a β -expansion of x . Sidorov [13] showed that for $\beta \in (1, 2)$ Lebesgue almost every $x \in I_\beta$ has a continuum of β -expansions. This is rather different from the case when $\beta = 2$ where every number in $I_2 = [0, 1]$ has a unique dyadic expansion except

for countably many points that have precisely two expansions. Given $x \in I_\beta$, among all of its β -expansions let

$$b(x, \beta) = (b_i(x, \beta))$$

be the *greedy* β -expansion of x , i.e., the lexicographically largest β -expansion of x . Such a sequence always exists and is generated by the orbit of x under the map T_β . Let σ be the *left-shift* on $\{0, 1\}^{\mathbb{N}}$ defined by $\sigma((c_i)) = (c_{i+1})$. Then $b(T_\beta(x), \beta) = \sigma(b(x, \beta))$ for any $x \in [0, 1)$. Similarly, for $x \in (0, 1/(\beta - 1)]$ let

$$a(x, \beta) = (a_i(x, \beta))$$

be the *quasi-greedy* β -expansion of x (cf. [3]), which is the lexicographically largest β -expansion of x not ending with 0^∞ . Here for a word \mathbf{c} we denote by $\mathbf{c}^\infty := \mathbf{c}\mathbf{c}\dots$ the periodic sequence with periodic block \mathbf{c} . Throughout the paper we will use the lexicographic order between sequences and words in the usual way. For example, for two sequences $(c_i), (d_i) \in \{0, 1\}^{\mathbb{N}}$ we write $(c_i) \prec (d_i)$ if $c_1 < d_1$, or there exists $n > 1$ such that $c_1 \dots c_{n-1} = d_1 \dots d_{n-1}$ and $c_n < d_n$. Furthermore, for two words \mathbf{c}, \mathbf{d} we say $\mathbf{c} \prec \mathbf{d}$ if $\mathbf{c}0^\infty \prec \mathbf{d}0^\infty$.

For $\beta \in (1, 2]$ recall that

$$\delta(\beta) = \delta_1(\beta)\delta_2(\beta)\dots$$

is the quasi-greedy β -expansion of 1, i.e., $\delta(\beta) = a(1, \beta)$. The following lexicographic characterizations of $\delta(\beta)$ and the greedy expansion $b(x, \beta)$ are essentially due to Parry [9] (see also [4]).

Lemma 2.1. (i) *The map $\beta \mapsto \delta(\beta)$ is a strictly increasing bijection from $(1, 2]$ onto the set of sequences $(\delta_i) \in \{0, 1\}^{\mathbb{N}}$ not ending with 0^∞ and satisfying*

$$\sigma^n((\delta_i)) \preceq (\delta_i) \quad \forall n \geq 0.$$

(ii) *Let $\beta \in (1, 2]$. Then the map $x \mapsto b(x, \beta)$ is a strictly increasing bijection from $[0, 1)$ onto the set of all sequences $(b_i) \in \{0, 1\}^{\mathbb{N}}$ satisfying*

$$\sigma^n((b_i)) \prec \delta(\beta) \quad \forall n \geq 0.$$

(iii) *For any $\beta \in (1, 2)$ the sequence $b(1, \beta) = (b_i)$ satisfies $\sigma^n((b_i)) \prec \delta(\beta) \forall n \geq 1$.*

For $\beta \in (1, 2]$ let $[t_L, t_R)$ be a β -Lyndon interval generated by a β -Lyndon word $s_1 \dots s_m$. Then by Definition 1.2 and Lemma 2.1 (ii) it follows that

$$b(t_L, \beta) = s_1 \dots s_m 0^\infty \quad \text{and} \quad b(t_R, \beta) = (s_1 \dots s_m)^\infty.$$

Lemma 2.2. *For any $\beta \in (1, 2]$ the β -Lyndon intervals are pairwise disjoint.*

Proof. Let $[t_L, t_R)$ and $[t'_L, t'_R)$ be two β -Lyndon intervals generated by the β -Lyndon words $s_1 \dots s_p$ and $s'_1 \dots s'_q$, respectively. Suppose on the contrary that $[t_L, t_R) \cap [t'_L, t'_R) \neq \emptyset$. Without loss of generality we assume $t_L < t'_L < t_R$. Then by Definition 1.2 and Lemma 2.1(ii) it follows that

$$s_1 \dots s_p 0^\infty \prec s'_1 \dots s'_q 0^\infty \prec (s_1 \dots s_p)^\infty.$$

This implies

$$q > p, \quad s'_1 \dots s'_p = s_1 \dots s_p \quad \text{and} \quad s'_{p+1} \dots s'_q 0^\infty \prec (s_1 \dots s_p)^\infty.$$

Write $q = Np + r$ with $N \geq 1$ and $0 < r \leq p$. So, either there exists $1 \leq k < N$ such that

$$s'_{p+1} \dots s'_{kp} = (s_1 \dots s_p)^{k-1} \quad \text{and} \quad s'_{kp+1} \dots s'_{(k+1)p} \prec s_1 \dots s_p,$$

or

$$s'_{p+1} \dots s'_{Np} = (s_1 \dots s_p)^{N-1} \quad \text{and} \quad s'_{Np+1} \dots s'_q \preceq s_1 \dots s_{q-Np}.$$

Using $s'_1 \dots s'_p = s_1 \dots s_p$ we conclude in both cases that

$$s'_{j+1} \dots s'_q \preceq s'_1 \dots s'_{q-j} \quad \text{for some } j \in \{p, p+1, \dots, q-1\}.$$

This is not possible by the definition of a β -Lyndon word. \square

To describe the Hausdorff dimension of the survivor set

$$K_\beta(t) = \{x \in [0, 1] : T_\beta^n(x) \geq t \quad \forall n \geq 0\},$$

we recall from [8, Chapter 4] the definition of topological entropy for a symbolic set. For a set $X \subset \{0, 1\}^{\mathbb{N}}$, its *topological entropy* is defined to be

$$h(X) = \liminf_{n \rightarrow \infty} \frac{\log \#B_n(X)}{n},$$

where $B_n(X)$ is the set of all length n prefixes of sequences from X .

The following characterization of the set-valued bifurcation set \mathcal{E}_β was implicitly given in [14] (see also [6, Proposition 2.3]). Furthermore, the Hausdorff dimension of $K_\beta(t)$ was implicitly given by Raith in [10], and was recently explicitly presented in [6, Equation (2.6)].

Proposition 2.3. (i) *Let $\beta \in (1, 2]$. Then*

$$\mathcal{E}_\beta = \{t \in [0, 1] : T_\beta^n(t) \geq t \quad \forall n \geq 0\}.$$

(ii) *Let $\beta \in (1, 2]$ and $t \in [0, 1]$. Then the Hausdorff dimension of $K_\beta(t)$ is given by*

$$\dim_H K_\beta(t) = \frac{h(\tilde{K}_\beta(t))}{\log \beta},$$

where $\tilde{K}_\beta(t) := \{(x_i) \in \{0, 1\}^{\mathbb{N}} : b(t, \beta) \preceq \sigma^n((x_i)) \preceq \delta(\beta) \quad \forall n \geq 0\}$. Furthermore, the dimension function $\eta_\beta : t \mapsto \dim_H K_\beta(t)$ is a Devil's staircase, i.e., η_β is a non-constant, decreasing and continuous function which is locally constant almost everywhere in $[0, 1]$.

3. PROOF OF THEOREM 1

In this section we will prove Theorem 1. First we show that the dimension bifurcation set \mathcal{B}_β coincides with the set-valued bifurcation set \mathcal{E}_β , we then derive a complete characterization of these sets via the β -Lyndon intervals. The proof heavily relies upon the transitivity of the symbolic survivor set $\tilde{K}_\beta(t)$ (see Lemma 3.2 below).

Proposition 3.1. *Let $\beta \in (1, 2)$ be a multinacci number. Then*

$$\mathcal{B}_\beta = \mathcal{E}_\beta = \left[0, 1 - \frac{1}{\beta}\right) \setminus \bigcup [t_L, t_R),$$

where the union is taken over all β -Lyndon intervals.

Observe by Lemma 2.2 that the β -Lyndon intervals are pairwise disjoint. In fact the closed β -Lyndon intervals $\{[t_L, t_R]\}$ are also pairwise disjoint. So by Proposition 3.1 it follows that each closed β -Lyndon interval is a maximal interval where the dimension function η_β is constant.

The proof of Proposition 3.1 will be split into several lemmas. We fix a multinacci number $\beta \in (1, 2)$ with $\delta(\beta) = (1^m 0)^\infty$ for some $m \geq 1$. In view of Proposition 2.3 it is necessary to investigate the symbolic survivor set

$$\tilde{K}_\beta(t) = \left\{ (x_i) \in \{0, 1\}^{\mathbb{N}} : b(t, \beta) \preceq \sigma^n((x_i)) \preceq \delta(\beta) \ \forall n \geq 0 \right\}.$$

Lemma 3.2. *Let $\beta \in (1, 2)$ with $\delta(\beta) = (1^m 0)^\infty$, and let $[t_L, t_R] \subset [0, 1 - 1/\beta)$ be a β -Lyndon interval. Then the set-valued map $t \mapsto \tilde{K}_\beta(t)$ is constant on $[t_L, t_R]$, and the set $\tilde{K}_\beta(t_R)$ is a transitive subshift of finite type.*

Proof. Suppose $[t_L, t_R]$ is a β -Lyndon interval generated by $s_1 \dots s_p$. First we claim that

$$(3.1) \quad \sigma^n((x_i)) \succcurlyeq s_1 \dots s_p 0^\infty \ \forall n \geq 0 \iff \sigma^n((x_i)) \succcurlyeq (s_1 \dots s_p)^\infty \ \forall n \geq 0.$$

Since $(s_1 \dots s_p)^\infty \succ s_1 \dots s_p 0^\infty$, the implication ‘ \Leftarrow ’ in (3.1) is obvious. For the reverse implication we assume $\sigma^n((x_i)) \prec (s_1 \dots s_p)^\infty$ for some $n \geq 0$. Then there exists $\ell \geq 0$ such that

$$x_{n+1} \dots x_{n+\ell p} = (s_1 \dots s_p)^\ell \quad \text{and} \quad x_{n+\ell p+1} \dots x_{n+(\ell+1)p} \prec s_1 \dots s_p.$$

This yields $\sigma^{n+\ell p}((x_i)) \prec s_1 \dots s_p 0^\infty$, completing the proof of ‘ \Rightarrow ’ in (3.1).

Take $t \in [t_L, t_R]$. Then by Lemma 2.1(ii) it follows that

$$\tilde{K}_\beta(t_R) \subseteq \tilde{K}_\beta(t) \subseteq \tilde{K}_\beta(t_L).$$

Observe that $\delta(\beta) = (1^m 0)^\infty$ for some $m \in \mathbb{N}$. Then

$$(3.2) \quad \begin{aligned} \tilde{K}_\beta(t_L) &= \{(x_i) : s_1 \dots s_p 0^\infty \preceq \sigma^n((x_i)) \preceq (1^m 0)^\infty \ \forall n \geq 0\} \\ &= \{(x_i) : (s_1 \dots s_p)^\infty \preceq \sigma^n((x_i)) \preceq (1^m 0)^\infty \ \forall n \geq 0\} = \tilde{K}_\beta(t_R). \end{aligned}$$

So, the set-valued map $t \mapsto \tilde{K}_\beta(t)$ is constant on $[t_L, t_R]$. Furthermore, $\tilde{K}_\beta(t_R)$ is a subshift of finite type with the set of forbidden blocks given by

$$\mathcal{F} = \left\{ c_1 \dots c_k \in \{0, 1\}^k : c_1 \dots c_k 0^\infty \prec s_1 \dots s_p 0^\infty \text{ or } c_1 \dots c_k 0^\infty \succ (1^m 0)^\infty \right\},$$

where $k = \max\{p, m+1\}$. It remains to prove the transitivity of $\tilde{K}_\beta(t_R)$.

Since $[t_L, t_R] \subset [0, 1 - \frac{1}{\beta})$, by Lemma 2.1 (ii) it follows that $b(t_R, \beta) \prec b(1 - \frac{1}{\beta}, \beta)$, which gives

$$(3.3) \quad (s_1 \dots s_p)^\infty \prec 01^m 0^\infty.$$

Arbitrarily fix an admissible word $\varepsilon = \varepsilon_1 \dots \varepsilon_k$ and an admissible sequence $\gamma = \gamma_1 \gamma_2 \dots$ in $\tilde{K}_\beta(t_R)$. We will construct a word ν such that $\varepsilon \nu \gamma \in \tilde{K}_\beta(t_R)$. Observe that $\sigma^n((s_1 \dots s_p)^\infty) \prec (1^m 0)^\infty$ for all $n \geq 0$. Thus, there exists a large integer N such that

$$(3.4) \quad \sigma^n((s_1 \dots s_p)^\infty) \prec (1^m 0)^N 0^\infty \quad \text{for all } n \geq 0.$$

Denote by $(\delta_i) := \delta(\beta) = (1^m 0)^\infty$. Note that $\varepsilon_{i+1} \dots \varepsilon_k \preceq \delta_1 \dots \delta_{k-i}$ for all $0 \leq i < k$. Let $i_0 \in \{0, 1, \dots, k-1\}$ be the smallest index such that

$$\varepsilon_{i_0+1} \dots \varepsilon_k = \delta_1 \dots \delta_{k-i_0}.$$

If such an index i_0 does not exist, then we put $i_0 = k$. In either case there exists a word μ such that $\varepsilon \mu = \varepsilon_1 \dots \varepsilon_{i_0} (1^m 0)^N$. Since $\gamma \preceq (1^m 0)^\infty$, there exists $q \in \{0, 1, \dots, m\}$ such that

γ begins with $\gamma_1 \dots \gamma_{q+1} = 1^q 0$. We emphasize here that if $q = 0$ then γ begins with digit 0. Now we claim that

$$\varepsilon \mu 1^{m-q} \gamma = \varepsilon_1 \dots \varepsilon_{i_0} (1^m 0)^{N+1} \gamma_{q+2} \gamma_{q+3} \dots \in \tilde{K}_\beta(t_R),$$

or equivalently,

$$(3.5) \quad (s_1 \dots s_p)^\infty \preceq \sigma^n(\varepsilon \mu 1^{m-q} \gamma) \preceq (1^m 0)^\infty \quad \text{for all } n \geq 0.$$

First we prove the second inequality in (3.5). By the definition of i_0 it follows that $\sigma^n(\varepsilon \mu 1^{m-q} \gamma) \prec \delta(\beta) = (1^m 0)^\infty$ holds for all $0 \leq n < i_0$. Furthermore, since $\gamma \in \tilde{K}_\beta(t_R)$, the second inequality in (3.5) also holds for $n \geq |\varepsilon| + |\mu| + m - q$. Here for a word \mathbf{c} we denote its length by $|\mathbf{c}|$. For the remaining n we observe that $\sigma^{i_0}(\varepsilon \mu 1^{m-q} \gamma) = (1^m 0)^{N+1} \gamma_{q+2} \gamma_{q+3} \dots$ and $\gamma_{q+2} \gamma_{q+3} \dots \in \tilde{K}_\beta(t_R)$. So it is easy to verify that

$$\sigma^n(\varepsilon \mu 1^{m-q} \gamma) \preceq (1^m 0)^\infty \quad \text{for all } i_0 \leq n < |\varepsilon| + |\mu| + m - q.$$

This proves the second inequality in (3.5).

For the first inequality in (3.5) we observe that $\varepsilon \mu 1^{m-q} \gamma = \varepsilon_1 \dots \varepsilon_{i_0} (1^m 0)^N 1^m \gamma_{q+1} \gamma_{q+2} \dots$ and $\gamma_{q+1} \gamma_{q+2} \dots \in \tilde{K}_\beta(t_R)$. Then by (3.3) it follows that

$$\sigma^n(\varepsilon \mu 1^{m-q} \gamma) \succeq (s_1 \dots s_p)^\infty \quad \text{for all } n \geq i_0.$$

If $i_0 = 0$, then we are done. Otherwise, we take $0 \leq n < i_0$. Since $\varepsilon_1 \dots \varepsilon_{i_0}$ is an admissible word in $\tilde{K}_\beta(t_R)$, we have

$$\varepsilon_{n+1} \dots \varepsilon_{i_0} \succ t_1 \dots t_{i_0-n},$$

where $(t_i) := (s_1 \dots s_p)^\infty$. The first inequality in (3.5) now holds by (3.4), which tells us that

$$(1^m 0)^N 1^m \gamma_{q+1} \gamma_{q+2} \dots \succ t_{i_0-n+1} t_{i_0-n+2} \dots$$

This completes the proof of our claim.

Since ε and γ are chosen arbitrarily, it follows that $\tilde{K}_\beta(t_R)$ is transitive. \square

Remark 3.3. • The fact that $\tilde{K}_\beta(t_R)$ is a subshift of finite type can also be deduced from [7].

- The proof of Lemma 3.2 can be adjusted to prove the more general case with $\beta > 2$ with $\delta(\beta) = (M^m k)^\infty$, where $M = \lceil \beta \rceil - 1$ and $k \in \{0, 1, \dots, M - 1\}$. The transitivity property of $\tilde{K}_\beta(t_R)$ holds only for t_R sufficiently close to 0.

To prove the coincidence of \mathcal{B}_β and \mathcal{E}_β we still need the following inequalities.

Lemma 3.4. *Let $(t_1 \dots t_N)^\infty \in \{0, 1\}^\mathbb{N}$ be a periodic sequence with period $N \geq 2$. If*

$$\sigma^n((t_1 \dots t_N)^\infty) \succeq (t_1 \dots t_N)^\infty \quad \forall n \geq 0,$$

then

$$t_{j+1} \dots t_N \succ t_1 \dots t_{N-j} \quad \forall 1 \leq j < N.$$

Proof. Note that $N \geq 2$ is the period of $(t_1 \dots t_N)^\infty$, and

$$(3.6) \quad \sigma^n((t_1 \dots t_N)^\infty) \succeq (t_1 \dots t_N)^\infty \quad \forall n \geq 0.$$

Then $t_1 = 0$ and $t_N = 1$. Taking the reflection on both sides of (3.6) it follows that

$$\sigma^n(\overline{(t_1 \dots t_N)^\infty}) \preceq \overline{(t_1 \dots t_N)^\infty} \quad \text{for all } n \geq 0.$$

Here for a word $c_1 \dots c_k \in \{0, 1\}^k$ its reflection is defined by $\overline{c_1 \dots c_k} := (1 - c_1)(1 - c_2) \dots (1 - c_k)$. By Lemma 2.1(i) it follows that $(t_1 \dots t_N)^\infty$ is the quasi-greedy expansion of 1 for some base $\beta' \in (1, 2]$, i.e., $\delta(\beta') = (t_1 \dots t_N)^\infty$. Since N is the period of the sequence $\delta(\beta')$, the greedy β' -expansion of 1 is given by

$$b(1, \beta') = \overline{t_1 \dots t_{N-1}} 10^\infty.$$

So, by Lemma 2.1 (iii) it follows that

$$\overline{t_{j+1} \dots t_N} \prec \overline{t_{j+1} \dots t_{N-1}} 1 \preccurlyeq \overline{t_1 \dots t_{N-j}} \quad \text{for all } 1 \leq j < N.$$

Then the lemma follows by taking the reflection in the above equation. \square

Now we prove the coincidence of the two bifurcation sets.

Lemma 3.5. *Let $\beta \in (1, 2)$ with $\delta(\beta) = (1^m 0)^\infty$. Then $\mathcal{E}_\beta = \mathcal{B}_\beta$.*

Proof. By the definition of the two bifurcation sets it is easy to see that $\mathcal{B}_\beta \subset \mathcal{E}_\beta$. So in the following we prove $\mathcal{E}_\beta \subset \mathcal{B}_\beta$.

Let $t \in \mathcal{E}_\beta$ with its greedy β -expansion $b(t, \beta) = (t_i)$. Then by Theorem 1.1 we have $t \leq 1 - 1/\beta < 1/\beta$. This gives $t_1 = 0$. By Lemmas 2.1 (ii) and Proposition 2.3 (i) it follows that

$$\sigma^n((t_i)) \succcurlyeq (t_i) \quad \text{for all } n \geq 0.$$

Let $N \geq 1$ be the smallest index such that $\sigma^N((t_i)) = (t_i)$. If such an integer N does not exist, then we set $N = \infty$. In the following we will prove $t \in \mathcal{B}_\beta$ by considering the following two cases: (I) $N < \infty$; and (II) $N = \infty$.

Case (I). $N < \infty$. We claim that $t_1 \dots t_N$ is a β -Lyndon word. If $N = 1$, then $(t_i) = t_1^\infty = 0^\infty$. It is easy to check that $t_1 = 0$ is a β -Lyndon word. In the following we assume $N \geq 2$. Since $\sigma^N((t_i)) = (t_i)$, we have $(t_i) = (t_1 \dots t_N)^\infty$. Note that (t_i) is the greedy β -expansion of t . Then by Lemma 2.1 (ii) it follows that

$$\sigma^n((t_1 \dots t_N)^\infty) \prec \delta(\beta) \quad \text{for all } n \geq 0.$$

Note that $\sigma^n((t_1 \dots t_N)^\infty) \succcurlyeq (t_1 \dots t_N)^\infty$. Then by Lemma 3.4 and the definition of N , it follows that

$$t_{j+1} \dots t_N \succ t_1 \dots t_{N-j} \quad \text{for all } 1 \leq j < N.$$

So by Definition 1.2 we establish the claim.

Hence, $t = ((t_1 \dots t_N)^\infty)_\beta = t_R$ is the right endpoint of a β -Lyndon interval generated by $t_1 \dots t_N$. By Lemma 3.2 it follows that $\tilde{K}_\beta(t)$ is a transitive subshift of finite type. Observe that for any $t' > t$ we have

$$\tilde{K}_\beta(t') \subset \tilde{K}_\beta(t) \quad \text{and} \quad (t_1 \dots t_N)^\infty \in \tilde{K}_\beta(t) \setminus \tilde{K}_\beta(t').$$

Recall by [8, Corollary 4.4.9] that for any transitive subshift of finite type, any proper subshift has strictly smaller topological entropy. Therefore,

$$h(\tilde{K}_\beta(t')) < h(\tilde{K}_\beta(t)) \quad \text{for any } t' > t.$$

By Proposition 2.3 (ii) this yields $\eta_\beta(t') < \eta_\beta(t)$ for any $t' > t$. So $t \in \mathcal{B}_\beta$.

Case (II). $N = \infty$. Then $\sigma^n((t_i)) \succcurlyeq (t_i)$ for all $n \geq 1$. So (t_i) is not periodic. Observe that (t_i) begins with digit 0, and

$$\sigma^n((t_i)) \prec (1^m 0)^\infty \quad \text{for all } n \geq 0.$$

So there exists a subsequence (m_k) of positive integers such that for any $k \geq 1$ we have $t_{m_k} = 0$, and the word $t_1 \dots t_{m_k}^+ := t_1 \dots t_{m_k-1}1$ does not contain $m+1$ consecutive ones. Then by noting $t_1 = 0$ it follows that

$$\sigma^n((t_1 \dots t_{m_k}^+)^\infty) \prec (1^m 0)^\infty \quad \forall n \geq 0.$$

Since $\sigma^n((t_i)) \succ (t_i)$ for all $n \geq 0$, by Definition 1.2 it follows that $t_1 \dots t_{m_k}^+$ is a β -Lyndon word for any $k \geq 1$. Let $s_k := ((t_1 \dots t_{m_k}^+)^\infty)_\beta$. Then s_k is the right endpoint of a β -Lyndon interval generated by $t_1 \dots t_{m_k}^+$. Furthermore, s_k strictly decreases to $t = ((t_i))_\beta$ as $k \rightarrow \infty$.

So, for any $t' > t$ we can find k such that $s_k \in (t, t')$. By the same arguments as in the proof of Case (I) for s_k we conclude that

$$\eta_\beta(t') < \eta_\beta(s_k) \leq \eta_\beta(t).$$

So $t \in \mathcal{B}_\beta$, completing the proof. \square

Finally, we describe the bifurcation sets via the β -Lyndon intervals.

Lemma 3.6. *Let $\beta \in (1, 2]$ with $\delta(\beta) = (1^m 0)^\infty$. Then*

$$\left[0, 1 - \frac{1}{\beta}\right) \setminus \bigcup [t_L, t_R) \subset \mathcal{E}_\beta.$$

Proof. Take $t \in [0, 1 - 1/\beta) \setminus \mathcal{E}_\beta$ with its greedy β -expansion (t_i) . Then $t_1 = 0$. Since $t \notin \mathcal{E}_\beta$, by Proposition 2.3 (i) there exists a smallest positive integer N such that $T_\beta^N(t) < t$, which implies

$$(3.7) \quad t_{N+1}t_{N+2} \dots \prec (t_i).$$

We claim that $t_1 \dots t_N$ is a β -Lyndon word. Clearly, if $N = 1$ then $t_1 = 0$ is a β -Lyndon word. In the following we assume $N \geq 2$. By Definition 1.2 it suffices to prove

$$(3.8) \quad t_{j+1} \dots t_N \succ t_1 \dots t_{N-j} \quad \text{for all } 1 \leq j < N,$$

and

$$(3.9) \quad \sigma^n((t_1 \dots t_N)^\infty) \prec (1^m 0)^\infty \quad \text{for all } n \geq 0.$$

First we prove (3.8). By the definition of N in (3.7) it follows that

$$(3.10) \quad t_{j+1}t_{j+2} \dots \succ (t_i) \quad \text{for all } 1 \leq j < N,$$

which implies $t_{j+1} \dots t_N \succ t_1 \dots t_{N-j}$ for all $1 \leq j < N$. Suppose $t_{j+1} \dots t_N = t_1 \dots t_{N-j}$ for some $j \in \{1, 2, \dots, N-1\}$. Applying (3.7) and then (3.10) it follows that

$$t_{j+1}t_{j+2} \dots = t_1 \dots t_{N-j}t_{N+1}t_{N+2} \dots \prec t_1 \dots t_{N-j}t_1t_2 \dots \preceq (t_i),$$

leading to a contradiction with the minimality of N . This proves (3.8).

To prove (3.9) we observe that $\delta(\beta) = (1^m 0)^\infty$ and (t_i) is the greedy β -expansion of t . Then by Lemma 2.1 (ii) it follows that $t_1 \dots t_N$ cannot contain $m+1$ consecutive ones. Since $t_1 = 0$, we have

$$\sigma^n((t_1 \dots t_N)^\infty) \preceq (1^m 0)^\infty \quad \text{for all } n \geq 0.$$

So to prove (3.9) it remains to prove that $\sigma^n((t_1 \dots t_N)^\infty) \neq (1^m 0)^\infty$ for any $n \geq 0$. Suppose the equality $\sigma^n((t_1 \dots t_N)^\infty) = (1^m 0)^\infty$ holds for some $n \geq 0$. Then by using $t_1 = 0$ it follows that

$$t_1 \dots t_{m+1} = 01^m.$$

This implies $b(t, \beta) = (t_i) \succ 01^m 0^\infty = b(1 - 1/\beta, \beta)$. By Lemma 2.1 (ii) we have $t \geq 1 - 1/\beta$, leading to a contradiction. This establishes (3.9).

By the claim there exists a β -Lyndon interval $[t_L, t_R)$ generated by $t_1 \dots t_N$. Furthermore, by (3.7) it follows that

$$\begin{aligned} (t_i) &= t_1 \dots t_N t_{N+1} t_{N+2} \dots \prec t_1 \dots t_N t_1 t_2 \dots = (t_1 \dots t_N)^2 t_{N+1} t_{N+2} \dots \\ &\prec (t_1 \dots t_N)^2 t_1 t_2 \dots = (t_1 \dots t_N)^3 t_{N+1} t_{N+2} \dots \\ &\dots \\ &\preccurlyeq (t_1 \dots t_N)^\infty. \end{aligned}$$

Therefore, $t_1 \dots t_N 0^\infty \preccurlyeq (t_i) \prec (t_1 \dots t_N)^\infty$, which gives $t \in [t_L, t_R)$ by Lemma 2.1 (ii). This completes the proof. \square

Proof of Proposition 3.1. By Lemmas 3.5 and 3.6 it suffices to prove

$$\mathcal{B}_\beta \subset \left[0, 1 - \frac{1}{\beta}\right) \setminus \bigcup [t_L, t_R).$$

Note by Lemma 3.5 and Theorem 1.1 that $\mathcal{B}_\beta = \mathcal{E}_\beta \subset [0, 1 - 1/\beta]$. In fact we have $\mathcal{E}_\beta \subset [0, 1 - 1/\beta)$. Observe that $b(1 - 1/\beta, \beta) = 01^m 0^\infty$. Then $T_\beta^{m+1}(1 - 1/\beta) < 1 - 1/\beta$. By Proposition 2.3 (i) this implies $1 - 1/\beta \notin \mathcal{E}_\beta$. Hence, $\mathcal{E}_\beta \subset [0, 1 - 1/\beta)$.

In the following it remains to prove $\mathcal{B}_\beta \cap \bigcup [t_L, t_R) = \emptyset$. Take a β -Lyndon interval $[t_L, t_R)$. If $t \in [t_L, t_R)$, then by (3.2) it follows that

$$\tilde{K}_\beta(t) = \tilde{K}_\beta(t_L) = \tilde{K}_\beta(t_R),$$

which gives $\eta_\beta(t') = \eta_\beta(t) = \eta_\beta(t_L)$ for all $t' \in (t, t_R)$. So, $t \notin \mathcal{B}_\beta$. \square

As a consequence of Proposition 3.1 and Theorem 1.1 it follows that for $\beta \in (1, 2]$ a multinacci number the β -Lyndon intervals cover $[0, 1 - 1/\beta)$ up to a Lebesgue null set.

Corollary 3.7. *Let $\beta \in (1, 2]$ be a multinacci number.*

- (i) *The union of all β -Lyndon intervals covers $[0, 1 - 1/\beta)$ up to a Lebesgue null set. Furthermore, for any $t \in \mathcal{B}_\beta$ and any $r > 0$ the interval $(t, t + r)$ contains infinitely many β -Lyndon intervals.*
- (ii) *$\eta_\beta(t) > 0$ if and only if $t < 1 - 1/\beta$.*

Proof. Note that \mathcal{E}_β is a Lebesgue null set which, by Theorem 1.1, has no isolated points. Then (i) follows from Proposition 3.1 which tells us that $\bigcup [t_L, t_R) = [0, 1 - 1/\beta) \setminus \mathcal{E}_\beta$. For (ii) it can be deduced from Proposition 3.1 and Theorem 1.1 that $\sup \mathcal{B}_\beta = 1 - 1/\beta$ and $1 - 1/\beta \notin \mathcal{B}_\beta$. \square

Now we turn to investigate the local dimension of the bifurcation set \mathcal{B}_β .

Lemma 3.8. *Let $\beta \in (1, 2]$ with $\delta(\beta) = (1^m 0)^\infty$. Then*

$$\lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t, t + r)) = \dim_H K_\beta(t) > 0 \quad \forall t \in \mathcal{B}_\beta.$$

Proof. Take $t \in \mathcal{B}_\beta$. By Proposition 3.1 we have $t < 1 - 1/\beta$, and then by Corollary 3.7 (ii) it gives $\eta_\beta(t) = \dim_H K_\beta(t) > 0$. Note by Proposition 3.1 and Proposition 2.3 (i) that

$$\mathcal{B}_\beta \cap (t, t+r) = \mathcal{E}_\beta \cap (t, t+r) \subseteq K_\beta(t) \quad \text{for any } r > 0.$$

Then $\lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t, t+r)) \leq \eta_\beta(t)$. So it remains to prove

$$(3.11) \quad \lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t, t+r)) \geq \eta_\beta(t).$$

We prove this now by considering the following two cases: (I) $t = t_R$ is the right endpoint of a β -Lyndon interval; (II) $t \in [0, 1 - 1/\beta) \setminus \bigcup [t_L, t_R]$.

Case (I). Suppose $t = t_R$ is the right endpoint of a β -Lyndon interval. Let $(t_i) = (t_1 \dots t_p)^\infty$ be the greedy β -expansion of t_R . Note that $t_R \in \mathcal{B}_\beta$. Then by Corollary 3.7 (i) there exists a sequence $(t_R^{(n)}) \subset \mathcal{B}_\beta$ such that each $t_R^{(n)}$ is a right endpoint of a β -Lyndon interval and $t_R^{(n)} \searrow t_R$ as $n \rightarrow \infty$. Fix $r > 0$. Then we can find a large integer N satisfying

$$t_R^{(n)} \in (t_R, t_R + r) \quad \text{for all } n \geq N.$$

Furthermore, since $b(t_R, \beta) = (t_1 \dots t_p)^\infty$, by Lemma 2.1 (ii) it follows that for each $n \geq N$ there exists an integer k_n such that the greedy β -expansion $b(t_R^{(n)}, \beta)$ of $t_R^{(n)}$ satisfies

$$(3.12) \quad b(t_R^{(n)}, \beta) \succ (t_1 \dots t_p)^{k_n} 1^\infty.$$

Observe by Proposition 3.1 and Proposition 2.3 (i) that

$$\mathcal{B}_\beta = \mathcal{E}_\beta = \{((s_i))_\beta : (s_i) \preceq \sigma^n((s_i)) \prec (1^m 0)^\infty \forall n \geq 0\}.$$

So by using $t_R \in \mathcal{B}_\beta$, (3.12) and Lemma 2.1 (ii) it follows that for any $n \geq N$,

$$(3.13) \quad \begin{aligned} & \left\{ ((t_1 \dots t_p)^{k_n} x_1 x_2 \dots)_\beta : x_1 \dots x_p = t_1 \dots t_p, (x_i) \in \tilde{K}_\beta(t_R^{(n)}) \right\} \\ & \subseteq \mathcal{B}_\beta \cap [t_R, t_R^{(n)}) \\ & \subseteq \mathcal{B}_\beta \cap [t_R, t_R + r). \end{aligned}$$

Note by Lemma 3.2 that $\tilde{K}_\beta(t_R^{(n)})$ is a transitive subshift of finite type. Then by (3.13) it follows that

$$\dim_H(\mathcal{B}_\beta \cap (t_R, t_R + r)) \geq \dim_H K_\beta(t_R^{(n)}) = \eta_\beta(t_R^{(n)}) \quad \text{for all } n \geq N.$$

Letting $n \rightarrow \infty$ and by the continuity of η_β (see Proposition 2.3 (ii)) we obtain that

$$\dim_H(\mathcal{B}_\beta \cap (t_R, t_R + r)) \geq \eta_\beta(t_R).$$

Since $r > 0$ was given arbitrary, letting $r \rightarrow 0$ we conclude that

$$(3.14) \quad \lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t_R, t_R + r)) \geq \eta_\beta(t_R).$$

Case (II). $t \in [0, 1 - 1/\beta) \setminus \bigcup [t_L, t_R]$. Then by Corollary 3.7 (i) there exists a sequence $(t_R^{(k)})$ such that each $t_R^{(k)}$ is the right endpoint of a β -Lyndon interval, and $t_R^{(k)} \searrow t$ as $k \rightarrow \infty$. So, for any $r > 0$ there exists a sufficiently large integer k such that $t_R^{(k)} \in (t, t+r)$. By (3.14) with t_R replaced by $t_R^{(k)}$ it follows that for any $\varepsilon > 0$ there exists $r_k > 0$ such that $(t_R^{(k)}, t_R^{(k)} + r_k) \subset (t, t+r)$ and

$$\dim_H(\mathcal{B}_\beta \cap (t, t+r)) \geq \dim_H(\mathcal{B}_\beta \cap (t_R^{(k)}, t_R^{(k)} + r_k)) \geq \eta_\beta(t_R^{(k)}) - \varepsilon.$$

Letting $r \rightarrow 0$, and then $t_R^{(k)} \rightarrow t$, we conclude by the continuity of η_β that

$$\lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t, t+r)) \geq \eta_\beta(t) - \varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, we obtain $\lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t, t+r)) \geq \eta_\beta(t)$. This, together with (3.14), proves (3.11). \square

Proof of Theorem 1. Let $\beta \in (1, 2)$ with $\delta(\beta) = (1^m 0)^\infty$. By Lemma 2.2, Proposition 3.1 and Lemma 3.8 it suffices to prove

$$(3.15) \quad \left\{ t \in [0, 1) : \lim_{r \rightarrow 0} \dim_H(\mathcal{B}_\beta \cap (t, t+r)) = \eta_\beta(t) > 0 \right\} \subset \mathcal{B}_\beta.$$

Take $t \in [0, 1) \setminus \mathcal{B}_\beta$. Then by Proposition 3.1 we have $t \in [1 - 1/\beta, 1)$ or $t \in [t_L, t_R)$ for some β -Lyndon interval. If $t \geq 1 - 1/\beta$, then $\eta_\beta(t) = 0$ by Corollary 3.7 (ii). If $t \in [t_L, t_R)$, then by Proposition 3.1 there exists $r > 0$ such that $\mathcal{B}_\beta \cap (t, t+r) = \emptyset$. This completes the proof. \square

Proof of Corollary 2. Note by Proposition 3.1 that $\mathcal{E}_\beta \subset [0, 1 - 1/\beta)$. So if $t \geq 1 - 1/\beta$, then clearly the result holds by Corollary 3.7 (ii). Now let $t \in [0, 1 - 1/\beta)$. Observe by Proposition 2.3 (i) that $\mathcal{E}_\beta \cap [t, 1] \subset K_\beta(t)$. So it suffices to prove

$$(3.16) \quad \dim_H(\mathcal{E}_\beta \cap [t, 1]) \geq \dim_H K_\beta(t).$$

If $t \in [0, 1 - 1/\beta) \setminus [t_L, t_R)$, then (3.16) follows by Lemma 3.8. If $t \in [t_L, t_R)$, then we still have (3.16) by using Lemma 3.8 that

$$\dim_H(\mathcal{E}_\beta \cap [t, 1]) \geq \dim_H(\mathcal{E}_\beta \cap [t_R, 1]) \geq \dim_H K_\beta(t_R) = \dim_H K_\beta(t),$$

where the last equality holds by (3.2). \square

4. FINAL REMARKS

The main results obtained in this paper can be easily modified to study the following analogous bifurcation sets:

$$\begin{aligned} \mathcal{E}'_\beta &:= \{t \in [0, 1) : K_\beta(t') \neq K_\beta(t) \ \forall t' \neq t\}, \\ \mathcal{B}'_\beta &:= \{t \in [0, 1) : \dim_H K_\beta(t') \neq \dim_H K_\beta(t) \ \forall t' \neq t\}. \end{aligned}$$

If $\beta \in (1, 2]$ is a multinacci number, one can show that

$$\begin{aligned} \mathcal{B}'_\beta &= \mathcal{E}'_\beta = \left[0, 1 - \frac{1}{\beta}\right) \setminus \bigcup [t_L, t_R] \\ &= \left\{ t \in [0, 1) : \lim_{r \rightarrow 0} \dim_H(\mathcal{E}_\beta \cap (t-r, t)) = \lim_{r \rightarrow 0} \dim_H(\mathcal{E}_\beta \cap (t, t+r)) = \dim_H K_\beta(t) > 0 \right\}, \end{aligned}$$

where the union is taken over all pairwise disjoint closed β -Lyndon intervals.

Observe that the main result Theorem 1 holds under the assumption that $\beta \in (1, 2]$ is a multinacci number, i.e., $\delta(\beta) = (1^m 0)^\infty$ for some $m \in \mathbb{N}$. The method used in this paper can be adapted to show that Theorem 1 still holds for $\beta \in (1, 2]$ with $\delta(\beta) = (10^m)^\infty$. It is worth mentioning that in [6] Kalle et al. considered a general Farey word base β , i.e., $\delta(\beta) = (s_1 \dots s_p)^\infty$ with $s_p s_{p-1} \dots s_2 s_1$ a non-degenerate Farey word. They showed that for a general Farey word base $\beta \in (1, 2)$, the set-valued bifurcation set \mathcal{E}_β has no isolated points and Theorem 1.1 holds. We finish by posing the following conjecture.

Conjecture 4.1. Let $\beta \in (1, 2]$. Then $\mathcal{B}_\beta = \mathcal{E}_\beta$ if and only if \mathcal{E}_β has no isolated points.

ACKNOWLEDGEMENTS

The authors were supported by an LMS Scheme 4 grant. The first author was supported by EPSRC grant EP/M001903/1. The second author was supported by NSFC No. 11971079 and the Fundamental and Frontier Research Project of Chongqing No. cstc2019jcyj-msxmX0338. He wishes to thank the Mathematical Institute of Leiden University.

REFERENCES

- [1] C. Carminati and G. Tiozzo. The local Hölder exponent for the dimension of invariant subsets of the circle. *Ergodic Theory Dynam. Systems*, 37(6):1825–1840, 2017.
- [2] L. Clark. The β -transformation with a hole. *Discrete Contin. Dyn. Syst.*, 36(3):1249–1269, 2016.
- [3] Z. Daróczy and I. Kátai. Univoque sequences. *Publ. Math. Debrecen*, 42(3-4):397–407, 1993.
- [4] M. de Vries and V. Komornik. Unique expansions of real numbers. *Adv. Math.*, 221(2):390–427, 2009.
- [5] G. Fuhrmann, M. Gröger, and A. Passeggi. The bifurcation set as a topological invariant for one-dimensional dynamics. *arXiv:1903.05172*.
- [6] C. Kalle, D. Kong, N. Langeveld, and W. Li. The β -transformation with a hole at 0. *arXiv:1803.07338*. To appear in *Ergodic Theory and Dynamical Systems*.
- [7] B. Li, T. Sahlsten, T. Samuel, and W. Steiner. Denseness of intermediate β -shifts of finite-type. *Proc. Amer. Math. Soc.*, 147(5):2045–2055, 2019.
- [8] D. Lind and B. Marcus. *An introduction to symbolic dynamics and coding*. Cambridge University Press, Cambridge, 1995.
- [9] W. Parry. On the β -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.*, 11:401–416, 1960.
- [10] P. Raith. Hausdorff dimension for piecewise monotonic maps. *Studia Math.*, 94(1):17–33, 1989.
- [11] A. Rényi. Representations for real numbers and their ergodic properties. *Acta Math. Acad. Sci. Hungar.*, 8:477–493, 1957.
- [12] J. Schmeling. Symbolic dynamics for β -shifts and self-normal numbers. *Ergodic Theory Dynam. Systems*, 17(3):675–694, 1997.
- [13] N. Sidorov. Almost every number has a continuum of β -expansions. *Amer. Math. Monthly*, 110(9):838–842, 2003.
- [14] M. Urbański. On Hausdorff dimension of invariant sets for expanding maps of a circle. *Ergodic Theory Dynam. Systems*, 6(2):295–309, 1986.
- [15] M. Urbański. Invariant subsets of expanding mappings of the circle. *Ergodic Theory Dynam. Systems*, 7(4):627–645, 1987.

(S. Baker) [SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, BIRMINGHAM, B15 2TT, UK](#)

Email address: simonbaker412@gmail.com

(D. Kong) [COLLEGE OF MATHEMATICS AND STATISTICS, CHONGQING UNIVERSITY, 401331, CHONGQING, P.R.CHINA](#)

Email address: derongkong@126.com