



International Journal of Phytomedicine 9 (2017) 655-672

<http://www.arjournals.org/index.php/ijpm/index>

Original Research Article



ISSN: 0975-0185

Phytochemistry investigation and antioxidant activity of four edible Verbenaceae of Burkina Faso.

Bangou Mindiédba Jean^{1*}, Reyes-Martínez Alfonso², Couliadiati Tangbadioa Hervé³, Megalizzi Véronique⁴, Nacoulma Odile Germaine⁵

*Corresponding author:

Bangou Mindiédba Jean

¹Laboratoire de Biochimie et de Chimie Appliquées (LABIOCA), Unité de Formation et de Recherche en Sciences et Techniques. Université Ouaga I, 09 BP 848 Ouagadougou 09, Burkina Faso.

²Instituto de Silvicultura e Industria de la Madera, Universidad Juárez del Estado de Durango, Boulevard del Guadiana 501, Fraccionamiento Ciudad Universitaria, 34160, Durango, Dgo., México.

³Section SVT, UFR/Sciences et Technologies, Université de Koudougou, 01 BP 376 Koudougou 01.

⁴Université Libre de Bruxelles. Faculté de Pharmacie, Service de Chimie Analytique Instrumentale et Bio Electrochimie et de Toxicologie cancéreuse Expérimentale. Campus de la Plaine-205/6. Boulevard du Triomphe B-1050 Bruxelles – Belgium.

⁵Laboratoire de Biochimie et de Chimie Appliquées (LABIOCA), Unité de Formation et de Recherche en Sciences et Techniques. Université Ouaga I, 09 BP 848 Ouagadougou 09, Burkina Faso

Abstract

Lantana rhodesiensis Moldenke, *Lippia chevalieri* Moldenke, *Vitex diversifolia* Bak and *Vitex doniana* Sweet are four species of Verbenaceae family widely used in traditional medicines in Burkina. The aim of this investigation was to study the chromatographic profile of phenols acids and that of the flavonoids as well as the antioxidants activities. Then, three types of extraction were made: acetonic (100%), decoction, ethanol-water(80/20 + 20/80); and fractions of ethanol-water which are hexane fraction (HF), ethyl acetate fraction (EAF), butanol fraction (BF) and aqueous fraction (AqF) of each species. These evaluations were supplemented by the quantification of the metal ions and the characterization by atomic absorption and HPLC-DAD methods of total phenols acids and total flavonoids. The whole of these proportioning was directed against the antioxidant activity through the anti-DPPH* (free radical scavenging activity), anti-TAC (total antioxidant capacity) and anti-IRP (iron reducing power). 36 flavonoids on 41 were highlighted by the HPLC-DAD method. A derivative glycosylate of quercetin was detected in the *V. diversifolia* extracts. *L. chevalieri* presented the strongest value in terms of ions content with a 55.88 g/kg of calcium. In terms of antioxidant activity, *L. rhodesiensis* and *L. chevalieri* presented the best activities on total antioxidant activity and anti-DPPH, and iron reducing power respectively. It's arise that decoction extract have the best's activities and can justify the traditional uses.

Keywords: chromatographic profile, antioxidant, phenols, flavonoids, metals, spectrometry.

Received: 21 Sep 2017

Accepted: 01 Dec 2017

Published: 28 Dec 2017

Introduction

Verbenaceae family contains around the world 100 genus and 2600 species. It's represented in intertropical Africa by 100 species

divided into 10 genus [1],[2]. Species from three genus of this family such as *Lippia*, *Lantana* and *Vitex* were the subject of our study. *L. rhodesiensis*, *L. chevalieri*, *V. diversifolia* and *V. doniana* are widely used traditionally in the management of several diseases including cancer, inflammatory diseases, arterial

DOI:10.5138/09750185.2192



This article is distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

hypertension, arterial high blood pressure, tumors, liver pathologies, nervous hepatitis, tooth aches, cutaneous diseases, intestinal bilharziose, bucco-anal and digestive candidiasis (Table 1). These plants are as well-known and used in traditional medicines in Burkina Faso [3],[4],[5],[6]. Phytochemistry and biological screening of these plants form integral part is our research plan. Several studies were undertaken on these plants: polyphenolic compound contents and chromatographic profiles of phenols acids and flavonoids [1],[4],[7], enzymatic activity [4], [8], antioxidant and antibacterial activities [1],[7],[9],[10]. The most elucidated studies on these plants were directed towards their essential oils contents [8],[11],[12],[13]. According to Nwachukwu and Uzoeto [14] acetone, ethanol, methanol, hot and cold water extracts of leaves of *Vitex doniana* have antimicrobial properties which can be pharmaceutically exploited on *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Bacillus subtilis* and *Staphylococcus aureus*. The antinociceptive activity of the ethanolic extract of *V. doniana* has been evaluated [13]. The extract demonstrated

significant antinociceptive activity in a dose dependent manner compared to control [15]. Recently, *Vitex diversifolia* exhibited the highest antibacterial activity against clinically isolated human pathogenic bacteria [16]. One of our prospects was to evaluate the polyphenolic compound contents and biological activity of these plants. Thus three types of extracts were used in this present study and the best extracts (ethanol-water) was to be selected for the sifting of the phenols acids, flavonoids and the evaluation of their antioxidant potential.

We also aimed to evaluate the antioxidant activity of these different extracts and fractions against free radical scavenging activity (DPPH*), total antioxidant capacity and iron reducing power. Specifically, in total extracts and fractions we have, (1) measured the phenolic and flavonoid compositions, (2) estimated their metals content, (3) studied their chromatographic profile through phenols acids and flavonoids (4) evaluated their antioxidants activities through three methods.

Table 1: Traditional uses of four (04) medicinal plant species (Verbenaceae)

Species	Traditional uses	Part used	References
<i>Lantana rhodesiensis</i> Moldenke	Arterial hypertension, cancer, diabetes mellitus arrhythmia, parasitic diseases, congestive heart failure, anti-proliferative, malaria, smallpox, lithiases	Fruits, leaves, stem-leaves	[3] [4] [8] [37] [58]
<i>Lippia chevalieri</i> Moldenke	Malaria, diarrheas, arterial high blood pressure, pathologies of the liver, bucco-anal and digestive candidiasis, menorrhagia of the menopause, nervous hepatitis, gout and the mental disorders	Stem-leaves, leaves,	[4] [5] [11] [25] [59]
<i>Vitex diversifolia</i> Bak	Antibacterial, tooth aches, cutaneous diseases, intestinal bilharziose, weaning, anti-inflammatory, arterial hypertension, Astringents	Fruits, bark, leaves, Stem-leaves	[6] [60] [61] [62]
<i>Vitex doniana</i> Sweet	Affection of respiratory tracks, tumors, dysentery, diarrhea, painful rules, chicken pox, small pox, evil of chronic stomach, madness, rheumatic pains, inflammatory disorders, Nutritive Sweetener	Fruits, bark, leaves, stem bark	[9] [12] [63] [64] [65] [66]

Experimental

Plant material

Stems leaves of *L. rhodesiensis* and *L. chevalieri* were collected to Boulbi (Saponé), leaves of *V. diversifolia* collected of Komadougou (11 Km North of Fada N'Gourma) and *V. doniana* (leaves) collected of Gampéla (25 Km East from Ouagadougou). All these were collected in Burkina Faso between May and October 2013. The plants were botanically identified by Professor Millogo-Rasolodimby from the Department of Plant Biology at the University

of Ouagadougou. The Voucher specimens were deposited in the OUA herbarium of the CIB (Centre d'Information sur la Biodiversité), UFR/SVT of the University of Ouagadougou.

Reagents and standards

Acetonitrile (HPLC grade), water (HPLC grade), ethanol (HPLC grade), ethyl acetate (analytical grade), hexane (analytical grade), sulfuric acid (analytical grade), and sodium phosphate were purchased from J. T. Baker (Xalostoc, Mexico). 2,2-Diphenyl-1-



picrylhydrazyl (DPPH^{*}), aluminum chloride, ammonium molybdate, and the references: quercetin, quercitrin (quercetin-3-rhamnoside), caffeic acid, and ascorbic acid were purchased from Sigma-Aldrich (St. Louis Missouri, USA). The standards kaempferol-3, 7-*O*-diglucoside, quercetin-3-*O*-[rhamnosyl-(1-6)-galactoside], kaempferol-3-*O*-[rhamnosyl-(1-6)-glucoside] came from Apin Chemicals Limited (Abingdon, Oxon, UK). Trichloroacetic acid and ferric chloride were obtained from Merck (Darmstadt, Germany). Potassium ferricyanide was purchased from Fermont (Monterrey, Mexico).

Preparation of extracts

Polyphenols and flavonoids were extracted from 4 g of dry powder from leaves, bark or stem-leaves were extracted by maceration in 40 mL of acetone 100% (acetone extract), ethanol 80% (v/v), or by decoction (decoction extract), for 24 hours in darkness, and at room temperature. The extracts were centrifuged (5000 rpm) for 10 min, at room temperature, and the supernatants separated. The pellets (only for ethanol 80% extract) were re-extracted in 100 mL of 20% ethanol (v/v) for 3 hours, centrifuged under the same conditions, and the supernatant decanted. Both supernatant were combined to form the total extracts (ethanol-water extract). The total extracts were concentrated to dry for polyphenolic, flavonoids and antioxidants activities. The ethanol-water extract was concentrated to half the volume and then fractioned twice with ethyl acetate (EAF). The pellets were also macerated in 100 ml hexane for 24 hours in darkness (HF). The two organic fractions (ethyl acetate and hexane), the aqueous fractions (AqF), and the separated aliquots of the total extracts, were individually concentrated under vacuum to dryness and then re-dissolved in 5 mL ethanol, aliquots were taken to be used in the determination of flavonoid content, in the HPLC-DAD analysis. The phenolic extractions were prepared from three different pools of samples. The ethanol-water extract was also subjected to sequential liquid-liquid extraction with n-hexane (HF), ethyl acetate (EAF), and n-butanol (n-BF).

Determination of phenolic content

Folin-Ciocalteu method was used for measurement of total content of phenolic compounds according to Nurmi *et al.* [17], by linear regression analysis from the standard curve of gallic acid ($y = 0.003x + 0.016$; $r = 0.997$). A 250 μ L extract was mixed with 2.5 ml of desionized water. Afterwards, 125 μ L Folin-Ciocalteu reagents were added and the mixture was allowed to stand for 5 min. Finally 375 μ L of 20% Na₂CO₃ was added. After 2 hours incubation at room temperature, the absorbance was measured at 760 nm on a Spectroscopic Analysis Mecasys (Optizen). The phenol contents were expressed as mg gallic acid equivalents g⁻¹ dry tissue (mg GAE g⁻¹ dry tissue). Three replicates of each sample were analyzed.

Determination of total flavonoid content

Flavonoid content was determined according to Lauranson-Broyer and Lebreton [18] by linear regression analysis from the following standard curve of quercetin: $Abs_{425nm} = 0.025x + 0.014$ [Quercetin], correlation coefficient $r = 0.998$. The curve was registered after the addition of 60 μ L of a freshly prepared 5% (w/v) aluminum chloride solution to 1 mL of quercetin solution (four different concentrations in the range of 100 to 1400 μ g/mL). The absorbance was immediately registered after the addition of aluminum chloride, at 425 nm, using a Spectronic Genesys 2 espectrophotometer (Rochester, New York, USA). The flavonoid content in each sample was also registered after the addition of aluminum chloride and the flavonoid content in each sample was expressed as mg of quercetin equivalents g⁻¹ dry tissue (mg QE g⁻¹ dry tissue). The addition of aluminum chloride produces bathochromic shifts (which can be perceived by a yellow coloration) in flavonoids containing orthodihydroxyl groups, due to the formation of complexes between the aluminium and C-4 keto group and either the C-3 or C-5 hydroxyl group of flavones and flavonols; complexes also are formed between the aluminum and the orthodihydroxyl groups in A or B-ring of flavonoids [19]. The addition of aluminium chloride represents a standard procedure for reproducibility [18]. The determination of flavonoid content was estimated individually for three pools of samples.

HPLC-DAD Analysis

To determine the flavonoid profiles, aliquots of each extract (concentrated to dryness and re-dissolved in 5 mL ethanol, as mentioned in the section Preparation of extracts) were analyzed as previously described [20] on a Perkin Elmer Series 200 HPLC system (Shelton, Connecticut, USA) and a Perkin Elmer Brownlee Analytical C18 column (4.6 x 250 mm, 5 μ m) (Shelton, Connecticut, USA), by an acidified acetonitrile-water gradient. Water adjusted to pH 2.5 with orthophosphoric acid was the solvent A, and acetonitrile was the solvent B, mixed according to the following gradient; starting with 100% A, decreasing to 91% over the next 12 mn, to 87% over next 8 mn, to 67% over the next 12 mn, to 57% over the next 10 mn, and held at this level until the end of the 60 mn analysis. Standard chromatograms were plotted at 260 and 340 nm. Spectral data for all peaks were accumulated in the range 220-400 nm using diode array detection (Perkin Elmer Series 200). Structural identification was obtained by direct comparisons of retention times and UV spectra of resolved compounds with those of standard; two kaempferol glycosides, identified as kaempferol-3,7-*O*-diglucoside and kaempferol-3-*O*-[rhamnosyl-(1-6)-glucoside], and one quercetin glycoside, identified as quercetin-3-*O*-[rhamnosyl-(1-6)-galactoside]. The structural information of compounds, for which standards were not available, was obtained from their spectral parameters according to the compilations of Mabry, Markhan and Thomas and Campos and Markham [19], [20].

Quantitative determinations were made by an external standard method, with the commercial reference (quercitrin), by area measurements, using the following standard curve: $\text{Area} = -0.0046x + 0.0278$ [Quercitrin], correlation coefficient $r = 0.992$. The content of each compound was expressed as μg of quercitrin equivalents g^{-1} dry tissue (g QiE g^{-1} dry tissue). The HPLC-DAD profiles were individually obtained and analyzed for the extracts from three pools of samples.

Total Antioxidant Capacity

The total antioxidant capacity (TAC) of each samples was evaluated through the method developed by Prieto *et al.* [21], in which the reduction of Mo (VI) to Mo (V) is carried out by the antioxidant, forming a green phosphate/Mo (V) complex at acidic pH. Aliquots (100 μL) of each sample (containing 100 $\mu\text{g}\cdot\text{mL}^{-1}$ of flavonols, respective concentrations of flavonols calculated from the standard curve of quercetin) were prepared and combined with 1 mL of a solution constituted of sulfuric acid (0.6 M), sodium phosphate (28 mM), ammonium molybdate (4mM) and incubated at 95 C for 90 min. After reaching room temperature, the absorbance of each sample was registered at 695 nm against a blank prepared as indicated for the samples, but adding ethanol instead of the sample. The reference quercetin was analyzed in the same manner. TAC was expressed as mg ascorbic acid equivalents mL^{-1} (mg AAE mL^{-1}). Ascorbic acid curve: $A_{695} = 3.678x - 0.092$ [ascorbic acid], correlation coefficient $r = 0,998$, constructed with ascorbic acid between 1.0 and 30.0 $\text{mg}\cdot\text{mL}^{-1}$. The analysis was done for independence aliquots of the samples from three pools of samples.

Free radical scavenging activity

The DPPH* method reported by Campos *et al.* [22] was used to evaluate the free radical scavenging activity. Four to five flavonoid concentrations of each sample were individually added to a DPPH* solution (40 $\mu\text{g}\cdot\text{mL}^{-1}$ in ethanol) in such a way so as to maintain a final volume of 1 mL. The decrease in absorbance was determined at 523 nm after 10 min. The DPPH* concentrations of samples were plotted to determine by linear regression, the efficient concentration at 50 %, defined as the amount of antioxidant needed to decrease by 50 % the initial DPPH* concentration (EC_{50}). The following calibration curve, made with DPPH* between 6.25 and 100 $\mu\text{g}\cdot\text{mL}^{-1}$, was used to calculate the DPPH* concentration ($\mu\text{g}\cdot\text{mL}^{-1}$) in the reaction medium: $A_{523} = 0,030x + 0,001$ [DPPH*], correlation coefficient $r = 0,999$. Antioxidant activities were expressed in terms of EC_{50} in $\mu\text{g}\cdot\text{mL}^{-1}$. The analysis was separately done for the samples from three pool samples.

Iron Reducing Power

The iron reducing power (IRP) method reported by Yanget *al.* [23] was used to evaluate the iron reducing power of each simple. Aliquots (1 mL) of each sample were combined with 2.5 mL (phosphate buffer, 0.2 M, pH 6.6), 2.5 mL (potassium ferricyanide, 30mM) and incubated at 50 C for 20 min. After, 2.5 mL trichloroacetic acid (0.6 M) was added and the mixture was centrifuged (2000 rpm for 10 min). From the upper layer, 2.5 mL of solution was removed and distilled water (2.5 mL) and ferric chloride (0.5 mL, 6 mM) were added to it. The absorbance at 700 nm of the formation of ferrous ions (Fe^{2+}) was registered after 10 min. The highest absorbance values indicated the greatest capacity of reducing ferric (Fe^{3+}) to ferrous (Fe^{2+}) ions. Four flavonol concentrations (10-400 μL combined with the proper volume of ethanol to reach 1 mL as final volume) of each sample (respective concentrations of flavonols calculated from standard curve of quercetin) were evaluated. The reducing power was expressed in terms of EC_{50} ($\text{mg}\cdot\text{mL}^{-1}$). The evaluation was separately done for the samples from three pools of samples.

Atomic absorption (*Samples preparation*)

Typical analytical procedure of 975.03 to AOAC was used adapted by Perkin-Elmer [24]. Take 1 g of ground dried plant sample and place it in a small beaker. Add 10 mL of concentrated HNO_3 and allow standing overnight. Heat carefully on a hot plate until the production of red NO_2 fumes has ceased. Cool the beaker and add a small amount (2 – 4 mL) of 70 % HClO_4 . Heat again and allow evaporating to a small volume. Transfer the sample to a 50 mL flask and dilute to volume with distilled water. Caution: always add HNO_3 to the tissue samples and allow this mixture to digest before adding HClO_4 . HClO_4 can react explosively with untreated organic matter.

Statistical analysis

All assays were carried out in triplicates and results are expressed as Means \pm Standard Deviation (SD) calculated with Excel 2007. Statistical comparisons were done with the XLSTAT7.5.2, using Spearman correlation. Differences were considered to be significant at $p < 0.05$.

Results and discussion

Biological investigation

Antioxidants activities

Several methods were developed to evaluate antioxidant potential of plants extracts. They are based on the determination of products resulting from oxidation or, on the contrary, measure the effectiveness of a substance to trap radicals, often by giving a form



H* [25].Recent studies showed that there can be differences in the determination of antioxidant activity according to the method used [25],[26]. Thus for better evaluating of antioxidant potential of plant extracts it is necessary to use at least two methods. In the present study we used three methods: total antioxidant capacity ($y = 3.6781x - 0.092$, $r = 0.9985$), iron reducing power ($y = 2.6486x + 0.0017$, $r = 0.9988$), free radical scavenging activity ($y = 0.0309x + 0.0019$, $r = 0.9996$) to evaluate the antioxidant activity of acetone, decoction, ethanol-water extracts and four fractions of ethanol-water of these plants.

Total Antioxidant Capacity

The results of the reduction of molybdate (VI) to molybdate (V) are presented in the Table 2. By taking account of the whole tests (total extracts and fractions), the results are included between 0.85 and 9.34 mg EAA/mL. In a general way the extracts of ethanol-water presented the best reducing activities with an average of 7 mg EAA/mL. More specifically the butanolic fraction of *V. diversifolia*

gave the best activity (9.34 Mg EAA/mL). The weakest activities are held by decoction extracts with an average of 0.9 mg EAA/mL. The low value being held by *L. chevalieri* (0.850 mg EAA/mL). At this level we observe a controversy correlation between polyphenolic compound content and molybdate reducing activity. Indeed sometimes a correlation seems existed such as polyphenolic compound content case of ethanol-water and decoctions extracts on the one hand and on the other hand with the reducing activities (Table 2). Often not correlation existing in a similar situation. In fact, with a total phenolic content of 44.74 mg EAG/g extracts (butanolic fraction of *V. diversifolia*), we observed a best activity on TAC than ethyl acetate fraction which gave 131.32 mg EAG/g extracts. These results corroborate with those of Barriada-Bernal *et al.* [27] who evaluated antioxidant activities against the total flavonoids contents. They don't found a clear correlation between total antioxidant capacity and the total flavonoids content. Indeed, 1210.48µg/g of total flavonoid content gave 4.646 mgEAA. However 120.48µg/g < 1210.48µg/g gave 6.113EAA > 4.646 mgEAA. The same results were obtained by Falleh *et al.* [28].

Table 2: Polyphenolic compounds contents and antioxidants activities of the plants

Species	Type of extracts/Fractions	Phenol contents (mg EAG/g dry extracts)	Flavonoid contents mg EQ/g dry extract)	Free radical scavenging activity EC ₅₀ (µg/mL)	Total antioxidant capacity (mg EAA/mL)	Iron reducing power EC ₅₀ (µg/mL)	
<i>L. rhodesiensis</i>	Acetone	24.85 ± 1.57 ^a	1.31 ± 0.09 ^b	2.31 ± 0.05 ^b	1.447 ± 0.047 ^b	0.24 ± 0.02 ^d	
		<i>L. chevalieri</i>	7.86 ± 0.42 ^c	0.90 ± 0.03 ^c	3.49 ± 0.08 ^a	1.337 ± 0.023 ^c	0.08 ± 0.00 ^f
		<i>V. diversifolia</i>	4.00 ± 1.03 ^d	1.89 ± 0.13 ^a	1.45 ± 0.02 ^d	1.277 ± 0.036 ^{cd}	0.10 ± 0.00 ^{ef}
		<i>V. doniana</i>	18.84 ± 1.34 ^b	2.08 ± 0.14 ^a	1.72 ± 0.04 ^c	1.496 ± 0.048 ^b	0.12 ± 0.00 ^e
<i>L. rhodesiensis</i>	Decoction	7.03 ± 0.46 ^c	0.48 ± 0.02 ^c	0.05 ± 0.00 ^d	1.049 ± 0.021 ^f	0.15 ± 0.00 ^d	
		<i>L. chevalieri</i>	16.10 ± 1.05 ^a	0.77 ± 0.02 ^b	1.30 ± 0.08 ^c	0.850 ± 0.008 ^g	0.06 ± 0.00 ^e
		<i>V. diversifolia</i>	12.14 ± 1.62 ^b	1.19 ± 0.03 ^a	8.12 ± 0.20 ^a	0.954 ± 0.028 ^f	0.07 ± 0.00 ^e
		<i>V. doniana</i>	18.70 ± 3.05 ^a	1.17 ± 0.06 ^a	2.95 ± 0.06 ^b	0.966 ± 0.004 ^f	0.07 ± 0.00 ^e
<i>L. rhodesiensis</i>	Ethanol-water	69.48 ± 8.07 ^a	3.27 ± 0.40 ^a	12.21 ± 0.95 ^b	8.936 ± 0.646 ^a	0.64 ± 0.01 ^c	
		<i>L. chevalieri</i>	19.48 ± 1.68 ^d	1.62 ± 0.33 ^b	13.93 ± 0.39 ^a	6.555 ± 0.096 ^c	0.23 ± 0.01 ^e
		<i>V. diversifolia</i>	33.48 ± 3.50 ^c	3.23 ± 0.43 ^a	6.07 ± 0.22 ^c	6.357 ± 0.122 ^c	0.31 ± 0.01 ^d
		<i>V. doniana</i>	54.14 ± 4.35 ^b	3.58 ± 0.20 ^a	5.04 ± 0.03 ^d	7.441 ± 0.267 ^b	0.28 ± 0.00 ^d
<i>(L. rhodesiensis)</i>	n-HF	91.76 ± 1.39 ^d	7.80 ± 0.04 ^g	0.33 ± 0.02 ^g	7.14 ± 0.18 ^e	0.35 ± 0.02 ⁱ	
	n-BF	90.52 ± 11.2 ^d	12.71 ± 0.02 ^d	0.57 ± 0.02 ^e	6.17 ± 0.15 ^f	2.21 ± 0.01 ^a	
	AqF	102.43 ± 14.55 ^{bc}	5.80 ± 0.04 ⁱ	1.21 ± 0.03 ^a	3.95 ± 0.01 ^j	0.89 ± 0.01 ^d	
	EAF	126.07 ± 10.74 ^a	25.71 ± 0.05 ^b	0.05 ± 0.00 ^j	7.34 ± 0.07 ^d	1.20 ± 0.03 ^b	
<i>(L. chevalieri)</i>	n-HF	40.20 ± 1.34 ^h	6.49 ± 0.08 ⁱ	0.11 ± 0.01 ⁱ	4.59 ± 0.15 ^h	0.09 ± 0.00 ^m	
	n-BF	73.00 ± 0.34 ^e	6.59 ± 0.12 ^j	0.28 ± 0.01 ^h	4.15 ± 0.05 ⁱ	0.26 ± 0.00 ^k	
	AqF	99.32 ± 0.56 ^{cd}	7.09 ± 0.06 ^h	ND	3.98 ± 0.03 ^j	0.35 ± 0.02 ⁱ	
	EAF	79.23 ± 1.48 ^e	4.96 ± 0.00 ^j	0.60 ± 0.02 ^d	3.98 ± 0.13 ^j	0.26 ± 0.00 ^k	
<i>(V. diversifolia)</i>	n-HF	41.803 ± 1.97 ^h	5.45 ± 0.18 ^{jk}	0.38 ± 0.00 ^f	2.86 ± 0.02 ^l	0.14 ± 0.01 ^l	
	n-BF	44.74 ± 2.17 ^{gh}	5.39 ± 0.42 ^k	0.99 ± 0.01 ^b	9.34 ± 0.12 ^a	0.30 ± 0.00 ^j	
	AqF	57.54 ± 3.64 ^{fg}	9.23 ± 0.24 ^{ef}	1.01 ± 0.04 ^b	7.53 ± 0.14 ^{cd}	0.36 ± 0.00 ⁱ	
	EAF	131.32 ± 4.43 ^a	35.25 ± 0.74 ^a	0.12 ± 0.01 ⁱ	8.61 ± 0.16 ^b	0.74 ± 0.02 ^f	



Table 2: Continued

Species	ETOH-H ₂ O fractions	Phenol contents (mg EAG/g dry extracts)	Flavonoid contents mg EQ/g dry extract)	Free radical scavenging activity EC ₅₀ (µg/mL)	Total antioxidant capacity (mg EAA/mL)	Iron reducing power IC ₅₀ (µg/mL)
<i>(V. doniana)</i>	n-HF	110.16 ± 9.51 ^b	8.04 ± 0.08 ^g	1.21 ± 0.02 ^a	3.92 ± 0.12 ^j	0.25 ± 0.00 ^k
	n-BF	60.12 ± 1.32 ^f	17.51 ± 0.05 ^c	0.34 ± 0.01 ^g	6.07 ± 0.13 ^f	0.59 ± 0.02 ^h
	AqF	10.87 ± 0.31 ⁱ	9.80 ± 0.11 ^e	0.72 ± 0.04 ^c	3.40 ± 0.01 ^k	0.36 ± 0.04 ⁱ
	EAf	93.54 ± 6.23 ^{cd}	17.81 ± 0.10 ^c	0.34 ± 0.01 ^g	5.23 ± 0.02 ^g	0.93 ± 0.00 ^c
(References)	Gallic acid				1.26 ± 0.03 ^o	0.92 ± 0.001 ^c
	Cafeic acid				1.73 ± 0.06 ^m	0.84 ± 0.01 ^e
	Ascorbic acid				1.78 ± 0.02 ^m	0.66 ± 0.03 ^g
	Quercetin				1.51 ± 0.01 ⁿ	0.65 ± 0.04 ^g

Freeradical scavenging activity

The anti-DPPH* activity of the vegetable extracts translates their aptitude to trap the free radicals of the organism. To this end we evaluated three types of extracts and four fractions coming from the ethanol-water total extract (Table 2). Results are evaluated in EC₅₀ and were between 0.05 (*L. rhodesiensis*) and 13.93 (*L. chevalieri*) EQ/g dry extracts. The EC₅₀ of a curve dose-effect represents the concentration of a compound where 50% of the maximum effect is observed. The antioxidant capacity of a compound is higher as its EC₅₀ is small. Among these results, the strongest value was observed with *L. rhodesiensis* in decoction extract and its ethyl acetate fraction (0.05 EQ/g dry extracts). These results are consigned in Table 2 and are similar to those obtained by Reyes-Martinez et al. [29]. These species also presented the strongest total phenolic and flavonoids contents in their fractions (ethyl acetate fraction). While looking at close the results related to the fractions, one notes that the ethyl acetate fractions presented the best anti-DPPH* activities. We couldn't establish a correlation of these activities with the total phenolic and flavonoid contents. Indeed on the level of total phenolic contents and ethanol-water extract, *L. rhodesiensis* which gave the strongest content (69.48 mg EAG/dry extract) also presented the second weaker DPPH* activity (12.21 µg/mL). Contrary to its decoction content which is 7.03 mg EAG/dry extract but it presented the best activity (0.05 EQ/g dry extracts). Our results are in disagreement with those of Barriada-Bernal et al. [27]. According to our former investigation [9], it was difficult to establish a clear correlation between the flavonoids rate and the DPPH* reduction. Several authors tried to justify this fact [30],[31],[32],[33],[34]. Confer relationship between antioxidants activities and polyphenolic compounds estimated part.

Iron Reducing Power

In the mechanism of phenolic antioxidant activity, the reduction of Fe (III) is often used like an indicator of donor of electrons [35]. Iron reducing power method makes it possible to measure the ability of phenolic compounds to reduce Fe (III) in Fe (II). The ascorbic acid was used as standard for the development of the calibration curve

whose regression equation is: $y = 2.6486x + 0.0017$; $r = 0.9988$. The results of this study are indicated in Table 2. Taken account of the three types of extraction the results give values ranging between 0.06 and 0.92 µg/mL. In terms of ions ferric reduction in ions ferrous the strongest values were obtained with *L. chevalieri* extracts with 0.06 µg/mL (decoction) > 0.08 µg/mL (acetone) > 0.23 µg/mL (ethanol-water). The low values are obtained with *L. rhodesiensis* with respectively 0.15 µg/mL (decoction) > 0.24 µg/mL (acetone) > 0.64 µg/mL (ethanol-water). According the results of this study there are not a correlation between total phenolic contents and ion ferric reducing activity. Indeed *L. rhodesiensis* which gave the best total phenolic content (69.48 mgEAG/g dry extracts) also presented the weakest ions ferric reduction (0.64 µg/mL). That is confirmed by the results obtained with the fractions. Firstly, fractions of *L. rhodesiensis* presented the lowest results (range between 0.35 and 2.21 µg/mL). However this plant gave the best total phenol contents of fraction with an average of 102.70 EAG/g dry extracts. Secondly, for 131.32 mg EAG/g dry extracts of total phenolic, one has a reducing activity of 0.74 µg/mL. On the other hand with a content of 10.87 mg EAG/g one obtained 0.36 µg/mL of ion ferric reduction. That corresponds to the higher half of the first activity. Finally, we note that ethyl acetate fraction gave the weakest results. Concerning total flavonoid content, we note a correlation between their contents and ion ferric reducing activity. These results are corroborated [27], [36]. While looking of close ions proportioning using atomic absorption method, we realize that among the four plants species, *L. chevalieri* presented the strongest content of ions ferrous (5.661 g/kg). This result is followed by that of *V. diversifolia* (1.145 g/kg). We can suppose that the ions ferrous reducing activity is correlated by the content of the plants.

Phytochemistry investigation

Phenolic and flavonoid contents

Three types of extraction were used in this study to evaluate phenolic and flavonoids contents. With the continuation we selected the type of extract having given the best contents of

polyphenolic to evaluate the fractions contents. Four types of fractions were evaluated such as hexanic fractions (n-HF), butanolic fractions (n-BF), ethyl acetate fractions (EAF) and aqueous fractions (AqF). Total extracts contents and their fractions were used for the evaluation of the antioxidant activities and the HPLC-DAD analysis.

In a general way, the results of phenolic and flavonoids quantification are decreasing in the order ethanol-water, acetone and decoction. The totals phenolic varied between 4.00 and 69.48 mg EAG/g dry extracts (Table 2). The best values are obtained with the ethanol-water extracts. At this level *L. rhodesiensis* holds the strong value (69.48 mg EAG/g), followed *V. doniana* (54.14 mg EAG/g). The low value is obtained with *L. chevalieri* (4.00 mg EAG/g) of acetone extracts. This value is followed of that of *L. rhodesiensis* (7.03 mg EAG/g) of decoction extracts.

In terms of total flavonoids content, the values vary between 0.48 and 3.58 mg EQ/g dry extracts. The strongest values are obtained with ethanol-water extracts of *V. doniana* = *V. diversifolia* = *L. rhodesiensis* (3.58 mg EQ/g). The weak values are held by the decoction extracts with 0.48 mg EQ/g for *L. rhodesiensis* and 0.77 mg EQ/g for *L. chevalieri*. However in terms of flavonoid contribution to total phenolics, the best contribution was observed with acetone extracts such as *V. diversifolia* with 47.25% > *V. doniana* = *L. rhodesiensis* (11.45%). These values are followed so far by the decoction extracts with 9.80%. *L. rhodesiensis* and *L. chevalieri* presented the best activities on total antioxidant activity, anti-DPPH*, and iron reducing power respectively. Former investigation of totals phenolic and flavonoid contents of *L. rhodesiensis* in methanolic extract showed 21.55 ± 0.75 mg GAE/100 mg and 5.09 ± 0.19 mg QE/100 mg extract, respectively [4]. Piero *et al.* [37] quantified in aqueous extracts of *L. rhodesiensis* (leaves) phenols content and find 685.25 ± 30.77 mg/g gallic acid equivalent, while flavonoids was 187.33 ± 54.97 mg/g. Our results are much lower than those of Piero *et al.* [37] who used aqueous extracts. However, they are better than those of Bangou *et al.* [4] who used methanolic extracts. Lagnika *et al.* [38] investigated antibacterial, antifungal, antioxidant activities and toxicity of *Vitex doniana* leaves against dichloromethane, methanol and hydroethanolic extracts. Their results supported the utilization of this plant in infectious diseases and also showed this plant as good sources for

antioxidants. Agbafor *et al.* [10] showed clearly that this best antioxidant activity of *Vitex doniana* is due to its vitamin C content.

Ions quantification

The analysis is related to eight minerals such as Ca²⁺, Fe²⁺, K⁺, Mg²⁺, Na⁺, Zn²⁺, Pb and As. No arsenic trace was found in our various plants. With the exception of lead and arsenic, we evaluated minerals for their implications of human physiological activity at low dose. The proportion of arsenic and lead wanted to be evaluating the risk of contamination of the grounds by the mining activities. The care by traditional medicine is generally related to the combinations of plants (phytotherapy), fragment of animals (opotherapy) and minerals (mineral therapy). Then, some oligo elements (Ca²⁺, Fe²⁺, K⁺, Mg²⁺, Na⁺, Zn²⁺) have an essential role in physiological balance and especially for the correct operation of the neurotransmitters [6],[25]. In general way their absences involve disturbances [6],[25]. Thus for example, magnesium deficiency will involve syndromes of neuromuscular hyperirritability, cardiovascular and thrombosis accidents([6],[25]). Its metabolism is also related to those of Ca²⁺ and P. In terms of ions quantification the values are included between 1.1x10⁻⁴ and 55.88 g/kg of dry weight. Calcium presented the best contents with 55.88 g/kg (*L. chevalieri*) > 23.36 g/kg (*L. rhodesiensis*) > 16.87 g/kg for *V. doniana* and *V. diversifolia*. Traces of lead are found on *Vitex* species. In terms of daily need for the ions highlighted, sodium comes at the head with 2g/day > 1.5 g/day (magnesium) and of potassium (0.5 g/day). Lead occupies the last place with 0.0005 g/day. Vunchi *et al.*[39], analyzed the proximate, vitamins and mineral compositions of *Vitex doniana* fruit pulp. They found importance values of minerals: potassium (16.5 mg/100g), sodium (10.4 mg/100g), calcium (30.27 mg/100g), iron (5.2 mg/100g), Copper (2.7 mg/100g), magnesium (20.10 mg/100g) and phosphorus (16.5 mg/100g). Confrontation of our results shows that the mineral values are higher in the fruits than in the leaves or stem-leaves (Table 3). Indeed, this study concluded that *Vitex doniana* fruit pulp could be promoted as: carbohydrate and lipid supplements for cereal-based diets in rural communities [39]. Sawadogo *et al.* [3] have quantified some metal ions (Na⁺ and K⁺) and showed their implication in the Na⁺-K⁺-pump action.

Table 3: Atomic absorption of seven oligo elements detected in the four edible plants

Species	Ca ²⁺	Fe ²⁺	K ⁺	Mg ²⁺	Na ⁺	Zn ²⁺	Pb
<i>L. rhodesiensis</i>	23.36 ± 1.92	0.51 ± 0.00	19.32 ± 0.56	3.23 ± 0.03	0.079 ± 0.001	0.035 ± 0.000	--
<i>L. chevalieri</i>	55.88 ± 1.38	5.66 ± 0.36	16.34 ± 0.79	6.76 ± 0.02	0.057 ± 0.001	0.05 ± 0.00	--
<i>V. diversifolia</i>	16.87 ± 0.05	1.145 ± 0.008	7.20 ± 0.41	5.32 ± 0.12	0.058 ± 0.001	0.037 ± 0.000	(1.1 ± 0.35)*10 ⁻⁴
<i>V. doniana</i>	16.58 ± 0.23	0.27 ± 0.00	10.11 ± 0.11	4.29 ± 0.11	0.06 ± 0.01	0.045 ± 0.001	(1.9 ± 0.43)*10 ⁻⁴

HPLC-DAD analysis

A total of 41 polyphenolic compounds (Figure 1: 1/7-7/7) was detected while excluding the aromatic acids from bases which

seem to be transverse to all plant extracts. The HPLC-DAD analysis related to the three types of extracts and fractions made on the basis of ethanol-water extract which presented the best contents of phenolic and flavonoids. The ethanol-water extract

presented more compounds (12 compounds). Only 8 compounds were detected on the fractions levels. Luteolin was highlighted in the two species of *Vitex* and on the whole types of extracts. In the same way a derivative of quercetin-3-O-arabioside (Dvidi6 and Avidi28) was highlighted in the decoction extract and in the acetone extract of *V. diversifolia* (Table 4)

Former studies showed that flavonoids are characterized by two bands maximum of absorbance of cycles A and B [1],[19]. More specifically flavones and flavonols are characterized by a band I (cycles A) in the areas 320 - 385 nm and a band II (cycles B) in interval 250 - 285 nm (Mabry *et al.* 1970). Band I of flavones being increasingly shorter from 20 to 30 nm with that of equivalent flavonols. Those of the phenols acids lie between 210 - 273 nm. Any increase in the number of hydroxyls grouping and/or glycosylation and O-methylation inducing bathochromic (displacement of maximum wavelengths towards the right) and hypsochromic (displacement of maximum wavelengths towards the left) effects, respectively [1], [19].

These observations associated at the protocol with elucidation [20] (Structure information from HPLC and on-line measured absorption spectra: Flavones, Flavonols and Phenolic Acids) enable us to distinguish 05 phenolsacids and 36 flavonoids (Table 4 & Figure 1: 1/7-7/7). Former investigations of the methanolic extracts of *L. chevalieri* highlighted 20 composed which have primarily acids phenols [1].

Thus we couldn't find the derivative luteolin and rutin [1] in our extracts. Other researchers showed in the dichloromethane fractions of *L. rhodesiensis*, two flavonoids such as 5,6,7,3',4',5'-hexamethoxyflavone and its analogue, 5-hydroxy-6,7,3',4',5'-pentamethoxyflavone [3]. The most elucidated studies on *L. chevalieri*, *L. rhodesiensis*, *V. diversifolia* and *V. doniana* especially related to their content of essential oils [1],[3], [5],[8].

Which could be the implication of these compounds highlighted in the biological activities observed?

Table 4: Wavelength characteristic of the phenols acids and flavonoids detected (1/2).

Number of compound	Compound	RT (min)	λ_{\max} (nm)
Dvido1	Acacetin-7-O-(rhamnosyl(1-2)glucoside)	35.248	269, 330
Dvido2	Luteolin	47.612	253, 267sh, 291sh, 347
Dvido3	chlorogenic acid	25,812 ± 0,034	245sh, 296sh, 323
Dvidi4	Derivative of Acacetin-7-O-[rhamnosyl (1-2) glucoside]	32,122 ± 0,143	271, 330
Dvidi5	Isoorientin	32.358	254sh, 268, 290sh, 348
Dvidi6	Derivative of quercetin -3-O-arabioside	33,187 ± 0,093	255, 267sh, 294sh, 346
Dvidi7	Luteolin	36.142	253, 267sh, 291sh, 347
Dlic 8	Chrysoeriol-6,8-di-C-glucoside	41,633 ± 0,066	253sh, 267, 343
Dlic9	Scutellarein-6,4'-dimethylether	41.217	274, 332
Dlic10	Acacetin-7-O-(rhamnosyl(1-2)glucoside)	29,579 ± 0,052	267, 333
Evidi11	Neocarlinoside	30.569	256sh, 271, 348
Evidi12	Acacetin-7-O-(rhamnosyl(1-2)glucoside)	31,845 ± 0,146	271, 332
Evidi 13	Isoorientin	32.153	245, 268sh, 290sh, 348
Evidi14	Orientin	33,114 ± 0,093	254, 268sh, 290sh, 350
Evidi15	Luteolin-7-O-rutinoside	36.201	254, 267sh, 348
Evidi 16	Luteolin-7-O-(glucuronosyl(1-2)glucuronide)	36.274	255, 267sh, 345
Evidi 17	Chlorogenic acid	39.09	245sh, 296sh, 324
Elar18	Dof 7-hydroxy-5,8,3',4'-tetramethoxyflavone	54,102 ± 0,052	255sh, 273, 342
Elar19	Genkwanin	31.126	270, 331
Elar20	2,4-dihydroxybenzoic acid	23,844 ± 2,333	240sh, 296sh, 326
Elar21	dof Luteolin-7,3'-di-O-glucuronide	36,795 ± 0,923	250, 278, 337
Elar22	Rosmarinic acid	36.201	250sh, 290sh, 328
Alar23	Vitexin	29.674	271, 331
Alar24	Acacetin-7-O-(rhamnosyl(1-2)glucoside)	30,54 ± 1,12	271, 332
Alic25	Scutellarein-6,4'-dimethylether	53.17	274, 332
Avidi26	Luteolin	47,949 ± 0,062	252sh, 266sh, 290sh, 348
Avidi27	Luteolin-7-O-glucoside	36,89 ± 0,06	254, 266sh, 348

D: decoction, E: ethanol-water, A: acetone, LAR: *L. rhodesiensis*, LIC: *L. chevalieri*, VID: *V. diversifolia*, VIDO: *V. doniana*.






Figure 1: UV spectra of 41 phenols acids flavonoids found in the four edible plants (1/7). Dvido: decoction extract of *Vitex doniana*, Dvidi: decoction extract of *Vitex diversifolia*.




Figure 1: UV spectra of 41 phenols acids flavonoids found in the four edible plants (2/7). Dvidi: decoction extract of *Vitex diversifolia*, Dlic: decoction extract of *Lippia chevalieri*, Evido: decoction extract of *Vitex doniana*.






Figure 1: UV spectra of 41 phenols acids flavonoids found in the four edible plants (3/7). Evidi: ethanol-water extract of *Vitex diversifolia*, Elar: ethanol-water extract of *Lantana rhodesiensis*.




Figure 1: UV spectra of 41 phenols acids flavonoids found in the four edible plants (4/7). Elar: ethanol-water extract of *Lantana rhodesiensis*, Alar: acetone extract of *Lantana rhodesiensis*.






Figure 1: UV spectra of 41 phenols acids flavonoids found in the four edible plants (5/7). Alic: ethanol-water extract of *Lippia chevalieri*, Avidi: acetone extract of *Vitex diversifolia*.




Figure 1: UV spectra of 41 phenols acids flavonoids found in the four edible plants (6/7). Avido: acetone extract of *Vitex doniana*, HFvido: hexane fraction of *Vitex doniana*, EAFlar: ethyl acetate fraction of *Lantana rhodesiensis*,






Figure 1: UV spectra of 41 phenols acids flavonoids found in the four edible plants (7/7). HFvido: hexane fraction of *Vitex doniana*, BFvido: butanolic fraction of *Vitex doniana*, EAFvido: ethyl acetate fraction of *Vitex doniana*

Relationship between Antioxidants Activities And Polyphenolic Compounds Estimated

The interest of the relation investigation is sought to understand the real implication of each group of compounds implied in the observed activities. It emerges from our investigations that total phenolic are implied in the antioxidant activities (Table 2). In fact, 16.10 mg EAG/g dry extract (decoction extract) of *L. chevalieri* produced the best total antioxidant activity (0.850 mg EAA/mL) against 4.78% of flavonoids extract contribution. However one expected that activity is held by *L. rhodesiensis* (ethanol-water extract) with 69.48 mg EAG/g dry extract. If the activities were related to the flavonoids contribution, then the best activity in this case would have been that of *V. diversifolia* of acetone extract (47.58% of contribution). That is not the case. It is the same for iron reducing power (contribution of total flavonoids in the best activity is 4.78%) and free radical scavenging activity (6.82% of total flavonoids contribution). It's thus difficult to establish a clear correlation between biological activity and polyphenolic compounds contents. It generally arises this controversy as for the implication of flavonoids in the antioxidant activities [9]. These results corroborate with those of Meda *et al.* [40]. Former research established that this controversy would be due to several

possibilities: (i) either the compounds existing in the extracts have large molecular weights or are heterosidic [9],[33], (ii) or the majority of the flavonoids are not antioxidants [9],[31], (iii) or there is under estimate of the flavonoids by the method of $AlCl_3$ [30],[34], [40].

The flavonoids inactive and stabilize the free radicals thanks to their hydroxyl grouping (C3-OH) strongly reactive. Indeed, the glycosylation of the 3-OH group of quercetin (case of rutin) or its suppression (case of the luteolin) decreases antioxidant activity [8]. That could explain the weak activity of *V. diversifolia* extracts where we highlighted a derivative of quercetin (quercetin-3-O-arabinoside). The spectral study enabled us to characterize some compounds (Table 4). With through the literature we know that 6-hydroxyflavones are much more stable than their 8-isomers [41]. According to these authors [41], flavonols like acacetin, genkwanin, chrysoeriol and diosmetin cannot be separate well on HPLC, but can be distinguished by their different colors reactions in UV light. According to the same authors, an additional effect existed between 6-hydroxyl group and apigenin which make the compound more polar and shorten the retention times [41]. That could explain in the same order the difference between retention times that we obtained (Table 4) and those of the method used [20]. Thus, hydroxylation decrease the retention time and methylation increase it [41],[42]. Concerning the antioxidant effect of these compounds, it's well known that the polyhydroxy substituted flavonoids

commonly found in most plants present the highest antioxidant activity. Specifically the C-ring plays an important role in this activity ([8],[42]). Traditional medicine being a science of natural substances synergy, we will be able to justify these activities observed by this synergy of plant compounds.

What happens exactly of antioxidants mechanism in the cellular metabolism? According to several authors, much time oxygen oxidizing take place in mitochondria, cytochrome P450, peroxisomes with some enzymes like NADPH oxidase and xanthine oxidase [43],[44],[45],[46]. Reactive oxygen species (ROS) include free radical of oxygen: superoxide anion, hydroxide radical and some derivative non radical form like hydrogen peroxide [25]. Figure 2 give some detail of oxygen share take origin probably in the mitochondria. Seifried et al. [45], tried to establish opposition between excess of oxidants and excess of antioxidants, and then the consequences on human body (Figure 3). It resort in this case, that excess of both of them have negative effect. The best situation is to establish balance in our cells [40]. But like any things, excess harms. The same authors were demonstrated that the excess of antioxidants (enzymatic antioxidant and non-enzymatic antioxidant) can involve the loss of their property [43],[45], [46],[47], [48],[49], [50]. In general the role of primary enzymes (superoxide dismutase, catalase and glutathione peroxidase), help to the detoxication of xenobiotics compounds [51]. The superoxide dismutases are the first defensive lines against the ROS, while the catalases prevent

the oxidative damages by ensuring the passage of H₂O₂ into H₂O and in O₂ [25]. In the case of cells proliferation of cancer, the conjugaison of glutathione (GSH) to the electrophilic compound permit to protect the element. It is well-known the great relationship between glutathione-s-transferase (GST) and glutathione. Indeed, all high concentration of glutathione present in the intracellular middle is always occupied, and it's a powerful detoxicant of xenobiotics compounds and free radicals [25]. And, the glutathione-s-transferases are implied in the resistance of tumorals cells [52],[53].

In a normal healthy human body, the generation of pro-oxidants in the form of ROS is effectively kept in check by the various levels of antioxidant defense [54]. Human body exposure to adverse physicochemical, environmental or pathological agents such as pollutions and overnutrition etc... maintained balance shifted in favor of pro-oxidants resulting in "oxidative stress" [54]. Natural compounds like dietary sources (fruits and leaves of *L. chevalieri*, *L. rhodesiensis*, *V. doniana*) provide a large number of antioxidants, which can prevent many human diseases [8],[54]. Some beverages such as tea (*L. chevalieri*) are also rich sources of antioxidants [8],[29], [54],[55], [56],[57]. According to Tilak et al. [54], a growing body of evidence suggests that moderate consumption of tea and dietary may protect against several forms of cancer, Alzheimer's disease, cardiovascular diseases, and formation of kidney stones, bacterial infections, and dental cavities.




Figure 2: cellular oxidative interactions [45]






Figure 3: defence mechanism against ROS [44]. **GSH:** glutathione reduced, **GSSG:** oxidized glutathione, **SOD:** superoxide dismutase, **NADPH:** nicotinamide adenine dinucleotide phosphate.

Conclusion

Antioxidant activity was investigated together with polyphenolic compounds, using acetone extracts, decoction extracts and ethanol-water extracts. The strongest total phenolic and flavonoids contents were obtained with ethanol-water extracts. Four types of fractions of ethanol-water were evaluated for polyphenolic quantification. Among the four edible species study, *L. rhodesiensis* and *L. chevalieri* presented the best activities on total antioxidant activity, anti-DPPH*, and iron reducing power respectively. By means of the literature we established the importance of balance between oxidant and antioxidant in the alive cells. Indeed it was demonstrated that the excess of antioxidants or oxidants can involve negative effect on human body. With the sight of polyphenolic compounds quality highlighted, one could justify the various activities observed thus the many uses in traditional medicine. Further analysis of these species should be focused to (1) the isolation and characterization, (2) the evaluation of the

capacity of the extracts and/or compounds isolated from its extracts to increase the rate of the superoxide dismutase, the catalase and of the glutathione in the human organism.

Acknowledgments

The authors thank The World Academic Sciences (TWAS) and the Consejo Nacional de Ciencia y Tecnologia (CONACYT) for the financial support through the training and excellence research fellowship. Award of 2012 CONACYT-TWAS Postdoctoral Fellowship. FR number: 3240267272. Authors also gratefully acknowledge the technical support of Professor Almaraz-Abarca Norma, director of «Laboratorio de Biotecnología», to have welcomed me in your laboratory and have allowed me to familiarize myself with its laboratory equipment and especially the HPLC-DAD. We cannot overlook, the support of the various laboratories of Instituto Politécnico Nacional, Unidad Durango.

References

- [1]. Bangou MJ, Almaraz-Abarca N, Méda NTR, Zeba B, Kiendrebéogo M, Millogo-Rasolodimby J, Nacoulma OG. Polyphenolic composition of *Lantana camara* and *Lippia chevalieri*, and their Antioxidant and Antimicrobial Activities, International Journal of Phytomedicine. 2012; 4(1):115–124.
- [2]. Ouattara FO. Treatment of sexually transmitted infections in Mali: Study of Phytochemistry and Biological Activities of *Annona senegalensis*

- (Annonaceae) and *Stachytarpheta val angustifolia* (Verbenaceae). Université de Bamako. 2005.
- [3]. Sawadogo WR, Cerella C, Al-Mourabit A, Moriou C, Teiten MH, Guissou IP, Dicato M, Diederich M. Cytotoxic, antiproliferative and pro-apoptotic effects of 5-Hydroxyl-6,7,31,41,51-Pentamethoxyflavone isolated from *Lantana ukambensis*, *Nutrients*. 2015; 7:10388–10397.
- [4]. Bangou MJ, Kiendrebeogo M, Compaoré M, Coulibaly AY, Méda NTR, Almaraz-Abarca N, Zeba B, Millogo-Rasolodimby J, Nacoulma OG. Enzyme Inhibition Effect and Polyphenolic Content of Medicinal Plant Extracts from Burkina Faso. *Journal of Biological Sciences*. 2011; 11(1):31-38
- [5]. Pascual ME, Slowing K, Carretero E, Sanchez Mata D, Villar A. *Lippia*: traditional uses, chemistry and pharmacology: a review, *J. Ethnopharmacol.* 2001; 76: 201 – 214.
- [6]. Nacoulma OG. Plantes médicinales et Pratiques médicales Traditionnelles au Burkina Faso: cas du plateau central T1&T2. Thèse Doctorat d'Etat ès Sciences Nat. Université de Ouagadougou, tome I et II. 1996.
- [7]. Taiwo O, Xu HX, Lee SF. Antibacterial activities of extracts from Nigerian chewing sticks, *Phytother. Res.* 1999; 13:675– 679.
- [8]. Bangou MJ, Meda N-TR, Kiendrebeogo M, Nacoulma OG, Almaraz-Abarca N. A review on *Lantana rhodesiensis* Moldenke: traditional uses, phytochemical constituents and pharmacological activities, *Int. J. of Phytomedicine*, 2017; 9(1): 1-10.
- [9]. Bangou MJ, Méda N-TR, Thiombiano AME, Kiendrebéogo M, Zeba B, Millogo-Rasolodimby J, Couliadiat YD, Yougbaré-Ziébrrou M, Nacoulma OG. Antioxidant and antibacterial activities of five verbenaceae species from Burkina Faso, *Curr. Res. J. Biol. Sci.* 2012; 4: 665 – 672.
- [10]. Agbafor KN, Nwachukwu N. Phytochemical analysis and antioxidant property of leaf extracts of *Vitex doniana* and *Mucuna pruriens*, *Biochem. Res. Int.* 2011; 459839:4
- [11]. Bangou MJ, Almaraz-Abarca N, Meda N-TR, Yougbaré-Ziébrrou M, Millogo-Rasolodimby J, Nacoulma OG. *Lippia chevalieri* Moldenke: A brief review of traditional uses, phytochemistry and pharmacology, *Int. J. Drug Delivery*, 2012; 4(3):289– 296.
- [12]. Agunu A, Yusuf S, Andrew GO, Zezi AU, Abdurahman EM. Evaluation of five medicinal plants used in diarrhoea treatment in Nigeria, *J. Ethnopharmacol.* 2005;101: 27 – 30.
- [13]. Nebie RHC, Yameogo RT, Belanger A, Sib FS. Chemical composition of essential oils of *Vitex diversifolia* Bark, *J. Essent. Oil Res.* 2005;17: 276 – 277.
- [14]. Nwachukwu E, Uzoeto HO. Antimicrobial activities of leaf of *Vitex doniana* and *Cajanus cajan* on some bacteria, *Researcher*. 2010; 2: 37 – 47.
- [15]. Mustapha AT, Fanna IA, Irfan ZK, Sandabe UK. The effects of ethanolic extract of *Vitex Doniana* Stem Bark on peripheral and central nervous system of laboratory animals, *JAPS*, 2012; 2: 74 – 79.
- [16]. Bibi A, Thangamani A, Venkatesalu V. Some endemic medicinal plants of Andamans with antimicrobial potential, *JAAR*. 2016; 1: 10 – 15.
- [17]. Nurmi K, Vladimir O, Erkki H, Kalevi P. Variation of total phenolic content and individual low-molecular-weight phenolics in foliage of mountain birch trees (*Betula pubescens* ssp. *Tortuosa*), *J. Chem. Ecol.* 1996; 22: 2023 – 2040.
- [18]. Lauranson-Broyer J, Lebreton P. Flavonoids and morphological traits of needles, as markers of natural hybridization between *Pinus uncinata* Ram and *Pinus sylvestris* L, *Biochem. Syst. Ecol.* 1993; 21: 241 – 247.
- [19]. Mabry TJ, Markham KR, Thomas MB. The systematic identification of flavonoids. Springer-Verlag, New York, 1970.
- [20]. Campos MG, Markham KR. Structure information from HPLC and on-line measured absorption spectra-flavone, flavonols and phenolic acids. Coimbra University Press, Coimbra, Portugal, 2007.
- [21]. Prieto P, Pineda M, Aguilar M. Spectrophometric quantification of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific to the determination of vitamin E, *Anal. Biochem.* 1999;269:337– 341.
- [22]. Campos MG, Cunha PA, Navarro MC, Utrilla MP. Free radical scavenger activity of bee pollen, *Polyphenols*. 1994; 94: 415 – 426.
- [23]. Yang J, Guo J, Yuan J. In vitro antioxidant properties of rutin, *LWT Food Sci. Technol.* 2008; 41:1060 – 1066.
- [24]. Perkin-Elmer, Analysis of Plant Tissue, AY-5, United States of America. 1982; :141–143.
- [25]. Bangou MJ. Etude phytochimique et activités biologiques des tiges feuillées de *Lantana camara* L. et de *Lippia chevalieri* Moldenke: deux Verbenaceae du Burkina Faso, Thèse Unique de Doctorat, Université de Ouagadougou. 2012.
- [26]. Nsimba RY, Kikuzaki H, Konishi Y. Antioxidant activity of various extracts and fractions of *Chenopodium quinoa* and *Amaranthus spp.* seeds. *Food Chem.* 2008;106:760 – 766.
- [27]. Barriada-Bernal LG, Almaraz-Abarca N, Delgado-Alvarado EA, Gallardo-Velázquez T, Avila-Reyes JA, Torres-Moran MI, González-Elizondo MS, Herrera-Arrieta Y. Flavonoid composition and antioxidant capacity of the edible flowers of *Agave durangensis* (Agavaceae), *CyTA-Journal of Food*. 2013; 12:105 – 114.
- [28]. Falleh H, Ksouri R, Medini F, Guyot S, Abdelly C, Magné C. Antioxidant activity and phenolic composition of the medicinal and edible halophyte *Mesembryanthemum edule* L, *Ind. Crop. Prod.* 2011; 34: 1066 – 1071.
- [29]. Reyes-Martínez A, Almaraz-Abarca N, Gallardo-Velázquez T, González-Elizondo MS, Herrera-Arrieta Y, Pajarito-Ravelero A, Alanís-Bañuelos RE, Torres-Morán MI. Evaluation of foliar phenols of 25 Mexican varieties of common bean (*Phaseolus vulgaris*

- L.) as antioxidants and varietal markers, *Nat. Prod. Res.* 2014; 28: 2158 – 2162
- [30]. Compaore M, Lamien-Meda A, Mogosan C, Lamien CE, Kiendrebeogo M, Vostinaru O, Vlase L, Ionescu C, Nacoulma OG. Antioxidant, diuretic activities and polyphenol content of *Stereospermum kunthianum* Cham.(Bignoniaceae), *Nat. Prod. Res.* 2011; 25: 1777 – 1788.
- [31]. Gursoy N, Sarikurkcü C, Cengiz M, Solak MH. Antioxidant activities, metal contents, total phenolics and flavonoids of seven *Morchella* species, *Food Chem. Toxicol.* 2009;47: 2381 – 2388.
- [32]. Lamien-Meda A, Lamien CE, Compaoré MMY, Meda RN-T, Kiendrebeogo M, Zeba B, Millogo JF, O.G. Nacoulma OG. Polyphenol content and antioxidant activity of fourteen wild edible fruits from Burkina Faso, *Molecules*, 2008;13:581 – 594.
- [33]. Manzi P, Marconi S, Aguzzi A. Commercial mushrooms: Nutritional quality and effect of cooking, *Food Chem.*, 2004; 84:201 – 206.
- [34]. Meda A, Lamien CE, Romito M, Millogo J, Nacoulma OG. Determination of the total phenolic, flavonoid and proline contents in Burkina Faso honey, as well as their radical scavenging activity, *Food Chem.* 2005;91:571– 577.
- [35]. Hinneburg I, Dorman HJD, Hiltunen R. Antioxidant activities of extracts from selected culinary herbs and spices, *Food Chem.* 2006; 97: 122 – 129.
- [36]. Tai Z, Cai L, Dai L, Dong L, Wang M, Yang Y, Ding Z. Antioxidant activity and chemical constituents of edible flower of *Sophora viciifolia*, *Food Chem.* 2011; 126: 1648 – 1654.
- [37]. Piero NM, Kimuni NS, Ngeranwa NJ, Orinda OG, Njagi MJ, Maina D, Agyirifo SD, Gathumbi K, Kinge WS, Njagi Eliud EN. Antidiabetic and safety of *Lantana rhodesiensis* in Alloxan induced diabetic rats, *J. Develop. Drugs.* 2015; 4(1): 1-10.
- [38]. Lagnika L, Amoussa M, Adjovi Y, Sanni A. Antifungal, antibacterial and antioxidant properties of *Adansonia digitata* and *Vitex doniana* from Bénin pharmacopeia, *J. Pharmacognosy Phytother.* 2012; 4: 44 – 52.
- [39]. Vunchi MA, Umar AN, King MA, Liman AA, Jeremiah G, Aigbe CO. Proximate, Vitamins and Mineral Composition of *Vitex doniana* (black plum) Fruit Pulp, *Nig. J. Basic Appl. Sci.* 2011; 19:97 – 101.
- [40]. Meda N-TR, Bangou MJ, Bakasso S, Millogo-Rasolodimby J, Nacoulma OG. Antioxidant activity of phenolic and flavonoid fractions of *Cleome gynandra* and *Maerua angolensis* of Burkina Faso, *J. Appl. Pharm. Sci.* 2013; 3:36 – 42.
- [41]. Greenham J, Harborne JB, Williams CA. Identification of Lipophilic Flavones and Flavonols by Comparative HPLC, TLC and UV Spectral Analysis. *Phytochemical Analysis*, 2003; 14:100-118.
- [42]. Tsimogiannis DI, Oreopoulou V. Free radical scavenging and antioxidant activity of 5, 7, 3V, 4V-hydroxy-substituted flavonoids. *Innovative Food Science and Emerging Technologies*, 2004; 5: 523-528.
- [43]. Carochó M, I.C.F.R. Ferreira ICFR. A review on antioxidants, prooxidants and related controversy: Natural and synthetic compounds, screening and analysis methodologies and future perspectives, *Food Chem. Toxicol.* 2013;51: 15 – 25.
- [44]. Lascarrou JB. Comparaison de deux méthodes de dosage de la capacité antioxydante totale du sérum au cours de la chirurgie thoracique. Université de Nantes. 2010.
- [45]. Seifried HE, Anderson DE, Fisher EI, JA. Milner JA. A review of the interaction among dietary antioxidants and reactive oxygen species, *J. Nutr. Biochem.* 2007; 18:567–579.
- [46]. Sies H. Oxidative Stress, Oxidants and Antioxidants, *Exp. Physiol.* 1997; 82: 291 – 295.
- [47]. Flora SJS. Structural, chemical and biological aspects of antioxidants for strategies against metal and metalloids exposure, *Oxid. Med. Cell. Longev.* 2009; 2:191 – 206.
- [48]. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health, *Pharmacogn. Rev.* 2010; 4:119 - 126.
- [49]. Pham-Huy LA, He H, Pham-Huy C. Free radicals, antioxidants in disease and health, *Int. J. Biomed. Sci.* 2008;4: 89 – 96.
- [50]. Rahman K. Studies on free radicals, antioxidants, and co-factors, *Clin. Interv. Aging*, 2007;2: 219 – 236.
- [51]. Thamotharan G, Sekar G, Ganesh T, Sen S, Chakraborty R, Kumar NS. Antiulcerogenic effects of *lantana camara* linn. leaves on *in vivo* test models in rats, *Asian J. Pharm. Clin. Res.* 2010;3: 57 – 60.
- [52]. Feng QL, Davey KG, Pang ASD, Ladd TR, Retnakaran A, Tomkins BL, Zheng S, Palli SR. Developmental expression and stress induction of glutathione S-transferases in the spruce budworm, *choristoneura fumiferana*, *J. Insect. Physiol.* 2001;47:1 – 10.
- [53]. Van Zanden JJ, Hamman OR, van Iersel MLPS, Boeren S, Cnubben PJ, I.M.C. Rietjens IMC. Inhibition of human glutathione S-transferases P1-1 by the flavonoid quercetin, *Chem. Biol. Interac.* 2003;145: 139 – 148.
- [54]. Tilak JC, Devasagayam TPA, Lele RD. Antioxidant Activities From Indian Medicinal Plants: A Review, *Current status and future prospects.* 2001; 21-24.
- [55]. Ayoola GA, Coker HAB, Adesegun SA, Adepoju-Bello AA, Obaweya K, Ezennia EC, Atangbayila TO. Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria, *Trop. J. Pharm. Res.* 2008; 7:1019 – 1024.
- [56]. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer, *Life Sci.* 2004; 74: 2157–2184.
- [57]. Djeridane A, Yousfi M, Nadjemi B, Boutassouna D, Stocker P, Vidal PN. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds, *Food Chem.* 2006;97:654–660.

- [58]. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. *Afri. J. Biotechnol.* 2005; 4: 685 – 688.
- [59]. Mevy JP, Bessiere JM, Dherbomez M, Millogo J, Viano J. Chemical composition and some biological activities of the volatile oils of a chemotype of *Lippia chevalieri* Moldenke, *Food Chem.* 2007;101:682 – 685.
- [60]. Suksamrarn A, Yingyongnarongkul B, Promrangsam N. Naturally Occurring 20, 26-Dihydroxyecdysone Exists As Two C-25 Epimers Which Exhibit Different Degrees Of Moulting Hormone Activity, *Tetrahedron*, 1998; 54:14565 – 14572.
- [61]. Aphajitt S, Nimgirawath K, Suksamrarn A, Tooptakong U. Isolation and Crystal Structure of Limonidilactone-A Labdane Diterpene From *Vitex limoniifolia*, *Aust. J. Chem.* 2006;48:133 – 137.
- [62]. Kannathasan K, Senthikumar A, Venkatesalu V. In vitro potential of some *Vitex* species against human pathogenic bacteria. *Asian Pac. J. Trop. Med.* 2011; 4:645 – 648.
- [63]. Egbekun MK, Akowe JI RJ, Ede RJ. Physico-chemical And Sensory Properties of Formulated Syrup from Black Plum (*Vitex doniana*) Fruit, *Plant Foods Hum. Nutr.* 1996; 49:301 – 306.
- [64]. Ladeji O, Udoh FV, Okoye ZSC. Activity of Aqueous Extract of The Bark of *Vitex doniana* On Uterine Muscle Response To Drugs, *Phytotherapy Res.* 2004;19:804 – 806.
- [65]. Ochieng CO, Ishola IO, Opiyo SA, Manguro LA, Owuor PO, Wong KC. Phytoecdysteroids from the stem bark of *Vitex doniana* and their anti-inflammatory effects, *Planta Med.* 2013; 79:52 – 59.
- [66]. Ajiboye TO, Standardized TO. Extract of *Vitex doniana* Sweet stalls protein oxidation, lipid peroxidation and DNA fragmentation in acetaminophen-induced hepatotoxicity. *J. Ethnopharmacol.* 2015; 164:273 – 282.

