
Nutrient Use Efficiency of Two Improved Cultivars of Taro (*Colocasia esculenta*) under Screen House Conditions in Samoa

Sanjay Anand^{1*} and Danilo Flores Guinto²

DOI: <http://doi.org/10.9734/bpi/npacs/v1>

ABSTRACT

Taro (*Colocasia esculenta* (L.) Schott) is a staple crop of many of the South Pacific nations with an ever increasing export demand. In recent years, yields of taro have increased dramatically through breeding and selection. However, selections of improved lines are often entirely based on final yield. There are many physiological pathways by which increased potential yield may be achieved. Factors such as the accumulation of dry matter and nutrient use efficiency, merit investigation. Two improved (blight resistant) taro cultivars were planted and harvested for biomass measurements on a monthly basis for a total of eight months (30-240 days after planting) through destructive sampling. At each harvest, plants were separated into various plant parts and their dry matter accumulation and nutrient content were determined. Comparatively, cultivar Samoa 2 showed significantly higher uptake of N (25%), P (37.5%), K (33%), Mg (36.4%), Mn (22.7%) and Zn (48.3%) than cultivar Samoa 1. Even though maximum levels of total plant uptake of nutrients by the two cultivars did not differ between the cultivars, cultivar Samoa 1 plants absorbed 17% less N, 26% less P and 20% less K than those of cultivar Samoa 2 with the uptake uniformly distributed over the entire life cycle of the crop. Although cultivar Samoa 2 resulted in higher total plant (19.6%) and corm dry matter (10.4%) productions, cultivar Samoa 1 had a higher nutrient use efficiency, (kg of edible dry matter produced per kg of nutrient taken up), for N, P, K, Mg, Mn and Cu over cultivar Samoa 2. However, for Ca, Fe and Zn. Cultivar Samoa 2 had a higher nutrient use efficiency over cultivar Samoa 1. Based on nutrient use efficiency of the cultivars, Samoa 1 is recommended for marginal to rich soils while Samoa 2 for good to rich soils.

Keywords: Dry matter accumulation; nutrient uptake; nutrient use efficiency; destructive sampling.

1. INTRODUCTION

Root and tuber crops are the major sources of dietary energy for many people in the Pacific island countries. In the Pacific Islands, taro has always been richly woven into the fabric of life [1,2]. Taro is postulated to have originated in southern or south-east Asia, and to have been dispersed to Oceania through the Island of New Guinea centuries ago [1,2]. The crop has evolved with the cultures of the people of the Asia/Pacific region. Not surprisingly, it has acquired considerable socio-cultural importance for the people. Among the food crops in Oceania region, the adulation and prestige attached to taro is equalled only by yam in certain localities [1,3].

In Samoa, taro is mostly grown with minimum use of tillage and agro-inputs, on small fragmented farm holdings averaging 2.4 ha in size. From this, the average allocation to taro is about 0.48 ha/holding. There are only a few commercial growers with farm sizes greater than 5 ha. The total number of taro holdings was 15,106 [4] with a total production area of 7,346 ha of which 65% is cultivated as monocrop while the remaining 35% is grown in intercropped and agro-forestry systems. The average local yield is 6.2 mt/ha and annual production ranges from 40,000 to 50,000 metric tonnes [4]. The annual production

¹Faculty of Business and Economics, School of Agriculture and Food Technology, The University of the South Pacific, Alafua Campus, Apia, Samoa.

²Ballance Fertilisers, Tauranga, New Zealand.

*Corresponding author: E-mail: anand_s@usp.ac.fj, sanjay.anand@samoa.usp.ac.fj;

of the Oceania region is around 300,000 metric tonnes with the leading producers being Papua New Guinea, Samoa, Tonga, Solomon Islands and Fiji [5].

Although the roots are the most widely consumed and important parts of the plant in the Pacific region, the leaves, petioles and corms of taro are all consumed as fresh vegetables depending on the cultivar and the culture [1,2]. Young taro leaves and stems can be eaten after boiling twice to remove the acrid flavor and the leaves are a good source of vitamins A and C and contain more protein than the corms [6]. Taro corms and cormels are good sources of essential mineral nutrients that contribute to growth as well as health maintenance and general well-being [7]. The major mineral nutrient in taro is K [5] and it is also rich in Fe, Zn and Ca [8,9]. Variable mineral nutrient levels between different cultivars of taro were observed in Papua New Guinea [10]. The corms contained K (250- 480 mg/100 g), Mg (19-37 mg/100 g), Ca (11-45 mg/100 g), Zn (0.2-6.3 mg/100 g), Fe (0.6-1.8 mg/100 g) and Na (0-3 mg/100 g). Taro is a good source of Na, K, Mg and Ca whose salts regulate the acid-base balance of the body [11]. Wide variations observed among different cultivars of taro have been attributed to differences in genetic background as well as climate, soil, season and agronomic factors [11,12].

Variation in mineral composition among the accessions of taro is probably due to differences in the genetic potential of each accession to obtain nutrients from the soil since different taro genotypes have different nutrient-use efficiencies [13,14]. As was found in the same study, regarding mineral content, high levels of variability in South East Asia and Oceania taro germplasm were also found with regards to chemical composition for minerals but also for lipids, proteins, amylose, glucose, fructose and saccharose [13,15]. Availability of N, P, K and S fertilizers increase yield as well as nutritional quality of root and tuber crops [16].

In most studies with food crops in the Pacific, biomass production and nutrient uptake receive little attention, particularly due to the tedious and difficult nature of the quantification process [12]. This has led to a scarcity of basic information regarding dry matter accumulation and nutrient uptake for the taro crop, particularly under intensive cropping systems which are aimed at satisfying the crop demand of a growing population and supplying corms for export markets. An essential step to increase the efficiency of fertilizers in order to improve yields is an understanding of nutrient uptake and allocation within the taro plant during the growing season [14]. These data are essential for the development of technological packages, especially involving nutrient inputs, growth simulation models, and decision support system [14]. This information is also critical for the establishment of taro breeding programs aimed at raising the yield potential of taro [2].

The objective of this research was to investigate the dry matter accumulation, nutrient uptake and nutrient use efficiency of two improved taro cultivars, *Samoa 1* and *Samoa 2*, grown under semi-controlled screen house environment, through regular temporal destructive sampling. These two new cultivars were bred locally for resistance to taro leaf blight and have adapted well to the local growing conditions. In addition, these two cultivars have excellent taste characteristics over other newly bred lines and have been successfully accepted in the local market and have been endorsed for the export markets as well as for value addition. Therefore, it is imperative to ascertain the nutrient uptake data which reflects on the nutritional value data for these two new local cultivars in order to realize their full economic potential.

2. MATERIALS AND METHODS

2.1 Description of the Trial

2.1.1 Study area

This experiment was conducted at the University of the South Pacific's Alafua Campus in Samoa (15-17° S and 171-173° W). The location experiences a humid tropical climate with an annual rainfall varying from 2,500 to 3,500 mm with a strong seasonality of distribution. The months from April to October are the driest times of the year. The average annual air temperature range is 20–33°C [3].

2.1.2 Soil characterization

The soil used was a well-drained inceptisol, oxic humitropept, clayey-skeletal oxidic isohyperthermic. The initial chemical characterization of the soil is given in Table 1 below. The soil was air dried and sieved through a 1 cm mesh. The potting bags were filled with 10 kg of soil each.

Table 1. Chemical characterization of the trial soil

Chemical and nutritional composition	
pH (H ₂ O)	6.0
CEC (cmol _c /kg)	20.83
EC (dS/m)	0.22
OC (%)	3.2
Total N (%)	0.43
Olsen P (mg/kg)	3.85
Exchangeable K (cmol _c /kg)	0.32
Exchangeable Ca (cmol _c /kg)	9.74
Exchangeable Mg (cmol _c /kg)	1.07
DTPA Extractable Fe (mg/kg)	38.97
DTPA Extractable Mn (mg/kg)	49.46
DTPA Extractable Cu (mg/kg)	3.45
DTPA Extractable Zn (mg/kg)	2.86

2.1.3 Nutrient supplementation and incubation

The entire package of macro and micronutrient elements, based on the soil pH, was included for nutrient supplementation to each pot, carried out at recommended levels as described by Asher et al., (Table 2) [17]. An incubation time of two weeks was allowed before the planting of the two taro cultivars.

2.1.4 Plant culture and experimental layout

Suckers of two improved taro cultivars, *Samoa 1* and *Samoa 2*, were planted in pots and laid out in a factorial arrangement, using randomised complete block design with five replications. Each replication consisted of plots randomly assigned to the two cultivars which were to accommodate eight randomly assigned monthly biomass harvests, sampled for dry matter accumulation and nutrient uptake at different stages of plant growth. There were six data plants of each variety from each block for each of the eight harvests totaling to 240 plants for each cultivar (480 plants for the whole experiment). The cultivars and harvest periods were completely randomized within a block.

2.2 Data Collection

Six taro plants of each cultivar from a block were harvested at 30, 60, 90, 120, 150, 180, 210, and 240 days after planting (DAP), to ascertain the dry matter measurements and total chemical analysis of individual plant parts. Plants in the sub-plots were harvested, washed and separated into petioles, corms, roots and sucker components. Samples of the various plant parts were oven dried to a constant weight at 65°C for dry matter determination. The dried samples were ground to pass through a 1.0-mesh screen and analysed for total N, P, K, Ca, Mg, Fe, Mn, Cu and Zn. The third most upper leaf lamina was also analysed for these elements at 30, 60, 90, 120, 150, 180, 210, and 240 days after planting (DAP). Nitrogen was determined by the micro-Kjeldahl procedure [18], P by molybdovanadophosphoric acid [18], and K, Ca, Mg, Zn, Fe, Mn, and Cu by atomic absorption spectrophotometry [19,20].

Table 2. Typical rates of nutrient supplementation for soils with a pH of 6.0

Element	Application rates (kg/ha)
N	100
P	60
K	80
Ca	35
Mg	30
Fe	5
Mn	5
Cu	3
Zn	4

2.3 Nutrient Uptake Calculations

Nutrient uptake and accumulation were calculated as the product of dry matter content and tissue nutrient concentrations for the various plant parts at various stages of growth over the entire growth cycle of the crop. The mean values from the six data plants, for each nutritional index and the number of plants per hectare were used to extrapolate nutrient uptake on a hectare basis. The nutrient use efficiency was calculated as the kg of corm dry matter produced per kg of nutrient taken up [14,15].

2.4 Statistical Analysis

All the data collected were subjected to two-way analysis of variance for differences between cultivars. Best-fit curves were determined using polynomial regression procedures of the Genstat Statistical Software package [21]. Only coefficients significant at $P < 0.05$ were retained in the model.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Dry matter accumulation by various plant organs

The accumulation of dry matter by various plant organs of the two cultivars is illustrated in Fig. 1a-f. The mean total dry matter yield showed cultivar Samoa 2 had 19.6% higher accumulation than cultivar Samoa 1 throughout the experimental period (Table 4). The first 90 days after planting (DAP) were characterized by low rates of total dry matter production by both the cultivars (Fig. 1a), however, statistically significant with cultivar Samoa 2 accumulating higher dry matter yield. During this period, leaves and petioles accounted for 58% of the total dry matter produced in each cultivar (Fig. 1a-c). Following 210 DAP, the dry matter content in the leaves and petioles declined to less than 25% of the total dry matter, but it increased significantly in corms and suckers (Fig. 1e and f). During the first 90 DAP, roots of cultivars Samoa 1 and Samoa 2 represented about 13% and 18% of the total dry matter content, respectively. Following 180 DAP, the dry matter content in the roots was never higher than 8% for Samoa 1 and 12% for Samoa 2. Cultivar Samoa 2 accumulated significantly higher root dry matter than Samoa 1 throughout the experimental period. It is noteworthy that, between 150 and 240 DAP, the suckers were a significant sink of dry matter in the taro plant. During this period, these organs accounted for 22% of the total plant dry matter in Samoa 1 and 13% in Samoa 2. Maximum significant dry matter accumulation in the corms of both cultivars was recorded between 210 and 240 DAP, accounting for about 46% of the total plant dry matter.

3.1.2 Nutrient uptake of the two taro cultivars

Two way analysis of variance revealed significantly higher uptake of N (25%), P (37.5%), K (33%), Mg (36.4%), Mn (22.7%) and Zn (48.3%) by cultivar Samoa 2. (Table 4). In general, the nutrient uptake was very similar between cultivars during the first 150 DAP; thereafter, the quantity of all the nutrients taken up by plants of cultivar Samoa 1 was lower than that of cultivar Samoa 2. The only exception was for Fe uptake where uptake by cultivar Samoa 1 was higher than cultivar Samoa 2, however, this was not significant (Fig. 2a-e and 3a-d, Table 3).




Fig. 1. Dry weights of plant organs of the two taro cultivars as influenced by age

3.1.3 Maximum levels of total plant uptake of nutrients by the two cultivars (kg/ha)

There were no statistical difference between the two cultivars for the maximum levels of total plant uptake of nutrients (Table 4). However, it is noteworthy that cultivar Samoa 1 plants absorbed 17% less N, 26% less P and 20% less K than those of cultivar Samoa 2 with the uptake uniformly distributed over the entire life cycle of the crop (Table 4).

3.1.4 Leaf nutrient concentrations

The nutritional analyses of the third uppermost leaf lamina of the two cultivars revealed only significantly higher concentrations of Mg (12.5%) and Zn (22.2%) in cultivar Samoa 2. In general, the concentrations of all the nutrients except Fe in the leaf lamina of cultivar Samoa 1 plants had greater concentrations than cultivar Samoa 2 plants though not statistically significant (Table 5).

3.1.5 Nutrient use efficiencies

There were significant differences in the total and corm dry matter productions between the cultivars throughout their entire growth period (Table 3). Cultivar Samoa 1 had a higher nutrient use efficiency (kg of edible dry matter produced per kg of nutrient taken up), for N, P, K, Mg, Mn and Cu over cultivar Samoa 2. However, for Ca, Fe and Zn, cultivar Samoa 2 had a higher nutrient use efficiency over cultivar Samoa 1 (Figs. 4 and 5).

3.2 Discussion

Taro exhibits continuous partitioning (a balance between vegetative growth and storage organ growth is maintained throughout the growing cycle) with an almost linear increase in fresh and dry weights [22,23]. The dynamics of dry matter accumulation, nutrient uptake and partitioning by two taro cultivars with fertilization under natural open field conditions showed similar patterns from a research carried out

in Isabella, Puerto Rico [14]. In a separate investigation involving comparisons between natural field conditions and 50% shade conditions, the corm yields were not affected by shade but the total biomass increased under shade as opposed to full sunlight [24-26]. Corm percentage dry matter, which reflects quality, was also reported to be higher under shade. These were attributed to greater photosynthetic efficiency resulting from increased stomatal and chlorophyll densities as aroids are shade tolerant crops [24,27].

The findings of this study showed that the dry matter accumulation by various plant organs followed analogous sigmoid patterns over the crop life cycle as reported by other authors [14,15]. Towards senescence, the suckers were the principal sink of dry matter for both the cultivars. This result is of particular importance because, when taro is grown under upland conditions, cormels of suckers seldom reach a marketable size; and they may compete for assimilates with the marketable main corm. This finding may influence such decisions as to remove the competing suckers at later stages of crop growth [13].

The comparatively higher nutrient uptake of cultivar Samoa 2 can be ascribed to the genotypic variations as reported by various other researchers who worked with taro [15,28]. Other studies on the N, P and K content of different plant parts at various growth stages revealed that the nutrient content changes with increase in age of the crop. The N and K contents in the foliage of taro were reported to be at its highest after 150 DAP; thereafter, decreased with maturity. The N content of root, tuber and pseudo-stem decreased towards maturity of the crop [14,15]. This was in agreement with the findings of this study with days after planting highly significant across all the nutrients analyzed.




Fig. 2. Macronutrient contents of the two taro cultivars as influenced by plant age
Table 3. Mean dry matter yield (kg/ha) and plant uptake (kg/ha) of nutrients by the two cultivars various across the 8 monthly biomass harvests

Cultivar	Mean (kg/ha)									
	Total dry matter**	N**	P***	K***	Ca	Mg****	Fe	Mn	Cu	Zn***
Samoa 1	664	78.7	13.6	123.6	47.3	10.7	5.6	0.43	0.040	0.178
Samoa 2	809	98.4	18.7	164.4	55.4	14.6	4.7	0.53	0.047	0.264
LSD (5%)	106.3	13.30	2.13	24.36	11.12	1.71	1.5	0.09	0.010	0.031

Table 4. Maximum levels of nutrient uptake by the two cultivars (kg/ha)

Nutrient	Samoa 1	Samoa 2	P (0.05)	LSD (5%)
N	146	176	0.310	62.11
P	35	47	0.051	12.06
K	259	321	0.257	111.35
Ca	165	183	0.477	40.88
Mg	20	28	0.087	9.02
Fe	21	15	0.156	7.80
Mn	0.9	1.1	0.237	0.39
Cu	0.07	0.08	0.102	0.39
Zn	0.39	0.54	0.095	0.175

Nutrient concentration of the two taro cultivars




Fig. 3. Micronutrient contents of the two taro cultivars as influenced by plant age




Fig. 4. Relationship between corm dry matter yield and macronutrient contents of the two cultivars




Fig. 5. Relationship between corm dry matter yield and micronutrient contents of the two cultivars

Table 5. Percent nutrient concentration in the lamina of the third uppermost leaf of the two taro cultivars at various stages of growth

Days after planting (DAP)	Cultivar (CV)	Nutrient Content (%)								
		N	P	K	Ca	Mg	Fe	Mn	Cu	Zn
30	Samoa 1	4.73	0.33	3.85	0.97	0.18	0.104	0.068	0.003	0.014
	Samoa 2	4.87	0.34	2.87	0.78	0.30	0.120	0.060	0.004	0.019
60	Samoa 1	4.21	0.44	3.33	1.82	0.16	0.425	0.059	0.005	0.017
	Samoa 2	4.39	0.46	3.11	1.54	0.18	0.489	0.075	0.005	0.028
90	Samoa 1	3.89	0.41	2.93	2.30	0.37	0.625	0.049	0.006	0.028
	Samoa 2	4.01	0.44	2.89	2.21	0.45	0.513	0.050	0.007	0.036
120	Samoa 1	4.37	0.41	3.75	2.23	0.37	0.675	0.053	0.010	0.024
	Samoa 2	4.53	0.43	3.71	1.76	0.37	0.733	0.045	0.011	0.040
150	Samoa 1	3.94	0.41	4.01	1.42	0.35	1.052	0.069	0.007	0.031
	Samoa 2	4.29	0.39	4.00	1.17	0.29	0.940	0.067	0.007	0.036
180	Samoa 1	2.94	0.39	3.78	12.11	0.31	1.103	0.085	0.006	0.033
	Samoa 2	3.43	0.38	3.63	10.82	0.31	1.364	0.090	0.006	0.036
210	Samoa 1	3.14	0.47	3.35	2.40	0.27	1.649	0.071	0.006	0.031
	Samoa 2	3.18	0.40	3.35	1.97	0.29	0.645	0.074	0.005	0.036
240	Samoa 1	4.18	0.56	3.34	1.95	0.36	0.575	0.066	0.006	0.029
	Samoa 2	3.41	0.48	2.91	2.41	0.34	0.050	0.064	0.005	0.038
P-value (5%)	CV	0.440	0.417	0.067	0.751	0.016	0.154	0.595	0.958	<0.001
	DAP	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
	CV x DAP	0.637	0.125	0.626	0.015	0.002	0.028	0.336	0.994	0.071
LSD (5%)	CV	0.271	0.023	0.205	0.463	0.1003	0.1067	0.0058	0.0010	0.0019
	DAP	0.543	0.045	0.411	0.926	0.2005	0.2134	0.0116	0.0019	0.0038
	CV x DAP	0.767	0.064	0.581	1.310	0.2836	0.3018	0.0164	0.0027	0.0053

Both cultivars exhibited higher levels of K uptake relative to N. This suggests that, as with most root crops, taro has a high requirement for K relative to N. Analogous findings were reported with the total plant as well as corm being characterized by high concentrations of K [24,29,30]. Potassium application resulted in greater leaf area and leaf area duration and exerted a profound influence in diverting greater proportion of dry matter into corms than N and increased the dry matter accumulation in corms, corm size, and yield. The increase in corm yield due to K was attributed partly to its effect in bringing about slightly earlier corm initiation and partly to an increase in bulking rate [31].

The variations in the leaf tissue nutrient concentrations can be attributed to genetic differences between the cultivars [24]. Higher plant vigour and sucker production was observed by cultivar Samoa 2 relative to cultivar Samoa 1 [27]. Among the different plant portions, leaf was found to be the richest in N (4-5%). Comparable findings were reported by other researchers [10,31]. This is of high nutritional significance, since leaves are consumed as fresh vegetable in the Pacific island communities.

Furthermore, the nutrient use efficiencies, computed as the weight of edible dry matter produced for every kg of nutrient taken up, revealed that though cultivar Samoa 2 had higher nutrient uptake, it required greater quantities of N, P, K, Mg, Mn and Cu to produce one kg of dry matter as compared to cultivar Samoa 1. Conversely, Ca, Fe and Zn were required in relatively higher amounts by cultivar Samoa 1 as opposed to cultivar Samoa 2, to produce one unit of corm dry matter. In another separate field trial, the effect of the taro genotype was significant for more than half of the analyzed minerals (i.e., Mg, Ca, Zn, Fe, Mn) [23]. Efficiency ratios can be influenced by the duration of the crop, fertilization, amount of solar radiation and drought [14,29]. Therefore, comparison of ratios among species or cultivars and across environments or management packages should be conducted with caution [30,31]. In this experiment, the cultivars were grown under identical conditions; however, future experimentation using a more limited nutrient supply would confirm the superior nutrient efficiency of cultivars under field conditions.

4. CONCLUSIONS

There has been limited number of experiments in the Pacific characterizing the inter-relationship between growth, development and nutrient uptake of the taro crop. However, as the demand for taro increases in the local, processing and export markets, the required volume will only be met through extensive plantings using modern management packages.

Implementation of such technological packages will require readdressing the current cultural and management practices and basic research to achieve higher yields.

The results of this study exhibited the inherent cultivar differences in relation to patterns of dry matter accumulation in various components of the taro plant.

The results of this study also revealed that both of the locally bred taro cultivars from Samoa are capable of absorbing a wide range of minerals with relevance to human dietary allowances and health. A complete information package on the nutritional composition of local taro germplasm would help to guide policy makers, nutritionist and researchers in incorporating the crop cultivars into the various diversification programs.

This investigation revealed that overall, cultivar Samoa 1 had had a relatively better nutrient use efficiency than cultivar Samoa 2. On the basis of this finding, Samoa 1 is better adapted for marginal to rich soils while Samoa 2 for moderate to rich soils.

Results from this investigation can be valuable for breeding programs dealing with improvements in taro nutrient use efficiency as well as nutritional composition.

ACKNOWLEDGEMENTS

The authors would hereby like to extend their profound gratitude to the following organization/ people for their generous assistance:

1. Australian Centre for International Agricultural Research (ACIAR) for funding the research.
2. Mr. Ami Chand Sharma and Mr. Samuelu Saulia for assisting in data collection and laboratory analyses work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Guinto DF, Lauga S, Dauara L, Walasi E, Autufuga D, Perera H, Seuoti D, Tauati S. Soil health assessment of taro (*Colocasia esculenta*) farms in Samoa. In: Moving farm systems to improved nutrient attenuation. (Eds. LD Currie, LL Burkitt) Occasional Report No. 28. Fertilizer and Lime Research Centre, Massey University, Palmerston North, New Zealand. 2015;7. Available: <http://flrc.massey.ac.nz/publications.html>
2. Tuivalalagi NS, Hunter DJ, Amosa F. Tackling land degradation and unsustainable land use in Samoa – with emphasis on agriculture sector. Proceedings of the National Environmental Forum held in Apia, Samoa. 2004;32-36.
3. Maathuis K, van Meer, H. Spatial evaluation of saturated hydraulic conductivity and soil erosion in Alafua catchment Samoa. MSc. thesis submitted to the Wageningen University; 2003.
4. Samoa Bureau of Statistics. Agriculture Census Tabulation Report. Econ. Stats. Div. Govt. of Samoa; 2009.
5. FAO. Roots, tubers, plantain and bananas in human nutrition. Food and Nutrition Series, No. 24. Food and Agriculture Organisation of the United Nations, Rome; 2007. ISBN: 9789251028629, P182.

6. Mcgee H. On food and cooking. Scribner; 2014.
ISBN 978-0-684-80001-1
7. South Pacific Commission. Taro: South Pacific foods leaflet 1. Community Educational Material Training Centre, Suva, Fiji; 2013.
8. Bradbury JH, Holloway WD. Chemistry of tropical root crops: Significant for nutrition and agriculture in pacifics. ACIAR Monograph. 2001;6:201.
9. Englberger L, Schierle J, Kraemer K, Aalbersberg W, Dolodolotawake, U. Carotenoid and mineral content of micronesian giant swamp taro (*Cyrtosperma*) cultivars. J. Food Compos. Anal. 2008;21:93-106.
10. Wills RBH, Lim JSK, Greenfield H, Bayliss-Smith T. Nutrient composition of taro (*Colocasia esculenta*) cultivars from the Papua New Guinea highlands. J. Sci. Food Agric. 2003;34:1137-1142.
11. Njoku PC, Ohia CC. Spectrophometric estimation studies of mineral nutrient in three cocoyam cultivars. Pak. J. Nutr. 2007;6:616-619.
12. Lebot V, Prana M, Kreik N, V. van Heck V, Pardales J. Characterisation of taro (*Colocasia esculenta* (L.) Schott) genetic resources in Southeast Asia and Oceania Genetic Resources. Crop Evol. 2004;51: 381-392.
13. Guchhait S, Bhattacharya A, Pal S, Mazumdar D, Chattopadhyay A, Das AK. Quality evaluation of cormels of new germplasm of taro. Int. J. Vegetable Sci. 2008;14:304-321.
14. Goenaga R, Chardon U. Nutrient, uptake, growth and yield performance of three taniar (*Xanthosoma* spp.) cultivars grown under intensive management. J. Agri. Univ. P.R. 1995;78:78-98.
15. Goenaga R, Chardon, U. Growth, yield and nutrient of taro grown under upland conditions. J. of Plant Nutr. 2008;18:1037-1048.
16. Wang ZH, Li SX, Malhi S. Effects of fertilization and other agronomic measures on nutritional quality of crops. J. Sci. Food Agric. 2008;88:7-23.
17. Asher C, Grundon N, Menzies N. How to unravel and solve soil fertility problems. ACIAR Monograph No. 83. 2002;139.
18. IBSNAT. Field and laboratory methods for IBSNAT. Department of Agronomy and Soil Science, College of Tropical Agriculture and Human Resources, University of Hawaii, Honolulu, Hawaii. Technical Report 2. 1987;37-38.
19. Chapman HD, Pratt PF. Methods for analysis of soils, plants and water. Division of Agricultural Sciences, University of California, Riverside, California; 1961.
20. Prasad M, Spiers M. Comparative study of ashing techniques for the digestion of horticultural plant samples. Agri. and Food Chem. 1978;26:824-827.
21. VSN International Ltd. Genstat discovery edition. Rothamsted Experimental Station; 2011.
Available: <http://discovery.genstat.co.uk>
22. Osorio NW, Shuai X, Miyasaka, Wang B, Shirey RL, Wigmore WJ. Nitrogen level and form affect taro growth and nutrition. Hort. Science. 2003;38:36-40.
23. Anand S, Developing a taro (*Colocasia esculenta*) production system based on genotype and fallow system for economic and environmental sustainability under local conditions in Samoa. The University of the South Pacific. Doctoral dissertation; 2016.
24. Norman MJT, Pearson CJ, Searle PGE. Sweet potato (*Ipomea batatas*), In: The Ecol. of Trop. Food Crops. Cambridge University Press, Cambridge, Great Britain. 1994;245-257.
25. Mergedus A, Kristl J, Ivancic A, Soba A, Sustar V, Krizan, T, Lebot V. Variation of mineral composition in different parts of taro (*Colocasia esculenta*) corms. Food Chem. 2014;170:37-46.
26. Mwenye OJ, Labuschagne MT, Herselman L, Benesi IRM. Mineral composition of Malawian cocoyam (*Colocasia esculenta* and *Xanthosoma sagittifolium*) Genotypes. J. of Bio Sci. 2011;11:331-335.
27. Hartemink AE, Johnstone M, O'Sullivan JN, Bloma S. Nitrogen use efficiency of taro and sweet potato in the humid lowlands of Papua New Guinea. Agri. Ecol. and Environ. 2000;79:271-280.
28. Saud BK, Alam S, Narzary BD. Integrated nutrient management of upland taro in Assam. Journal of Root Crops. 2013;39(2):38-47.
29. Chianu JN, Tsujii N. Integrated nutrient management in the farming systems of the savannas of northern Nigeria: What future? Outlook Agriculture. 2005;34:19- 202.

30. Lebot V. Tropical Root and Tuber Crops: Cassava, Sweet Potato, Yams and Aroids. CABI. Wallingford, Oxfordshire. 2009;234.
31. John KS. Soil Fertility Management Strategies in Edible Yams and Aroids: A Review. J. of Root Crops. 2011;37:3-18.

© Copyright 2019 The Author(s), Licensee Book Publisher International, This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

DISCLAIMER

This chapter is an extended version of the article published by the same authors in the following journal with CC BY license. Journal of Agriculture and Ecology Research International, Vol 11, Issue 4, page numbers 1-11.

Reviewers' Information

- (1) Wafula Nelson, Kenyatta University, Kenya.
- (2) Bucagu Charles, University of Rwanda, Rwanda.
- (3) Suqing Li, Shanxi University, China.
- (4) Ibok Oduro, Kwame Nkrumah University of Science and Technology, Ghana.
- (5) Mary O. Iwuagwu, Abia State University, Uturu, Nigeria.