

## SYSTEMATICS AND PHYLOGENY

# Contemporary integrative taxonomy for sexually deprived protists: A case study of *Trachelomonas* (Euglenaceae) from western Ukraine

Anže Žerdoner Čalasan,<sup>1,2</sup>  Juliane Kretschmann<sup>1</sup> & Marc Gottschling<sup>1</sup> 

<sup>1</sup> Department Biologie, Systematische Botanik und Mykologie, GeoBio-Center, Ludwig-Maximilians-Universität München, Menzinger Str. 67, 80 638 Munich, Germany

<sup>2</sup> Fachbereich 5: Biologie/Chemie, Botanik, Universität Osnabrück, Barbarastr. 11, 49 076 Osnabrück, Germany

Address for correspondence: Marc Gottschling, [gottschling@bio.lmu.de](mailto:gottschling@bio.lmu.de)

DOI <https://doi.org/10.1002/tax.12206>

**Abstract** As many other protist groups, euglenophytes are prone to false identification based solely on morphology because of a limited amount of morphological features and cryptic speciation. One of the supposedly completely asexual groups within the freshwater phototrophic representatives of euglenophytes is *Trachelomonas*, capable of forming an inorganic shell around its cell (i.e., the lorica). *The International Code of Nomenclature for algae, fungi, and plants* regulates the taxonomy not only of flowering plants, but explicitly also of phototrophic protists, and provides powerful tools to resolve various taxonomic challenges. To exemplify some of the problems and potential solutions, a number of *Trachelomonas* strains were collected from the muddy, lake-rich region of Dobrostan and cultivated under stable laboratory conditions. Being a type locality of 58 unclarified *Trachelomonas* names, this region in western Ukraine is of great taxonomic importance. Based on light and electron microscopy, and on RAxML and MrBayes phylogenetics using multiple loci and a representative taxon sample, a detailed description of investigated strains and their systematic placement is provided. Morphologically, the strains differed slightly but consistently in minute characters such as size, lorica shape and ornamentation. The presently most comprehensive molecular tree of the Euglenaceae indicated to the existence of at least five different species present in the newly investigated samples, although they were collected from localities in very close vicinity to each other and at the same date. Based on morphological comparisons with type illustrations of species validly described 100 or more years ago, biological material was used to epitypify three names of *Trachelomonas*, eternally linking morphology with reliable genetic information. This taxonomic application is one of the powerful methods to clarify ambiguous scientific names, which has particular importance in character-poor protists such as the euglenophytes.

**Keywords** asexual organism; epitypification; Euglenida; Euglenophyta; molecular phylogenetics; morphology; ribosomal RNA; type locality

**Supporting Information** may be found online in the Supporting Information section at the end of the article.

## ■ INTRODUCTION

Unambiguous scientific names are the prerequisite for proper identification and, therefore, any subsequent application of biological species. The taxonomy not only of flowering plants, but also of phototrophic protists, is regulated by the *International Code of Nomenclature for algae, fungi, and plants* (ICN; Turland & al., 2018). Unicellular organisms, which can only be identified under the microscope, are particularly prone to taxonomic confusion (De Clerck & al., 2013; Manoylov, 2014), and corresponding “standard” (i.e., type) material is often poorly preserved or entirely lost. Diverging methodological approaches have led to some deviations of phycological research within botany, which is also expressed by explicitly disregarding Rec. 46A of the *Code* (i.e., author abbreviations of scientific names) in a number of distinguished phycological journals. However, effective, sustainable

and approved algal taxonomy can be assured only by a powerful federal organisation such as the International Association for Plant Taxonomy (IAPT) with its periodical *Taxon*.

The exemplary subject of this study to demonstrate a good practice to resolve taxonomic challenges in protists are euglenophytes. These represent an ecologically and economically important group of unicellular plankton organisms, as they include numerous primary producers but can cause massive algal blooms, resulting in fish culture losses (Kulczycka & al., 2018). *Trachelomonas* Ehrenb. comprises of not less than 1700 uncritically listed scientific names at the species level and below (Guiry & Guiry, 2017). Together with *Colacium* Ehrenb., *Cryptoglena* Ehrenb., *Discoplastis* Triemer, *Euglena* Ehrenb., *Euglenaformis* M.S.Bennett & Triemer, *Euglenaria* Karnkowska-Ishikawa & al., *Eutreptia* Perty, *Eutreptiella* A.G.Cunha, *Lepocinclis* Perty, nom. cons. (Senn in Briquet, 1930; Silva, 1960; Ross, 1966), *Monomorphina*

**Article history:** Received: 10 Jun 2019 | accepted: 16 Dec 2019 | published online: 10 Apr 2020 | **Associate Editor:** Øjvind Moestrup | © 2020 The Authors.

TAXON published by John Wiley & Sons Ltd on behalf of International Association for Plant Taxonomy.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Mereschk., *Phacus* Dujard., nom. cons. (Silva, 1960; Ross, 1966) and *Strombomonas* Deflandre, *Trachelomonas* represents an integral element of the freshwater Euglenophyceae as inferred from molecular phylogenetics (Kim & al., 2015; Bicudo & Menezes, 2016).

The striking apomorphy of all euglenophytes is the pellicle. It is a unique surface structure, comprised of a microtubular system, endoplasmic reticulum cisterns and a plasma membrane, connected with protein layers (Arnott & Walne, 1967). In addition, euglenophytes are capable of producing and secreting an additional mucilage layer (Dunlap & Walne, 1985). In some cases, such as *Strombomonas* and *Trachelomonas*, this extracellular matrix becomes thicker and gets impregnated with inorganic substances, forming an inorganic shell termed lorica (Pringsheim, 1953; Leedale, 1975). Chemical analyses show that the main inorganic compounds found in loricae of *Trachelomonas* are iron and manganese (Rino & Pereira, 1991), which are together with a great deal of other elements incorporated in the lorica via non-active physical processes (Pereira & al., 2003). Depending on its mineral composition, the colour of the lorica varies from translucent to brown or reddish-brown (Pringsheim, 1953).

Within Euglenaceae, state reconstruction indicates that a lorica is a synapomorphic feature shared by the sister groups *Strombomonas* and *Trachelomonas* (Karnkowska & al., 2015). *Trachelomonas* can be distinguished from *Strombomonas* based on lorica ontogeny: *Strombomonas* loricae namely always develop progressively from the anterior towards the posterior part of the cell, whereas the formation of *Trachelomonas* loricae takes place evenly across the entire cell (Brosnan & al., 2005). Additionally, *Strombomonas* loricae do not possess pores, whereas *Trachelomonas* loricae are always ornamented (i.e., possessing pores and/or spines) at least at one stage of their development (Brosnan & al., 2003). A diversity of spines and pores on the lorica surface, and their arrangement, have been thought to be of great taxonomic importance, not only at species but also on infraspecific levels of *Trachelomonas* (Rino & Pereira, 1988). However, environmental factors can alter lorica traits significantly (Nudelman & al., 2003), and its morphology also depends on the protoplast metaboly, as it starts to develop in later ontogenetic stages (Brosnan & al., 2005). Thus, such morphological characteristics should be treated with great caution when describing a certain species, as the classification based on lorica morphology alone is often inadequate and misleading (Pringsheim, 1953; Ciugulea & al., 2008).

Biodiversity assessment of euglenophytes and *Trachelomonas* is perplexing due to several reasons. Combining both phototrophic and heterotrophic representatives, this group is an example of the ambireginal taxonomic treatment under both, the *ICN* (Turland & al., 2018) and the *International Code of Zoological Nomenclature* (*ICZN*; Ride, 2003). The diversity of biological phenomena further prevents us from an unambiguous definition of what a biological species actually is. Euglenophyte reproduction, and thus also of *Trachelomonas*, is considered to be almost entirely asexual (Leedale, 1967), and such organisms may be prone to greater mutability because of the missing

regulatory system of crossing over. Nevertheless, asexual clones still possess a consistent set of traits, which are the result of natural selection, and in this manner resemble organisms with sexual reproduction (Gottschling, 2008). However, asexually propagating organisms such as cyanobacteria and bdelloid rotifers are still equally ranking in taxonomy with “regular” biological species.

During the past decades, molecular DNA sequence data have gained great importance for taxonomic delimitations in the microbial world including *Trachelomonas*. As in many other protist species, cryptic speciation is also observed here (Kim & al., 2013; Przybos & Tarcz, 2016), resulting in a number of morphologically indistinguishable species. In turn, it is also highly plausible that size variations and different ecological forms of the same species have been described as several distinct species due to their plasticity (Bicudo & Menezes, 2016). Thus, diversity of *Trachelomonas* has been continuously under- and overestimated at the same time.

Because of cryptic speciation, it is important to link DNA sequence data to a type specimen, but the original material of microscopic organisms consists in many (particularly older) cases of illustrations only. Since protists frequently lack diagnostic morphological features (Pawlowski & al., 2012), type specimens (including illustrations) cannot provide for unambiguous identification of certain species. As one is not able to connect a type illustration to a DNA sequence, the same problem arises, as if the type material is in poor condition. In such cases, it is recommended to designate an epitype via epitypification (Hyde & Zhang, 2008). This powerful tool has already been applied to solve taxonomic ambiguities in various protistan groups such as green algae (Demchenko & al., 2012) and dinophytes (Zinßmeister & al., 2011; John & al., 2014; Kretschmann & al., 2015, 2018a,b; Tillmann & al., 2017) but also in euglenophytes (Marin & al., 2003; Kosmala & al., 2005; Bennett & Triemer, 2012; Karnkowska-Ishikawa & al., 2012; Kim & Shin, 2014).

For all the above reasons, one is faced with big taxonomic confusion in euglenophytes (Brosnan & al., 2003, 2005). In this study, we provide a robust molecular phylogeny of Euglenaceae and clarify the taxonomy of three names within *Trachelomonas* (all of which have rarely been applied in the past: Starmach, 1983; Borisova & al., 2006). Fifty-eight taxa at the species level and below have been described from the lake district near Dobrostan west of Lviv (suppl. Table S1), making it a taxonomic hot spot of great importance. In the beginning of the 20th century, this lake district has comprised primarily of three small ponds, namely Wolicki pond in the North, centrally located Dobrostan pond and Białogórski pond in the South (Koczwara, 1915). Already available data of Euglenaceae are thus further enriched based on the material collected in western Ukraine, the most southern region ever affected by the maximum Scandinavian glaciations (Lindner & al., 2004). The utmost western part of the country is part of a so-called forest-steppe region, characterised by grey soils and covered with either steppe vegetation or mixed forests, dominated by oak (Borisova & al., 2006). Due

to massive human impact, the majority of water bodies are highly eutrophic or eutrophic-polytrophic (Konenko & al., 1965), representing suitable living conditions for euglenophytes. The region is also characterised by radioactive contamination of water bodies, last but not least because of the accident at the Chernobyl Nuclear Power Plant (Kuzmenko, 2000). Euglenophytes are resistant to moderate radiation (Margulis & al., 1990), and, therefore, we expected the investigated species to still be present on their type localities, from which they have been described as long as a century ago.

## ■ MATERIALS AND METHODS

**Sampling and cultivation.** — During a field trip, water tow samples were collected with a plankton net (with pores' diameter 20  $\mu\text{m}$ ) at several different localities – in Poland on 10 September 2012, and in Ukraine on 15 and 16 September 2012. Single, motile, greenish through brownish cells were isolated using a glass micropipette under a CKX41 inverted microscope (Olympus; Hamburg, Germany). To avoid contaminations, the isolated cells were rinsed at least three times in WC-Medium, prior to cultivation (Guillard & Lorenzen, 1972). Single or few cells were then transferred in six-well microplates (Zefa; Munich, Germany), containing 5 ml of modified WC-Medium (Guillard & Lorenzen, 1972) and 1 ml of soil extract (McFadden & Melkonian, 1986). Plates were stored in a climate chamber Percival I-36VL (CLF Plant Climatics; Emersacker, Germany) at 18°C, at a light intensity of 80  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$  and under a 12 : 12 h light : dark photoperiod.

To maintain the established strains, they were supplied with fresh medium every four weeks by being transferred into a new well. Prior to adding 2 ml of fresh medium to the strain, the same volume of the old medium was discarded using a syringe equipped with a disposable sterile filter and a pore size of 0.2  $\mu\text{m}$  (Sartorius; Göttingen, Germany). In order to avoid further contamination with possible bacteria and fungi, medium transfer was done under a clean bench (ET 130/H, Ehret; Emmendingen, Germany). Once dense material of *Trachelomonas* strains was obtained, a part of the grown strain was transferred into a new 1.5 ml collection Eppendorf tube (Eppendorf; Hamburg, Germany), centrifuged and stored in the freezer at –20°C for DNA isolation.

**Light and electron microscopy.** — For morphological analysis, 80 cells of every strain were separated and individually measured (length and width) with a CKX41 inverted microscope (Olympus) equipped with a DP73 digital camera (Olympus). Measurements were represented by mean values and standard errors and were statistically evaluated by one-way analysis of variance (ANOVA), using SigmaPlot v.11.0 (Systat Software; San Jose, California, U.S.A.). Significant differences between mean values were determined using the post-hoc Dunn's test (Dunn, 1961). Only mature cells exhibiting lorica were measured, to minimise the measuring errors as much as possible.

For Scanning Electron Microscopy (SEM), cultivated material was fixed in 4% OsO<sub>4</sub> (1 : 1 ratio, resulting in a final

concentration of 2% OsO<sub>4</sub>) for at least 1 h. The cells were then filtered onto an Omnipore-membrane filter (5  $\mu\text{m}$ ; Merck; Darmstadt, Germany), which was placed in a Swinnex filter holder (Merck, Millipore; Darmstadt, Germany). The preparation liquids were exchanged with the aid of a plastic syringe connected to the above-mentioned filter holder. Fixed cells were rinsed four times; first in sodium cacodylate buffer for 15 min and then three times in distilled water (the first time for 5 min, second time for 15 min and third time for 30 min). Next step included subsequent dehydration in a graded acetone series of 15 min in 10%, 30%, 50%, 70% and 90% acetone, followed by three additional dehydration steps in 100% acetone (the first time for 5 min and two times for 30 min). The membrane filters were critical-point dried in liquid CO<sub>2</sub> and afterwards glued to a SEM-stub with double-adhesive carbon disks. Additionally, small drops of Planocarbon N650 (Plano; Wetzlar, Germany) were applied on the edges of the filter to improve the electrical contact between the filter and the stub. The filter was coated with platinum (BAL-TEC SCD 050 sputter coater; Schalksmühle, Germany) for 4 min and examined in a LEO 435VP SEM (LEO Electron Microscopy; Cambridge, U.K.).

For epitype preparation, 1 ml of the corresponding strain was taken and centrifuged in a 1.5 ml Eppendorf-tube at a maximum speed of 200g (Z320, Hermile; Gosheim, Germany) for 30 min to make the sample denser. The cells were then fixed in 2.5% glutaraldehyde in 75 mM NaCacodylate-buffer (C<sub>2</sub>H<sub>6</sub>AsNaO<sub>2</sub>·3H<sub>2</sub>O) containing 2 mM MgCl<sub>2</sub> for at least 1 h. Afterwards, the cells were stained in 0.5% astra blue and 2% tartaric acid dissolved in WC-Medium for 30 min. After staining, a cleaning step in WC-Medium and dehydration in a graded ethanol series (30%, 50%, 70% and 90%, followed by two changes to 100%; each time for 15 min) was performed. A second staining with 0.1% (ethanol-based) eosin was carried out during the incubation in 90% ethanol for 30 min. Ethanol-based Technovit 7100 (Heraeus; Wehrheim, Germany) was used for embedding, following the manufacturer's instructions. For the final preparation, 40  $\mu\text{l}$  aliquots of the Technovit mixture including the embedded samples were transferred to three microscopic glass slides and were enclosed by a cover slide.

**Molecular phylogenetics.** — Genomic DNA was extracted from fresh material using the Nucleo Spin Plant II Kit (Machery-Nagel; Düren, Germany). Two nuclear regions of the rRNA were amplified, using primers and following the standard protocols (Hansen & al., 2000; Kim & al., 2010). The taxon sample included all Euglenaceae taxa (excluding *Euglena* itself, see Discussion), whose sequences were available for at least two out of three nuclear loci (or parts of loci; Appendix 1). The only exception is *Trachelomonas* cf. *volvocinopsis* Svirenko (only SSU), which was used as a reference sequence (based on a BLAST search: Altschul & al., 1990) for the strain GeoM\*521 established in this study. The sequences were separately aligned, using MAFFT v.7.304 (Katoh & Standley, 2013; freely available at <http://mafft.cbrc.jp/alignment/software/>) under default settings, and concatenated afterwards (Appendix S1).

Phylogenetic analyses were carried out with maximum likelihood (ML) and Bayesian approaches, as described in detail

previously (Gottschling & al., 2012), using the resources available from the CIPRES Science Gateway (Miller & al., 2010). The Bayesian analysis was performed using MrBayes v.3.2.6 (Ronquist & al., 2012; freely available at <http://mrbayes.sourceforge.net/download.php>) under the GTR +  $\Gamma$  substitution model and the random-addition-sequence method with 10 replicates. Two independent analyses of four chains (one cold and three heated) with 15 million cycles were run, sampled every 1000th cycle, with an appropriate burn-in (10%) as inferred from the evaluation of the trace files using Tracer v.1.6 (freely available at <http://tree.bio.ed.ac.uk/software/tracer/>). For the ML calculation, the MPI version of RAxML v.8.0.24 (Stamatakis, 2014; freely available at <http://www.exelixis-lab.org/>) was applied by using the GTR +  $\Gamma$  substitution model. To determine the best-fitted ML tree, 10 tree searches from distinct random-stepwise-addition-sequence maximum parsimony starting trees were executed and 1000 non-parametric bootstrap replicates were performed. Statistical support values (LBS: ML bootstrap support, BPP: Bayesian posterior probabilities) were drawn on the resulting, best-scoring tree. The phylogenetic tree was rooted posteriorly with *Cryptoglena* and *Monomorphina*.

## ■ RESULTS

**General morphological observations.** — All six investigated strains differed morphologically between each other based on lorica shape and ornamentation. Another morphological trait that was taken into consideration is size, since some strains were significantly larger than others, considering length and width (Figs. 1, 2, suppl. Tables S2, S3). Generally, the motile stage was predominant in all investigated strains not showing a considerable contamination, but immotile stages were more frequent in heavily contaminated sub-strains. Cell replication and lorica formation were enhanced with an addition of a soil extract. Dividing motile cells formed big cell clumps, which were either moving extremely slowly or were lying on the bottom of the cultivation plates.

Three developmental stages were observed in the life-history of a strain, namely small and transparent cells, bigger and partially transparent cells and non-transparent brown cells, which did not differ greatly in size in comparison to the partially transparent cells. For clearer terminology, the cells were given terms “immature cells” (Figs. 3A,E,I, 4M,Q,U), “mature naked cells” (Figs. 3B,F,J, 4N,R,V) and “mature loricate cells” (Figs. 3C,G,K, 4O,S,W); empty loricae are depicted in the fourth column of every strain illustration (Figs. 3D,H,L, 4P,T, X). These terms are not used in the strains’ descriptions, as they present an artificial classification and as the description itself is built on more than one stage of life-history.

Regardless of the ontogenetic stage, the majority of the cells possessed a clearly distinctive eye spot, which was located always at the anterior end of the cell, near the flagellar apparatus. This structure also served as an anchor point to orientate the cells into the same direction. Length of the visible flagellum varied greatly between strains. Moreover, the cells were densely filled with

granules of varying sizes and chloroplasts, which were hard to distinguish from the rest of granular structures, in mature loricate cells.

**Strain descriptions.** — *GeoM 520* (Figs. 3A–D, 5A–F). — Cell colour yellowish to greenish-brown; lorica ellipsoid, 19.88 ( $\pm 1.66$ )  $\mu\text{m}$  long, 16.01 ( $\pm 1.19$ )  $\mu\text{m}$  wide, the surface finely punctate, with uniformly and somewhat densely covered short, conical spines; apical pore without a collar or with a short, slightly raised margin; flagellum at least twice the length of the cell; chloroplasts up to 5, plate-like, green, including paramylon cups accompanying the diplopyrenoids; eye spot orange, visible in most cells, except in mature loricate cells.


*GeoM\* 524* (Figs. 3E–H, 6A). — Cell colour yellowish to brownish, greenish only during division; lorica spherical through ellipsoid, 24.85 ( $\pm 3.71$ )  $\mu\text{m}$  long, 21.45 ( $\pm 2.83$ )  $\mu\text{m}$  wide, the surface densely, deeply and somewhat uniformly reticulate without spines (reticulation could be a result of an uncompleted lorica formation); apical pore without a collar (in SEM images) and with a distinctive collar (in some LM images); flagellum length variable; chloroplasts up to 5, plate-like, green, including paramylon cups accompanying the diplopyrenoids; eye spot orange to orange reddish, visible in most cells, except in mature loricate cells.

*GeoM 526* (Figs. 3I–L, 6B). — Cell colour yellow-greenish to greenish-brown; lorica broadly ellipsoid, 24.23 ( $\pm 2.45$ )  $\mu\text{m}$  long, 19.53 ( $\pm 1.79$ )  $\mu\text{m}$  wide, the surface punctate; apical pore with a distinctive diagonally ended collar (seen only in naked cells, but not in loricate ones); flagellum at least twice as long as the cell; chloroplasts up to 5, plate-like, green, including paramylon cups accompanying the diplopyrenoids; eye spot bright-orange to orange-dark reddish, visible in most cells, regardless of cell maturity.


*GeoM 527* (Figs. 4M–P, 5G–L). — Cell colour dark greenish to dark greenish-brown; lorica very widely ellipsoid through almost spherical, 19.98 ( $\pm 2.19$ )  $\mu\text{m}$  long, 16.08 ( $\pm 1.78$ )  $\mu\text{m}$  wide, the surface irregularly and sparsely punctate; apical pore without a collar; flagellum at least three times as long as the cell, at least in younger cell stages; chloroplasts up to 5, plate-like, yellowish-green including paramylon cups accompanying the diplopyrenoids; eye spot red to reddish, visible in all cells, regardless of cell maturity.

*GeoM 529* (Figs. 4Q–T, 6C). Cell colour bright greenish, greenish through dark greenish-brownish; lorica ellipsoid, 24.46 ( $\pm 2.27$ )  $\mu\text{m}$  long, 20.79 ( $\pm 2.24$ )  $\mu\text{m}$  wide, the surface uniformly, densely and deeply reticulate, covered sparsely with sharp spines; apical pore without collar or with short, slightly raised margin; flagellum at least twice as long as the cell; chloroplasts up to 5, plate-like, green, including paramylon cups accompanying the diplopyrenoids (hard to observe, since the granular structures in the mature naked cells have almost the same colour as the chloroplasts); eye spot big, orange to dark-reddish, visible in all cells, except in mature loricate cells.


*GeoM 540* (Figs. 4U–X, 5M–R). Cell colour yellow-brownish to almost grey-brownish; lorica ellipsoid, 18.79 ( $\pm 1.76$ )  $\mu\text{m}$  long, 15.57 ( $\pm 1.50$ )  $\mu\text{m}$  wide, the surface uniformly and densely reticulate; apical pore without collar or




**Fig. 1.** Box plot display of cell length of investigated mature naked cells and mature loricate cells of *Trachelomonas* strains. Statistically significant clusters are indicated with letters a or b and green or red colour, corresponding to the phylogenetic tree (see Fig. 7).



**Fig. 2.** Box plot display of cell width of investigated mature naked cells and mature loricate cells of *Trachelomonas* strains. Statistically significant clusters are indicated with letters a or b and green or red colour, corresponding to the phylogenetic tree (see Fig. 7).



**Fig. 3.** Different ontogenetic stages and empty loricae of selected *Trachelomonas* strains (LM; all at the same scale). **A–D**, Young immature cell, mature naked cell, mature loricate cell and empty lorica of *Trachelomonas hispida* var. *volicensis* GeoM 520; **E–H**, Young immature cell, mature naked cell, mature loricate cell and empty lorica of *Trachelomonas* sp. GeoM\*524; **I–L**, Young immature cell, mature naked cell, mature loricate cell and empty cell and empty lorica of *Trachelomonas* sp. GeoM 526.




**Fig. 4.** Different ontogenetic stages and empty loricae of selected *Trachelomonas* strains (LM; all at the same scale). **M–P**, Young immature cell, mature naked cell, mature loricate cell and empty lorica of *Trachelomonas teres* var. *minor* GeoM 527; **Q–T**, Young immature cell, mature naked cell, mature loricate cell and empty lorica of *Trachelomonas hispida* var. *irregularis* GeoM 529; **U–X**, Young immature cell, mature naked cell, mature loricate cell and empty lorica of *Trachelomonas teres* var. *granulata* GeoM 540.

with short, slightly raised margin (seen only in mature naked cells under SEM); flagellum at least twice as long as the cell; chloroplasts up to 5, plate-like, green, including paramylon cups accompanying the diplopyrenoids; eye spot orange through reddish, visible in most cells.

**Molecular phylogenetics.** — Newly obtained sequences were deposited in the NCBI GenBank under MK894259–MK894271. The alignment comprised 4150 parsimony-informative positions, and the proportion of gaps and ambiguous characters was 57.12% (for additional alignment traits, see suppl. Table S3). Our dataset provided a well-resolved backbone phylogeny of a representative taxon sample of selected photosynthetic euglenacean clades (Fig. 7). Maximum

likelihood and Bayesian analyses recovered phylogenetic trees with congruent topologies. Five monophyletic groups were recovered: *Colacium* (100 LBS, 1.00 BPP), *Cryptoglana* (100 LBS, 1.00 BPP), *Monomorphina* (100 LBS, 1.00 BPP), *Strombomonas* (100 LBS, 1.00 BPP) and *Trachelomonas* (89 LBS, 1.00 BPP). Both loricate taxa *Strombomonas* and *Trachelomonas* constituted sister groups (97 LBS, 1.00 BPP).

*Trachelomonas* segregated into a number of well-supported lineages at high taxonomic level. As such lineages could not be correlated with non-DNA sequence traits, they are informally indicated with Greek letters (Fig. 7). The type of *Trachelomonas*, *T. volvocina* (Ehrenb.) Ehrenb., is nested with clade  $\alpha$  (99 LBS, 1.00 BPP), which also one of our strains



**Fig. 5.** LM, epitype, SEM and protologue images of mature *Trachelomonas* cells from selected strains. **A–F**, Mature loricate cells (**A & B**), epitype images (**C & D**), SEM image (**E**) and protologue (**F**) of *Trachelomonas hispida* var. *volucensis* GeoM 520; **G–L**, Mature loricate cells (**G & H**), epitype images (**I & J**), SEM image (**K**) and protologue (**L**) of *Trachelomonas teres* var. *minor* GeoM 527; **M–R**, Mature loricate cells (**M & N**), epitype images (**O & P**), SEM image (**Q**) and protologue (**R**) of *Trachelomonas teres* var. *granulata* GeoM 540.

(GeoM\*521) was assigned to. However, this strain represented a phylogenetically isolated branch shared with strain M1422 (determined as *T. cf. volvocinopsis*). All other our strains were nested within clade  $\beta$  (100 LBS, 1.00 BPP), which included both significantly small and large representatives. Strains GeoM\*524, GeoM 526 and GeoM 529 clustered together with other large species such as *T. hispida* (Perty) F.Stein, *T. pertyi* E.G.Pringsh. and *T. variabilis* Kam.P.Singh, while strains GeoM 520, GeoM 527 and GeoM 540 build a clade together with small-celled *T. echinata* A.M.Cunha (100 LBS, 1.00 BPP). Notably, strains GeoM 520 and GeoM 540 (and also SAG 1283-22, determined as *T. echinata*) shared identical SSU and LSU sequences, but possessed different morphological features (see above).

## ■ DISCUSSION


**Molecular phylogenetics and at least a few correlations to morphology.** — The study of the euglenophytes in general and *Trachelomonas* in particular is challenging because of various biological phenomena such as elusive diagnostic traits, intraspecific modifications because of environmental influence, cryptic speciation and last but not least due to the absence of sexual reproduction (Leedale, 1967; Kim & al., 2013; Przybos & Tarcz, 2016; Wołowski & al., 2016). Subsequently, depicting driving forces and dynamics of the molecular evolution in *Trachelomonas* is extremely problematic, causing an extensive taxonomic mess still persisting within numerous protist lineages. In such cases, an integrative approach considering morphological and molecular data combined (as applied in the present study) is advised.

The necessary prerequisite to infer evolutionary dynamics is a well-supported phylogenetic tree, and the concatenation of loci generally improves the reconstructions (not only in euglenophytes: Brosnan & al., 2003, Kim & al., 2015, but also in other protist groups such as the dinophytes: Gottschling & al., 2012, 2020; Tillmann & al., 2014; Kretschmann & al., 2018a,b). This approach reveals statistically highly supported

topologies, which is due to a proportionally high share of parsimony-informative sites (suppl. Table S4). Anyhow, a persistent limitation hindering phylogenetic reconstructions in the microbial world is the taxon sample, being biased towards phototrophic and therefore easily cultivatable taxa. Only a single (of few) heterotrophic presenter is included into the phylogenetic analysis, namely *Trachelomonas reticulata* G.A.Klebs, whose phototrophy is believed to have been lost secondarily (Ciugulea & al., 2008). We do not know yet how many times phototrophy has been lost in *Trachelomonas* (or Euglenaceae in general), but corresponding species (groups) are presumably small and nested within their phototrophic relatives. Thus, our analysis still illustrates relationships based on a representative taxon sample of *Trachelomonas* (Kim & al., 2015; Bicudo & Menezes, 2016).


Four *Trachelomonas* subclades are recovered in our analysis, in comparison to five in a previous study (Ciugulea & al., 2008). There is no obligatory criterion for subclade delimitation due to the complex biology of *Trachelomonas*. Our choice to delimitate individual subclades within the phylogeny is based on branch length and bootstrap support, and not on differences in lorica morphology as in Ciugulea & al. (2008). Our reasoning is that lorica morphology is evidently more influenced by the environment than the phylogenetic signal, making the latter one a more reliable source for subclade delimitation. As *Trachelomonas*, *Euglena* also includes phototrophic and heterotrophic presenters as well (Marin & al., 2003; Triemer & al., 2006). Including *Euglena* to the taxon sample resulted in the same topology as shown in Fig. 7, but with lower statistical support of the backbone. *Euglena* is set on a clade exhibiting long branches (phylogenetic tree not shown), which is indicative for strong rate heterogeneity perturbing phylogenetic reconstructions in general. Moreover, *Euglena* possesses a high number of autapomorphic sequence positions, which are impossible to align reliably with the rest of euglenophytes' genetic features.

Based on our well-resolved phylogenetic tree of Euglenaceae, an unexpected phylogenetic signal has been recovered between the DNA tree and cell size. Due to plasticity, size



**Fig. 6.** Loricae showing different shape and ornamentation (SEM; all at the same scale). **A**, Lorica of *Trachelomonas* sp. GeoM\*524; **B**, Lorica of *Trachelomonas* sp. GeoM 526; **C**, Lorica of *Trachelomonas hispida* var. *irregularis* GeoM 529; **D**, Protologue of *Trachelomonas hispida* var. *irregularis*.





**Fig. 7.** Maximum likelihood-phylogenetic tree of Euglenaceae, derived from concatenated 18S rRNA and partial 28S rRNA loci. Major clades are indicated, and strains of newly obtained sequences are pointed out in bold font. Epitypified taxa (or those corresponding to reference material) are marked in bold. Branch lengths are drawn to scale, with the scale bar indicating the number of nt substitutions per site. The numbers on the branches show statistical support values (above: ML bootstrap values, values <50 are not shown; below: Bayesian posterior probabilities, values <.90 are not shown). Asterisks indicate maximal support. Significantly smaller taxa are shaded in green and significantly larger taxa in red. Well-supported lineages that could not be correlated with non-DNA sequence traits are informally indicated with Greek letters.

does not usually play an important role in taxon delimitation in protists, but three (out of six investigated) strains forming significantly smaller cells, cluster together (shaded in light green in Fig. 7). As inferred from outgroup comparison, smaller size appears of an apomorphic nature within *Trachelomonas*. It should be pointed out, however, that cell biometry is reliable only for the strains investigated in this study. Assessments of other material for which sequence data are available are inferred from the names introduced in the original literature (Swirensko, 1914; Koczwara, 1915; Dreżepolski, 1923, 1925; Skvortzov, 1925; Deflandre, 1926). In addition, the considerable amount of biometric data is missing, as most of the *Trachelomonas* accessions (as well as those of other Euglenaceae) included in the phylogenetic analysis are missing their names.

#### Towards clarified taxonomy in *Trachelomonas*. —

Presenting a polyphyletic group, algae are extremely diverse in terms of their biology. This heterogeneity could be at least partially controlled via a robust taxonomy. However, researchers are not obliged to follow the *ICN* regulations, which further hinders us from understanding of the fundamental issues in phylogenetic research (as for example in our case of *Trachelomonas*). Because of the complex biology, loricate euglenophytes represent also a taxonomically very challenging organismal group. Merely, 18% of all described species are currently accepted (Guiry & Guiry, 2017), but the unadjusted synonymy includes likewise many historical names dating back to 100 years ago or more. Essentially, all scientific names in the microbial world dating prior to the DNA era are ambiguous in the sense of *ICN* Art. 9.9, as original material mostly comprises of illustrations (frequently of a single cell) with no DNA sequence information. It is therefore of even greater importance to link the DNA sequence of newly collected strains to original material, since individual species in cryptic complexes cannot be distinguished in any other way. For all the above reasons, any application of historical names is particularly unreliable in *Trachelomonas*, and thus each clarified name is an important step towards a stable taxonomy (Romeikat & al., 2019).

Fifty-eight unclarified *Trachelomonas* names described from either Dobrostrany lakes or from nearby vicinity of Lviv were compiled from the literature in the course of our study (suppl. Table S1). Of such names, approximately half correspond to large taxa (i.e., more than 20 µm in length), but we could not assign any of our larger strains to them due to incompatibilities in shape and ornamentation of the lorica. Excluding criteria for the smaller taxa (i.e., maximally 20 µm in length) have been specialised structures such as a distinct collar and extreme spherical or cylindrical shapes, all absent in our investigated strains. Consequently, we apply *ICN* Art. 9.9. while epitypifying the remaining three such historical names, namely *Trachelomonas hispida* var. *volicensis* Dreżep., *Trachelomonas teres* var. *granulata* Dreżep. and *Trachelomonas teres* var. *minor* Dreżep. (see Taxonomy). In no single case, contradictions to the corresponding protocols have been found based on the present integrative study of the material collected at Ukrainian type localities. A fourth

name can be assigned to the collected material (namely *T. hispida* var. *irregularis* Dreżep.: Fig. 6C,D), but the strain did not originate from the corresponding type locality, and we thus refrain from its epitypification. However, the DNA sequence information assigned to the taxon can be regarded as reference until the final clarification of the name.

Sequences of strain GeoM 527 (corresponding to *T. teres* var. *minor*) are distinct from any other GenBank record, but we refrain from elevating the taxon to species level as long as the identity of *Trachelomonas teres* Maskell from New Zealand is unclarified. Surprisingly, DNA sequence data alone are not always discriminative for species identification in *Trachelomonas*. The two strains GeoM 520 (corresponding to *T. hispida* var. *volicensis*) and GeoM 540 (corresponding to *T. teres* var. *granulata*) share identical SSU and partial LSU sequences. However, both strains possess distinct lorica morphologies, which considerably alter from each other, despite insignificant size differences (Figs. 1, 2, 3A–E, 4U–X, 5A–F, M–R, suppl. Tables S2, S3). Both of the strains have been cultivated in the same medium, during the same time, under the same conditions and have been added soil extract at the same time; therefore, it is unlikely that the morphological differences are of ecological and/or ontogenetic nature, a cultivation artefact or expression of intraspecific variability. We refrain from taxonomic conclusions here as well (i.e., elevation to species rank or synonymisation under, for example, *Trachelomonas echinata* A.M.Cunha) because we cannot irrefutably conclude on its taxonomic status at this moment in time. Those two taxa may present a similar dinophyte “species pair” as brackish *Apocalathium malmogiense* (G.Sjöstedt) Craveiro & al. and freshwater *Apocalathium aciculiferum* (Lemmerm.) Craveiro & al., which are molecularly indistinguishable, yet possess morphological and ecological differences (Gottschling & al., 2005; Kremp & al., 2005; Annenkova & al., 2015).

Considerable effort has been made in the past years to clarify the complex taxonomy of euglenophytes (e.g., Shin & Triemer, 2004; Kosmala & al., 2005, 2007; Łukomska-Kowalczyk & al., 2015). However, there have been several cases in which Art. 7.11 of the *ICN* was violated [“... designation of a type is achieved only ..., if the typification statement includes the phrase ‘designated here’ (hic designatus) or an equivalent”], and subsequently, all such typifications should thus remain neglected (though effectively published from the nomenclatural point of view). Moreover, many typification statements do not provide locality information, which is an indication that those authors doubt its relevance. Thus, it seems like they are supporters of the “everything is everywhere” hypothesis (Finlay, 2002), which has been challenged numerous times in the past and is nowadays rather outdated (Bass & Boenigk, 2011; Bates & al., 2013; Žerdoner Čalasan & al., 2019). We are certain – and this is exemplified by the approach pled in the present study – that taxa, for example, from Ukraine or Germany, should not (readily) be epitypified with material collected in far distant places, such as in England, Portugal or even in the U.S.

The questionable taxonomic approaches are also illustrated by, for example, the typifications of *Phacus longicauda* var. *torta* Lemmerm.: Łukomska-Kowalczyk & al. (2015) chose an illustration as lectotype that was published 16 years after the protologue (Lemmermann, 1910; Skuja, 1926) and is thus not original material in the sense of the *ICN* (Art. 9.4). Lemmermann (1910) referred to an image of Stein (1878), which Łukomska-Kowalczyk & al. (2015) chose for typification purposes of another taxon, namely *Phacus helikoides* Pochm. More thought should be given to the taxonomy of euglenophytes, and our study might stimulate a discussion about more appropriate ways to disentangle the existing historical complexities. The fact that numerous cases of taxonomic activity are not in accordance with the rules of the *ICN* provides also an opportunity, frankly speaking, to revise former, inaccurate decisions regarding typifications in euglenophytes.

In conclusion, three historical names of *Trachelomonas* are taxonomically clarified, and meaningful in-depth studies of such epitypified taxa are from this point onward possible. Based on our taxonomic decisions, we are also able to take intraspecific variability of lorica traits into consideration, confirming a much higher morphological variability than previously thought (Wołowski & al., 2016). This is once more a strong argument for epitypification particularly for the microbial world, as the names clarified here have been rarely applied in the past (Starmach, 1983; Borisova & al., 2006). This study furthermore gives an example of successful taxonomic clarification within protists, putting them side-by-side with macro-organismic groups, in which taxonomy does not represent such a topical issue. Anyhow, we live in an era, where the newly produced amount of data is getting out of hand, and scientific publications in favour of discarding all long-forgotten names, which could cause taxonomic instability (Smith & al., 2016), are a matter of great concern. Nevertheless, not only that authors of such papers simply ignore the fact that taxonomy of different groups of organisms is facing different obstacles; with such a radical approach, they are sabotaging their own research by discarding the work of taxonomists of previous centuries, not realising that their work could be treated in the same way after a period of time. Despite being time consuming, epitypification (if done properly) is still one of the most powerful methods to clarify ambiguous scientific names, which is of particular importance especially in character-poor protists such as the euglenophytes. Integrating both morphological and phylogenetic methods, taking type locality into consideration and carrying out epitypification, this study represents a considerable step forward towards stable and reliable taxonomy also in protists. As epitypification is only available under the *ICN*, we furthermore urge fellow researchers to carry out any kind of taxonomic treatments of ambiregnal taxa under the *ICN*.

## ■ TAXONOMY

*Trachelomonas hispida* var. *volicensis* Drezep. in Kosmos (Lvov) 50: 216, t. 1, fig. 35. 1925 – **Lectotype (designated here):**

[illustration] in Kosmos (Lvov) 50: t. 1, fig. 35. 1925! [showing a non-fossil individual from Republic of Ukraine. Lviv; exact locality and collecting date unknown: *R. Dreżepolski s.n.*] – **Epitype (designated here):** Republic of Ukraine. Lviv, Yavoriv, Dobrostan, 15 Sep 2012 [non-fossil]: *M. Gottschling, N.H. Filipowicz & C. Zinßmeister P57* [J. Kretschmann GeoM 520] (KW No. KW-A-221!; isoeptypes: B barcode B 40 0043801!, M barcodes M-0299991! & M-0299992!).

The nomenclatural act has been registered in PhycoBank under <http://phycobank.org/100580>.

*Other original elements.* – Non-fossil specimens from Lviv and Wolicki pond (Ukraine), without exact dates (January, March), collected by R. Dreżepolski, none preserved.

*Note.* – Average length, width and shape of strain GeoM 520 correspond to the protologue data. Additionally, absence of collar and overall punctated surface, which is more pronounced at the anterior part of the cell (seen in SEM images), is also in correspondence with the protologue.

*Trachelomonas teres* var. *granulata* Drezep., nom. corr. (*ICN* Art. 60.1.), in Kosmos (Lvov) 50: 223, t. 1, fig. [32]. 1925 – **Lectotype (designated here):** [illustration] in Kosmos (Lvov) 50: t. 1, fig. [32]. 1925! [not numbered on the plate; showing a non-fossil individual from Republic of Ukraine; Lviv, Yavoriv, Dobrostan, Jul (without year): *R. Dreżepolski s.n.*] – **Epitype (designated here):** Republic of Ukraine. Lviv, Yavoriv, Dobrostan, 15 Sep 2012 [non-fossil]: *M. Gottschling, N.H. Filipowicz & C. Zinßmeister P56* [J. Kretschmann GeoM 540] (KW No. KW-A-224!; isoeptypes: B barcode B 40 0043804!, M barcodes M-0299987! & M-0299988!).

The nomenclatural act has been registered in PhycoBank under <http://phycobank.org/100568>.

*Note.* – Average length and width of the strain GeoM 540 more or less correspond to the protologue data. Cell shape and overall ornamentation (seen in LM and SEM images) are in correspondence with the protologue. Additionally, thickened edge on the flagellar pore region (seen in LM images) corresponds to the original description. Finally, samples were collected in summer, which also correlates with the original description.

*Trachelomonas teres* var. *minor* Drezep. in Kosmos (Lvov) 50: 223, t. 1, fig. 26. 1925 – **Lectotype (designated here):** [illustration] in Kosmos (Lvov) 50: t. 1, fig. 26. 1925! [showing a non-fossil individual; exact locality and collecting date unknown: *R. Dreżepolski s.n.*] – **Epitype (designated here):** Republic of Ukraine. Lviv, Lubień Mały, 15 Sep 2012 [non-fossil]: *C. Zinßmeister, N.H. Filipowicz & M. Gottschling P39* [J. Kretschmann GeoM 527] (KW No. KW-A-222!; isoeptypes: B barcodes B 40 0043802! & B 40 0043803!, KW No. KW-A-223!, M barcodes M-0299989! & M-0299990!).

The nomenclatural act has been registered in PhycoBank under <http://phycobank.org/100569>.

*Other original elements.* – Non-fossil specimens from Dobrostan and Lviv (Ukraine), Łuków (Poland) and Tuhanowicze (Belarus), without exact date (May, Jul, Sep, Nov), collected by R. Dreżepolski, none preserved.

*Note.* – Average length, width and shape of the strain GeoM 527 correspond to the protologue data. Double-husked pyrenoids are not seen in all the cells, yet the subclade within strain GeoM 527 clustered is known to possess diplopyrenoids (Ciugulea & al., 2008). Additionally, absence of collar (seen in SEM and LM images) and overall smooth surface (seen in LM images) are also in correspondence with the protologue. Finally, samples were collected in September, which also correlates with the original description.

## ■ AUTHOR CONTRIBUTIONS

AZC: morphological study, molecular phylogenetics, led the writing; JK: strain establishment, morphological study; MG: scientific concept, field work, taxonomy; all authors critically read and approved the written text. — AZC, <https://orcid.org/0000-0003-2081-2076>; MG, <https://orcid.org/0000-0002-4381-8051>

## ■ ACKNOWLEDGEMENTS

We sincerely thank Natalia H. Filipowicz (Gdańsk) and Carmen Zinßmeister (Wilhelmshaven) for the collection of algae in Ukraine. Furthermore, we would also like to thank Anina Neumann (Munich), who devotedly curated the established euglenophyte strains.

## ■ LITERATURE CITED

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. 1990. Basic local alignment search tool. *J. Molec. Biol.* 215: 403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
- Annenkova, N.V., Hansen, G., Moestrup, Ø. & Rengefors, K. 2015. Recent radiation in a marine and freshwater dinoflagellate species flock. *ISME J.* 9: 1821–1834. <https://doi.org/10.1038/ismej.2014.267>
- Arnott, H.J. & Walne, P.L. 1967. Observations on the fine structure of the pellicle pores of *Euglena granulata*. *Protoplasma* 64: 330–344. <https://doi.org/10.1007/BF01253711>
- Bass, D. & Boenigk, J. 2011. Everything is everywhere: A twenty-first century de-/reconstruction with respect to protists. Pp. 88–110 in: Fontaneto, D. (ed.), *Biogeography of microscopic organisms: Is everything small everywhere?* Cambridge: Cambridge University Press. <https://doi.org/10.1017/CBO9780511974878.007>
- Bates, S.T., Clemente, J.C., Flores, G.E., Walters, W.A., Parfrey, L.W., Knight, R. & Fierer, N. 2013. Global biogeography of highly diverse protistan communities in soil. *ISME J.* 7: 652–659. <https://doi.org/10.1038/ismej.2012.147>
- Bennett, M.S. & Triemer, R.E. 2012. A new method for obtaining nuclear gene sequences from field samples and taxonomic revisions of the photosynthetic euglenoids *Lepocinlis (Euglena) helicoideus* and *Lepocinlis (Phacus) horridus* (Euglenophyta). *J. Phycol.* 48: 254–260. <https://doi.org/10.1111/j.1529-8817.2011.01101.x>
- Bicudo, C.E.D.M. & Menezes, M. 2016. Phylogeny and classification of Euglenophyceae: A brief review. *Frontiers Ecol. Evol.* 4: 1–15. <https://doi.org/10.3389/fevo.2016.00017>
- Borisova, O.V., Dogadina, T.V., Gorbulin, O.S., Kovalenko, O.V., Krakhmalny, O.F., Levantets, A.A., Lillitska, G.G., Massjuk, N.P., Petlovany, O.A., Skyba, Y.A., Tsarenko, O.M., Tsarenko, P.M., Vinogradova, O.M. & Wasser, S.P. 2006. *Algae of Ukraine: Diversity, nomenclature, taxonomy, ecology and geography*. Ruggell: Gantner.
- Briquet, J. 1930. *Recueil synoptique des documents destinées à servir de base aux débats de la sous-section de nomenclature du Vme Congrès international de Botanique Cambridge*. Berlin: Friedländer.
- Brosnan, S., Shin, W., Kjer, K.M. & Triemer, R.E. 2003. Phylogeny of the photosynthetic euglenophytes inferred from the nuclear SSU and partial LSU rDNA. *Int. J. Syst. Evol. Microbiol.* 53: 1175–1186. <https://doi.org/10.1099/ijs.0.02518-0>
- Brosnan, S., Brown, P.J.P., Farmer, M.A. & Triemer, R.E. 2005. Morphological separation of the euglenoid genera *Trachelomonas* and *Strombomonas* (Euglenophyta) based on lorica development and posterior strip reduction. *J. Phycol.* 41: 590–605. <https://doi.org/10.1111/j.1529-8817.2005.00068.x>
- Ciugulea, I., Nudelman, M.A., Brosnan, S. & Triemer, R.E. 2008. Phylogeny of the euglenoid loricate genera *Trachelomonas* and *Strombomonas* (Euglenophyta) inferred from nuclear SSU and LSU rDNA. *J. Phycol.* 44: 406–418. <https://doi.org/10.1111/j.1529-8817.2008.00472.x>
- De Clerck, O., Guiry, M.D., Leliaert, F., Samyn, Y. & Verbruggen, H. 2013. Algal taxonomy: A road to nowhere? *J. Phycol.* 49: 215–225. <https://doi.org/10.1111/jpy.12020>
- Deflandre, G. 1926. Monographie du genre *Trachelomonas* Ehr. *Rev. Gén. Bot.* 38: 358–380, 449–469, 518–528, 646–658, 687–706.
- Demchenko, E., Mikhailyuk, T., Coleman, A.W. & Pröschold, T. 2012. Generic and species concepts in *Microglena* (previously the *Chlamydomonas monadina* group) revised using an integrative approach. *Eur. J. Phycol.* 47: 264–290. <https://doi.org/10.1080/09670262.2012.678388>
- Dreżepolski, R. 1923. Eugleniny wolnożyjące ze zbioru glonów podlaskich i litewskich dr J. Grochmalickiego. *Rozpr. Wiadom. Muz. Dzieduszyckich* 7–8: 1–17.
- Dreżepolski, R. 1925. Przyczynek do znajomości polskich Euglenin. *Kosmos (Lvov)* 50: 173–270. <https://archive.lib.msu.edu/DMC/Euglena/pdfs/drez0173.pdf>
- Dunlap, J.R. & Walne, P.L. 1985. Fine-structure and biomineralization of the mucilage in envelopes of *Trachelomonas lefevrei* (Euglenophyceae). *J. Protozool.* 32: 437–441. <https://doi.org/10.1111/j.1550-7408.1985.tb04040.x>
- Dunn, O.J. 1961. Multiple comparisons among means. *J. Amer. Statist. Assoc.* 56: 54–64. <https://doi.org/10.1080/01621459.1961.10482090>
- Finlay, B.J. 2002. Global dispersal of free-living microbial eukaryote species. *Science* 296: 1061–1063. <https://doi.org/10.1126/science.1070710>
- Gottschling, M. 2008. *Aktuelle Herausforderungen für Diversitätserfassung und Systematik: Blütenpflanzen, kalkige Dinoflagellaten und Papillomviren im Vergleich*. Habilitation thesis, Freie Universität Berlin, Germany.
- Gottschling, M., Keupp, H., Plötner, J., Knop, R., Willems, H. & Kirsch, M. 2005. Phylogeny of calcareous dinoflagellates as inferred from ITS and ribosomal sequence data. *Molec. Phylog. Evol.* 36: 444–455. <https://doi.org/10.1016/j.ympev.2005.03.036>
- Gottschling, M., Söhner, S., Zinßmeister, C., John, U., Plötner, J., Schweikert, M., Aligizaki, K. & Elbrächter, M. 2012. Delimitation of the Thoracosphaeraceae (Dinophyceae), including the calcareous dinoflagellates, based on large amounts of ribosomal RNA sequence data. *Protist* 163: 15–24. <https://doi.org/10.1016/j.protis.2011.06.003>
- Gottschling, M., Chacón, J., Žerdoner Čalasan, A., Neuhaus, S., Kretschmann, J., Stibor, H. & John, U. 2020. Phylogenetic placement of environmental sequences using taxonomically reliable databases helps to rigorously assess dinophyte biodiversity in Bavarian lakes (Germany). *Freshwater Biol.* 65: 193–208. <https://doi.org/10.1111/fwb.13413>

- Guillard, R.R.L. & Lorenzen, C.J. 1972. Yellow-green algae with chlorophyllide *c*. *J. Phycol.* 8: 10–14. <https://doi.org/10.1111/j.1529-8817.1972.tb03995.x>
- Guiry, M.D. & Guiry, G.M. 2017. AlgaeBase: World-wide electronic publication. <http://www.algaebase.org/> (accessed 14 May 2017).
- Hansen, G., Daugbjerg, N. & Henriksen, P. 2000. Comparative study of *Gymnodinium mikimotoi* and *Gymnodinium aureolum*, comb. nov. (= *Gyrodinium aureolum*) based on morphology, pigment composition, and molecular data. *J. Phycol.* 36: 394–410. <https://doi.org/10.1046/j.1529-8817.2000.99172.x>
- Hyde, K.D. & Zhang, Y. 2008. Epitypification: Should we epitypify? *J. Zhejiang Univ. Sci. B* 9: 842–846. <https://doi.org/10.1631/jzus.B0860004>
- John, U., Litaker, R.W., Montresor, M., Murray, S., Brosnahan, M.L. & Anderson, D.M. 2014. Formal revision of the *Alexandrium tamarense* species complex (Dinophyceae) taxonomy: The introduction of five species with emphasis on molecular-based (rDNA) classification. *Protist* 165: 779–804. <https://doi.org/10.1631/jzus.B0860004>
- Karnkowska, A., Bennett, M.S., Wata, D., Kim, J.I., Zakryś, B. & Triemer, R.E. 2015. Phylogenetic relationships and morphological character evolution of photosynthetic euglenids (*Excavata*) inferred from taxon-rich analyses of five genes. *J. Eukar. Microbiol.* 62: 362–373. <https://doi.org/10.1111/jeu.12192>
- Karnkowska-Ishikawa, A., Milanowski, R., Triemer, R.E. & Zakryś, B. 2012. Taxonomic revisions of morphologically similar species from two euglenoid genera: *Euglena* (*E. granulata* and *E. velata*) and *Euglenaria* (*Eu. anabaena*, *Eu. caudata*, and *Eu. clavata*). *J. Phycol.* 48: 729–739. <https://doi.org/10.1111/j.1529-8817.2012.01140.x>
- Katoh, K. & Standley, D.M. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molec. Biol. Evol.* 30: 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kim, J.I. & Shin, W. 2014. Molecular phylogeny and cryptic diversity of the genus *Phacus* (Phacaceae, Euglenophyceae) and the descriptions of seven new species. *J. Phycol.* 50: 948–959. <https://doi.org/10.1111/jpy.12227>
- Kim, J.I., Shin, W. & Triemer, R.E. 2010. Multigene analyses of photosynthetic euglenoids and new family, Phacaceae (Euglenales). *J. Phycol.* 46: 1278–1287. <https://doi.org/10.1111/j.1529-8817.2010.00910.x>
- Kim, J.I., Shin, W. & Triemer, R.E. 2013. Cryptic speciation in the genus *Cryptoglena* (Euglenaceae) revealed by nuclear and plastid SSU and LSU rRNA gene. *J. Phycol.* 49: 92–102. <https://doi.org/10.1111/jpy.12032>
- Kim, J.I., Linton, E.W. & Shin, W. 2015. Taxon-rich multigene phylogeny of the photosynthetic euglenoids (Euglenophyceae). *Frontiers Ecol. Evol.* 3: 1–11. <https://doi.org/10.3389/fevo.2015.00098>
- Koczwara, M. 1915. Fytoplankton stawów dobrostańskich. *Kosmos (Lvov)* 40: 231–275.
- Konenko, G.D., Pidgajko, M.L. & Radzimovskij, D.O. 1965. *Ponds of wood-steppe, steppe and mountain regions of Ukraine*. Kiev: Naukova dumka.
- Kosmala, S., Karnkowska, A., Milanowski, R., Kwiatowski, J. & Zakryś, B. 2005. Phylogenetic and taxonomic position of *Lepocinclis fusca* comb.nov. (= *Euglena fusca*) (Euglenaceae): Morphological and molecular justification. *J. Phycol.* 41: 1258–1267. <https://doi.org/10.1111/j.1529-8817.2005.00141.x>
- Kosmala, S., Bereza, M., Milanowski, R., Kwiatowski, J. & Zakryś, B. 2007. Morphological and molecular examination of relationships and epitype establishment of *Phacus pleuronectes*, *Phacus orbicularis*, and *Phacus hamelii*. *J. Phycol.* 43: 1071–1082. <https://doi.org/10.1111/j.1529-8817.2007.00386.x>
- Kremp, A., Elbrächter, M., Schweikert, M., Wolny, J.L. & Gottschling, M. 2005. *Woloszynskia halophila* (Biecheler) comb. nov.: A bloom-forming cold-water dinoflagellate co-occurring with *Scrippsiella hangoei* (Dinophyceae) in the Baltic Sea. *J. Phycol.* 41: 629–642. <https://doi.org/10.1111/j.1529-8817.2005.00070.x>
- Kretschmann, J., Elbrächter, M., Zinßmeister, C., Söhner, S., Kirsch, M., Kusber, W.H. & Gottschling, M. 2015. Taxonomic clarification of the dinophyte *Peridinium acuminatum* Ehrenb., = *Scrippsiella acuminata*, comb. nov. (Thoracosphaeraceae, Peridinales). *Phytotaxa* 220: 239–256. <https://doi.org/10.11646/phytotaxa.220.3.3>
- Kretschmann, J., Żerdoner Čalasan, A. & Gottschling, M. 2018a. Molecular phylogenetics of dinophytes harbouring diatoms as endosymbionts (Kryptoperidiniaceae, Peridinales), with evolutionary interpretations and a focus on the identity of *Durinskia oculata* from Prague. *Molec. Phylogen. Evol.* 118: 392–402. <https://doi.org/10.1016/j.ympev.2017.10.011>
- Kretschmann, J., Żerdoner Čalasan, A., Kusber, W.-H. & Gottschling, M. 2018b. Still curling after all these years: *Glenodinium apiculatum* Ehrenb. (Peridinales, Dinophyceae) repeatedly found at its type locality in Berlin (Germany). *Syst. Biodivers.* 16: 200–209. <https://doi.org/10.1080/14772000.2017.1375045>
- Kulczycka, A., Łukomska-Kowalczyk, M., Zakryś, B. & Milanowski, R. 2018. PCR identification of toxic euglenid species *Euglena sanguinea*. *J. Appl. Phycol.* 30: 1759–1763. <https://doi.org/10.1007/s10811-017-1376-z>
- Kuzmenko, M.I. 2000. Radio ecological problems of waterbodies in Ukraine. *Hydrobiol. J.* 36: 3–32.
- Leedale, G.F. 1967. *Euglenoid flagellates*. Englewood Cliffs: Prentice Hall.
- Leedale, G.F. 1975. Envelope formation and structure in the euglenoid genus *Trachelomonas*. *Brit. Phycol. J.* 10: 17–41. <https://doi.org/10.1080/00071617500650031>
- Lemmermann, E.J. 1910. *Algen I (Schizophyceen, Flagellaten, Peridinee)*. Leipzig: Bornträger. <https://doi.org/10.5962/bhl.title.4953>
- Lindner, L., Gozhik, P., Marciniak, B., Marks, L. & Yelovicheva, Y. 2004. Main climatic changes in the quaternary of Poland, Belarus and Ukraine. *Geol. Quart.* 48: 97–114.
- Łukomska-Kowalczyk, M., Karnkowska, A., Milanowski, R., Łach, Ł. & Zakryś, B. 2015. Delimiting species in the *Phacus longicauda* complex (Euglenida) through morphological and molecular analyses. *J. Phycol.* 51: 1147–1157. <https://doi.org/10.1111/jpy.12352>
- Manoylov, K.M. 2014. Taxonomic identification of algae (morphological and molecular): Species concepts, methodologies, and their implications for ecological bioassessment. *J. Phycol.* 50: 409–424. <https://doi.org/10.1111/jpy.12183>
- Margulis, L., Corliss, J.O., Melkonian, M. & Chapman, D.J. 1990. *Handbook of Protoctista*. Boston: Jones & Bartlett.
- Marin, B., Palm, A., Klingberg, M. & Melkonian, M. 2003. Phylogeny and taxonomic revision of plastid-containing euglenophytes based on SSU rDNA sequence comparisons and synapomorphic signatures in the SSU rDNA secondary structure. *Protist* 154: 99–145. <https://doi.org/10.1078/143446103764928521>
- McFadden, G.I. & Melkonian, M. 1986. Use of Hepes buffer for microalgal culture media and fixation for electron microscopy. *Phycologia* 25: 551–557. <https://doi.org/10.2216/i0031-8884-25-4-551.1>
- Miller, M.A., Pfeiffer, W. & Schwartz, T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Pp. 45–52 in: *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, Louisiana, 14 Nov 2010. Piscataway: IEEE. <https://doi.org/10.1109/GCE.2010.5676129>
- Nudelman, M.A., Rossi, M.S., Conforti, V. & Triemer, R.E. 2003. Phylogeny of Euglenophyceae based on small subunit rRNA sequences: Taxonomic implications. *J. Phycol.* 39: 226–235. <https://doi.org/10.1046/j.1529-8817.2003.02075.x>
- Pawlowski, J., Audic, S., Adl, S., Bass, D., Belbahri, L., Berney, C., Bowser, S.S., Cepicka, I., Decelle, J., Dunthorn, M.,

- Fiore-Donno, A.M., Gile, G.H., Holzmann, M., Jahn, R., Jirku, M., Keeling, P.J., Kostka, M., Kudryavtsev, A., Lara, E., Lukes, J., Mann, D.G., Mitchell, E.A., Nitsche, F., Romeralo, M., Saunders, G.W., Simpson, A.G., Smirnov, A.V., Spouge, J.L., Stern, R.F., Stoeck, T., Zimmermann, J., Schindel, D. & de Vargas, C. 2012. CBOL protist working group: Barcoding eukaryotic richness beyond the animal, plant, and fungal kingdoms. *PLoS Biol.* 10: e1001419. <https://doi.org/10.1371/journal.pbio.1001419>
- Pereira, M.J., Azeiteiro, U.M.M., Goncalves, F. & Soares, A.M.V.M. 2003. Inorganic composition of the envelopes of *Trachelomonas* Ehr. (Euglenophyta). *Acta Oecol.* 24: S317–S324. [https://doi.org/10.1016/S1146-609X\(03\)00027-4](https://doi.org/10.1016/S1146-609X(03)00027-4)
- Pringsheim, G. 1953. Observations on some species of *Trachelomonas* grown in culture. *New Phytol.* 52: 93–113, 238–266. <https://doi.org/10.1111/j.1469-8137.1953.tb07514.x>
- Przybos, E. & Tarcz, S. 2016. *Paramecium jenningsi* complex: Existence of three cryptic species confirmed by multi-locus analysis and strain crosses. *Syst. Biodivers.* 14: 140–154. <https://doi.org/10.1080/14772000.2015.1134701>
- Ride, W.D.L. 2003. The International Code of Zoological Nomenclature, 4th edition — what next? Pp. 673–682 in: Legakis, A., Sfenthourakis, S., Polymeni, R. & Thessalou-Legaki, M. (eds.), *The new panorama of animal evolution: Proceedings, XVIII International Congress of Zoology*. Sofia: Pensoft.
- Rino, J.A. & Pereira, M.J. 1988. Euglenophyta da Região Centro de Portugal, I. Género *Trachelomonas* Ehrenb. 1833 emend. Defl. 1926. *Revista Biol. Univ. Aveiro* 2: 129–161.
- Rino, J.A. & Pereira, M.J. 1991. Estrutura e composição química da lórica de *Trachelomonas nigra* Swirenko (Algae, Euglenophyta). *Revista Biol. Univ. Aveiro* 4: 95–107.
- Romeikat, C., Izquierdo López, A., Tietze, C., Kretschmann, J. & Gottschling, M. 2019. Typification for reliable application of subspecific names within *Peridinium cinctum* (Peridinales, Dinophyceae). *Phytotaxa* 424: 147–157. <https://doi.org/10.11646/phytotaxa.424.3.2>
- Ronquist, F., Teslenko, M., Van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A. & Huelsenbeck, J.P. 2012. MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Ross, R. 1966. General Committee 1964–1969, Report No. 1. *Taxon* 15: 286–287. <https://doi.org/10.1002/j.1996-8175.1966.tb01995.x>
- Shin, W. & Triemer, R.E. 2004. Phylogenetic analysis of the genus *Euglena* (Euglenophyceae) with particular reference to the species *Euglena viridis*. *J. Phycol.* 40: 759–771. <https://doi.org/10.1111/j.1529-8817.2004.03142.x>
- Silva, P.C. 1960. Remarks on algal nomenclature, III. *Taxon* 9: 18–25. <https://doi.org/10.2307/1217351>
- Skuja, H. 1926. Vorarbeiten zu einer Algenflora von Lettland. I. *Acta Horti Bot. Univ. Latv.* 1: 33–54.
- Skvortzov, B.V. 1925. Die Euglenaceengattung *Trachelomonas* Ehrenberg: Eine systematische Übersicht. *Trudy Sungar. Rechn. Biol. Stantsii* 1(2): 1–101.
- Smith, G.F., Figueiredo, E. & Moore, G. 2016. Increasing nomenclatural stability by preventing the introduction of long-forgotten names that will compete with ones in use: A solution must be found, and soon. *Taxon* 65: 1385–1390. <https://doi.org/10.12705/656.11>
- Stamatakis, A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Starmach, K. 1983. *Euglenophyta—Eugleniny*. Warsaw: Państwowe Wydawnictwo Naukowe.
- Stein, F. von 1878. *Der Organismus der Infusionsthier nach eigenen Forschungen in systematischer Reihenfolge bearbeitet*, 3(1). Leipzig: Engelmann. <https://doi.org/10.5962/bhl.title.3933>
- Swirenko, D.O. 1914. Zur Kenntnis der russischen Algenflora. I. Die Euglenaceen der Gattung *Trachelomonas*. *Arch. Hydrobiol. Planktonk.* 9: 630–647.
- Tillmann, U., Gottschling, M., Nézan, E., Krock, B. & Bilien, G. 2014. Morphological and molecular characterization of three new *Azadinium* species (Amphidomataceae, Dinophyceae) from the Irminger Sea. *Protist* 165: 417–444. <https://doi.org/10.1016/j.protis.2014.04.004>
- Tillmann, U., Hoppenrath, M., Gottschling, M., Kusber, W.-H. & Elbrächter, M. 2017. Plate pattern clarification of the marine dinophyte *Heterocapsa triquetra* sensu Stein (1883) collected at the Kiel Fjord (Germany). *J. Phycol.* 53: 1305–1324. <https://doi.org/10.1111/jpy.12584>
- Triemer, R.E., Linton, E., Shin, W., Nudelman, A., Monfils, A., Bennett, M. & Brosnan, S. 2006. Phylogeny of the Euglenales based upon combined SSU and LSU rDNA sequence comparisons and description of *Discoplastis* gen. nov (Euglenophyta). *J. Phycol.* 42: 731–740. <https://doi.org/10.1111/j.1529-8817.2006.00219.x>
- Turland, N.J., Wiersma, J.H., Barrie, F.R., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Kusber, W.-H., Li, D.-Z., Marhold, K., May, T.W., McNeill, J., Monro, A.M., Prado, J., Price, M.J. & Smith, G.F. (eds.) 2018. *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017*. Glashütten: Koeltz. <https://doi.org/10.12705/Code.2018>
- Wolowski, K., Poniewozik, M. & Jurán, J. 2016. Morphological variability of loricae in *Trachelomonas caudata* complex (Euglenophyta). *Cryptog. Algol.* 37: 97–108. <https://doi.org/10.7872/crya/v37.iss2.2016.97>
- Žerdoner Čalasan, A., Kretschmann, J., Filipowicz, N.H., Irimia, R.-E., Kirsch, M. & Gottschling, M. 2019. Towards global distribution maps of unicellular organisms such as calcareous dinophytes based on DNA sequence information. *Mar. Biodivers.* 49: 749–758. <https://doi.org/10.1007/s12526-018-0848-y>
- Zinßmeister, C., Söhner, S., Facher, E., Kirsch, M., Meier, K.J.S. & Gottschling, M. 2011. Catch me if you can: The taxonomic identity of *Scrippsiella trochoidea* (F.Stein) A.R.Loeb. (Thoracosphaeraceae, Dinophyceae). *Syst. Biodivers.* 9: 145–157. <https://doi.org/10.1080/14772000.2011.586071>

#### Appendix 1. Voucher list for the multilocus-based alignment of a representative taxon set of selected photosynthetic euglenoid clades.

Taxon, strain number, locality, collector plus collection number, GenBank accession numbers for 18S rRNA and partial 28S rRNA loci. The latter locus occasionally under two different GenBank accession numbers.

*Colacium calvum* F.Stein, MI 107, locality unknown, collector and coll. number unknown, EF999907, EF999910; *Colacium mucronatum* Bourr. & Chadeff., SAG 1211-1, United Kingdom, England, Cambridge, epizoic on *Daphnia*. *E.G. Pringsheim, s.n.*, AJ532441, EF999906; *Colacium vesiculosum* Ehrenb., MI 105, locality unknown, collector and coll. number unknown, EF999905, EF999912; *Cryptoglena pigra* Ehrenb., CCAP1212/1, United Kingdom, England,

## Appendix 1. Continued.

Cambridge, Jesus Ditch, collector and coll. number unknown, AJ532437, DQ140101; *Cryptoglena skujae* B. Marin & Melkonian, SAG 10.88, Austria, Neusiedler See at Rust, *E. Kusel-Fetzmann*, *s.n.*, AY014998, AY130236; *Monomorpha aenigmatica* (Drezip.) Nudelman & Triemer, UTEX 1284, United Kingdom, England, Lyme Regis, *M.R. Droop*, *s.n.*, AF190814, DQ140117; *Monomorpha inaequalis* (Deflandre) B. Marin & Melkonian, ACOI 1295, Portugal, Castelo Branco, Penha Garcia, *L. Santos*, *s.n.*, DQ140129, DQ140111; *Monomorpha pseudopyrum* Kosmala, Milanowski, Brzóska, Pékala, Kwitowski & Zakryś, CCAC 0093, The Netherlands, Ameland (West), (Nassau) Entenkooi, *Klingberg*, *M. s.n.*, AJ532433, EF999909; *Monomorpha pyrum* (Ehrenb.) Mereschk., UTEX 2354, locality unknown, collector and coll. number unknown, AF112874, AY130238; *Strombomonas balvayi* Bourr. & Couté, CP1D, locality unknown, collector and coll. number unknown, EF999897, AY359916, KT304970; *Strombomonas borystheniensis* (Y. V. Roll) T.G. Popova, S10, locality unknown, collector and coll. number unknown, DQ140131, AY359920, KT304971; *Strombomonas costata* Deflandre, ACOI 1273, Portugal, Quiaios, Lagoa das Braças, *M. F. Santos*, *s.n.*, DQ140152, AY359915, KT304972; *Strombomonas eurystoma* (F. Stein) T.G. Popova, S600, locality unknown, collector and coll. number unknown, DQ140132, AY359918, KT304974; *Strombomonas ovalis* (Playfair) Deflandre, S115, locality unknown, collector and coll. number unknown, DQ140133, AY359919, KT304975; *Strombomonas triquetra* (Playfair) Deflandre, S604, locality unknown, collector and coll. number unknown, DQ140153, AY359917, KT304978; *Strombomonas verrucosa* (Daday) Deflandre, S5C, locality unknown, collector and coll. number unknown, AF445461, AY359911, KT304979; *Trachelomonas abrupta* Svirenko, T 801, USA, collector and coll. number unknown, DQ140134, AY359941, KT304980; *Trachelomonas armata* (Ehrenb.) F. Stein, ACOI 1323, Portugal, Casal Novo do Rio, canal, *M.F. Santos*, *s.n.*, EF999904, KT304981; *Trachelomonas bernardinensis* Vischer, ACOI 1103, Portugal, Castelo Branco, Penamacor, *L. Santos*, *s.n.*, EF999908, AY359950, KT304982; *Trachelomonas echinata* A.M. Cunha, SAG 1283-22, United States, Indiana, Griffy Lake at Bloomington, *R.C. Starr*, *s.n.*, AY015001, AY130242, KT304985; *Trachelomonas ellipsoidalis* Kam.P. Singh, ST1, locality unknown, collector and coll. number unknown, DQ140135, AY359935; *Trachelomonas grandis* Kam.P. Singh, SAG 204.80, United States, Tennessee, quarry in Nashville, *K.P. Singh (reisolated by E.G. Pringsheim)*, *s.n.*, AJ532446, AY359936; *Trachelomonas hispida* (Perty) F. Stein, UTEX 1325, United Kingdom, England, Cambridge, Landbeach, *E.G. Pringsheim*, *s.n.*, AF445462, AY130817; *Trachelomonas hispida* (Perty) F. Stein, UTEX 1326, United Kingdom, England, Debden, *E.G. Pringsheim*, *s.n.*, AF090377, AY130817; *Trachelomonas hispida* var. *coronata* Lemmerm., Jigock 121309C, Republic of Korea, Jigock, collector and coll. number unknown, KT304850, KT304989; *Trachelomonas hispida* var. *volicensis* Drezip., GeoM 520 (≡ CCCM7069, CCAC 6780 B), Republic of Ukraine, Lviv, Yavoriv, Dobrostan, 15 Sep 2012: *M. Gottschling*, *N.H. Filipowicz* & *C. Zinßmeister* P56 [J. Kretschmann GeoM 520] (epitype: KW No. KW-A-224!; isoeotypes: B barcode B 40 0043804!, M barcodes M-0299987! & M-0299988!), MK894269\*, MK894263\*; *Trachelomonas lefevrei* Deflandre, Pungcheon 093006B, Republic of Korea, Pungcheon, collector and coll. number unknown, KT304852, KT304993; *Trachelomonas lefevrei* Deflandre, SAG 1283-10, United Kingdom, England, Shelford/Cambridge, *E.G. Pringsheim*, *s.n.*, DQ140136, AY359949; *Trachelomonas magdaleniana* Deflandre, 472057, Argentina, collector and coll. number unknown, EF999903, EF999911; *Trachelomonas oblonga* Lemmerm., T 516, locality unknown, collector and coll. number unknown, DQ140137, AY359947, KT304994; *Trachelomonas pertyi* E.G. Pringsh., Posudun 091809D, Republic of Korea, Posudun, collector and coll. number unknown, KT304853, KT304995; *Trachelomonas planctonica* Svirenko, WT1, locality unknown, collector and coll. number unknown, DQ140138, AY359954, KT304996; *Trachelomonas reticulata* G.A. Klebs, SAG 239.80, Germany, Göttingen, mud from a road puddle, *O. Pringsheim*, *s.n.*, DQ140139, AY359934; *Trachelomonas rugulosa* F. Stein, TS2, locality unknown, collector and coll. number unknown, DQ140140, AY359942, KT304997; *Trachelomonas scabra* Playfair, T 235, United States, collector and coll. number unknown, DQ140141, AY359951; *Trachelomonas similis* A. Stokes, SAG 1283-14, United Kingdom, England, pond near Debden, *E.G. Pringsheim*, *s.n.*, DQ140142, AY359948, KT304999; *Trachelomonas sp.*, Bakunji 101709F, Republic of Korea, Bakunji, collector and coll. number unknown, KT304858, KT305004; *Trachelomonas sp.*, Bidduk 050909G, Republic of Korea, Bidduk, collector and coll. number unknown, KT304859, KT305005; *Trachelomonas sp.*, Bokheungje 092609A, Republic of Korea, Bokheungje, collector and coll. number unknown, KT304860, KT305006; *Trachelomonas sp.*, Chosan 090509B, Republic of Korea, Chosan, collector and coll. number unknown, KT304861, KT305007; *Trachelomonas sp.*, GeoM\*521, Republic of Ukraine, Lviv, Zhovkva, Dublyany, 16 Sep 2012: *N.H. Filipowicz*, *C. Zinßmeister* & *M. Gottschling*, P58 [J. Kretschmann GeoM\*521], MK894271\*; *Trachelomonas sp.*, GeoM\*524, Republic of Ukraine, Lviv, Lubień Mały, 15 Sep 2012: *C. Zinßmeister*, *N.H. Filipowicz* & *M. Gottschling*, P38 [J. Kretschmann GeoM\*524], MK894265\*, MK894259\*; *Trachelomonas sp.*, GeoM 526, Republic of Poland, Włocławek, Chodecz, Chodeckie lake, 10 Sep 2012: *C. Zinßmeister*, *N.H. Filipowicz* & *M. Gottschling*, P1 [J. Kretschmann GeoM 526], MK894267\*, MK894261\*; *Trachelomonas sp.*, GeoM 529 (≡ CCCM7132), Republic of Ukraine, Lviv, Lubień Mały, 15 Sep 2012: *C. Zinßmeister*, *N.H. Filipowicz* & *M. Gottschling*, P37 [J. Kretschmann GeoM 529], MK894266\*, MK894260\*; *Trachelomonas sp.*, Geumdong 090304A, Republic of Korea, Geumdong, collector and coll. number unknown, KT304865, KT305011; *Trachelomonas sp.*, Hoisan 032709F, Republic of Korea, Hoisan, collector and coll. number unknown, KT304866, KT305012; *Trachelomonas sp.*, Hongseong 091706C, Republic of Korea, Hongseong, collector and coll. number unknown, KT304867, KT305013; *Trachelomonas sp.*, Jakeun 052407A, Republic of Korea, Jakeun, collector and coll. number unknown, KT304868, KT305014; *Trachelomonas sp.*, Jinal 030207BT, Republic of Korea, Jinal, collector and coll. number unknown, KT304870, KT305016; *Trachelomonas sp.*, Juam 072909B, Republic of Korea, Juam, collector and coll. number unknown, KT304871, KT305017; *Trachelomonas sp.*, Kwoanam 102007E, Republic of Korea, Kwoanam, collector and coll. number unknown, KT304872, KT305018; *Trachelomonas sp.*, LnE082603E, locality unknown, collector and coll. number unknown, KT304864, KT305010; *Trachelomonas sp.*, Mulryang 090509E, Republic of Korea, Mulryang, collector and coll. number unknown, KT304873, KT305019; *Trachelomonas sp.*, Nogok 101407B, Republic of Korea, Nogok, collector and coll. number unknown, KT304875, KT305021; *Trachelomonas sp.*, Ojung 110506C, Republic of Korea, Ojung, collector and coll. number unknown, KT304876, KT305022; *Trachelomonas sp.*, Pungcheon 093006A, Republic of Korea, Pungcheon, collector and coll. number unknown, KT304877, KT305023; *Trachelomonas sp.*, Silo081903A, United States, collector and coll. number unknown, KT304880, KT305026; *Trachelomonas sp.*, Suckhyun 092606A, Republic of Korea, Suckhyun, collector and coll. number unknown, KT304885, KT305031; *Trachelomonas sp.*, T 101, locality unknown, collector and coll. number unknown, EF999898, AY359943; *Trachelomonas sp.*, T 201, locality unknown, collector and coll. number unknown, EF999899, AY359922; *Trachelomonas sp.*, T 307, locality unknown, collector and coll. number unknown, EF999900, AY359923; *Trachelomonas sp.*, T 502, locality unknown, collector and coll. number unknown, EF999901, AY359939; *Trachelomonas sp.*, T 603, locality unknown, collector and coll. number unknown, DQ140143, AY359940; *Trachelomonas sp.*, T 812, United States, collector and coll. number unknown, KT304886, AY359945; *Trachelomonas sp.*, T 815, locality unknown, collector and coll. number unknown, EF999902, AY359946; *Trachelomonas sp.*, T 900, United States, collector and coll. number unknown, KT304887, AY359927; *Trachelomonas sp.*, Tukuba 080509E, Japan, Tukuba, collector and coll. number unknown, KT304888, KT305034; *Trachelomonas sp.*, Yeonhwaji 091109F, Republic of Korea, Yeonhwaji, collector and coll. number unknown, KT304890, KT305036; *Trachelomonas teres* var. *granulata* Drezip., GeoM 540 (≡ CCAC 6782 B), Republic of Ukraine, Lviv, Yavoriv, Dobrostan, 15 Sep 2012: *M. Gottschling*, *N.H. Filipowicz* & *C. Zinßmeister*, P56 [J. Kretschmann GeoM 540] (epitype: KW No. KW-A-224!; isoeotypes: B barcode B 40 0043804!, M barcodes M-0299987! & M-0299988!), MK894270\*, MK894264\*; *Trachelomonas teres* var. *minor* Drezip., GeoM 527 (≡ CCCM7129, CCAC 6781 B), Republic of Ukraine, Lviv, Lubień Mały, 15 Sep 2012: *C. Zinßmeister*, *N.H. Filipowicz* & *M. Gottschling*, P39 [J. Kretschmann GeoM 527] (epitype: KW No. KW-A-222!; isoeotypes: B barcodes B 40 0043802! & B 40 0043803!, KW No. KW-A-223!, M barcodes M-0299989! & M-0299990!), MK894268\*, MK894262\*; *Trachelomonas variabilis* Kam.P. Singh, SAG 1283-24, United States, Tennessee, Nashville, pond in greenhouse of Vanderbilt University, *E.G. Pringsheim*, *s.n.*, KT304891, KT305038; *Trachelomonas volvocina* (Ehrenb.) Ehrenb., UTEX 1327, United Kingdom, England, Debden, *E.G. Pringsheim*, *s.n.*, KT304892, AY359953; *Trachelomonas* cf. *volvocinopsis* Svirenko, M1422, Germany, Muenster castle, small waterfall, *U. Powlowski*, *s.n.*, AJ532452; *Trachelomonas volvocinopsis* Svirenko, SAG 1283-16, United Kingdom, England, Trumpington/Cambridge, *E.G. Pringsheim*, *s.n.*, DQ140144, AY359944, KT305040; *Trachelomonas volvocinopsis* var. *spiralis* E.G. Pringsh., UTEX 1313, locality unknown, collector and coll. number unknown, AY015004, AY130816; *Trachelomonas zorensis* Deflandre, UTEX 1331, locality unknown, collector and coll. number unknown, DQ140145, AY359952, KT305037.