

Paul, Bartok, et al. Published in *Advances in Pediatrics*, 56(1), 2009. 107-133.

Opportunities for the Primary Prevention of Obesity during Infancy

Ian M. Paul, MD, MSc^{a,*}, Cynthia J. Bartok, PhD^b,
Danielle S. Downs, PhD^b, Cynthia A. Stifter, PhD^b,
Alison K. Ventura, PhD^c, Leann L. Birch, PhD^b

^aPediatrics, Penn State College of Medicine, HS83, 500 University Drive, Hershey, PA 17033, USA

^bPenn State College of Health and Human Development, University Park, PA, USA

^cMonell Chemical Senses Center, Philadelphia, PA, USA

WEIGHT GAIN DURING INFANCY AND LONG-TERM EFFECTS

Are chubby babies healthy babies? Whereas most seem well during infancy, evidence is increasing that heavier babies have a poorer long-term health trajectory than their trimmer counterparts. Data have emerged over the past 2 decades that early life growth patterns and behaviors play an important role in the etiology of obesity, yet there has been very little focus on the primary prevention of obesity during infancy by the medical, behavioral health, and public health communities. A recent report from the National Health and Nutrition Examination Survey (NHANES) highlighted the need for very early intervention when it revealed that between 2003 and 2006, a staggering 24.4% of children aged 2 to 5 years *already* were overweight or obese (body mass index [BMI; calculated as the weight in kilograms divided by height in meters squared] 85th–94th and ≥ 95 th percentiles, respectively) [1]. NHANES data also have described obesity (weight-for-length/height ≥ 95 th percentile) among infants younger than 2 years (Fig. 1). Between the late 1970s and 2000, the prevalence of obesity among infants 6 to 23 months old increased by more than 60% [2]. Reports from the Centers for Disease Control and Prevention (CDC) Pediatric Nutrition Surveillance System [3] and a Massachusetts Health Maintenance Organization [4] similarly showed significant increases in the prevalence of overweight for infants and toddlers for all age groups since the 1980s.

The Institute of Medicine publication, “Preventing Childhood Obesity: Health in the Balance,” stated that the prevention of obesity in children should

This work was supported by grants DK72996 and DK075867 from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Additional support was received from the Penn State Children, Youth and Families Consortium. Financial Disclosure and Conflicts of Interest: None.




Fig. 1. Trends in infant (weight-for-length ≥ 95 th percentile) and toddler (BMI ≥ 95 th percentile) obesity. (Data from Ogden CL, Carroll MD, Flegal KM. High body mass index for age among US children and adolescents, 2003–2006. *JAMA* 2008;299:2401–05; Ogden CL, Flegal KM, Carroll MD, et al. Prevalence and trends in overweight among US children and adolescents, 1999–2000. *JAMA* 2002;288:1728–32; and Koplan JP, Liverman CT, Kraak VI, editors. Preventing childhood obesity: health in the balance. Washington, DC: The National Academies Press; 2005.)

be a national public health priority [5]. More specific to younger children was the summary of the “Conference on Preventing Childhood Obesity,” where it was remarked, “The prenatal period, infancy, and early childhood may be stages of particular vulnerability to obesity development because they are unique periods for cellular differentiation and development. This unique vulnerability might make it possible for actions taken at these stages to determine the future course of adiposity [6].” This statement has been magnified by the numerous studies demonstrating the association between rapid or accelerated infant weight gain and subsequent obesity [7–24], as well as hypertension [25–28], coronary heart disease [29,30], and type 2 diabetes mellitus [31,32]. Further, numerous studies have now shown that overweight infants and toddlers are at increased risk of staying overweight as they age [9,16,33–46]. It has been theorized that overnutrition in infancy adversely “programs” the components of the metabolic syndrome and the way energy is stored [47,48]. These relationships may be especially true for those born to overweight parents, as genetic and familial influences, combined with pregnancy weight gain, are strongly associated with obesity in offspring [19,34,35,49,50].

Whereas all of the concerns about infant growth and subsequent morbidity make a strong case for very early intervention, there is little evidence regarding what, if anything, works to prevent the development of obesity during the first

years of life although the extant literature provides some suggestions regarding potentially promising approaches [5,51]. Early intervention and prevention hold great promise for interrupting the vicious cycle of obese children becoming obese adults who subsequently have obese offspring themselves. The following sections summarize numerous aspects of infant life that affect weight status, the way information on this subject should be communicated with parents, and interventions that can be suggested to families to prevent the development of obesity, based on the currently available evidence.

OBESITY PREVENTION DURING THE NEWBORN PERIOD AND EARLY INFANCY

Role of clinicians in addressing infant weight gain

Many parents, grandparents, and clinicians propagate the belief that “a chubby baby is a healthy baby” despite evidence even in the short term to the contrary [52–55], and substantial long-term evidence as described earlier. During infancy, growth charts are typically used by health care providers to ensure adequate and proportional growth with respect to weight, length, and head circumference, but information is usually communicated to parents without significant explanation so long as the child does not (a) raise concern for failure to thrive, or (b) demonstrate disproportionate or very excessive growth on 1 of the 3 measurements.

In addition, there is often a disconnect between health care provider definitions of overweight and obesity and parents’ interpretation of these terms [53,56–65]. Many parents believe heavier infant weight and appearance indicates good infant health and higher levels of parenting competence, particularly parents from poor or minority backgrounds [57,65–70]. In contrast, parents often perceive their children as picky eaters even when their weight gain is progressing normally [71], and infants and children perceived as too small often are given developmentally inappropriate nutrition, including the early introduction of solids or table foods [68,70]. The association of food with love in some cultures may also contribute to higher infant weight [72].

Potential intervention

Given the childhood obesity epidemic and the evidence that early upward crossing of major percentile lines on the growth curve is associated with later obesity [7–11,13–15,73], clinicians must pay closer attention to patterns of growth during early childhood and the way parents interpret infant growth. Providers must better understand healthy infant growth patterns and communicate this information regularly and accurately to parents. In addition, they must be familiar with early interventions that can prevent unhealthy patterns of weight gain in infancy and corrective interventions when problems are identified (Table 1).

Early feeding mode

Epidemiologic and experimental evidence has consistently indicated that breastfeeding offers modest protection against obesity later in life compared

Table 1**Potential interventions to prevent or address obesity during infancy**

Opportunity	Intervention
<i>Early infancy</i> Growth monitoring Infant feeding mode	Educate parents about growth charts, percentiles, and their meaning <ul style="list-style-type: none">– Support of breastfeeding as the preferred source of infant nutrition– Plot breastfed infants on the WHO growth chart, particularly when there are weight gain concerns– Educate parents on satiety cues– For bottle-fed infants, emphasize to parents that the volume of formula consumed should be child, not parent driven
Sleep	Educate parents on methods to lengthen sleep duration and soothe at night without feeding as a first response to nocturnal crying
Parental regulation of distress	<ul style="list-style-type: none">– Respond quickly to crying early in infancy, but use alternative methods to soothe than feeding– Use nonfood items as rewards later in infancy
Introduction of solid foods	<ul style="list-style-type: none">– Delay introduction of complementary foods until at least age 4 months– Avoid placing cereal into a bottle; complementary foods should only be fed with a spoon– Use repeated exposure to healthy foods as a response to normal infant neophobia
<i>Later infancy</i> Parent feeding style Transition to cow's milk Sweet beverage consumption	Avoid coercive or restrictive feeding styles Use low-fat cow's milk <ul style="list-style-type: none">– Do not give juice to children ≤ 6 months old– Limit daily consumption of 100% fruit juice to ≤ 6 ounces (170 g) per day– Give 100% juice only in a cup, never in a bottle– Do not allow children to easily transport juice so that they will not steadily consume it throughout the day
Transitional feeding and table foods	<ul style="list-style-type: none">– Completely avoid fruit drinks and soft drinks– Emphasize healthy dietary choices that have high nutritional value and low energy density such as fresh fruits, cooked vegetables, cheese, yogurt, whole-grain breads and crackers, and cereals– Avoid foods with added salt or sugar
Physical activity and sedentary behaviors	<ul style="list-style-type: none">– Choose physical activities that are interactive, stimulating, easy to do, and incorporated into daily routines– Avoid television watching for children younger than 2 years– Keep televisions out of bedroom

with formula feeding [74–80], and both exclusivity and duration of breastfeeding strengthen this association [75,77,78,80–83]. There are several reasons or mechanisms by which this protection may occur. First, breastfeeding promotes self-regulation of intake by the infant, and breastfed infants regulate the volume of feeds in response to the energy density of breast milk [84]. In contrast, formula feeding is a more parent-driven feeding activity, with the regulation of intake directed by the parents rather than the infant. Compared with nursing infants, bottle-fed infants are fed on a more regular schedule and the volume of feeds is very consistent, suggesting that parents are driving intake patterns [85]. Subsequent research has shown that common bottle feeding practices, such as “emptying the bottle” and serving larger volumes of formula at feedings, are associated with excess weight gain in the first 6 months of life [86].

The composition of breast milk may also contribute to the protective effects of breastfeeding. Human breast milk contains hundreds of components serving nutritive and nonnutritive functions within the infant [87], both of which may affect short- and long-term growth patterns of children [88]. Interspecies comparisons suggest that the high lactose and cholesterol content of human milk supports growth of the central nervous system, whereas the high protein and mineral content of other species’ milk (eg, cow’s milk) supports substantial and rapid gains in physical size [88,89]. Recent experimental research in humans suggest that the high levels of protein and minerals in formula may stimulate excess physical growth later in infancy, with persistent effects even at 2 years of age [90].

Other “bioactive” components of breast milk may have potential roles in the regulation of growth and development of the infant [91]. Human milk contains growth-regulating components such as leptin, ghrelin, insulinlike growth factor-1, and adiponectin [92–94], and blood leptin levels in breastfed infants are comparatively higher to formula-fed infants [92]. Subsequent research focused solely on breastfed infants has shown that maternal milk leptin levels are negatively associated with weight gain during early infancy and through 2 years of age [95,96].

The sum of this early feeding experience for breastfed and formula-fed infants produces clear growth differences by feeding mode that persists for at least the first 2 years after birth [97]. Limited research assessing body composition has shown that growth differences are likely due to increases in adiposity in formula-fed infants between 6 months and 24 months old [98]. These growth differences were viewed historically as a sign that breastfed infants were not thriving, and the widely used CDC 2000 growth reference has been criticized as inadequate for monitoring the growth of breastfed infants because it is based predominantly on data from formula-fed infants in the United States. The creation of the World Health Organization (WHO) International Growth Standards, based on an international sample of healthy breastfed infants, has helped to support the perception of breastfed infants’ growth as the “reference” growth pattern and formula-fed infants’ growth as deviant from this reference [99–101]. Of note, when comparing how breastfed infants’ growth

trajectories track along the CDC versus the WHO charts across the first 2 years after birth, substantial differences in expected growth are apparent [101]. With this comparison, when using the WHO chart as the growth reference, a normal deceleration of growth for breastfed infants is easily apparent after age 2 months and a slower rate of growth is evident through the first year after birth. In contrast, when using the CDC chart as the growth reference, the average breastfed infant would cross below the 50th percentile at age 7 months and proceed further below the 50th percentile through the first year. Without reference to a growth chart for breastfed infants, some infants might come to attention for failure to thrive when in fact their growth is normal. Further highlighting this is the fact that at 12 months of age, the WHO median weight for age is 1.2 kg lower for females and 1.5 kg lower for males in comparison with CDC charts.

Potential intervention

The CDC Guide to Breastfeeding Interventions outlines evidence-based practices for promoting breastfeeding, and improving breastfeeding duration and exclusivity [102]. Optimal breastfeeding practices can be promoted through a variety of avenues, including the health care system, places of work/employment, the community, and broader society. Among the interventions that can improve breastfeeding rates are several changes that can be made within the health care system. Prenatal, intrapartum, and postpartum education to improve breastfeeding knowledge and skills is an integral part of the promotion of breastfeeding. In addition, access to professional support (eg, lactation consultants, nurses, physicians) when families experience problems is a critical component of a supportive health care system. Institutional changes within the hospital or clinical setting have been shown to improve breastfeeding initiation and duration rates. The changes may be discrete, such as not handing out formula promotion gift packs to families, or they may be comprehensive, such as becoming a designated Baby Friendly Hospital Initiative hospital [102].

For physicians who monitor the growth of infants, practice guidelines should emphasize the expected, natural, and health-promoting aspects of “slower” growth in breastfed infants during the second 6 months after birth and beyond. Physicians should reassure parents who may be concerned about their infant’s performance on CDC growth charts that the growth patterns of breastfed infants are healthy. Plotting children on the appropriate WHO growth standard may provide both physicians and parents with needed reassurance.

For families choosing to formula feed their infant, parents should be given specific education aimed at reducing problematic bottle-feeding behaviors. For example, parents should be encouraged to feed their infants when they are hungry, rather than on a set schedule outside of the immediate newborn period. Parents should be instructed to be responsive to infant cues for satiety, rather than ensuring their infant finishes the bottle contents. Age-specific guidelines for how much formula should be dispensed at a feeding would help parents start the feed with an appropriate portion size for their infant. Finally,

encouraging parents to discern whether an infant is hungry or needing alternative soothing may reduce overfeeding of the bottle-fed infant.

Sleep

Short sleep duration may have other health effects other than fatigue. There is some evidence that it may be a contributor to the development of obesity. During the past 40 years, sleep duration in the United States has decreased by 1 to 2 hours per day whereas the prevalence of obesity has markedly increased [103–105]. It is estimated that children, a group with a rapid increase in the prevalence of obesity, are currently sleeping 1 to 2 hours less than they require, and that approximately 15 million American children are affected by inadequate sleep [106,107].

The link between short sleep duration and childhood obesity was first shown by a study of French 5-year olds, in which investigators found a significant risk for overweight among children who slept less than 11 hours per day [108]. Since the publication of that study, several investigations have shown that short sleep duration during early childhood (age 3–5 years) is associated with overweight, obesity, and higher body fat during school age [9,11,74,109,110]. Most recently, Taveras and colleagues demonstrated that sleep duration of less than 12 hours during infancy is a risk factor for overweight and adiposity in preschool-aged children [111].

There are several mechanisms by which shorter sleep duration may lead to overweight even among the youngest of children. The first 2 months after birth represent a critical period in the development of sleep patterns, a period during which feeding and sleeping are inextricably linked with infants waking every 2 to 4 hours, typically to feed [112–115]. These first months are also central for the development of normal circadian rhythms [116]. As a result of these rhythms, infants have periods of arousal from sleep, and how parents handle the infant's night waking represents a source of variability in infants' developing nighttime sleep patterns [117,118].

To evaluate different parenting styles and the impact on sleep, St. James-Roberts evaluated a “proximal care” model of parenting infants characterized by prolonged holding, frequent breastfeeding, rapid response to infant frets and cries, and cosleeping with infants at night with other approaches to infant care that have less parent-infant contact per day [117]. He found that the proximal care group of infants had more frequent night waking and crying at age 12 weeks. In another investigation, infants whose parents were present when their child fell asleep were more likely to wake at night than infants whose parents were not present, suggesting that infants who were able to self-soothe in the absence of feeding were more likely to sleep through the night [119].

The early development of sleeping through the night and its association with subsequent weight status is based on several findings. First, children who are unable to achieve a sleep duration of 6 hours by age 5 months have a much greater risk of short sleep duration and sleep problems later in childhood [120,121]. Second, relative sleep duration for age compared with norms for

age was shown to remain constant for approximately 90% of children in a recent longitudinal study of sleep in children [122].

Next, to understand a potential physiologic mechanism for a relationship between sleep and obesity, one might consider adult data that have demonstrated that sleep restriction results in a significant reduction in the anorexigenic hormone, leptin, and an increase in the appetite stimulating peptide, ghrelin [123]. Reduced leptin and increased ghrelin were associated with a significant increase in hunger and appetite. The relationship between short sleep duration, reduced leptin, and increased ghrelin was also found in another investigation with more than 1000 participants whereby the links were shown to exist independent of BMI [124]. Although limited data on this subject exist for infants and children, lower cord blood ghrelin levels have been linked to slower weight gain from age 0 to 3 months [125]. These findings suggest that efforts to increase sleep duration for children could result in lower ghrelin levels, which could limit rapid weight gain during infancy. The importance of research studying potential links between sleep and obesity in children is becoming apparent, and was emphasized in a recent editorial in the *Archives of Internal Medicine* [126]. There, Bass and Turek wrote, “It is now critical to determine the importance of a lack of sleep during the early formative years in putting our youth on a trajectory toward obesity and the metabolic syndrome—a trajectory that could be altered if sleep loss is indeed playing a role in this epidemic.” In summary, the roots of short sleep duration can be found in infancy, and are linked to parenting practices surrounding sleeping and feeding with potential long-term consequences for weight status.

Two seemingly conflicting theories regarding the prevention of obesity intersect in the discussion of sleep. Whereas prolonged sleep duration may be protective for obesity, breastfeeding, which is also protective for obesity, is associated with shorter sleep segments, increased night waking, and reduced total daily sleep [121,127–130]. This shorter sleep duration can persist even after the child has been weaned [121,127]. Breastfed infants also may be more easily aroused from sleep than those that are bottle fed [131]. Further, more frequent night waking has been described as a source of distress for mothers and a cause for them changing from breastfeeding to formula feeding [121,132]. Clearly the relationship between sleep duration, breastfeeding, and weight status is complex, but interventions to promote longer sleep duration such as those now described could achieve objectives of breastfeeding promotion and obesity prevention.

Potential intervention

Because short sleep duration in infancy and childhood is linked to childhood obesity, interventions designed increase sleep duration during infancy may have long-term protective effects against obesity. Of note, some research suggests that sleep duration during infancy can be significantly lengthened with appropriate interventions, even for breastfed infants. An intervention developed by Pinilla and Birch was successful in increasing nocturnal sleep

duration in breastfed infants, who were taught to sleep for at least 5 consecutive hours by age 8 weeks [133]. In this study, parents in the treatment group were given a simple set of instructions to gradually lengthen intervals between middle-of-the-night feedings by performing alternative caregiving behaviors before feeding (eg, reswaddling, diapering, rocking). By age 3 weeks, treatment infants showed significantly longer sleep episodes at night, and by 8 weeks 100% of treatment infants were sleeping a 5-hour duration overnight, compared with 23% of control infants. Of note, Infants made up for the reduced nocturnal milk consumption with a larger early morning feed. One of the interventions described in the study, swaddling, has been shown to calm infants, reduce arousals during sleep, improve sleep efficiency, and reduce spontaneous awakenings [134,135]. In addition, other pediatric practitioners have expanded on these techniques in an attempt to calm and soothe infants, and improve infant sleep through the addition of “white noise,” gentle rocking, and nonnutritive sucking [136,137]. However, like the use of pacifiers, the techniques described in this study should not be attempted until breastfeeding success is well established.

Whereas those who advocate for breastfeeding “on demand” may initially be averse to attempts to lengthen sleep duration, it is important to recognize that there is no uniform definition of “on demand” and mothers may interpret “feeding on demand” in a highly variable fashion. Thus, teaching mothers sleep-lengthening techniques may potentially enhance their ability to distinguish hunger cues from other distress cues, improve feelings of parental competence, reduce formula use, and increase the duration of breastfeeding.

Parental regulation of distress, temperament, and self-regulation of emotion

The ability to regulate behavior and emotion are important developmental tasks [138]. Moreover, the *inability* to successfully regulate one’s emotional state characterizes many childhood problem behaviors [139]. The self-regulation of emotion may also have important implications for other areas of development such as physical health, more specifically weight gain. There are several possible reasons why emotion regulation may be important in understanding and potentially preventing abnormal weight gain. First, infant difficultness, a temperamental quality characterized by fussiness and difficulty soothing, has been related to rapid weight gain [140] and later body composition [141] in childhood, whereas negative child emotionality has been linked to adult body mass [142]. Next, studies of emotion and eating in adults show emotions, particularly negative emotions, to be related to increased eating, prompting theorists to hypothesize that obese individuals eat to reduce discomfort [143]. Finally, there is increasing evidence that mental health problems such as depression and conduct disorder are strong predictors of adulthood obesity [144]. This link may be biologic, as several central nervous system processes responsible for feeding regulation also are involved in regulating emotion

[145]. Taken together, these data suggest that the ability to self-regulate emotion may be an important factor in preventing obesity.

Developmental theorists agree that the soothing environment provided by the parent serves not only to alleviate immediate distress but also facilitates the infant's development of self-regulation [138,146]. Soothing a distressed infant models emotion regulatory strategies and demonstrates the effectiveness of various behaviors for reducing distress. Previous research has shown parental sensitivity, or contingent responsiveness to infant cues, predict infant crying and behavior [147]. That is, parents exhibiting high sensitivity had infants who cried less both concurrently and longitudinally. Furthermore, whereas some researchers have examined the effect of certain soothing techniques on reducing infant distress or supporting concurrent infant regulation [148,149], only one study has demonstrated the carry-over effect of parental regulation to infant self-regulation. Jahromi and Stifter found that mothers who were most effective in soothing their infants at 2 months had infants who cried less with immunization injections 4 months later [148].

Unfortunately from an obesity prevention perspective, many parents use feeding as a method to soothe a distressed infant, and this practice may have negative consequences for weight gain [150]. Using food to reward or punish behavior in children has similarly been associated with binge eating and heavier weight status when those children become adults [151]. Thus, whereas parent regulation of distress is important to the development of the child's ability to self-regulate his or her emotion, the use of food to soothe or reward may contribute to unhealthy outcomes.

Potential intervention

As feeding in infancy is predominantly under the control of the parent, using food in circumstances unrelated to hunger and sustenance may lead to children's understanding that food has other 'rewardlike' qualities. Infants may learn to eat in response to cues other than hunger, such as the presence of food, or their own emotional distress. This compromised ability to self-regulate their food intake may put them at risk for overweight. Given this evidence and that parents report using food to soothe their children [67], parental regulation seems to be a critical variable in the development of children's overall self-regulation skills.

One simple strategy parents can employ is to use nonfood items to soothe during early infancy and nonfood rewards for good behaviors during later infancy. Alternative soothing strategies for young infants, as discussed earlier in relation to sleep, can be applied during the day as well. For older infants, praise or rewards with stickers, toys, book reading, singing songs, playing, or visits to special places may be good alternatives to rewarding good behaviors with food.

As for the core issue of improving parental regulation, the results of 2 studies suggest that the effect of improving parental sensitivity may be most important for the highly reactive infant. In one study, an intervention focused on mothers

of irritable infants found that training mothers to respond contingently and appropriately to the fussing and crying of the infant improved maternal interactive behavior and infant self-soothing, exploration, and attachment [152]. In the second study, an intervention aiming to improve mother-infant attachment found the effect to be greatest for those infants who were highest in negative emotionality [153]. Taken together, these studies suggest that parents should (1) recognize that infant distress is a signal to act, (2) respond quickly as waiting does not reduce crying, and (3) use strategies such as rocking, presenting alternative activities, and providing a pacifier to reduce the crying and fussing. Not only does this ameliorate the infant's distress but, if applied consistently, will enhance the parent-child relationship, and demonstrate how the child might self-regulate his or her own emotions in the future. Promoting improved parental regulation of infant distress as well as less intense negative responses from the infant seems to be essential to the development of the child's ability to self-regulate. Furthermore, interventions that provide the parents with alternative strategies to using food to soothe may not only be practical and economical, but may directly prevent childhood obesity.

Introduction of solid foods

Timing of solid food introduction

The American Academy of Pediatrics (AAP) suggests that the introduction of solid foods should begin between 4 and 6 months because, in general, exclusive milk feeding is adequate to support growth until this time, and supplementary foods are not needed [154,155]. Others add that solids should not be introduced until an infant has good head and neck control and can sit with support [156]. Recent NHANES data suggest, however, that about 30% of breastfed infants and about 50% of formula-fed infants are consuming some infant cereal by 2 to 3 months [157]. Further, data from the Early Childhood Longitudinal Study showed that in a nationally representative cohort, children placed in child care before age 3 months were nearly twice as likely to receive introduction of solids before turning 4 months [158]. Also, many parents continue to add cereal to their infant's bottle, a practice not supported by the AAP because it can lead to excessive weight gain [159]. Related to the previous discussion about sleep and regulation of distress, one commonly attempted approach to improve infant sleep is to give infant cereal at bedtime, a practice that has not been shown to be effective at helping infants sleep despite the widespread belief that the early addition of solids promotes sleeping through the night [160]. Of note, the time when solids are first introduced may be related to increased caloric intake, rapid weight gain, and the subsequent development of overweight.

When the topic of the relationship between the timing of introduction of complementary foods and subsequent obesity has been evaluated in large, population-based studies, there has been minimal to little influence of this variable on the development of obesity [79,80]. However, several more focused studies have revealed important associations. Shukla and colleagues [161] found that

overweight status at 13 weeks was related to the extra calories provided by solid foods early in life, and overweight status at age 13 weeks persisted through 1 year. Von Kries and colleagues [11] similarly showed a relationship between the introduction of solids before age 4 months and the risk of overweight status between the age of 5 and 6 years. Others have further characterized this relationship by showing the association between increased rate of weight gain in the first 6 to 12 months after birth and early introduction of solids, particularly for formula-fed infants [158,162–164]. The analysis of the Danish National Birth Cohort by Baker and colleagues revealed perhaps the most intriguing data on this subject. Their regression analyses showed that the relationship between maternal obesity and greater weight gain during the first year could be nearly eliminated by breastfeeding for more than 40 weeks and delaying the introduction of complementary foods until after age 20 weeks [165].

Method of solid food introduction and the use of repeated exposure

Across cultures, dietary diversity is vast, but all infants begin life consuming the same food: milk. However, as omnivores, they are prepared to learn to consume the diet of their culture. Infants “come equipped” to prefer sweet and salty tastes, to be “neophobic” and reject new foods when they are first offered (at least those that are not sweet or salty), and to learn to like foods of the adult diet of their culture and ethnicity, via various associative conditioning processes that involve the pairing of food with the aspects of the social contexts and physiologic consequences of eating [166]. Given this set of predispositions, all normal infants will readily accept sweet and salty foods such as French fries and sweetened drinks. In contrast, many healthy foods such as pureed vegetables, infant cereals, meats, and dairy products, which are not high in sugar or salt, will be initially rejected by infants. These findings have revealed that infants typically need several opportunities to sample these new foods before intake will increase [167,168]. The liking for complex food flavors that are not dominated by sweet or salty tastes must be learned.

Potential intervention

In addition to delaying the introduction of complementary foods until age 4 to 6 months, parents should be instructed on the infant neophobic response to new foods. Parents often interpret the initial rejection of new foods as “she doesn’t like it,” and the food is not offered again. However, the infant’s initial neophobic response is usually transient, and that infants’, toddlers’, and preschoolers’ intake of new healthy foods will typically increase if the infant has repeated opportunities to taste them [167–170]. As few as 1 or 2 exposures [168] or as many as 5 or 10 exposures [169] may be needed, depending on the age of the child; and all foods will not be accepted by some children, despite repeated exposure. However, parents must be instructed that such repeated exposure is necessary if not sufficient for acceptance of new foods that are not very sweet or salty. Most of the healthy foods that are developmentally appropriate for infants and toddlers fall into this category, including most

vegetables, complex carbohydrates, and meat and dairy products. Although exceptions to these instructions are cases in which food aversions and dislikes develop when consumption is followed by nausea or vomiting, or when eating is coerced or unpleasant, parents need guidance on how to introduce new foods, to increase the likelihood that the child will learn to consume a variety of foods that will constitute a balanced diet promoting healthy weight gain.

In addition to repeated exposure to healthy foods, parents should be alerted to cues for fullness when feeding complementary foods. Pursued lips, closed mouth, spitting out food, turning of the head, and leaning back are examples of cues to stop feeding [156].

OBESITY PREVENTION DURING LATER INFANCY AND THE EARLY TODDLER YEARS

Parent feeding style

During the first years of life, infants and young children are learning an enormous amount about food and eating as they are being introduced to the adult diet of their culture. Growing children learn when to eat, what is food and what is not, how much to eat, and as a result of their experiences with food and eating, are developing food preferences and dislikes. The period from birth to 3 years is also a crucial one for the development of the controls of food intake, and the development of food preferences and eating behaviors [166]. There is evidence that early feeding practices are linked to patterns of food acceptance and the developing controls of food intake [166,171,172]. A feeding style using coercion is unlikely to be successful in the short term and can result in unhealthy weight status in the long term, as recent data suggest that maternal control of feeding during the first years after birth moderates infant weight gain [173,174]. Coercive feeding practices include tactics such as (1) pressuring children to finish their vegetables or (2) restricting their access to sweets and junk food, or using such foods as rewards [71]. In addition to promoting dislikes for healthy foods, and increased liking and wanting of restricted junk foods, coercive practices can also promote dysregulation of intake by promoting over-eating, and learning to eat in response to the presence of food on the plate (as in “finish your vegetables”) [175–177].

Potential intervention

When introducing solids, parents should be aware that new foods may be initially rejected, and should be advised to be patient, as learning to like new foods takes time. If parents offer new foods repeatedly, over a series of 5 or 10 days, many of these foods will eventually be accepted and even preferred [167–169]. However, pressuring or coercive feeding practices engender resistance and foster dislikes, and should be discouraged in favor of providing repeated experience in positive contexts. During the transition to table foods, parents can be effective positive models, promoting infant acceptance and liking for healthy foods by consuming healthy foods and avoiding “junk foods” at family meals. The use of positive feeding practices such as repeated exposure

and modeling are more likely to promote the establishment of healthy diets than the use of coercive feeding practices [178]. Parents should be advised to avoid pressure and restriction in feeding, and encouraged to put their efforts into promoting the liking and acceptance of table foods that are part of a healthy diet.

Transition from human milk or formula to cow's milk

Although consumption of human milk or infant formula may continue for over 1 year, most infants transition to cow's milk at age 1 year. As opposed to children 2 years or older, however, many do not recommend the use of nonfat or low-fat milk for children 12 to 23 months old [179–182]. This recommendation likely stems from concerns from years ago that infants consume a sufficient quantity of dietary fat and essential fatty acids for normal growth and development. However, as opposed to the evidence for infants younger than 12 months, the recommendation for 12- to 23-month-old infants is not evidence based [183].

The question of milk fat intake is important because the preference for high-fat foods develops early in life, and children still consume 20% to 25% of their calories from cow's milk between the ages of 1 and 2 years [184,185]. Further, a majority of preschool children drink whole milk, particularly those who are from minorities or low-income households [186,187].

Potential intervention

Reducing milk fat intake has been previously identified as a target for dietary fat reduction and obesity prevention in toddlers and preschool children [188–190], and reducing fat intake among infants and toddlers has been shown to be well tolerated without adverse effects on growth, nutrient intake, or development [191–193]. In contrast, high fat intake between 12 and 23 months of age has been associated with inhibition of the normal decrease in body fat between the ages of 2 and 5 years [194]. Given that contemporary diets for children older than 1 year typically contain a significant amount of dietary fat from nonmilk sources [195], the common advice to only use whole milk for children 12 to 23 months old is no longer necessary for infants who consume a well-balanced diet [186]. In fact, the 2008 AAP policy statement “Lipid Screening and Cardiovascular Health in Childhood” includes as its first recommendation the use of reduced fat milk for 12- to 23-month-old children [196]. Although studies of older children have suggested that consumption of dairy fat is not associated with weight gain [197], because each cup of whole milk contains 146 kcal compared with only 86 kcal per cup of nonfat milk [198], a change to low-fat or even nonfat milk could reduce energy consumption without adverse effects for infants.

Sweet beverage consumption

Over the past 3 decades, the percentage of daily calories obtained from soft drinks, fruit drinks, sweetened beverages, and fruit juice each have steadily increased among American children [199]. Among children 12 to 24 months

old, 100% juice and sweetened beverages account for the second and third greatest source of dietary calories, respectively, exceeded only by milk [185]. Although data have in general not supported a link between consumption of 100% juice and obesity unless consumed in excess [200], there is a strong association between sweetened beverages and soft drinks with obesity [201]. Further, as suggested by Skinner and colleagues [202], because early life diet preferences predict later preferences, the development of beverage consumption patterns before age 2 years is important for long-term consumption.

Regarding infant and toddler consumption of juice, the AAP has recommended that juice (a) not be given to infants younger than 6 months, (b) only be given to infants that can drink from a cup and never given in a bottle, (c) not be given in a fashion that allows for a child to easily transport it with easy consumption throughout the day, and (d) not be given in quantities greater than 6 ounces (170 g) per day [203]. Data from the Feeding Infants and Toddlers Study (FITS), a national dietary survey, demonstrated that many infants consume fruit juice before 6 months of age, and exceed the recommended amount of juice later during infancy and the toddler years [202].

Unlike fruit juice, the AAP has no formal policy statement or recommendation regarding the consumption of other sweetened beverages and soft drinks prior to school entry, but the Institute of Medicine has recommended the avoidance of high-calorie, nutrient-poor beverages [5]. Of note, consumption of sweet drinks has been associated with obesity even among preschool children [204,205]. In addition, in FITS, sweetened drinks and soda also were being consumed at the expense of milk in this sample of infants and toddlers [203]. Further, several investigators have suggested that reducing intake of sweetened beverages is one of the most important and promising obesity prevention strategies [204–207].

Potential intervention

Given the current evidence, adherence to the AAP policy regarding fruit juice is appropriate. Further, because they have no nutritional benefit and are associated with obesity and other morbidities, the use of sweetened beverages, soft drinks, and fruit drinks should be totally discouraged for infants and toddlers. Given the alternatives of milk, water, and the now increasingly available low-calorie flavored water, there are sufficient options for parents to give their young children.

Transitional feeding and table foods

The transition from the exclusive milk diet of infancy to a modified adult diet is completed by age 3 years, and during this early period children already have begun acquiring food preferences and aversions [166]. Children are learning a great deal about when, what, and how much to eat. Unfortunately, recent data from FITS revealed that energy intakes among typical infants and toddlers exceed requirements by 20% to 30% [208]. Of note, in FITS this excess already was apparent in infants aged 4 to 6 months, in whom intake exceeded energy requirements by 10%, suggesting that patterns of intake promoting excessive

weight gain and obesity were being established at an early age. In addition to consuming too much energy, from an early age (4 to 24 months) children consume significant amounts of developmentally inappropriate foods, high in energy density and low in nutrients, while consuming too few of the foods that should form the basis of a healthy weaning diet [195]. The high energy density of foods offered appeared to be the same ones that contribute to energy intakes that were in excess of energy requirements in other age groups studied [208]. For example, in children 7 and 24 months old, 18% and 33%, respectively, consumed no servings of vegetables during a given 24-hour period. Twenty-three percent of 7-month-old and 33% of 24-month-old children did not consume any fruits. Further, parents reported that French fries were the third most common vegetable consumed by infants 9 to 11 months old, and by 15 to 18 months were the most common vegetable consumed. The diets of infants in this age group may, not surprisingly, be deficient of key nutrients [209].

Potential intervention

The FITS findings underscore the need to provide parents with anticipatory guidance regarding the transition to a modified adult diet. The Start Healthy Feeding Guidelines for Infants and Toddlers provides excellent guidance on the nutritional needs of children younger than 2 years [210]. As opposed to giving children French fries, parents should be encouraged to meet these nutritional requirements with fresh fruits, cooked vegetables, cheese, yogurt, whole-grain breads and crackers, and cereals [156,211,212]. All of these foods should not have added sugar or salt.

Physical activity and sedentary behaviors

There is substantial epidemiologic evidence that regular physical activity is essential for good physical and psychological health and disease prevention among children and adults. However, to date the recommendations for infant and toddler physical activity (from birth to age 5 years) have not been evidence based. Rather, the guidelines developed by the National Association for Sport and Physical Education [213] were adapted from evidence accumulated among older children and adolescents [214]. The recommendations for infants suggest that they should be: (1) interacting with caregivers in daily physical activities that promote movement and exploration of their environment, (2) engaging in activities that promote the development of movement skills and large muscle activities, and (3) placed in safe settings that facilitate physical activity and do not restrict physical activity for prolonged periods of time. These recommendations suggest that toddlers should: (1) accumulate at least 30 minutes of daily structured physical activity, (2) engage in at least 60 minutes and up to several hours a day of unstructured physical activity and should not be sedentary for more than 60 minutes at a time except when sleeping, and (3) develop movement skills that are building blocks for more complex movement behaviors [213]. Nonetheless, because physical activity and sedentary patterns, much like feeding and sleeping behaviors, become established in the early years, there

is increasing support for promoting physical activity and reducing sedentary behaviors as soon as possible during the early infancy period [213,215].

Environmental risk factors associated with sedentary activities during the early infancy and toddler periods may predispose children to low levels of physical activity in later childhood. For example, restricting infants to car seats, swings, carriers, strollers, and small play spaces for long periods of time may limit motor development and delay physical activity such as crawling and walking [213,216]. Also, limited time for leisure activities and parent-child play, concern for neighborhood safety, and using television or computer games to occupy a child's attention may promote sedentary lifestyle during the first 3 years after birth [216,217]. Television viewing during the infancy and toddler years seems to be one particular sedentary practice that is an environmental risk factor for obesity development. It is reported that 82% of 1-year-old (11 average hours per week) and 96% of 2-year-old (15 average hours per week) children watch television or videos, and having a television in the bedroom elevates the risk of being overweight [218]. Also, Taveras and colleagues found that children who slept less than 12 hours per day and viewed 2 hours per day or more of television had an obesity probability at 3 years old of 17% [111].

Potential intervention

Although there is little debate that physical activity is important for obesity prevention, questions remain about how much and what types of physical activity during the infant and early toddler years is needed to prevent later obesity. In general, parents of infants should choose physical activities that are interactive, stimulating, easy to do, and incorporated into their daily routine, to reinforce the concept that physical activity is rewarding [213,215].

In addition, 2 existing programs offer possible blueprints for more extensive prevention efforts. The Infant Feeding Activity and Nutrition Trial is an early intervention to prevent childhood obesity that targets infants through age 18 months in Victoria, Australia. This program teaches parents about the development of positive diet and physical activity behaviors while reducing sedentary behaviors during infancy [219]. Parents in the program learn about age-appropriate physical activity behaviors, the risks associated with sedentary behaviors, and how parental modeling of physical activity and sedentary behaviors influences their child's behaviors.

The second program, Fighting Fit Tots, also aims to prevent obesity in the first years of life by promoting toddler physical activity and parent lifestyle education [220]. This program is currently being implemented in Lambeth, South London (United Kingdom) and consists of 11 weekly 2-hour sessions that include guided parent-child physical activity (eg, jumping, skipping, hopping, dancing, singing), snack (water and fruit), and parental education (eg, healthy lifestyle workshop) aimed at increasing physical activity behaviors, confidence, and decreasing BMI and waist circumference among children 18 to 30 months old.

Regarding sedentary behavior, one clear target for intervention is television viewing. Because most children watch television by age 2 years, educational efforts and interventions about limiting television/video viewing need to be implemented before this age to potentially impact overweight development. Of note, the AAP discourages television watching for children younger than 2 years, and suggests that televisions never be placed in children's bedrooms [221].

SUMMARY

Given the vast array of topics that are important to cover at infant health maintenance visits, extensive discussion about growth, growth charts, and healthy lifestyle may be challenging for providers. Nonetheless, obesity and its comorbidities threaten both individual patients and the health care system. To break the vicious cycle of obese children becoming obese adults who have obese offspring, preventing behaviors that lead to obesity must be implemented during the very earliest periods of life, the prenatal period and infancy. For pediatric care providers, there are numerous opportunities to intervene, and good communication with families about healthy growth and lifestyle are a promising beginning.

References

- [1] Ogden CL, Carroll MD, Flegal KM. High body mass index for age among US children and adolescents, 2003–2006. *JAMA* 2008;299:2401–5.
- [2] Ogden CL, Flegal KM, Carroll MD, et al. Prevalence and trends in overweight among US children and adolescents, 1999–2000. *JAMA* 2002;288:1728–32.
- [3] Mei Z, Scanlon KS, Grummer-Strawn LM, et al. Increasing prevalence of overweight among US low-income preschool children: the Centers for Disease Control and Prevention pediatric nutrition surveillance, 1983 to 1995. *Pediatrics* 1998;101:E12.
- [4] Kim J, Peterson KE, Scanlon KS, et al. Trends in overweight from 1980 through 2001 among preschool-aged children enrolled in a health maintenance organization. *Obesity (Silver Spring)* 2006;14:1107–12.
- [5] Koplan JP, Liverman CT, Kraak VI, editors. Preventing childhood obesity: health in the balance. Washington, DC: The National Academies Press; 2005.
- [6] Lederman SA, Akabas SR, Moore BJ. Editors' overview of the conference on preventing childhood obesity. *Pediatrics* 2004;114:1139–45.
- [7] Eid EE. Follow-up study of physical growth of children who had excessive weight gain in first 6 months of life. *Br Med J* 1970;2:74–6.
- [8] Ong KK, Ahmed ML, Emmett PM, et al. Association between postnatal catch-up growth and obesity in childhood: prospective cohort study. *BMJ* 2000;320:967–71.
- [9] Reilly JJ, Armstrong J, Dorosty AR, et al. Early life risk factors for obesity in childhood: cohort study. *BMJ* 2005;330:1358–60.
- [10] Stettler N, Zemel BS, Kumanyika S, et al. Infant weight gain and childhood overweight status in a multicenter, cohort study. *Pediatrics* 2002;109:194–9.
- [11] von Kries R, Toschke AM, Wurmser H, et al. Reduced risk for overweight and obesity in 5- and 6-y-old children by duration of sleep—a cross-sectional study. *Int J Obes Relat Metab Disord* 2002;26:710–6.
- [12] Cameron N, Pettifor J, De Wet T, et al. The relationship of rapid weight gain in infancy to obesity and skeletal maturity in childhood. *Obes Res* 2003;11:457–60.

- [13] Mellbin T, Vuille JC. Physical development at 7 years of age in relation to velocity of weight gain in infancy with special reference to incidence of overweight. *Br J Prev Soc Med* 1973;27:225–35.
- [14] Stettler N, Kumanyika SK, Katz SH, et al. Rapid weight gain during infancy and obesity in young adulthood in a cohort of African Americans. *Am J Clin Nutr* 2003;77:1374–8.
- [15] Ekelund U, Ong K, Linne Y, et al. Upward weight percentile crossing in infancy and early childhood independently predicts fat mass in young adults: the Stockholm Weight Development Study (SWEDES). *Am J Clin Nutr* 2006;83:324–30.
- [16] Monteiro PO, Victora CG, Barros FC, et al. Birth size, early childhood growth, and adolescent obesity in a Brazilian birth cohort. *Int J Obes Relat Metab Disord* 2003;27:1274–82.
- [17] Wells JC, Hallal PC, Wright A, et al. Fetal, infant and childhood growth: relationships with body composition in Brazilian boys aged 9 years. *Int J Obes (Lond)* 2005;29:1192–8.
- [18] Sachdev HS, Fall CH, Osmond C, et al. Anthropometric indicators of body composition in young adults: relation to size at birth and serial measurements of body mass index in childhood in the New Delhi birth cohort. *Am J Clin Nutr* 2005;82:456–66.
- [19] Blair NJ, Thompson JM, Black PN, et al. Risk factors for obesity in 7-year-old European children: the Auckland Birthweight Collaborative Study. *Arch Dis Child* 2007;92:866–71.
- [20] Dennison BA, Edmunds LS, Stratton HH, et al. Rapid infant weight gain predicts childhood overweight. *Obesity (Silver Spring)* 2006;14:491–9.
- [21] Hui LL, Schooling CM, Leung SS, et al. Birth weight, infant growth, and childhood body mass index: Hong Kong's children of 1997 birth cohort. *Arch Pediatr Adolesc Med* 2008;162:212–8.
- [22] Botton J, Heude B, Maccario J, et al. Postnatal weight and height growth velocities at different ages between birth and 5 y and body composition in adolescent boys and girls. *Am J Clin Nutr* 2008;87:1760–8.
- [23] Yliharsila H, Kajantie E, Osmond C, et al. Body mass index during childhood and adult body composition in men and women aged 56–70 y. *Am J Clin Nutr* 2008;87:1769–75.
- [24] Chomtho S, Wells JC, Williams JE, et al. Infant growth and later body composition: evidence from the 4-component model. *Am J Clin Nutr* 2008;87:1776–84.
- [25] Singhal A, Cole TJ, Fewtrell M, et al. Is slower early growth beneficial for long-term cardiovascular health? *Circulation* 2004;109:1108–13.
- [26] Huxley RR, Shiell AW, Law CM. The role of size at birth and postnatal catch-up growth in determining systolic blood pressure: a systematic review of the literature. *J Hypertens* 2000;18:815–31.
- [27] Law CM, Shiell AW, Newsome CA, et al. Fetal, infant, and childhood growth and adult blood pressure: a longitudinal study from birth to 22 years of age. *Circulation* 2002;105:1088–92.
- [28] Parker L, Lamont DW, Unwin N, et al. A lifecourse study of risk for hyperinsulinaemia, dyslipidaemia and obesity (the central metabolic syndrome) at age 49–51 years. *Diabet Med* 2003;20:406–15.
- [29] Eriksson JG, Forsen T, Tuomilehto J, et al. Catch-up growth in childhood and death from coronary heart disease: longitudinal study. *BMJ* 1999;318:427–31.
- [30] Barker DJ, Osmond C, Forsen TJ, et al. Trajectories of growth among children who have coronary events as adults. *N Engl J Med* 2005;353:1802–9.
- [31] Forsen T, Eriksson J, Tuomilehto J, et al. The fetal and childhood growth of persons who develop type 2 diabetes. *Ann Intern Med* 2000;133:176–82.
- [32] Bhargava SK, Sachdev HS, Fall CH, et al. Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N Engl J Med* 2004;350:865–75.
- [33] Fisch RO, Bilek MK, Ulstrom R. Obesity and leanness at birth and their relationship to body habitus in later childhood. *Pediatrics* 1975;56:521–8.
- [34] Charney E, Goodman HC, McBride M, et al. Childhood antecedents of adult obesity. Do chubby infants become obese adults? *N Engl J Med* 1976;295:6–9.

- [35] Whitaker RC, Wright JA, Pepe MS, et al. Predicting obesity in young adulthood from childhood and parental obesity. *N Engl J Med* 1997;337:869-73.
- [36] Guo SS, Wu W, Chumlea WC, et al. Predicting overweight and obesity in adulthood from body mass index values in childhood and adolescence. *Am J Clin Nutr* 2002;76:653-8.
- [37] Mei Z, Grummer-Strawn LM, Scanlon KS. Does overweight in infancy persist through the preschool years? An analysis of CDC Pediatric Nutrition Surveillance System data. *Soz Praventivmed* 2003;48:161-7.
- [38] Sayer AA, Syddall HE, Dennison EM, et al. Birth weight, weight at 1 y of age, and body composition in older men: findings from the Hertfordshire Cohort Study. *Am J Clin Nutr* 2004;80:199-203.
- [39] Freedman DS, Khan LK, Serdula MK, et al. The relation of childhood BMI to adult adiposity: the Bogalusa heart study. *Pediatrics* 2005;115:22-7.
- [40] Nader PR, O'Brien M, Houts R, et al. Identifying risk for obesity in early childhood. *Pediatrics* 2006;118:e594-601.
- [41] Vogels N, Posthumus DL, Mariman EC, et al. Determinants of overweight in a cohort of Dutch children. *Am J Clin Nutr* 2006;84:717-24.
- [42] Jouret B, Ahluwalia N, Cristini C, et al. Factors associated with overweight in preschool-age children in southwestern France. *Am J Clin Nutr* 2007;85:1643-9.
- [43] Serdula MK, Ivery D, Coates RJ, et al. Do obese children become obese adults? A review of the literature. *Prev Med* 1993;22:167-77.
- [44] Garn SM, LaVelle M. Two-decade follow-up of fatness in early childhood. *Am J Dis Child* 1985;139:181-5.
- [45] Rolland-Cachera MF, Deheeger M, Guillaud-Bataille M, et al. Tracking the development of adiposity from one month of age to adulthood. *Ann Hum Biol* 1987;14:219-29.
- [46] Muramatsu S, Sato Y, Miyao M, et al. A longitudinal study of obesity in Japan: relationship of body habitus between at birth and at age 17. *Int J Obes* 1990;14:39-45.
- [47] Singhal A, Lucas A. Early origins of cardiovascular disease: is there a unifying hypothesis? *Lancet* 2004;363:1642-5.
- [48] Gillman MW. The first months of life: a critical period for development of obesity. *Am J Clin Nutr* 2008;87:1587-9.
- [49] Oken E, Taveras EM, Kleinman KP, et al. Gestational weight gain and child adiposity at age 3 years. *Am J Obstet Gynecol* 2007;196(322):e1-8.
- [50] Whitaker RC. Predicting preschooler obesity at birth: the role of maternal obesity in early pregnancy. *Pediatrics* 2004;114:e29-36.
- [51] Gillman MW, Rifas-Shiman SL, Kleinman K, et al. Developmental origins of childhood overweight: potential public health impact. *Obesity (Silver Spring)* 2008;16:1651-6.
- [52] Jaffe M, Kosakov C. The motor development of fat babies. *Clin Pediatr (Phila)* 1982;21:619-21.
- [53] Shibli R, Rubin L, Akons H, et al. Morbidity of overweight (> or =85th percentile) in the first 2 years of life. *Pediatrics* 2008;122:267-72.
- [54] Wake M, Hardy P, Sawyer MG, et al. Comorbidities of overweight/obesity in Australian preschoolers: a cross-sectional population study. *Arch Dis Child* 2008;93:502-7.
- [55] Taveras EM, Rifas-Shiman SL, Camargo CA Jr, et al. Higher adiposity in infancy associated with recurrent wheeze in a prospective cohort of children. *J Allergy Clin Immunol* 2008;121:1161-6, e3.
- [56] Jain A, Sherman SN, Chamberlin LA, et al. Why don't low-income mothers worry about their preschoolers being overweight? *Pediatrics* 2001;107:1138-46.
- [57] Sherry B, McDivitt J, Birch LL, et al. Attitudes, practices, and concerns about child feeding and child weight status among socioeconomically diverse white, Hispanic, and African-American mothers. *J Am Diet Assoc* 2004;104:215-21.
- [58] Jackson J, Strauss CC, Lee AA, et al. Parents' accuracy in estimating child weight status. *Addict Behav* 1990;15:65-8.

- [59] Baughcum AE, Chamberlin LA, Deeks CM, et al. Maternal perceptions of overweight preschool children. *Pediatrics* 2000;106:1380–6.
- [60] Maynard LM, Galuska DA, Blanck HM, et al. Maternal perceptions of weight status of children. *Pediatrics* 2003;111:1226–31.
- [61] Jeffery AN, Voss LD, Metcalf BS, et al. Parents' awareness of overweight in themselves and their children: cross sectional study within a cohort (EarlyBird 21). *BMJ* 2005;330:23–4.
- [62] Carnell S, Edwards C, Croker H, et al. Parental perceptions of overweight in 3–5 y olds. *Int J Obes (Lond)* 2005;29:353–5.
- [63] Eckstein KC, Mikhail LM, Ariza AJ, et al. Parents' perceptions of their child's weight and health. *Pediatrics* 2006;117:681–90.
- [64] Campbell MW, Williams J, Hampton A, et al. Maternal concern and perceptions of overweight in Australian preschool-aged children. *Med J Aust* 2006;184:274–7.
- [65] Reifsnider E, Flores-Vela AR, Beckman-Mendez D, et al. Perceptions of children's body sizes among mothers living on the Texas-Mexico border (La Frontera). *Public Health Nurs* 2006;23:488–95.
- [66] Kramer MS, Barr RG, Leduc DG, et al. Maternal psychological determinants of infant obesity. Development and testing of two new instruments. *J Chronic Dis* 1983;36:329–35.
- [67] Baughcum AE, Burklow KA, Deeks CM, et al. Maternal feeding practices and childhood obesity: a focus group study of low-income mothers. *Arch Pediatr Adolesc Med* 1998;152:1010–4.
- [68] Bentley M, Gavin L, Black MM, et al. Infant feeding practices of low-income, African-American, adolescent mothers: an ecological, multigenerational perspective. *Soc Sci Med* 1999;49:1085–100.
- [69] Contento IR, Basch C, Zybert P. Body image, weight, and food choices of Latina women and their young children. *J Nutr Educ Behav* 2003;35:236–48.
- [70] Boyington JA, Johnson AA. Maternal perception of body size as a determinant of infant adiposity in an African-American community. *J Natl Med Assoc* 2004;96:351–62.
- [71] Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. *Pediatrics* 1998;101:539–49.
- [72] Bruss MB, Morris J, Dannison L. Prevention of childhood obesity: sociocultural and familial factors. *J Am Diet Assoc* 2003;103:1042–5.
- [73] Karaolis-Danckert N, Buyken AE, Bolzenius K, et al. Rapid growth among term children whose birth weight was appropriate for gestational age has a longer lasting effect on body fat percentage than on body mass index. *Am J Clin Nutr* 2006;84:1449–55.
- [74] Lederman SA, Akabas SR, Moore BJ, et al. Summary of the presentations at the conference on preventing childhood obesity, 2003. *Pediatrics* 2004;114:1146–73.
- [75] Owen CG, Martin RM, Whincup PH, et al. Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics* 2005;115:1367–77.
- [76] Owen CG, Martin RM, Whincup PH, et al. The effect of breastfeeding on mean body mass index throughout life: a quantitative review of published and unpublished observational evidence. *Am J Clin Nutr* 2005;82:1298–307.
- [77] Arenz S, Ruckerl R, Koletzko B, et al. Breast-feeding and childhood obesity—a systematic review. *Int J Obes Relat Metab Disord* 2004;28:1247–56.
- [78] Dewey KG. Is breastfeeding protective against child obesity? *J Hum Lact* 2003;19:9–18.
- [79] Hediger ML, Overpeck MD, Kuczmarski RJ, et al. Association between infant breastfeeding and overweight in young children. *JAMA* 2001;285:2453–60.
- [80] Gillman MW, Rifas-Shiman SL, Camargo CA Jr, et al. Risk of overweight among adolescents who were breastfed as infants. *JAMA* 2001;285:2461–7.
- [81] Bogen DL, Hanusa BH, Whitaker RC. The effect of breastfeeding with and without concurrent formula feeding on the risk of obesity at 4 years of age: a retrospective cohort study. *Obes Res* 2004;12:1527–35.

- [82] von Kries R, Koletzko B, Sauerwald T, et al. Breast feeding and obesity: cross sectional study. *BMJ* 1999;319:147–50.
- [83] Harder T, Bergmann R, Kallischnigg G, et al. Duration of breastfeeding and risk of overweight: a meta-analysis. *Am J Epidemiol* 2005;162:397–403.
- [84] Dewey KG, Heinig MJ, Nommsen LA, et al. Maternal versus infant factors related to breast milk intake and residual milk volume: the DARLING study. *Pediatrics* 1991;87:829–37.
- [85] Wright P, Fawcett J, Crow R. The development of differences in the feeding behaviour of bottle and breast fed human infants from birth to two months. *Behav Processes* 1980;5:1–20.
- [86] Dewey KG, Nommsen-Rivers LA, Lonnerdal B. Plasma insulin and insulin-releasing amino acid (IRAA) concentrations are higher in formula fed than breastfed infants at 5 months of age [abstract]. Experimental Biology meeting, 2004.
- [87] Jensen RG, editor. Handbook of milk composition. San Diego (CA): Academic Press; 1995.
- [88] Hambraeus L. Proprietary milk versus human breast milk in infant feeding. A critical appraisal from the nutritional point of view. *Pediatr Clin North Am* 1977;24:17–36.
- [89] Offedal OT, Iverson SJ. Comparative analysis of nonhuman milks: phylogenetic variation in the gross composition of milks. In: Jensen RG, editor. Handbook of milk composition. San Diego (CA): Academic Press; 1995. p. 749–89.
- [90] EU Childhood Obesity Programme. EU Childhood Obesity Programme press pack. Budapest, Hungary: Danone Institute; 2007.
- [91] Read LC, Penttila IA, Howarth GS, et al. Role and function of growth factors in infant nutrition. In: Raiha NCR, Rubaltelli FF, editors. Infant formula: closer to the reference. Philadelphia: Vevey/Lippincott Williams & Wilkins; 2002. p. 185–95.
- [92] Savino F, Fissore MF, Grassino EC, et al. Ghrelin, leptin and IGF-I levels in breast-fed and formula-fed infants in the first years of life. *Acta Paediatr* 2005;94:531–7.
- [93] Elmlinger MW, Hochhaus F, Loui A, et al. Insulin-like growth factors and binding proteins in early milk from mothers of preterm and term infants. *Horm Res* 2007;68:124–31.
- [94] Martin LJ, Woo JG, Geraghty SR, et al. Adiponectin is present in human milk and is associated with maternal factors. *Am J Clin Nutr* 2006;83:1106–11.
- [95] Dundar NO, Anal O, Dundar B, et al. Longitudinal investigation of the relationship between breast milk leptin levels and growth in breast-fed infants. *J Pediatr Endocrinol Metab* 2005;18:181–7.
- [96] Miralles O, Sanchez J, Palou A, et al. A physiological role of breast milk leptin in body weight control in developing infants. *Obesity (Silver Spring)* 2006;14:1371–7.
- [97] Dewey KG. Growth characteristics of breast-fed compared to formula-fed infants. *Biol Neonate* 1998;74:94–105.
- [98] Dewey KG, Heinig MJ, Nommsen LA, et al. Breast-fed infants are leaner than formula-fed infants at 1 y of age: the DARLING study. *Am J Clin Nutr* 1993;57:140–5.
- [99] American Academy of Pediatrics policy statement. breastfeeding and the use of human milk. *Pediatrics* 2005;115:496–506.
- [100] de Onis M, Garza C, Victora CG, et al. The WHO Multicentre Growth Reference Study: planning, study design, and methodology. *Food Nutr Bull* 2004;25:S15–26.
- [101] de Onis M, Garza C, Onyango AW, et al. Comparison of the WHO child growth standards and the CDC 2000 growth charts. *J Nutr* 2007;137:144–8.
- [102] Shealy K, Li R, Benton-Davis S, et al. The CDC guide to breastfeeding interventions. Atlanta: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2005.
- [103] National Sleep Foundation. “Sleep in America” poll. Washington, DC: National Sleep Foundation; 2000.
- [104] National Sleep Foundation. “Sleep in America” poll. Washington, DC: National Sleep Foundation; 2001.

- [105] National Sleep Foundation. "Sleep in America" poll. Washington, DC: National Sleep Foundation; 2002.
- [106] National Sleep Foundation. "Sleep in America" poll. Washington, DC: National Sleep Foundation; 2004.
- [107] Smaldone A, Honig JC, Byrne MW. Sleepless in America: inadequate sleep and relationships to health and well-being of our nation's children. *Pediatrics* 2007;119(Suppl 1): S29-37.
- [108] Locard E, Mamelle N, Billette A, et al. Risk factors of obesity in a five year old population. Parental versus environmental factors. *Int J Obes Relat Metab Disord* 1992;16:721-9.
- [109] Sekine M, Yamagami T, Hamanishi S, et al. Parental obesity, lifestyle factors and obesity in preschool children: results of the Toyama Birth Cohort study. *J Epidemiol* 2002;12:33-9.
- [110] Agras WS, Hammer LD, McNicholas F, et al. Risk factors for childhood overweight: a prospective study from birth to 9.5 years. *J Pediatr* 2004;145:20-5.
- [111] Taveras EM, Rifas-Shiman SL, Oken E, et al. Short sleep duration in infancy and risk of childhood overweight. *Arch Pediatr Adolesc Med* 2008;162:305-11.
- [112] Moore T, Ucko LE. Night waking in early infancy. *Arch Dis Child* 1957;32:333-42.
- [113] Parmelee AH, Wenner WH, Schulz HR. Infant sleep patterns: from birth to 16 weeks of age. *J Pediatr* 1964;65:576-82.
- [114] Anders TF, Keener M. Developmental course of nighttime sleep-wake patterns in full-term and premature infants during the first year of life. I. *Sleep* 1985;8:173-92.
- [115] James-Roberts IS, Conroy S, Hurry J. Links between infant crying and sleep-waking at six weeks of age. *Early Hum Dev* 1997;48:143-52.
- [116] Glotzbach SF, Edgar DM, Boeddiker M, et al. Biological rhythmicity in normal infants during the first 3 months of life. *Pediatrics* 1994;94:482-8.
- [117] St James-Roberts I, Alvarez M, Csipke E, et al. Infant crying and sleeping in London, Copenhagen and when parents adopt a "proximal" form of care. *Pediatrics* 2006;117:e1146-55.
- [118] Anders TF. Night-waking in infants during the first year of life. *Pediatrics* 1979;63:860-4.
- [119] Adair R, Bauchner H, Philipp B, et al. Night waking during infancy: role of parental presence at bedtime. *Pediatrics* 1991;87:500-4.
- [120] Touchette E, Petit D, Paquet J, et al. Factors associated with fragmented sleep at night across early childhood. *Arch Pediatr Adolesc Med* 2005;159:242-9.
- [121] Wolke D, Meyer R, Ohrt B, et al. Co-morbidity of crying and feeding problems with sleeping problems during infancy: concurrent and predictive associations. *Early Dev Parenting* 1995;4:191-208.
- [122] Jenni OG, Molinari L, Caflisch JA, et al. Sleep duration from ages 1 to 10 years: variability and stability in comparison with growth. *Pediatrics* 2007;120:e769-76.
- [123] Spiegel K, Tasali E, Penev P, et al. Brief communication: sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* 2004;141:846-50.
- [124] Taheri S, Lin L, Austin D, et al. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* 2004;1:e62.
- [125] James RJ, Drewett RF, Cheetham TD. Low cord ghrelin levels in term infants are associated with slow weight gain over the first 3 months of life. *J Clin Endocrinol Metab* 2004;89: 3847-50.
- [126] Bass J, Turek FW. Sleepless in America: a pathway to obesity and the metabolic syndrome? *Arch Intern Med* 2005;165:15-6.
- [127] Wright P, MacLeod HA, Cooper MJ. Waking at night: the effect of early feeding experience. *Child Care Health Dev* 1983;9:309-19.
- [128] Elias MF, Nicolson NA, Bora C, et al. Sleep/wake patterns of breast-fed infants in the first 2 years of life. *Pediatrics* 1986;77:322-9.
- [129] Eaton-Evans J, Dugdale AE. Sleep patterns of infants in the first year of life. *Arch Dis Child* 1988;63:647-9.

- [130] Scher A, Tirosh E, Jaffe M, et al. Sleep patterns of infants and young children in Israel. *Int J Behav Dev* 1995;18:701–11.
- [131] Horne RS, Parslow PM, Ferens D, et al. Comparison of evoked arousability in breast and formula fed infants. *Arch Dis Child* 2004;89:22–5.
- [132] Ball HL. Breastfeeding, bed-sharing, and infant sleep. *Birth* 2003;30:181–8.
- [133] Pinilla T, Birch LL. Help me make it through the night: behavioral entrainment of breast-fed infants' sleep patterns. *Pediatrics* 1993;91:436–44.
- [134] Gerard CM, Harris KA, Thach BT. Spontaneous arousals in supine infants while swaddled and unswaddled during rapid eye movement and quiet sleep. *Pediatrics* 2002;110:e70.
- [135] Franco P, Seret N, Van Hees JN, et al. Influence of swaddling on sleep and arousal characteristics of healthy infants. *Pediatrics* 2005;115:1307–11.
- [136] Karp H. *The happiest baby on the block*. New York: Bantam Books; 2002.
- [137] Karp H. The "fourth trimester": a framework and strategy for understanding and resolving colic. *Contemp Pediatr* 2004;21:94–116.
- [138] Kopp CB. Regulation of distress and negative emotions: a developmental review. *Dev Psychol* 1989;25:343–54.
- [139] Loeber R, Hay D. Key issues in the development of aggression and violence from childhood to early adulthood. *Annu Rev Psychol* 1997;48:371–410.
- [140] Carey W, Hegvik R, McDevitt S. Temperamental factors associated with rapid weight gain and obesity in middle childhood. *J Dev Behav Pediatr* 1988;9:197–8.
- [141] Wells JC, Stanley M, Laidlaw AS, et al. Investigation of the relationship between infant temperament and later body composition. *Int J Obes Relat Metab Disord* 1997;21:400–6.
- [142] Pulkki-Raback L, Elovainio M, Kivimaki M, et al. Temperament in childhood predicts body mass in adulthood: the Cardiovascular Risk in Young Finns Study. *Health Psychol* 2005;24:307–15.
- [143] Canetti L, Bachar E, Berry EM. Food and emotion. *Behav Processes* 2002;60:157–64.
- [144] Pine DS, Cohen P, Brook J, et al. Psychiatric symptoms in adolescence as predictors of obesity in early adulthood: a longitudinal study. *Am J Public Health* 1997;87:1303–10.
- [145] Kishi T, Elmquist JK. Body weight is regulated by the brain: a link between feeding and emotion. *Mol Psychiatry* 2005;10:132–46.
- [146] Thompson RA. Emotion regulation: a theme in search of definition. *Monogr Soc Res Child Dev* 1994;59:25–52.
- [147] Crockenberg S, Leerkes E. The family context of infant development. In: Zenah C, editor. *Handbook of infant mental health*. New York: Guilford; 1999. p. 60–90.
- [148] Jahromi LB, Putnam SP, Stifter CA. Maternal regulation of infant reactivity from 2 to 6 months. *Dev Psychol* 2004;40:477–87.
- [149] Crockenberg SC, Leerkes EM. Infant and maternal behaviors regulate infant reactivity to novelty at 6 months. *Dev Psychol* 2004;40:1123–32.
- [150] Baughcum AE, Powers SW, Johnson SB, et al. Maternal feeding practices and beliefs and their relationships to overweight in early childhood. *J Dev Behav Pediatr* 2001;22:391–408.
- [151] Puhl RM, Schwartz MB. If you are good you can have a cookie: how memories of childhood food rules link to adult eating behaviors. *Eat Behav* 2003;4:283–93.
- [152] van den Boom D. Behavioral management of early infant crying in irritable babies. In: Barr RG, StJames-Roberts I, Keefe MR, et al, editors. *New evidence on unexplained infant crying: its origins, nature, and management*. New Brunswick (NJ): Johnson & Johnson Pediatric Institute; 2001. p. 209–28.
- [153] Velderman MK, Bakermans-Kranenburg MJ, Juffer F, van IMH. Effects of attachment-based interventions on maternal sensitivity and infant attachment: differential susceptibility of highly reactive infants. *J Fam Psychol* 2006;20:266–74.
- [154] Chapter 6: Complementary feeding. In: Kleinman RE, editor. *Pediatric nutrition handbook*. Elk Grove (IL): American Academy of Pediatrics; 2004.

- [155] Gartner LM, Morton J, Lawrence RA, et al. Breastfeeding and the use of human milk. *Pediatrics* 2005;115:496–506.
- [156] Parents' survival guide to transitional feeding. The Institute of Pediatric Nutrition; 2003. Available at: http://abbottnutrition.com/home/breastfeeding/index.aspx?p=0.5.2_ForHealthCareProfessionals.
- [157] US Department of Health and Human Services, National Center for Health Statistics. Third National Health and Nutrition Survey, 1988–1994, NHANES III. Examination data file: CD-ROM series 11 No 1A and No 2A. In: Hyattsville (MD): Public use data file documentation number 76200; July 1997 and April 1998, respectively.
- [158] Kim J, Peterson KE. Association of infant child care with infant feeding practices and weight gain among US infants. *Arch Pediatr Adolesc Med* 2008;162:627–33.
- [159] Shelov SP, Hannemann RE, editors. *Caring for your baby and young child: birth to age 5*. 4th edition. New York: Bantam Books; 2004.
- [160] Macknin ML, Medendorp SV, Maier MC. Infant sleep and bedtime cereal. *Am J Dis Child* 1989;143:1066–8.
- [161] Shukla A, Forsyth HA, Anderson CM, et al. Infantile overnutrition in the first year of life: a field study in Dudley, Worcestershire. *Br Med J* 1972;4:507–15.
- [162] Ferris AG, Laus MJ, Hosmer DW, et al. The effect of diet on weight gain in infancy. *Am J Clin Nutr* 1980;33:2635–42.
- [163] Kramer MS, Barr RG, Leduc DG, et al. Determinants of weight and adiposity in the first year of life. *J Pediatr* 1985;106:10–4.
- [164] Ong KK, Emmett PM, Noble S, et al. Dietary energy intake at the age of 4 months predicts postnatal weight gain and childhood body mass index. *Pediatrics* 2006;117:e503–8.
- [165] Baker JL, Michaelsen KF, Rasmussen KM, et al. Maternal prepregnant body mass index, duration of breastfeeding, and timing of complementary food introduction are associated with infant weight gain. *Am J Clin Nutr* 2004;80:1579–88.
- [166] Birch LL. Development of food preferences. *Annu Rev Nutr* 1999;19:41–62.
- [167] Sullivan SA, Birch LL. Infant dietary experience and acceptance of solid foods. *Pediatrics* 1994;93:271–7.
- [168] Birch LL, Gunder L, Grimm-Thomas K, et al. Infants' consumption of a new food enhances acceptance of similar foods. *Appetite* 1998;30:283–95.
- [169] Birch LL, Marlin DW. I don't like it; I never tried it: effects of exposure on two-year-old children's food preferences. *Appetite* 1982;3:353–60.
- [170] Birch LL, McPhee L, Shoba BC, et al. What kind of exposure reduces children's food neophobia? Looking vs. tasting. *Appetite* 1987;9:171–8.
- [171] Birch LL. Development of food acceptance patterns in the first years of life. *Proc Nutr Soc* 1998;57:617–24.
- [172] Dietz WH, Franks AL, Marks JS. The obesity problem. *N Engl J Med* 1998;338:1157 [author reply 8].
- [173] Farrow C, Blissett J. Does maternal control during feeding moderate early infant weight gain? *Pediatrics* 2006;118:e293–8.
- [174] Farrow CV, Blissett J. Controlling feeding practices: cause or consequence of early child weight? *Pediatrics* 2008;121:e164–9.
- [175] Birch LL, McPhee L, Shoba BC, et al. "Clean up your plate". Effects of child feeding practices on the conditioning of meal size. *Learn Motiv* 1987;18:301–17.
- [176] Galloway AT, Fiorito L, Lee Y, et al. Parental pressure, dietary patterns, and weight status among girls who are "picky eaters". *J Am Diet Assoc* 2005;105:541–8.
- [177] Galloway AT, Fiorito LM, Francis LA, et al. 'Finish your soup': counterproductive effects of pressuring children to eat on intake and affect. *Appetite* 2006;46:318–23.
- [178] Ventura AK, Birch LL. Does parenting affect children's eating and weight status? *Int J Behav Nutr Phys Act* 2008;5:15.
- [179] Gidding SS, Dennison BA, Birch LL, et al. Dietary recommendations for children and adolescents: a guide for practitioners. *Pediatrics* 2006;117:544–59.

- [180] Green M, editor. Bright futures: guidelines for health supervision of infants, children, and adolescents. Arlington (VA): National Center for Education in Maternal and Child Health; 1994.
- [181] Dietz WH, Stern L, editors. American Academy of Pediatrics: guide to your child's nutrition. New York: Villard; 1999.
- [182] Kleinman RE, editor. Pediatric nutrition handbook. 5th edition. Elk Grove (IL): American Academy of Pediatrics; 2004.
- [183] American Academy of Pediatrics. Committee on Nutrition. Cholesterol in childhood. Pediatrics 1998;101:141-7.
- [184] Birch LL. Children's preferences for high-fat foods. Nutr Rev 1992;50:249-55.
- [185] Fox MK, Reidy K, Novak T, et al. Sources of energy and nutrients in the diets of infants and toddlers. J Am Diet Assoc 2006;106:S28-42.
- [186] Dennison BA, Erb TA, Jenkins PL. Predictors of dietary milk fat intake by preschool children. Prev Med 2001;33:536-42.
- [187] Kranz S, Lin PJ, Wagstaff DA. Children's dairy intake in the United States: too little, too fat? J Pediatr 2007;151:642-6, 6 e1-e2.
- [188] Basch CE, Shea S, Zybert P. Food sources, dietary behavior, and the saturated fat intake of Latino children. Am J Public Health 1992;82:810-5.
- [189] Sigman-Grant M, Zimmerman S, Kris-Etherton PM. Dietary approaches for reducing fat intake of preschool-age children. Pediatrics 1993;91:955-60.
- [190] Dixon LB, McKenzie J, Shannon BM, et al. The effect of changes in dietary fat on the food group and nutrient intake of 4- to 10-year-old children. Pediatrics 1997;100:863-72.
- [191] Friedman G, Goldberg SJ. An evaluation of the safety of a low-saturated-fat, low-cholesterol diet beginning in infancy. Pediatrics 1976;58:655-7.
- [192] Niinikoski H, Viikari J, Ronnema T, et al. Regulation of growth of 7- to 36-month-old children by energy and fat intake in the prospective, randomized STRIP baby trial. Pediatrics 1997;100:810-6.
- [193] Lagstrom H, Seppanen R, Jokinen E, et al. Influence of dietary fat on the nutrient intake and growth of children from 1 to 5 y of age: the Special Turku Coronary Risk Factor Intervention Project. Am J Clin Nutr 1999;69:516-23.
- [194] Karaolis-Danckert N, Gunther AL, Kroke A, et al. How early dietary factors modify the effect of rapid weight gain in infancy on subsequent body-composition development in term children whose birth weight was appropriate for gestational age. Am J Clin Nutr 2007;86:1700-8.
- [195] Fox MK, Pac S, Devaney B, et al. Feeding Infants and Toddlers Study: what foods are infants and toddlers eating? J Am Diet Assoc 2004;104:s22-30.
- [196] Daniels SR, Greer FR. Lipid screening and cardiovascular health in childhood. Pediatrics 2008;122:198-208.
- [197] Berkey CS, Rockett HR, Willett WC, et al. Milk, dairy fat, dietary calcium, and weight gain: a longitudinal study of adolescents. Arch Pediatr Adolesc Med 2005;159:543-50.
- [198] USDA National Nutrient Database for Standard Reference. Available at: http://www.nal.usda.gov/fnic/foodcomp/cgi-bin/nut_search_new.pl. Accessed February 7, 2008.
- [199] Nielsen SJ, Popkin BM. Changes in beverage intake between 1977 and 2001. Am J Prev Med 2004;27:205-10.
- [200] O'Connor TM, Yang SJ, Nicklas TA. Beverage intake among preschool children and its effect on weight status. Pediatrics 2006;118:e1010-8.
- [201] American Dietetic Association. Childhood overweight evidence analysis project: factors associated with overweight. Available at: <http://www.adaevidencelibrary.com/topic.cfm?cat=2792>. Accessed February 15, 2007.
- [202] Skinner JD, Ziegler P, Ponza M. Transitions in infants' and toddlers' beverage patterns. J Am Diet Assoc 2004;104:s45-50.
- [203] American Academy of Pediatrics. The use and misuse of fruit juice in pediatrics. Pediatrics 2001;107:1210-3.

- [204] Welsh JA, Cogswell ME, Rogers S, et al. Overweight among low-income preschool children associated with the consumption of sweet drinks: Missouri, 1999–2002. *Pediatrics* 2005;115:e223–9.
- [205] Warner ML, Harley K, Bradman A, et al. Soda consumption and overweight status of 2-year-old Mexican-American children in California. *Obesity (Silver Spring)* 2006;14:1966–74.
- [206] Murray R, Frankowski B, Taras H. Are soft drinks a scapegoat for childhood obesity? *J Pediatr* 2005;146:586–90.
- [207] James J, Kerr D. Prevention of childhood obesity by reducing soft drinks. *Int J Obes (Lond)* 2005;29(Suppl 2):S54–7.
- [208] Devaney B, Ziegler P, Pac S, et al. Nutrient intakes of infants and toddlers. *J Am Diet Assoc* 2004;104:s14–21.
- [209] Picciano MF, Smiciklas-Wright H, Birch LL, et al. Nutritional guidance is needed during dietary transition in early childhood. *Pediatrics* 2000;106:109–14.
- [210] Butte N, Cobb K, Dwyer J, et al. The Start healthy feeding guidelines for infants and toddlers. *J Am Diet Assoc* 2004;104:442–54.
- [211] Ryan C, Dwyer J. The toddler's smorgasbord. *Nutr Today* 2003;38:164–9.
- [212] McConahy KL, Picciano MF. How to grow a healthy toddler—12 to 24 months. *Nutr Today* 2003;38:156–63.
- [213] National Association for Sport and Physical Education. Active start: a statement of physical activity guidelines for children birth to five years. New York: AAHPERD Publications; 2002.
- [214] Corbin C, Pangrazi R, Beighle A, et al. Guidelines for appropriate physical activity for elementary school children 2003 update. A position statement. Council for Physical Education for Children (COPEC) of the National Association for Sport and Physical Education, an Association of the American Alliance for Health Physical Education and Recreation. New York: AAHPERD Publications; 2003.
- [215] Gunner KB, Atkinson PM, Nichols J, et al. Health promotion strategies to encourage physical activity in infants, toddlers, and preschoolers. *J Pediatr Health Care* 2005;19:253–8.
- [216] Dennison BA, Boyer PS. Risk evaluation in pediatric practice aids in prevention of childhood overweight. *Pediatr Ann* 2004;33:25–30.
- [217] Neumark-Sztainer D. Childhood and adolescent obesity: an ecological perspective. *Basic Pediatrics* 2003;101:12–20.
- [218] Dennison BA, Erb TA, Jenkins PL. Television viewing and television in bedroom associated with overweight risk among low-income preschool children. *Pediatrics* 2002;109:1028–35.
- [219] Campbell K, Hesketh K, Crawford D, et al. The Infant Feeding Activity and Nutrition Trial (INFANT) an early intervention to prevent childhood obesity: cluster-randomised controlled trial. *BMC Public Health* 2008;8:103.
- [220] Wolman J, Skelly E, Kolotourou M, et al. Tackling toddler obesity through a pilot community-based family intervention. *Community Pract* 2008;81:28–31.
- [221] American Academy of Pediatrics. Children, adolescents, and television. *Pediatrics* 2001;107:423–6.