

A sub-kpc-scale binary active galactic nucleus with double narrow-line regions

Jong-Hak Woo,^{1,2,3★†} Hojin Cho,¹ Bernd Husemann,⁴ S. Komossa,^{5,3} Daeseong Park¹ and Vardha N. Bennert⁶

¹*Astronomy Program, Department of Physics and Astronomy, Seoul National University, Seoul 151-742, Republic of Korea*

²*The Observatories of the Carnegie Institution for Science, 813 Santa Barbara Street, Pasadena, CA 91101, USA*

³*Kavli Institute for Theoretical Physics, Santa Barbara, CA 93106, USA*

⁴*Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, D-14482 Potsdam, Germany*

⁵*Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany*

⁶*Physics Department, California Polytechnic State University, San Luis Obispo, CA 93407, USA*

Accepted 2013 September 25. Received 2013 September 24; in original form 2013 August 27

ABSTRACT

We present the kinematic properties of a type-2 Quasi stellar object (QSO), SDSS J132323.33–015941.9 at $z \sim 0.35$, based on the analysis of Very Large Telescope integral field spectroscopy and *Hubble Space Telescope* (*HST*) imaging, which suggest that the target is a binary active galactic nucleus (AGN) with double narrow-line regions. The QSO features double-peaked emission lines ([O III] and H β) which can be decomposed into two kinematic components. The flux-weighted centroids of the blue and red components are separated by ~ 0.2 arcsec (0.8 kpc in projection) and coincide with the location of the two stellar cores detected in the *HST* broad-band images, implying that both stellar cores host an active black hole. The line-of-sight velocity of the blue component is comparable to the luminosity-weighted velocity of stars in the host galaxy, while the red component is redshifted by ~ 240 km s⁻¹, consistent with typical velocity offsets of two cores in a late stage of a galaxy merger. If confirmed, the target is one of the rare cases of sub-kpc-scale binary AGNs, providing a test-bed for understanding the binary AGN population.

Key words: galaxies: active – galaxies: interactions – galaxies: nuclei – quasars: supermassive black holes.

1 INTRODUCTION

Galaxy mergers represent an important stage in the evolution of galaxies. In the course of gas-rich mergers, accretion activity is triggered on to one or both black holes (BHs). As the merger evolves, a bound pair of supermassive black holes (SMBHs) is formed (Begelman, Blandford & Rees 1980), which will ultimately coalesce, emitting a giant burst of gravitational wave emission (see Centrella et al. 2010 for a review). Identifying pairs of accreting SMBHs in all stages of galaxy merging is therefore of great interest; hence, an active search has been ongoing. While a few *candidate* sub-parsec binary SMBHs have been reported based on semiperiodic light curves or characteristic structures in radio jets (e.g. Valtonen et al. 2011, see Komossa 2006 for a review), they lack spatially resolved observations, thus still awaiting confirmation. Spatially resolved spectroscopy is a powerful tool to reveal and confirm the wider pairs of SMBHs, and an intense search is currently ongoing. Nevertheless, only a few *spatially resolved* close

pairs [binary active galactic nuclei (AGNs)] at $\lesssim 1$ kpc separation have been confirmed so far through X-ray and radio observations (Komossa et al. 2003; Rodriguez et al. 2006; Fabbiano et al. 2011).

Recently, double-peaked narrow emission lines have been employed to search for binary AGNs, making use of large spectroscopic data bases, i.e. the Sloan Digital Sky Survey (SDSS; e.g. Zhou et al. 2004; Xu & Komossa 2009; Liu et al. 2010, 2013; Smith et al. 2010; Comerford et al. 2012; Fu et al. 2012; Ge et al. 2012; Barrows et al. 2013; Benítez et al. 2013). Apart from binary AGN, however, a number of other mechanisms can potentially produce double-peaked narrow emission lines at separations of a few hundred km s⁻¹, including jet–cloud interactions, bi-conical outflows, special narrow-line region (NLR) geometries or a single AGN illuminating the interstellar media of two galaxies (Xu & Komossa 2009, and references therein). Therefore, follow-up spatially resolved spectroscopy is essential in selecting true AGN pairs, by identifying the host galaxy (merger) itself and the two active cores. Follow-up observations of double-peaked emission line AGNs have shown that only a small fraction (2–10 per cent) of them likely harbours dual AGNs (e.g. Shen et al. 2011; Comerford et al. 2012; Fu et al. 2012). In the radio regime, for example, some sources turned

* E-mail: woo@astro.snu.ac.kr

† TJ Park Science Fellow.




Figure 1. *HST* images of the host galaxy of type-2 QSO, J1323 observed with three different filters, *F435W* (left), *F550M* (middle left) and *F775W* (middle right). The flux contours are plotted for the *F550M*-band image to show the flux contrast (right).

out to be dominated by radio jets (e.g. Rosario et al. 2010), while others are genuine dual radio-AGN (e.g. Fu et al. 2011).

Here, we present the discovery of a *sub-kpc*-scale binary AGN candidate in a type-2 Quasi stellar object (QSO), SDSS J132323.33–015941.9 (hereafter J1323), based on Very Large Telescope (VLT) spectroscopy combined with *Hubble Space Telescope* (*HST*) imaging. At the redshift of the QSO, $z = 0.35$, 1 arcsec corresponds to ~ 4.9 kpc, assuming $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.7$ and $\Omega_m = 0.3$. Very few such sub-kpc systems have been reported so far (Rodríguez et al. 2006; Fabbiano et al. 2011). Their identification is also important for studying the initial conditions which later define the rapidity of binary coalescence and the amplitude of gravitational wave recoil.

2 OBSERVATION AND REDUCTION

2.1 VLT IFU spectroscopy

Using the VLT VIMOS instrument (Le Fèvre et al. 2003), we obtained integral field unit (IFU) spectra of J1323 on 2006 February 24, with ~ 0.7 arcsec seeing according to the seeing monitor of the observatory. The total exposure time, 5445 s, was split into three exposures of 1815 s each. We chose the high-resolution (HR) grism of VIMOS with a spectral resolution of $R \sim 2650$, covering a spectral range of 5250–7400 Å, which includes the rest-frame [O III] $\lambda\lambda 4959, 5007$ and H β lines of the target. The VIMOS field of view was set to 13 arcsec \times 13 arcsec, providing a spatial resolution of 0.33 arcsec spaxel $^{-1}$.

The data were reduced by a customized pipeline based on the data reduction software developed for the Calar Alto legacy integral field area (CALIFA) survey (Sánchez et al. 2012; Husemann et al. 2013). Principal reduction steps include bias subtraction, cosmic ray detection with PYCOSMIC (Husemann et al. 2012), fibre tracing, optimal extraction of spectra, wavelength calibration using arc lamp exposures, fibre transmission correction, flux calibration using standard star exposures and correction for differential atmospheric refraction. VIMOS suffers from instrumental flexures that need to be properly taken into account during the calibration of each exposure. The flexure effect was characterized in each science exposure by measuring the offset of the fibre traces between the science and continuum exposures as a function of wavelength as well as measuring offsets from the expected centroid wavelength of prominent sky lines.

The slightly varying spectral resolutions of the four individual VIMOS spectrographs were characterized across each detector from the width of emission lines in the arc lamp exposures. With an adap-

tive Gaussian smoothing of the individual spectra, we homogenized the spectral resolution to 2 Å (full width at half-maximum) for each spectrum along the entire wavelength range.

2.2 HST images

J1323 was previously observed with the Wide-Field Channel of the Advanced Camera for Surveys on board *HST* (Zakamsk et al. 2006), selected as one of the high [O III] luminosity type-2 AGNs. Three filters (*F435W*, *F550M*, *F775W*) were chosen to cover the rest-frame UV, blue and yellow continuum of the host galaxy, avoiding strong AGN emission lines, i.e. [O III], H α and H β , of the QSO (see fig. 1 in Zakamska et al. 2006). In these images, we detect a merging pair with two distinct cores, which are separated by $0.20 \text{ arcsec} \pm 0.01$ with a north-to-east position angle $\text{PA} = 12:9 \pm 4.0$, based on the flux centroid calculations. As shown in Fig. 1, the flux ratio of the two cores varies depending on the adopted filters; however, the position of the centre of each core and the PA is unchanged. While the two cores have similar flux in the rest-frame blue-band image, the south core is more prominent in the rest-frame UV image. In contrast, the north core is brighter than the south core in the rest-frame yellow-band image. The host galaxy seems to be undergoing a merging process with the two cores separated by a sub-kpc-scale in a common envelope. As a consistency check for the separation and the PA of the two cores, we perform the decomposition analysis with three Sérsic components to fit the two cores and the envelope in the *F550M* image, using GALFIT (Peng et al. 2002). Although the image is relatively shallow to accurately determine the luminosity of each component, we confirm that the separation ($0.22 \text{ arcsec} \pm 0.02$) and the PA ($9:2 \pm 5.4$) based on the GALFIT analysis is consistent with those based on the flux centroid calculation within the uncertainties.

Since the weaker emission lines of the QSO were included in the broad-band photometry, we quantify the contribution of the narrow emission lines to the broad-band flux, by convolving the SDSS spectrum of the QSO with the bandpasses of the aforementioned three filters. We find that the AGN emission-line contribution to the broad-band flux is a few per cent, indicating that the *HST* images represent the stellar component.

3 ANALYSES AND RESULTS

3.1 SDSS spectroscopy

To determine the systemic velocity of the host galaxy, we use the SDSS spectrum of J1323 since the VIMOS IFU spectra have




Figure 2. Examples of the IFU spectra of [O III] (left) and H β (right), and their decomposition at the central 5×5 spaxels. The black solid lines represent the continuum-subtracted spectra, while green lines represent the best-fitting model, which is composed of two kinematic components (blue and red lines, respectively). The vertical dashed lines indicate the centres of [O III] and H β lines expected from the systemic velocity based on stellar absorption lines. Each panel represents a spaxel with a $0.33 \text{ arcsec} \times 0.33 \text{ arcsec}$ scale, centred on the spaxel where the [O III] line flux peaks.

low signal-to-noise ratio (S/N) on the continuum with no clear signs of stellar absorption lines. In contrast, the SDSS spectrum presents various stellar lines over the observed spectral range although the level of noise is relatively high. We fit the stellar lines at the rest-frame range, 5100–5350 Å, including the Mgb triplet (5172 Å) and Fe (5300 Å) lines, using the penalized pixel-fitting software (Cappellari & Emsellem 2004) and the synthesized stellar templates based on 180 stellar spectra from the MILES library (Falcón-Barroso et al. 2011). Based on the best-fitting template, we derive the line-of-sight (LOS) velocity. To account for the uncertainty of the LOS velocity, we perform Monte Carlo simulations using 200 mock spectra generated by adding random noise to the SDSS spectrum. Then, we fit the stellar lines and determine the LOS velocity for each spectrum and adopt the 1σ dispersion of the measurements, 24.9 km s^{-1} , as the uncertainty of the LOS velocity. Due to the low S/N on the stellar features, the LOS velocity is somewhat uncertain. Thus, we use the determined systemic velocity as a reference.

Since we used the 3 arcsec fibre SDSS spectrum, the derived LOS velocity represents the systemic velocity of the luminosity-weighted stellar component in the host galaxy. To quantify the luminosity contribution of the north core to the SDSS spectrum, we integrate the flux of the north core as well as the flux from the envelope within a 3 arcsec aperture, using the *HST* $F775W$ -band image. The contribution of the north core is only 10–20 per cent, indicating that the north core is not dominant in the luminosity weight and that the measured velocity represents the systemic velocity of the host galaxy.

3.2 Narrow-line regions

In Fig. 2, we present the observed spectra overplotted with the best-fitting model for the central 5×5 spaxels, covering a $1.65 \text{ arcsec} \times 1.65 \text{ arcsec}$ region, since emission lines are much weaker at outer spaxels. Here, the centre (0, 0) is defined as the location of the peak [O III] flux, corresponding to the location between the north and south cores in the *HST* images.

As shown in Fig. 2, the VLT IFU spectra provide spatial constraints of flux and profile changes of the emission lines, [O III] and H β . The [O III] and H β lines show a double-peaked profile, particularly at the central spaxels, and the flux ratio between the two peaks dramatically changes spatially. Note that the double-peaked profile was not detected in the SDSS spectrum due to its lower spectral resolution. As a consistency check, we combine all spectra from our IFU spaxels, then degrade the spectral resolution to match that of the SDSS spectrum. Indeed, the combined spectrum shows the same line profile as in the SDSS spectrum.

We kinematically decompose the double-peaked [O III] $\lambda 5007$ line by fitting the line profile with various models, including single, double and triple Gaussian, and double Gauss–Hermitian models, using the MPFIT (Markwardt 2009) Levenberg–Marquardt least-squares minimization routine. We find that these Gaussian models do not generate consistent fluxes between the nearest spaxels, making these models unfavourable since we expect similar fluxes in neighbouring spaxels due to the seeing. In contrast, when a pair of Lorentzian profiles was used to fit each emission line, we obtain the best results with spatially consistent fluxes. In the fitting process, we assume that [O III] $\lambda 5007$ and [O III] $\lambda 4959$ lines have the same




Figure 3. [O III] flux map (top), LOS velocity distribution (middle) and [O III]/H β ratio distribution of the blue (left-hand panels) and red components (right-hand panels). Each pixel represents a spaxel of $0.33 \text{ arcsec} \times 0.33 \text{ arcsec}$. Stellar contours measured from the *HST* *F550M*-band image are overlapped. The flux-weighted centroids of the blue and red components of [O III] are denoted with blue and red circles, respectively, with a 1σ error bar.

redshift and width and that the flux ratio between them is 3:1. The H β line was also fitted under the assumption that its LOS velocity coincides with that of [O III]. Although H β is much weaker than [O III] and kinematic decomposition can be done only for central several spaxels, the H β line seems to show a similar spatial flux change between blue and red components.

In Fig. 3, we present the kinematic analysis, showing the spatial distributions of the fluxes, LOS velocities and the [O III]/H β flux ratios for both kinematic components of [O III]. The [O III] flux maps may suggest that both components are extended over kpc scales. However, we were not able to confirm whether the [O III] line gas is extended over kpc-scale based on our point spread function (PSF) convolution test since the exact PSF at the time of observation is not known.

Although these two components cannot be spatially separated due to the seeing limit ($\sim 0.7 \text{ arcsec}$), the IFU data enable us to determine the luminosity-weighted centroid of each component as denoted with blue and red points, respectively, in Fig. 3. The two centroids are separated by $0.17 \text{ arcsec} \pm 0.01$, which is $\sim 0.8 \text{ kpc}$ in projection at the distance of the target, with a $\text{PA} = 13^\circ 9 \pm 0.5$. Note that the S/N of [O III] in each spaxel is very high, leading to relatively small measurement uncertainties of the flux centroids. To better quantify the uncertainty of the separation and the PA, we repeatedly measured the flux centroids using various sets of spaxels selected from the [O III] map and flux weighting schemes. We obtained an average separation 0.17 arcsec with a 0.01 arcsec standard deviation and a mean $\text{PA} = 13^\circ 9$ with a $1^\circ 8$ standard deviation. We also computed the separation and the PA using the triple Gaussian

models in fitting the [O III] line, in order to test the dependence of the line profile modelling on the separation measurement. While the PA remains consistent with that based on the double Lorentzian models, the separation becomes slightly larger, presumably due to the uncertainties in modelling the blended wings of [O III]. Assuming this discrepancy as the systematic uncertainty, we conservatively add ~ 0.06 arcsec uncertainty to the measurement error. By combining the measurement error (0.01 arcsec), the standard deviation due to the spaxel sampling and weighting schemes (0.01 arcsec) and the systematic uncertainty in the line profile modelling (~ 0.06 arcsec) in quadrature, we determine the uncertainty of the separation as 0.06 arcsec. Thus, we finalize the measurements as separation = 0.17 arcsec ± 0.06 and PA = $13^{\circ}.9 \pm 1.8$ between the two flux centroids.

Interestingly, the separation and the PA of the two centroids are consistent with those of the two stellar cores measured from the *HST* images. In Fig. 3, we overplot the contours of the host galaxy from the *F550M*-band image, after shifting the *HST* image to match the location of the [O III] flux centroids. Note that since the S/N of the continuum in the IFU spectra is too low to independently determine the exact location of the stellar cores, we shift the *HST* image in order to match the location of the stellar cores to that of the [O III] flux centroids. As a consistency check, we compare the flux map of the weak continuum detected in the IFU spectra with that of the *HST* image, after convolving the *HST* image with a typical PSF of the IFU. The two flux maps are overlapping each other, confirming that the two stellar cores are close to the centre of the [O III] flux distribution. The absolute spatial registration between VLT IFU image and *HST* image is uncertain given the current data; hence, future confirmation is required. However, since the separation and PA of the two components coincide between IFU and *HST* images, it is likely that the flux centroids of the blue and red components of [O III] indeed coincide with the locations of the north and south stellar cores. Thus, we interpret these results as indication of both stellar cores hosting an active BH with an associated NLR.

By measuring the flux centroid of the line profile of each [O III] component, we obtained the velocity offset ~ 211 km s $^{-1}$ between blue and red [O III] components. We also measured the LOS velocity of each [O III] component with respect to the systemic velocity measured from stellar absorption lines. The velocity of the blue component shows a small redshift of 30 ± 5 km s $^{-1}$ with respect to the stellar component, while that of the red component is redshifted by 241 ± 5 km s $^{-1}$ on average with a velocity gradient of ~ 8 km s $^{-1}$ kpc $^{-1}$ towards south-west. The velocity of the blue component is consistent with the system velocity of the luminosity-weighted stellar component within the measurement errors (~ 25 km s $^{-1}$), implying that the active BH in the south core and the accompanied NLR is close to the dynamical centre of the host galaxy, while the NLR of the active BH in the north core is dynamically offset (see Fig. 3). Assuming two BHs are hosting NLRs, we estimate the mass of each BH based on the M_{BH} -stellar velocity dispersions relationship adopted from McConnell & Ma (2013; see also Woo et al. 2013), using the width of the [O III] line as a proxy for the stellar velocity dispersion. Although these estimates are highly uncertain, we find that the BH mass ($\log M_{\text{BH}}/M_{\odot}$) of the blue and red components are 7.6 and 6.7, respectively.

The [O III]/H β flux ratios of both components are much larger than 3, indicating that the ionization source of each component is an AGN. The [O III]/H β flux ratio of each component does not vary significantly over ± 1 arcsec scale, within which we are able to measure the flux of each H β components (see Fig. 2). In contrast,

the [O III]/H β flux ratio between the two components is different by 0.2 dex (58 per cent).

4 DISCUSSIONS AND SUMMARY

The nature of the double-peaked [O III] lines detected with spatially integrated spectra, i.e. SDSS spectra, is generally ambiguous since it can be explained by various mechanisms: the projection of two unrelated AGNs, bi-conical outflows, the interaction of a two-sided jet with NLR clouds, special NLR geometries, a single AGN illuminating the interstellar media of two galaxies or a true binary AGN (e.g. Axon et al. 1998; Xu & Komossa 2009; Crenshaw et al. 2010; Shen et al. 2011). The uncertainty in interpreting the data arises mostly from the fact that the (SDSS) spectroscopic data do not come with spatially resolved kinematic information or well-resolved images of host galaxies, as demonstrated by follow-up studies to overcome the lack of spatial information (e.g. Shen et al. 2011; Fu et al. 2012; Liu et al. 2013).

In this paper, we presented a binary AGN candidate based on the spatial and kinematic decomposition of the double-peaked [O III] line profile using VLT IFU spectroscopy and *HST* imaging. Although the spatial resolution of the IFU image is limited to the seeing (~ 0.7 arcsec), the coincidence of the separation and the PA between the two stellar cores, with those between the two flux centroids of the [O III] kinematic components, suggests that a binary AGN harbours double NLRs. As the velocity separation of ~ 200 km s $^{-1}$ between blue and red components is typical of two cores in a galaxy merger, the sub-kpc-scale spatial separation may indicate a binary AGN in a late stage of a galaxy merger.

It is also possible that the double-peaked [O III] is due to jet-cloud interaction in a single active BH system. In this case, however, the jet orientation should coincide with the PA of the two stellar cores, instead of oriented randomly. There is only a small chance that the projected jet direction is close to the PA of the two stellar cores. Thus, localized jet-cloud interactions or outflows and special NLR geometries are less viable scenarios for J1323.

Another plausible scenario is that a single AGN photoionizes interstellar media centred on two stellar cores. In this scenario, we expect that the fluxes of [O III] in two NLRs are different due to the distance from an ionizing source, covering factor and gas properties. Given the similar strengths of blue and red peaks in the [O III] line, it is less likely that a *single* AGN ionizes *two* NLRs, which produces a stronger difference in line luminosity under most NLR geometries. High-resolution radio, optical (e.g. *HST*) and X-ray observations are essential to confirm that two NLRs are associated with two active nuclei. In the X-ray, despite recent improvements in sub-pixel imaging techniques with *Chandra* (e.g. Liu et al. 2013), even marginally resolving two faint cores with a ~ 0.2 arcsec separation will be extremely challenging. The two cores could be easily resolved in the radio, but only if both cores are bright radio emitters. In contrast, spatially resolved spectroscopy with *HST* would allow us to measure the size and kinematics structure of the two NLRs, confirming the binary nature of J1329.

If confirmed, J1323 is one of the rare cases of sub-kpc-scale binary AGNs. While a handful of binary AGNs have been confirmed with a separation of several kiloparsecs between two active BHs, only two binary cases show a sub-kpc-scale separation, i.e. ~ 10 and ~ 100 pc, respectively (Rodríguez et al. 2006; Fabbiano et al. 2011; see also Komossa et al. 2003 for NGC 6240), indicating that the discovery of close pairs of BHs in the late stage of galaxy merging is observationally difficult, partly due to the challenge of spatially resolving two active cores. Thus, confirming the binary nature of

J1323 will provide another test-bed for understanding the binary AGN population.

ACKNOWLEDGEMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) (No. 2012-006087 and 2010-0021558). JHW and SK acknowledge the support and hospitality by the Kavli Institute for Theoretical Physics during the programme ‘A Universe of Black Holes’. This research was supported in part by the National Science Foundation under grant no. NSF PHY11-25915.

REFERENCES

- Axon D. J., Marconi A., Capetti A., Maccetto F. D., Schreier E., Robinson A., 1998, *ApJ*, 496, L75
- Barrows R. S. et al., 2013, *ApJ*, 769, 95
- Begelman M. C., Blandford R. D., Rees M. J., 1980, *Nature*, 287, 307
- Benítez E. et al., 2013, *ApJ*, 763, 36
- Cappellari M., Emsellem E., 2004, *PASP*, 116, 138
- Centrella J., Baker J. G., Kelly B. J., van Meter J. R., 2010, *Rev. Mod. Phys.*, 82, 3069
- Comerford J. M., Gerke B. F., Stern D., Cooper M. C., Weiner B. J., Newman J. A., Madsen K., Barrows R. S., 2012, *ApJ*, 753, 42
- Crenshaw D. M., Schmitt H. R., Kraemer S. B., Mushotzky R. F., Dunn J. P., 2010, *ApJ*, 708, 419
- Fabbiano G., Wang J., Elvis M., Risaliti G., 2011, *Nature*, 477, 431
- Falcón-Barroso J., Sánchez-Blázquez P., Vazdekis A., Ricciardelli E., Cardiel N., Cenarro A. J., Gorgas J., Peletier R. F., 2011, *A&A*, 532, A95
- Fu H. et al., 2011, *ApJ*, 740, L44
- Fu H., Yan L., Myers A. D., Stockton A., Djorgovski S. G., Aldering G., Rich J. A., 2012, *ApJ*, 745, 67
- Ge J.-Q., Hu C., Wang J.-M., Bai J.-M., Zhang S., 2012, *ApJS*, 201, 31
- Husemann B., Kamann S., Sandin C., Sánchez S. F., García-Benito R., Mast D., 2012, *A&A*, 545, A137
- Husemann B. et al., 2013, *A&A*, 549, A87
- Komossa S., 2006, *Mem. Soc. Astron. Ital.*, 77, 733
- Komossa S., Burwitz V., Hasinger G., Predehl P., Kaastra J. S., Ikebe Y., 2003, *ApJ*, 582, L15
- Le Fèvre O. et al., 2003, in Iye M., Moorwood A. F. M., eds, *Proc. SPIE Conf. Ser. Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes*. SPIE, Bellingham, p. 1670
- Liu X., Greene J. E., Shen Y., Strauss M. A., 2010, *ApJ*, 715, L30
- Liu X., Civano F., Shen Y., Green P., Greene J. E., Strauss M. A., 2013, *ApJ*, 762, 110
- McConnell N. J., Ma C.-P., 2013, *ApJ*, 764, 184
- Markwardt C. B., 2009, in Bohlender D. A., Durand D., Dowler P., eds, *ASP Conf. Ser. Vol. 411, Astronomical Data Analysis Software and Systems XVIII*. Astron. Soc. Pac., San Francisco, p. 251
- Peng C. Y., Ho L. C., Impey C. D., Rix H.-W., 2002, *AJ*, 124, 266
- Rodríguez C., Taylor G. B., Zavala R. T., Peck A. B., Pollack L. K., Romani R. W., 2006, *ApJ*, 646, 49
- Rosario D. J., Shields G. A., Taylor G. B., Salviander S., Smith K. L., 2010, *ApJ*, 716, 131
- Sánchez S. F. et al., 2012, *A&A*, 538, A8
- Shen Y., Liu X., Greene J. E., Strauss M. A., 2011, *ApJ*, 735, 48
- Smith K. L., Shields G. A., Bonning E. W., McMullen C. C., Rosario D. J., Salviander S., 2010, *ApJ*, 716, 866
- Valtonen M. J., Lehto H. J., Takalo L. O., Sillanpää A., 2011, *ApJ*, 729, 33
- Woo J.-H., Schulze A., Park D., Kang W.-R., Kim S. C., Riechers D. A., 2013, *ApJ*, 772, 49
- Xu D., Komossa S., 2009, *ApJ*, 705, L20
- Zakamska N. L. et al., 2006, *ApJ*, 132, 1496
- Zhou H., Wang T., Zhang X., Dong X., Li C., 2004, *ApJ*, 604, L33

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.