

Dose-Response: An International Journal

Volume 12 | Issue 2

Article 10

6-30-2014

COMMENTARY: ETHICAL ISSUES OF CURRENT HEALTH-PROTECTION POLICIES ON LOW-DOSE IONIZING RADIATION

Yehoshua Socol

Academic Forum for Nuclear Awareness, Israel

Ludwik Dobrzynski

National Center for Nuclear Research, Poland

Mohan Doss

Fox Chase Cancer Center, USA

Ludwig E. Feinendegen

Heinrich-Heine University, Germany

Marek K. Janiak

Military Institute of Hygiene and Epidemiology, Poland

See next page for additional authors

Follow this and additional works at: https://scholarworks.umass.edu/dose_response

Recommended Citation

Socol, Yehoshua; Dobrzynski, Ludwik; Doss, Mohan; Feinendegen, Ludwig E.; Janiak, Marek K.; Miller, Mark L.; Sanders, Charles L.; Scott, Bobby R.; Ulsh, Brant; and Vaiserman, Alexander (2014) "COMMENTARY: ETHICAL ISSUES OF CURRENT HEALTH-PROTECTION POLICIES ON LOW-DOSE IONIZING RADIATION," *Dose-Response: An International Journal*: Vol. 12 : Iss. 2 , Article 10.

Available at: https://scholarworks.umass.edu/dose_response/vol12/iss2/10

This Article is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Dose-Response: An International Journal by an authorized editor of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

COMMENTARY: ETHICAL ISSUES OF CURRENT HEALTH- PROTECTION POLICIES ON LOW-DOSE IONIZING RADIATION

Authors

Yehoshua Socol, Ludwik Dobrzynski, Mohan Doss, Ludwig E. Feinendegen, Marek K. Janiak, Mark L. Miller, Charles L. Sanders, Bobby R. Scott, Brant Ulsh, and Alexander Vaiserman

COMMENTARY: ETHICAL ISSUES OF CURRENT HEALTH-PROTECTION POLICIES ON LOW-DOSE IONIZING RADIATION

Yehoshua Socol □ Academic Forum for Nuclear Awareness, Israel

Ludwik Dobrzyński □ National Center for Nuclear Research, Poland

Mohan Doss □ Fox Chase Cancer Center, USA

Ludwig E. Feinendegen □ Heinrich-Heine University, Germany

Marek K. Janiak □ Military Institute of Hygiene and Epidemiology, Poland

Mark L. Miller □ Sandia National Laboratories, USA

Charles L. Sanders □ USA

Bobby R. Scott □ Lovelace Respiratory Research Institute, USA

Brant Ulsh □ M.H. Chew & Associates, USA

Alexander Vaiserman □ Institute of Gerontology, Ukraine

□ The linear no-threshold (LNT) model of ionizing-radiation-induced cancer is based on the assumption that every radiation dose increment constitutes increased cancer risk for humans. The risk is hypothesized to increase linearly as the total dose increases. While this model is the basis for radiation safety regulations, its scientific validity has been questioned and debated for many decades. The recent memorandum of the International Commission on Radiological Protection admits that the LNT-model predictions at low doses are “speculative, unproven, undetectable and ‘phantom’.” Moreover, numerous experimental, ecological, and epidemiological studies show that low doses of sparsely-ionizing or sparsely-ionizing plus highly-ionizing radiation may be beneficial to human health (hormesis/adaptive response). The present LNT-model-based regulations impose excessive costs on the society. For example, the median-cost medical program is 5000 times more cost-efficient in saving lives than controlling radiation emissions. There are also lives lost: e.g., following Fukushima accident, more than 1000 disaster-related yet non-radiogenic premature deaths were officially registered among the population evacuated due to radiation concerns. Additional negative impacts of LNT-model-inspired radiophobia include: refusal of some patients to undergo potentially life-saving medical imaging; discouragement of the study of low-dose radiation therapies; motivation for radiological terrorism and promotion of nuclear proliferation.

Key Words: low-dose radiation, risk, hormesis, adaptive response

With the linear no-threshold (LNT) model of radiation-induced cancers it is assumed that each ionizing radiation dose increment, no matter

Address correspondence to Yehoshua Socol, Academic Forum for Nuclear Awareness, POB 3067 Karney Shomron, Israel; Phone +972-9-50-661-9622, E-mail socol@FalconAnalytics.com

Ethical issues of current policies

how small, constitutes an increase in the cancer risk to humans. The risk is assumed to increase linearly as total dose increases, with an adjustment made to the slope of the dose-response curve for the reduced risk at lower dose rates. Typically, the slope is scaled down by a factor of 2 for very low dose rates (e.g. for Fukushima down-winders) in comparison to the slope for high dose rates (e.g. Hiroshima and Nagasaki).

Where mixtures of different radiations are involved (e.g., alpha, beta, and gamma), special *radiation weighting factors* (RWFs) are used to obtain a weighted dose named *equivalent dose*. RWF values are based on relative biological effectiveness (RBE) and vary from 1 (X, beta, gamma) to 20 (alpha). The RBE values come from animal and in vitro studies and vary a lot for different conditions. Where different organs are involved, *tissue weighting factors* are also used, which relate to differing tissue sensitivities; the resulting overall dose assigned to an individual applies to the whole body and is called *effective dose*. Effective dose has the following property: if e.g., only the lung is irradiated and the risk of lung cancer is 0.01, then the effective dose is the hypothetical uniform gamma-ray dose to the total body that results in the same risk (0.01) of cancer, when all cancer types are considered. The partitioning of the risk between cancer types is based on LNT and assigned uncertain tissue weighting factors.

Both equivalent dose and effective dose are expressed in units of sievert (Sv). Small effective doses on average (e.g., 0.1 mSv = 0.0001 Sv) to each member of a large population (e.g., 1 million persons downwind of Fukushima) are added to obtain a large collective dose (e.g., 0.1 millisievert \times 1 million persons = 100,000 person-millisieverts), a hypothetical value which is then multiplied by a risk coefficient to predict hypothetical cancer cases or cancer deaths for the population. It is important to recognize that the risk coefficient makes sense and both equivalent dose and effective dose are directly related to cancer risk only when dose-response relationships of interest are of the LNT type. Thus, collective dose is a LNT-hypothesis-related hypothetical value.

The LNT model in a more complex form (e.g., weighted average of absolute and relative risk forms) is presently relied on for cancer risk assessment. The LNT model is also relied on by regulatory agencies, and as such it has become the basis for radiation safety regulations. Moreover, the LNT model is widely accepted by the general public. However, the scientific validity of this model has never been proven and has been seriously questioned and debated for many decades (Taylor 1980; Feinendegen 1991; Jaworowski 1999; Tanooka 2001; Sakai *et al.* 2003; Scott 2008; Tubiana *et al.* 2009; Cuttler 2010; Fornalski and Dobrzyński 2010; Sanders 2010; Feinendegen *et al.* 2013). The absence of scientific consensus has always been officially acknowledged, including by the US Congress Office of Technology Assessment (OTA 1979). The recent

Y. Socol and others

memorandum of the ICRP (International Commission on Radiological Protection) Task Group (Gonzalez *et al.* 2013) states that:

*“While **prudent for radiological protection**, the LNT model is not universally accepted as biological truth, and its influence and inappropriate use to attribute health effects to low dose exposure situations is often ignored...*

***Speculative, unproven, undetectable and ‘phantom’ numbers** are obtained by multiplying the nominal risk coefficients by an estimate of the collective dose received by a huge number of individuals theoretically incurring very tiny doses that are hypothesized from radioactive substances released into the environment.”* (Highlights are by the authors).

Thus, the Task Group of the ICRP, one of the main bodies promoting the LNT model, admits that LNT predictions at low doses (up to 100 mSv) are *“speculative, unproven, undetectable and ‘phantom’;*” raising the reasonable question of how such a model can be *“prudent for radiological protection”* and be justifiably used in low-dose radiation risk assessment. The supporters of the LNT model claim that its use is “conservative” and should be continued until the model is proven to be untrue. They claim that in the field of safety every risk factor should be considered hazardous until proven safe, like every firearm should be considered loaded until proven unloaded. The case of radiation protection is quite different, as discussed below.

Numerous studies (experimental, epidemiological, and ecological) have shown that low doses of ionizing radiation can be beneficial to health (Feinendegen *et al.* 2004; Jaworowski 2008; Tubiana *et al.* 2009; Sanders 2010; Thompson 2011). For example, in an epidemiological study of cancer among nuclear industry workers, the rate of cancer mortality (as well as overall mortality) among the workers was substantially lower than in the reference population (Sponsler and Cameron 2005). In an epidemiological study of lung cancer association with residential radon exposure, low doses of radiation were found to prevent the occurrence of some lung cancers (Thompson 2011). Also, the healing properties of radon from spas have been utilized for centuries before people heard the word “radiation” and radon treatment is widely accepted by both the medical community and patients in Europe (Erickson 2007). Radon therapy is also popular in Japan and to a lesser extent in the United States. The lack of popularity in the United States appears to relate at least in part to the claim by the U. S. Environmental Protection Agency that residential radon causes thousands of lung cancer deaths annually among U. S. citizens.

The low-dose radiation benefits mentioned above and numerous others (Mitsunobu *et al.* 2003; Boreham *et al.* 2007; Lacoste-Colin *et al.* 2007;

Ethical issues of current policies

Liu 2007; Cohen 2008; Nakatsukasa *et al.* 2008; Scott 2008, 2011; Scott *et al.* 2008; Sanders 2010, Thompson 2011; Doss 2012; Sanders 2012; Scott and Dobrzyński 2012; Ulsh 2012; Calabrese 2013; Feinendegen *et al.* 2013; Nomura *et al.* 2013) comprise emerging scientific support for the application of radiation hormesis/adaptive response for a variety of health benefits.

The present LNT-based regulations impose excessive costs to the society, effectively leading to loss, rather than saving, of life. For example:

- According to the researchers from the Harvard School of Public Health (Graham 1995), spending \$100,000,000 per year on controlling radiation emissions might save 1 life-year per year, if the LNT model were valid, while life-saving medical program median cost is \$19,000 per life-year saved. Another study concluded that costs of radiation protection are about 5000 times higher than the cost of protection of workers from all other and much more probable events (Inhaber 2001).
- At Chernobyl and Fukushima, compulsory relocation (ordered by the authorities on the basis of ICRP recommendations which are based on the LNT model predictions) led to social destruction, which caused significant emotional/psychological problems and life-shortening. After Fukushima alone, more than 1000 non-radiogenic disaster-related premature deaths were officially registered among the evacuated population during the first year after the accident (Saji 2013). If not evacuated, these people would have received low doses of radiation that would have led, according to the LNT model, to shortening of life expectancy by less than one week (Socol *et al.* 2013) – while even this estimation is “speculative, unproven, undetectable and ‘phantom’” according to the above-mentioned ICRP Task Group memorandum.

There are additional aspects of human cost because of the LNT model and the associated radiophobia – an irrational fear of radiation hazards:

- “*Predictions of hypothetical cancer incidence and deaths ... cause some patients and parents to refuse medical imaging procedures, placing them at substantial risk by not receiving the clinical benefits of the prescribed procedures*” (AAPM 2011).
- Present policy significantly dissuades the study of low-dose radiation therapies for beneficial effects in medicine, whereas animal studies have shown potential for treatment of diseases for which presently no treatments are available, e.g., treatment of Alzheimer’s disease using low-dose radiation (Wei *et al.* 2012).
- After Chernobyl, there were more than 100,000 unnecessary abortions of pregnancies among females that received negligible radiation doses

Y. Socol and others

(or no dose at all) associated with the reactor accident (Ketchum 1987).

- Finally, unrelated to medical treatment but related to ethics, radiophobia contributes to motivating radiological terrorism and promoting nuclear proliferation (Socol *et al.* 2013).

In light of the above we suggest that the scientific community address these questions:

1. Can the LNT model, whose predictions are “speculative, unproven, undetectable and ‘phantom’”, be “prudent for radiological protection” and “accurate for low-dose-risk estimation”?
2. Doesn’t the high human cost of LNT-model-based policy necessitate serious reconsideration of this policy?
3. Should the present approval procedure for using low-dose radiation in medical research/treatment be eased in cases of cancer, autoimmune disease, diabetes, bronchial asthma, Parkinson’s, Alzheimer’s and other presently-incurable diseases associated with major suffering?
4. Should the medical community attend to debunking radiophobia by *explaining the evidence against the LNT model*?
5. Should bio-medical research of low-dose radiation be given a priority in order to resolve the existing controversy about negative/zero/positive carcinogenic effect?

Note: This paper is an adaptation of a letter recently submitted to the Israeli Bioethics Commission by some of the authors (Yehoshua Socol, Ludwik Dobrzyński, Mohan Doss, Ludwig E. Feinendegen, Marek K. Janiak, Charles L. Sanders, Brant Ulsh, Alexander Vaiserman). All authors of this paper are members of **Scientists for Accurate Radiation Information (SARI)** whose mission is *to help prevent unnecessary, radiation-phobia-related deaths, morbidity, and injuries associated with nuclear/radiological emergencies through countering phobia-promoting misinformation spread by alarmists via the news and other media including journal publications.*

DISCLAIMER: This paper represents the professional opinions of the authors, and does not necessarily represent the views of their affiliated institutions.

REFERENCES

- AAPM (American Association of Physicists in Medicine). 2011. AAPM Position Statement on Radiation Risks from Medical Imaging Procedures, PP 25–A. Available at: <http://www.aapm.org/org/policies/details.asp?id=318&type=PP>
- Boreham DR, Dolling J-A, Somers C, Quinn J, and Mitchel REJ. 2006. The adaptive response and protection against heritable mutations. *Dose-Response* 4:317–326
- Calabrese EJ. 2013. Hormetic mechanisms. *Critical Reviews in Toxicology* 43(7):580–606

Ethical issues of current policies

- Cohen BL. 2008. The linear no-threshold theory of radiation carcinogenesis should be rejected. *J Am Physicians Surg* 13(3):70–76
- Cuttler JM. 2010. Commentary on using LNT for radiation protection and risk assessment. *Dose-Response* 8(3):378–384
- Doss M. 2012. Evidence supporting radiation hormesis in atomic bomb survivor cancer mortality data. *Dose-Response* 10(4):644–663
- Erickson BE. 2007. The therapeutic use of radon: a biomedical treatment in Europe; an “alternative” remedy in the United States. *Dose-Response* 5:48–62 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2477705>
- Feinendegen LE, 1991. Radiation risk of tissue late effect, a net consequence of probabilities of various cellular responses. *Europ. J. Nucl. Med.* 18, 740-751
- Feinendegen LE, Pollycove M, and Sondhaus CA. 2004. Responses to low doses of ionizing radiation in biological systems. *Nonlinearity in Biol. Toxicol. Med.* 2: 143-171
- Feinendegen LE, Pollycove M, and Neumann RD. 2013. Hormesis by low dose radiation effects: Low-dose cancer risk modeling must recognize up-regulation of protection. In: Baum RP. (Ed.) *Therapeutic Nuclear Medicine*, Springer 2013, 500 p. ISBN 978-3-540-36718-5
- Fornalski KW and Dobrzyński L. 2010. The healthy worker effect and nuclear industry workers. *Dose-Response* 8:125–147
- Gonzalez AJ, Akashi M, Boice JD, Chino M, Homma T, Ishigure N, Kai M, Kusumi S, Lee JK, Menzel HG, Niwa O, Sakai K, Weiss W, Yamashita S, and Yonekura Y. 2013. Radiological protection issues arising during and after the Fukushima nuclear reactor accident. *J Radiol Pro* 33:497–571
- Graham JD. 1995. Comparing Opportunities To Reduce Health Risks: Toxin Control, Medicine and Injury Prevention. NCPA Policy Report No. 192, National Center for Policy Analysis, Dallas TX. Available at: <http://www.ncpa.org/pdfs/st192.pdf>
- Inhaber H. 2001. Public and occupational risks of the Nevada (USA) test site. *Environmental Management*, 28(4): 505–517
- Jaworowski Z. 1999. Radiation risk and ethics. *Physics Today* (52):9 24–29
- Jaworowski Z. 2008. The paradigm that failed. *Int J Low Radiat* 5(2):151–155
- Ketchum LE. 1987. Lessons of Chernobyl: SNM members try to decontaminate the world threatened by fallout. *Newsline* 28(6):933–941
- Lacoste-Collin L, Jozan S, Cances-Lauwers V, Pipy B, Gasset G, Caratero C, and Courtade-Saidi M. 2007. Effect of continuous irradiation with a very low dose of gamma rays on lifespan and immune system in SJL mice prone to B-cell lymphoma. *Radiat Res* 168:725–732
- Liu SZ. 2007. Cancer control related to stimulation of immunity by low-dose radiation. *Dose-Response* 5:39–47
- Mitsunobu F, Yamaoka K, Hanamoto K, Kojima S, Hosaki Y, Ashida K, Sugita K, and Tanizaki Y. 2003. Elevation of antioxidants enzymes in the clinical effects of radon and thermal therapy for bronchial asthma. *J Radiat Res* 44:95–99
- Nakatsukasa H, Tsukimoto M, Oshima Y, Tago F, Masada A, and Kojima S. 2008. Suppressing effect of low dose gamma-ray irradiation on collagen-induced arthritis. *J Radiat Res* 49:381–389
- Nomura T, Sakai K, Ogata H, and Magay J. 2013. Prolongation of lifespan in accelerated aging *klotho* mouse model by low-dose-rate continuous gamma irradiation. *Radiat Res* 179:717–724
- OTA (Office of Technology Assessment). 1979. *The Effects of Nuclear War*. U.S. Government Printing Office, Washington, DC, 20402, p. 111. Available at: <http://www.fas.org/ota/reports/7906.pdf>
- Saji G. 2013. A post-accident safety analysis report of the Fukushima accident. Future direction of the evacuation: lessons learned. Proceedings of the 21st International Conference on Nuclear Engineering ICONE21 July 29 - August 2, 2013, Chengdu, China ICONE21-16526.
- Sakai K, Hoshi Y, Nomura T, Oda T, Iwasaki T, Fujita K, Yamada T, and Tanooka H. 2003. Suppression of carcinogenic process in mice by chronic low dose rate gamma-irradiation. *Int J Low Radiat* 1:142–146
- Sanders CL. 2010. *Radiation hormesis and the linear-no-threshold assumption*. 1st ed. Heidelberg, Dordrecht, London, New York: Springer
- Sanders CL. 2012. Potential treatment of inflammatory and proliferative diseases by ultra-low doses of ionizing radiation. *Dose-response* 10(4):610–625
- Scott BR. 2008. Low-dose risk extrapolation fallacy associated with the linear-no-threshold model. *Hum Exp Toxicol* 27:163–168

Y. Socol and others

- Scott BR, Sanders CL, Mitchel REJ, and Boreham DR. 2008. CT scans may reduce rather than increase the risk of cancer. *Journal of American Physicians and Surgeons*. 13(1):8–11
- Scott BR. 2011. Residential radon appears to prevent lung cancer. *Dose-Response* 9:444–464
- Scott BR and Dobrzyński L. 2012. Special issue introduction. *Dose-Response* 10(4):462–466
- Socol Y, Yanovskiy M, and Zatskovetsky I. 2013. Low-dose ionizing radiation: scientific controversy, moral-ethical aspects and public choice. *Int J of Nuclear Governance, Economy and Ecology*, 4:59–75
- Sponsler R and Cameron JR. 2005. Nuclear shipyard worker study (1980–1988): a large cohort exposed to low-dose-rate gamma radiation. *Int J of Low Radiation* 1: 463–478
- Tanooka H. 2001. Threshold dose response in radiation carcinogenesis: an approach for chronic beta-irradiation experiments and a review of non tumor doses. *Int J Radiat Biol* 77:541–551
- Taylor LS. 1980. Some non-scientific influences on radiation protection standards and practice – the 1980 Sievert lecture. *Health Phys* 39(6) 851–874
- Thompson RE. 2011. Epidemiological evidence for possible radiation hormesis from radon exposure: A case-control study conducted in Worcester, MA. *Dose-Response* 9(1):59–75
- Tubiana MF, Feinendegen LE, Yang C, and Kaminski JM. 2009. The linear no-threshold relationship is inconsistent with radiation biologic and experimental data. *Radiology* 251:13–22
- Ulsh BA. 2012. The new radiobiology: returning to our roots. *Dose-Response* 10(4):593–609
- Wei LC, Ding YX, Liu YH, Duan L, Bai Y, Shi M and Chen LW. 2012. Low-dose radiation stimulates Wnt/beta-catenin signaling, neural stem cell proliferation and neurogenesis of the mouse hippocampus in vitro and in vivo. *Curr Alzheimer Res* 9: 278–89