

12-2006

## RADIATION-INDUCED BYSTANDER EFFECTS: EVIDENCE FOR AN ADAPTIVE RESPONSE TO LOW DOSE EXPOSURES?

Carmel Mothersill

*McMaster University, Hamilton, ON, Canada*

Colin Seymour

*McMaster University, Hamilton, ON, Canada*

Follow this and additional works at: [https://scholarworks.umass.edu/dose\\_response](https://scholarworks.umass.edu/dose_response)

---

### Recommended Citation

Mothersill, Carmel and Seymour, Colin (2006) "RADIATION-INDUCED BYSTANDER EFFECTS: EVIDENCE FOR AN ADAPTIVE RESPONSE TO LOW DOSE EXPOSURES?," *Dose-Response: An International Journal*: Vol. 4 : Iss. 4 , Article 5. Available at: [https://scholarworks.umass.edu/dose\\_response/vol4/iss4/5](https://scholarworks.umass.edu/dose_response/vol4/iss4/5)

This Article is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Dose-Response: An International Journal by an authorized editor of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

*Dose-Response*, 4(4):283–290, 2006  
Formerly *Nonlinearity in Biology, Toxicology, and Medicine*  
Copyright © 2006 University of Massachusetts  
ISSN: 1559-3258  
DOI: 10.2203/dose-response.06-111.Mothersill



## RADIATION-INDUCED BYSTANDER EFFECTS: EVIDENCE FOR AN ADAPTIVE RESPONSE TO LOW DOSE EXPOSURES?

**Carmel Mothersill and Colin Seymour** □ Medical Physics and Applied Radiation Sciences Unit, McMaster University, Hamilton, Ontario, Canada, L8S 4K1

□ This paper reviews our current knowledge of the mechanisms underlying the induction of bystander effects by low dose, low-LET ionizing radiation and discusses how they may be related to observed adaptive responses or other protective effects of low dose exposures. Bystander effects appear to be the result of a generalized stress response in tissues or cells. The signals may be produced by all exposed cells, but the response appears to require a quorum in order to be expressed. The major response involving low LET radiation exposure discussed in the existing literature is a death response. This has many characteristics of apoptosis but is p53 independent. While a death response might appear to be adverse, the position is argued in this paper that it is in fact protective and removes damaged cells from the population. Since many cell populations carry damaged cells without being exposed to radiation, so called “background damage”, it is possible that low doses exposures cause removal of cells damaged by agents other than the test dose of radiation. This mechanism would lead to the production of “U-shaped” dose response curves. In this scenario, the level of “adaptive” or beneficial response will be related to the background damage carried by the cell population. This model may be important when attempting to predict the consequences of mixed exposures involving radiation and other environmental stressors.

### I. INTRODUCTION

Radiation-induced bystander signals appear to coordinate cellular responses even in cells not directly exposed or traversed by radiation. This work has led to a paradigm shift in radiobiology over the last 5-10 years<sup>1-4</sup>. Prior to this, it was held that DNA double strand breaks and cellular survival/damage were inextricably linked and that radiation damage could be defined as a function of DNA double strand breaks. This is now being challenged because of an increasing number of studies that demonstrate indirect (i.e., non-DNA related) effects and coordinated tissue responses<sup>5</sup>. These appear to saturate at low doses and lead to a breakdown of the dose response relationship that dominates at high doses<sup>6, 7</sup>. The low dose mechanisms may mitigate or exacerbate the direct effects

Address correspondence to Carmel Mothersill or Colin Seymour, , Medical Physics and Applied Radiation Sciences Unit, McMaster University, Hamilton, Ontario, Canada, L8S 4K1. Phone: 905 525 9140 ext 26227; fax: 905 522 5982; e-mail: mothers@mcmaster.ca, seymouc@mcmaster.ca

*C. Mothersill and C. Seymour*

of the dose and dominate the results at doses below 0.5 Gy<sup>7, 8</sup>. Signal production has been detected at doses as low as 5 mGy, although at these doses the recipient cell may transduce the signal in a different way<sup>9</sup>. Current conventional models of radiation dose response do not accommodate these new findings and as long as the mechanisms remain unclear, modelling low dose effects is difficult and uncertainty is high.

While there is obvious interest in general in this field, the key applications are likely to be in radiation protection and biotechnology. A novel mechanism for coordination of tissue responses is clearly being induced by radiation and probably by other substances. This offers new avenues for development of drugs aimed not at cell destruction but at restoring the tissues own control and coordination of response following DNA damage.

## **II. BYSTANDER EFFECTS AND ADAPTIVE RESPONSES**

Many of the newly recognized effects are similar to systemic stress or innate immune responses, in that there is no simple relationship between exposure and effect, and the outcome is not obviously dependent on dose or number of cells hit by radiation<sup>10-13</sup>. Mitochondria and reactive oxygen species appear to be important to the coordination and regulation of these effects<sup>14-17</sup>. So far, research by our group and by others has suggested that radiation causes hit cells to produce signals, which can be received by cells close to or distant from the targeted cell<sup>18-23</sup>. The recipient cells transduce the signals and appear to coordinate an appropriate (by definition ADAPTIVE) response. Responses recorded to date include initiation of apoptosis, differentiation or proliferation<sup>13, 24-26</sup>. These coordinated responses can be protective as, for example, an apoptotic response can remove an abnormal cell from the population, but the response can also involve fixation of mutations, induction of genomic instability or cellular transformation as pre-malignant responses. Which response predominates appears to depend on genetic and environmental influences and not to be related to dose<sup>27, 28</sup>.

## **III. WHAT ARE THE SIGNALING MECHANISMS?**

The nature of the signal(s) is (are) unknown, although the properties are becoming clearer. Much of the phenomenological data are suggestive of a very small (less than 1000 dalton) (lipo) peptide molecule or biogenic amine, but it is also possible to argue for long-lived radicals leading to peroxide or aldehyde release from cells<sup>22, 29-31</sup>. The mechanisms by which the cells coordinate their responses are also unknown, but signaling which leads to persistently increased ROS and modulation of biochemical pathways in mitochondria (particularly HMP shunt) have been demonstrated<sup>15, 32-34</sup>.

*Radiation-induced bystander effects***IV. MODELS TO STUDY BYSTANDER EFFECTS**

Many *in vitro* models to study these effects have been developed. These can involve irradiation using low doses of high or low LET radiation using microbeams or low fluences of alpha particles, where not all cells in the field are hit by a radiation traversal. Effects are looked for in “un-hit” cells<sup>35-40</sup>. A simple medium transfer protocol, which enables low dose, low LET radiation effects to be studied, has also been published by our own group<sup>41-42</sup>. This work has shown that medium from irradiated cells and from the distant progeny of irradiated cells contains a factor or factors, which can significantly alter survival of cells that were never irradiated and were never in contact with irradiated cells. Inhibitors of the production of the factor (or response to it) include the MAO inhibitor L-deprenyl and lactate<sup>15, 43</sup>. A major feature of current research in the field is aimed at dissecting out the relative importance of signal production and cellular response. Results to date suggest that these are independently modulated and that cell lines, which do not produce a signal, may respond to one. This clearly implicates a genetic component in the mechanism which is further indicated by the *in vivo* work available<sup>44, 45</sup>.

**V. BYSTANDER EFFECTS AND GENOMIC INSTABILITY**

One of the most interesting areas in this field is the link between bystander effects and the induction and perpetuation of genomic instability<sup>1-4</sup>. Radiation-induced genomic instability is characterized by the appearance, in cell populations, of progeny with higher than normal levels of NON-CLONAL cytogenetic abnormalities and cell death. The instability is persistent, but effects occur at a stable rate in the post-irradiation survivors for many generations. Affected progeny populations do not either die out or dominate—an apparent paradox, which is difficult to reconcile with the current “world view” of competitive natural selection of favourable genes. The mechanism of perpetuation is now thought to be epigenetic and to involve an excess generation of reactive oxygen species (ROS). This is “signaled” to neighbours and perpetuated in progeny via mechanisms similar to the bystander mechanisms discussed earlier. The transmissible factors are very likely to be related to “bystander factors”.

While knowledge about radiation-induced genomic instability and bystander effects has been growing in the radiation field for over 15 years, it has only recently become apparent that chemicals in the natural environment can also induce the state of genomic instability in cells and hence low dose chemical toxicity probably also involves bystander effects<sup>46-47</sup>. This widens the relevance of these indirect damage mechanisms to include environmental toxins other than radiation and makes it important to understand the mechanisms involved as they may contribute to mixed exposure responses in biota.

*C. Mothersill and C. Seymour*

## **VI. RELEVANCE OF BYSTANDER EFFECTS IN THE ADAPTIVE RESPONSE FIELD**

Evidence, which suggests that bystander mechanisms may be involved in adaptive responses, comes from published data and also from deductive reasoning. The published data show that signals produced by irradiated cells can induce protection against a real dose of ionizing radiation<sup>48, 49</sup>. These authors have also shown that intracellular calcium fluxes precede the induction of responses in bystander cells exposed to signals from irradiated cells<sup>24, 50-52</sup>. While the response that generally follows exposure to these bystander signals is cell death, this can be protective if it eliminates damaged cells from the population. Following low dose exposure, where few cells will have damage, it seems appropriate to remove them. At higher doses, where many cells are damaged and tissue integrity is at risk of collapse, such a bystander effect would be an added problem for an already compromised population. It is interesting to note here that repair deficient cells have larger death-inducing bystander effects than the corresponding repair proficient parent lines<sup>53, 54</sup>. This would be expected if the damaged cell couldn't be repaired, if the bystander effect is assumed to be protective. Many cell lines and most tumour explants do not produce death-inducing signals after exposure to radiation, and no calcium pulse is seen<sup>55-57</sup>. It is not known whether they produce no signals or whether different signals, not transduced through the calcium pulse-apoptotic death pathway, are involved.

## **VII. MODELS AND RELEVANCE TO RADIATION PROTECTION**

If we accept that bystander effects are the result of a generalized stress response in tissues or cells, what are the implications for radiation protection? Does the effect alter the acceptability of the Linear-No-Threshold hypothesis, upon which all radiation protection legislation is based? How can dose be used as a measure of effect or harm, if low doses (which are those experienced in the workplace) do not produce any type of linear dose-effect curve? There is clearly some very complex biology involved because the signals may be produced by all exposed cells, but the response appears to require a quorum in order to be expressed<sup>18, 58-60</sup>. The major response involving low LET radiation exposure discussed in the existing literature is a death response. This has many characteristics of apoptosis, but is p53 independent? While a death response might appear to be adverse, the position is argued in this paper that it is in fact protective and removes damaged cells from the population. Since many cell populations carry damaged cells without being exposed to radiation, so called "background damage", it is possible that low dose exposures cause removal of cells damaged by agents other than the test dose of radiation. This mechanism could lead to the production of U-shaped dose

*Radiation-induced bystander effects*

response curves so common in toxicology<sup>61-63</sup>. In this scenario, the level of “adaptive” or beneficial response will be related to the level of background damage carried by the cell population. These considerations may be important when attempting to predict the consequences of mixed exposures involving radiation and other environmental stressors.

**VIII. CONCLUSION**

To conclude, it is clear that adaptive responses, bystander effects and genomic instability belong to a suite of effects that predominately modulate the low dose response to radiation. These mechanisms are part of the cellular homeostatic response and, while we can detect low dose *effects*, there is little evidence that these translate into *harm*. It is likely that for many genotypes there is an operational threshold for harmful radiation damage that probably occurs at a point where the functional activity of the tissue is being compromised by the level of (protective) cell death. For genotypes where the bystander response, if there is one, does not involve coordinated cell death, it is likely that there is no operational threshold and that stochastic effects such as carcinogenesis have some very small probability of occurring at low doses. What this probability is, though, is not easy to determine. It is unlikely to be definable by extrapolation from high dose data because the underlying mechanisms are so different. Many of the current research efforts in this field are aimed at modulating the bystander effect using chemicals. This approach should, perhaps, not only look at preventing the bystander effect but also at causing or simulating it in tissues and cells which do not have the capacity to mount this response.

**ACKNOWLEDGMENTS**

We acknowledge support from the Science Foundation of Ireland and Saint Luke’s Institute of Cancer Research.

**REFERENCES**

1. Morgan WF. (2003) Non-targeted and delayed effects of exposure to ionizing radiation: I. Radiation-induced genomic instability and bystander effects in vitro. *Radiat Res* 159:567-580 Review.
2. Mothersill C, Seymour C. (2003) Low-dose radiation effects: experimental hematology and the changing paradigm. *J Exp Haemat* 31:437-445 Review.
3. Lorimore SA, Wright EG. (2003) Radiation-induced genomic instability and bystander effects: related inflammatory-type responses to radiation-induced stress and injury? A review. *Int J Radiat Biol* 79:15-25 Review.
4. Little JB, Morgan WF. (2003) Guest editors. *Oncogene* 13;22(45):6977.
5. Barcellos-Hoff MH, Brooks AL. (2001) Extracellular signaling through the microenvironment: a hypothesis relating carcinogenesis, bystander effects, and genomic instability. *Radiat Res* 156:618-627.
6. Prise KM, Folkard M, Michael BD. (2003) A review of the bystander effect and its implications for low-dose exposure. *Radiat Prot Dosimetry* 104:347-355.

## C. Mothersill and C. Seymour

7. Seymour CB, Mothersill C. (2000) Relative contribution of bystander and targeted cell killing to the low-dose region of the radiation dose-response curve. *Radiat Res* 153:508-511.
8. Zhou H, Randers-Pehrson G, Geard CR, Brenner DJ, Hall EJ, Hei TK. (2003) Interaction between radiation-induced adaptive response and bystander mutagenesis in mammalian cells. *Radiat Res* 160:512-516.
9. Keyes E, Howe O, Seymour CB, Lyng F, Mothersill C. (in press) Prolonged expression of delayed cell death in the progeny of cells exposed to direct irradiation or to culture medium from irradiated cells. *Int J Radiat Biol*.
10. Zhou H, Randers-Pehrson G, Suzuki M, Waldren CA, Hei TK. (2002) Genotoxic damage in non-irradiated cells: contribution from the bystander effect. *Radiat Prot Dosimetry* 99:227-232.
11. Azzam EI, DE Toledo SM, Little JB. (2003) Oxidative metabolism, gap junctions and the ionizing radiation-induced bystander effect. *Oncogene* 13:7050-7057.
12. Geard CR, Jenkins-Baker G, Marino SA, Ponnaiya B. (2002) Novel approaches with track segment alpha particles and cell co-cultures in studies of bystander effects. *Radiat Prot Dosimetry* 99:233-236.
13. Belyakov OV, Folkard M, Mothersill C, Prise KM, Michael BD. (2003) A proliferation-dependent bystander effect in primary porcine and human urothelial explants in response to targeted irradiation. *Br J Cancer* 88:767-774.
14. Limoli CL, Giedzinski E. (2003) Induction of chromosomal instability by chronic oxidative stress. *Neoplasia* 5:339-346.
15. Mothersill C, Stamato TD, Perez ML, Cummins R, Mooney R, Seymour CB. (2000) Involvement of energy metabolism in the production of 'bystander effects' by radiation. *Br J Cancer* 82:1740-1746.
16. Iyer R, Lehnert BE. (2002) Low dose, low-LET ionizing radiation-induced radioadaptation and associated early responses in unirradiated cells. *Mutat Res* 503:1-9.
17. Shao C, Furusawa Y, Kobayashi Y, Funayama T, Wada S. (2003) Bystander effect induced by counted high-LET particles in confluent human fibroblasts: a mechanistic study. *FASEB J* 17:1422-1427.
18. Gerashchenko BI, Howell RW. (2003) Cell proximity is a prerequisite for the proliferative response of bystander cells co-cultured with cells irradiated with gamma-rays. *Cytometry* 56A:71-80.
19. Lorimore SA, Coates PJ, Wright EG. (2003) Radiation-induced genomic instability and bystander effects: inter-related non-targeted effects of exposure to ionizing radiation. *Oncogene* 22:7058-7069.
20. Little JB, Azzam EI, De Toledo SM, Nagasawa H. (2002) Bystander effects: intercellular transmission of radiation damage signals. *Radiat Prot Dosimetry* 99:159-162.
21. Reznikov K, Kolesnikova L, Pramanik A, Tan-No K, Gileva I, Yakovleva T, Rigler R, Terenius L, Bakalkin G. (2000) Clustering of apoptotic cells via bystander killing by peroxides. *FASEB J* 14:1754-1764.
22. Emerit I, Oganessian N, Arutyunian R, Pogossian A, Sarkisian T, Cernjavski L, Levy A, Feingold J. (1997) Oxidative stress-related clastogenic factors in plasma from Chernobyl liquidators: protective effects of antioxidant plant phenols, vitamins and oligoelements. *Mutat Res* 377:239-246.
23. Faguet GB, Reichard SM, Welter DA. (1984) Radiation-induced clastogenic plasma factors. *Cancer Genet Cytogenet* 12:73-83.
24. Trosko JE. (1998) Hierarchical and cybernetic nature of biologic systems and their relevance to homeostatic adaptation to low-level exposures to oxidative stress-inducing agents. *Environ Health Perspect* 106 Suppl 1:331-339 Review.
25. Dent P, Yacoub A, Fisher PB, Hagan MP, Grant S. (2003) MAPK pathways in radiation responses. *Oncogene* 22(37):5885-5896 Review.
26. Belyakov OV, Folkard M, Mothersill C, Prise KM, Michael BD. (2002) Bystander-induced apoptosis and premature differentiation in primary urothelial explants after charged particle microbeam irradiation. *Radiat Prot Dosimetry* 99:249-251.
27. Baverstock K. (2000) Radiation-induced genomic instability: a paradigm-breaking phenomenon and its relevance to environmentally induced cancer. *Mutat Res* 254:89-109 Review.
28. Mothersill C, Rea D, Wright EG, Lorimore SA, Murphy D, Seymour CB, O'malley K. (2001) Individual variation in the production of a 'bystander signal' following irradiation of primary cultures of normal human urothelium. *Carcinogenesis* 22:1465-1471.

*Radiation-induced bystander effects*

29. Davies MJ. (2003) Singlet oxygen-mediated damage to proteins and its consequences. *Biochem Biophys Res Commun* 305:61-70 Review.
30. Balcer-Kubiczek EK, Harrison GH, Xu JF, Gutierrez PL. (2002) Coordinate late expression of trefoil peptide genes (pS2/TFF1 and ITF/TFF3) in human breast, colon, and gastric tumor cells exposed to X-rays. *Mol Cancer Ther* 6:405-415.
31. Azzam EI, De Toledo SM, Spitz DR, Little JB. (2002) Oxidative metabolism modulates signal transduction and micronucleus formation in bystander cells from alpha-particle-irradiated normal human fibroblast cultures. *Cancer Res* 62:5436-5442.
32. Lehnert BE, Iyer R. (2002) Exposure to low-level chemicals and ionizing radiation: reactive oxygen species and cellular pathways. *Hum Exp Toxicol* 21:65-69.
33. Iyer R, Lehnert BE. (2002) Low dose, low-LET ionizing radiation-induced radioadaptation and associated early responses in unirradiated cells. *Mutat Res* 503:1-9.
34. Burlakova EB, Mikhailov VF, Azurik VK. (2001) The redox homeostasis system in radiation-induced genomic instability. *Radiat Biol Radioecol* 41:489-499.
35. Osterreicher J, Prise KM, Michael BD, VOGT J, Butz T, Tanner JM. (2003) Radiation-induced bystander effects: Mechanisms, biological implications, and current investigations at the Leipzig LIPSION facility. *Strahlenther Onkol* 179:69-77 Review.
36. Prise KM, Belyakov OV, Newman HC, Patel S, Schettino G, Folkard M, Michael BD. (2002) Nontargeted effects of radiation: bystander responses in cell and tissue models. *Radiat Prot Dosimetry* 99:223-226.
37. Sawant SG, Zheng W, Hopkins KM, Randers-Pehrson G, Lieberman HB, Hall EJ. (2002) The radiation-induced bystander effect for clonogenic survival. *Radiat Res* 157:361-364.
38. Prise KM, Belyakov OV, Folkard M, Michael BD. (1998) Studies of bystander effects in human fibroblasts using a charged particle microbeam. *Int J Radiat Biol* 74:793-798.
39. Iyer R, Lehnert BE. (2002) Alpha-particle-induced increases in the radioresistance of normal human bystander cells. *Radiat Res* 157:3-7.
40. Nagasawa H, Little JB. (1992) Induction of sister chromatid exchanges by extremely low doses of alpha-particles. *Cancer Res* 52:6394-6396.
41. Mothersill C, Seymour C. (1997) Medium from irradiated human epithelial cells but not human fibroblasts reduces the clonogenic survival of unirradiated cells. *Int J Radiat Biol* 71:21-27.
42. Mothersill C, Seymour CB. (1998) Cell-cell contact during gamma irradiation is not required to induce a bystander effect in normal human keratinocytes: evidence for release during irradiation of a signal controlling survival into the medium. *Radiat Res* 149:256-262.
43. Seymour CB, Mothersill C, Mooney R, Moriarty M, Tipton KF. (2003) Monoamine oxidase inhibitors l-deprenyl and clorgyline protect nonmalignant human cells from ionising radiation and chemotherapy toxicity. *Br J Cancer* 89:1979-1986.
44. Coates PJ, Lorimore SA, Lindsay KJ, Wright EG. (2003) Tissue-specific p53 responses to ionizing radiation and their genetic modification: the key to tissue-specific tumour susceptibility? *J Pathol* 201:377-388.
45. Lorimore SA, Coates PJ, Scobie GE, Milne G, Wright EG. (2001) Inflammatory-type responses after exposure to ionizing radiation in vivo: a mechanism for radiation-induced bystander effects? *Oncogene* 20:7085-7095.
46. Mothersill C, Crean M, Lyons M, McSweeney J, Mooney R, O'Reilly J, Seymour CB. (1998) Expression of delayed toxicity and lethal mutations in the progeny of human cells surviving exposure to radiation and other environmental mutagens. *Int J Radiat Biol* 74:673-680.
47. Coen N, Mothersill C, Kadhim M, Wright EG. (2001) Heavy metals of relevance to human health induce genomic instability. *J Pathol* 195:293-299.
48. Mothersill C, Seymour CB. (2002) Bystander and delayed effects after fractionated radiation exposure. *Radiat Res* 158:626-633.
49. Maguire M, Mothersill C, Seymour CB and Lyng FM. An adaptive response following exposure of cultures to medium from cells irradiated to low doses of cobalt 60 gamma rays. *Radiation Res* (in press).
50. Lyng FM, Seymour CB, Mothersill C. (2002) Early events in the apoptotic cascade initiated in cells treated with medium from the progeny of irradiated cells. *Radiat Prot Dosimetry* 99:169-172.
51. Lyng FM, Seymour CB, Mothersill C. (2002) Initiation of apoptosis in cells exposed to medium from the progeny of irradiated cells: a possible mechanism for bystander-induced genomic instability? *Radiat Res* 157:365-370.



C. Mothersill and C. Seymour

52. Lyng FM, Seymour CB, Mothersill C. (2000) Production of a signal by irradiated cells which leads to a response in unirradiated cells characteristic of initiation of apoptosis. *Br J Cancer* 83:1223-1230.
53. Nagasawa H, Huo L, Little JB. (2003) Increased bystander mutagenic effect in DNA double-strand break repair-deficient mammalian cells. *Int J Radiat Biol* 79:35-41.
54. Mothersill, C., Seymour RJ, and Seymour CB. (2004) Bystander effects in repair deficient cell lines. *Radiat. Res* (in press).
55. Mothersill C, Seymour CB, Joiner MC. (2002) Relationship between radiation-induced low-dose hypersensitivity and the bystander effect. *Radiat Res* 157:526-532.
56. Nagar S, Smith LE, Morgan WF. (2003) Characterization of a novel epigenetic effect of ionizing radiation: the death-inducing effect. *Cancer Res* 63:324-328.
57. Mothersill C, Lyng F, O'Reilly S, Harney J, Seymour CB. (1996) Expression of lethal mutations is suppressed in neoplastically transformed cells and after treatment of normal cells with carcinogens. *Radiat Res* 145:714-721.
58. Mothersill C, Seymour C. (1997) Survival of human epithelial cells irradiated with cobalt 60 as microcolonies or single cells. *Int J Radiat Biol* 72:597-606.
59. Erickson AC, Barcellos-Hoff MH. (2003) The not-so innocent bystander: the microenvironment as a therapeutic target in cancer. *Expert Opin Ther Targets* 7:71-88.
60. Bishayee A, Hill HZ, Stein D, Rao DV, Howell RW. (2001) Free radical-initiated and gap junction-mediated bystander effect due to non-uniform distribution of incorporated radioactivity in a three-dimensional tissue culture model. *Radiat Res* 155:335-344.
61. Calabrese EJ, Baldwin LA. (2002) Applications of hormesis in toxicology, risk assessment and chemotherapeutics. *Trends Pharmacol Sci* 23:331-337 Review.
62. Calabrese EJ, Baldwin LA. (2001) The frequency of U-shaped dose responses in the toxicological literature. *Toxicol Sci* 62:330-338.
63. Calabrese EJ, Baldwin LA. (2001) Hormesis: U-shaped dose responses and their centrality in toxicology. *Trends Pharmacol Sci* 22:285-291 Review.