

University of Massachusetts Amherst  
**ScholarWorks@UMass Amherst**

---

International Conference on Engineering and  
Ecohydrology for Fish Passage

International Conference on Engineering and  
Ecohydrology for Fish Passage 2015

---

Jun 23rd, 5:00 PM - 5:15 PM

## Session A6: Timing, Frequency and Environmental Conditions Associated with Mainstem-Tributary Movement by a Lowland River Fish, Golden Perch

Wayne Koster

*Arthur Rylah Institute, DELWP, Victoria, Australia, [wayne.koster@delwp.vic.gov.au](mailto:wayne.koster@delwp.vic.gov.au)*

David Dawson

*Arthur Rylah Institute, DELWP, Victoria, Australia*

Follow this and additional works at: [https://scholarworks.umass.edu/fishpassage\\_conference](https://scholarworks.umass.edu/fishpassage_conference)



Part of the [Aquaculture and Fisheries Commons](#), and the [Hydraulic Engineering Commons](#)

---

Koster, Wayne and Dawson, David, "Session A6: Timing, Frequency and Environmental Conditions Associated with Mainstem-Tributary Movement by a Lowland River Fish, Golden Perch" (2015). *International Conference on Engineering and Ecohydrology for Fish Passage*. 7.

[https://scholarworks.umass.edu/fishpassage\\_conference/2015/June23/7](https://scholarworks.umass.edu/fishpassage_conference/2015/June23/7)

This Event is brought to you for free and open access by the Fish Passage Community at UMass Amherst at ScholarWorks@UMass Amherst. It has been accepted for inclusion in International Conference on Engineering and Ecohydrology for Fish Passage by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

# Patterns of connectivity between mainstem–tributary habitats for a lowland river fish

Wayne Koster and David Dawson

Arthur Rylah Institute, DELWP, Victoria, Australia  
Wayne.Koster@delwp.vic.gov.au



Department of  
Environment, Land,  
Water & Planning



# Background



- Connections between mainstem and tributaries important for range of ecological processes and functions

# Background



- Connections between mainstem and tributaries important for range of ecological processes and functions
- Mainstem–tributary movements important for maintaining fish populations in river networks

# Background



- Connections between mainstem and tributaries important for range of ecological processes and functions
- Mainstem–tributary movements important for maintaining fish populations in river networks
- Hydrological regimes major driver of river ecosystems and provide cues for a range of important behaviours in fishes

# Aims



1. To determine patterns of connectivity between a mainstem river and tributary for golden perch

# Aims



1. To determine patterns of connectivity between a mainstem river and tributary for golden perch
2. To determine whether the frequency and/or direction of mainstem–tributary movement change during the spawning period

# Aims



1. To determine patterns of connectivity between a mainstem river and tributary for golden perch
2. To determine whether the frequency and/or direction of mainstem–tributary movement change during the spawning period
3. To determine whether hydrology or temperature influences the occurrence of mainstem–tributary movements



# Methods




- Study area – Goulburn and Murray rivers, southeast

Australia

Goulburn River



Murray River




# Methods



- Study area – Goulburn and Murray rivers, southeast

Australia

- 79 adult fish tagged (2007-2009)




# Methods



- Study area – Goulburn and Murray rivers, southeast

Australia

- 79 adult fish tagged (2007-2009)
- 12 acoustic loggers deployed  
(data collected 2007-2012)



# Findings



- One quarter of Murray River (mainstem) fish moved into Goulburn (tributary)

# Findings





- One quarter of Murray River (mainstem) fish moved into Goulburn (tributary)
- Half of Goulburn River (tributary) fish moved into Murray (mainstem)

# Findings



- One quarter of Murray River (mainstem) fish moved into Goulburn (tributary)
- Half of Goulburn River (tributary) fish moved into Murray (mainstem)
- Mostly temporary occupation, but ~10% shift between rivers


# Findings





- Differences in temporal patterns of movement among rivers

*A – Fish tagged in Murray River (mainstem)*

*B – Fish tagged in Goulburn River (tributary)*





# Findings



- Differences in temporal patterns of movement among rivers



*A – Fish tagged in Murray River (mainstem)*

*B – Fish tagged in Goulburn River (tributary)*





# Findings




- Differences in temporal patterns of movement among rivers

*A – Fish tagged in Murray River  
(mainstem)*




*B – Fish tagged in Goulburn River  
(tributary)*



# Findings




- Movement from tributary to mainstem



# Findings



- Movement from mainstem to tributary




# Conclusion




- Spatially and temporally complex pattern of movement between mainstem and tributary locations

# Conclusion




1. Most fish remain in their original 'home' river

# Conclusion




1. Most fish remain in their original 'home' river
2. Some fish move between mainstem and tributary locations, characterised by temporary occupation, followed by fish returning to their original river

# Conclusion



1. Most fish remain in their original 'home' river
2. Some fish move between mainstem and tributary locations, characterised by temporary occupation, followed by fish returning to their original river
  - a) Movement of tributary fish into the mainstem is common during spring/summer, particularly during changes in flow


# Conclusion



1. Most fish remain in their original 'home' river
2. Some fish move between mainstem and tributary locations, characterised by temporary occupation, followed by fish returning to their original river
  - a) Movement of tributary fish into the mainstem is common during spring/summer, particularly during changes in flow
  - b) Movement of mainstem fish into tributaries is not concentrated during any one period



# Conclusion



1. Most fish remain in their original 'home' river
2. Some fish move between mainstem and tributary locations, characterised by temporary occupation, followed by fish returning to their original river.
3. Some fish shift between mainstem and tributary locations and do not return

# Conclusion



- Spatially and temporally complex pattern of movement between mainstem and tributary locations
- Fish populations don't conform to artificially constrained management units

# Conclusion



- Spatially and temporally complex pattern of movement between mainstem and tributary locations
- Fish populations don't conform to artificially constrained management units
- Important to consider flows to facilitate connectivity among rivers

# Conclusion



- Spatially and temporally complex pattern of movement between mainstem and tributary locations
- Fish populations don't conform to artificially constrained management units
- Important to consider flows to facilitate connectivity among rivers
- Environmental flow recommendations need to be developed interdependently across rivers

# Acknowledgements



- Dave Crook (Charles Darwin University)
- Paul Moloney (Arthur Rylah Institute)
- Simon Casanelia, Geoff Earl, Mark Turner, Wayne Tennant, Megan Judd (Goulburn Broken Catchment Management Authority)

OPEN ACCESS Freely available online

PLOS ONE

## Timing, Frequency and Environmental Conditions Associated with Mainstem–Tributary Movement by a Lowland River Fish, Golden Perch (*Macquaria ambigua*)

Wayne M. Koster<sup>1,2\*</sup>, David R. Dawson<sup>1</sup>, Damien J. O'Mahony<sup>1a</sup>, Paul D. Moloney<sup>1</sup>, David A. Crook<sup>1ab</sup>

<sup>1</sup> Arthur Rylah Institute for Environmental Research, Department of Environment and Primary Industries, Heidelberg, Victoria, Australia, <sup>2</sup> School of Life and Environmental Sciences, Deakin University, Warrambool, Victoria, Australia