

Utah State University

DigitalCommons@USU

---

Aspen Bibliography

Aspen Research

---


1977

## Biomass Estimation For Some Shrubs From Northeastern Minnesota

David F. Grigal

Lewis F. Ohmann

Follow this and additional works at: [https://digitalcommons.usu.edu/aspens\\_bib](https://digitalcommons.usu.edu/aspens_bib)

 Part of the [Agriculture Commons](#), [Ecology and Evolutionary Biology Commons](#), [Forest Sciences Commons](#), [Genetics and Genomics Commons](#), and the [Plant Sciences Commons](#)


---

### Recommended Citation

Grigal, David F. and Ohmann, Lewis F. 1977. Biomass estimation for some shrubs from northeastern Minnesota. USDA Forest Service. Research Note NC-226. North Central Forest Experiment Station, St. Paul, Minnesota

This Report is brought to you for free and open access by the Aspen Research at DigitalCommons@USU. It has been accepted for inclusion in Aspen Bibliography by an authorized administrator of DigitalCommons@USU. For more information, please contact [digitalcommons@usu.edu](mailto:digitalcommons@usu.edu).





## RESEARCH NOTE NC-226

NORTH CENTRAL FOREST EXPERIMENT STATION, FOREST SERVICE—U.S. DEPARTMENT OF AGRICULTURE  
Folwell Avenue, St. Paul, Minnesota 55101 1977

### BIOMASS ESTIMATION FOR SOME SHRUBS FROM NORTHEASTERN MINNESOTA

David F. Grigal, *Associate Professor*  
*Soil Science and Forest Resources Department*  
*University of Minnesota*  
*St. Paul, Minnesota*

and

Lewis F. Ohmann, *Principal Plant Ecologist*  
*St. Paul, Minnesota*

**ABSTRACT.**--Biomass prediction equations were developed for 23 northeastern Minnesota shrub species. The allometric function was used to predict leaf, current annual woody twig, stem, and total woody biomass (dry grams), using stem diameter class estimated to the nearest 0.25 cm class at 15 cm above ground level as the independent variable.

**OXFORD:** 182.46:182.5(776). **KEY WORDS:** wild-life habitat, nonlinear regression analysis, allometric relations, dry weight, browse estimation.

A recent study (unpublished) to quantitatively describe a series of pine plantations in northeastern Minnesota as white-tailed deer and snowshoe hare habitat placed special emphasis on woody plants as potential browse. The descriptive information included shrub density and basal area. To estimate the browse represented by these data, equations were needed to determine the biomass of leaf and current annual woody-twig growth for each species. Reported here are the data collected and the resulting regression equations developed to produce the needed estimates. These equations should be useful to land managers, game managers, and other researchers who need to estimate biomass values for these species within the northern Lake States region.

For each of the major shrub species found in the plantations, stems spanning the size range were collected during the last half of

August 1976. In the plantation study, shrub diameters had been determined to the nearest 0.25 cm at 15 cm above ground level. In sampling for the biomass estimation equations, we attempted to collect individuals from the lowest, highest, and middle size class for each species. Thus, if a species occurred in the study in size classes 1 (0 to 0.25 cm) through 7 (1.50 to 1.75 cm), we selected a minimum of three typical stems within each of three size classes spanning that range: for example, size class 1, 4 (0.75 to 1.00 cm), and 7. For shrubs with small diameters, such as *Rosa blanda*, we selected only stems within the lowest three size classes.

The stems were selected in the field, clipped at ground level, and taken to the laboratory where the leaves and current annual woody-twig growth were each separated from the remainder of the woody stem. Each component of each stem was bagged separately and dried at 70 C to constant weight. Each component was weighed to the nearest 0.01 gram immediately after removal from the oven.

Because of dry weather during the summer of 1976, some shrub stems had lost a portion of their leaves by late August. In those cases no leaf data were collected and the number of leaf biomass observations used in the leaf biomass regressions was reduced.

The independent variable, stem diameter size class; and each dependent variable-leaf,

current annual woody-twig, stem, and total woody biomass were summarized for each species (table 1).

The numerical value used for stem diameter was the upper end of the diameter class range. Stems 0.50 to 0.75 cm diameter, for example, were considered to be 0.75 cm diameter in the data summary.

In earlier work estimating biomass of

five shrub species from northeastern Minnesota, we found that shrub biomass data conformed to the allometric function as well as or better than they did to alternative functional relations (Ohmann, Grigal, and Brander 1976)<sup>1</sup>.

<sup>1</sup>Ohmann, L. F., D. F. Grigal, and R. B. Brander. 1976. Biomass estimation for five shrubs from northeastern Minnesota. USDA For. Serv. Res. Pap. NC-133, 11 p. North Cent. For. Exp. Stn., St. Paul, MN 55108.

Table 1.--Dimensional data for 23 shrub species

	<i>Acer</i>	<i>Acer</i>	<i>Alnus</i>	<i>Alnus</i>	<i>Amelanchier</i>	<i>Betula</i>	<i>Cornus</i>	<i>Corylus</i>	<i>Lonicera</i>	<i>Lonicera</i>	<i>Lonicera</i>	<i>Populus</i>
	<i>nigrum</i>	<i>spicatum</i>	<i>crispa</i>	<i>rugosa</i>	spp.	<i>papyrifera</i>	<i>rugosa</i>	<i>cornuta</i>	<i>canadensis</i>	<i>hirsuta</i>	<i>oblongifolia</i>	<i>tremula</i>
Independent variable												
Stem diameter class (cm)	10	27	26	15	27	10	9	43	9	13	4	28
Number of observations	10	27	26	15	27	10	9	43	9	13	4	28
Mean	1.17	1.17	1.14	.90	1.17	1.23	1.00	1.08	.75	.69	.44	1.14
Standard deviation	.51	.52	.52	.53	.52	.55	.43	.53	.43	.27	.13	.53
Standard error	.16	.10	.10	.14	.10	.17	.14	.08	.14	.38	.06	.10
Range	.50-	.50-	.50-	.50-	.50-	.50-	.50-	.50-	.25-	.25-	.25-	.50-
	1.75	1.75	1.75	1.75	1.75	1.75	1.50	1.75	1.25	1.00	.50	1.75
Dependent variables												
Leaf biomass (dry grams)												
Number of observations	---	11	25	12	14	4	9	9	4	10	4	14
Mean	---	4.33	9.51	6.04	8.62	3.11	11.41	15.46	2.33	2.34	.54	6.77
Standard deviation	---	11.79	9.18	8.64	6.75	4.33	10.35	15.75	1.52	2.86	.35	9.81
Standard error	---	3.56	1.84	2.49	1.80	2.16	3.45	5.25	.76	.91	.18	2.62
Range	---	.01-	.07-	.15-	.05-	.62-	1.32-	.11-	.16-	.18-	.08-	.16-
	---	39.44	27.00	29.08	19.95	9.57	27.17	36.21	3.68	9.29	.89	25.37
Current annual twig (dry g)												
Number of observations	10	27	26	15	27	10	9	43	9	13	4	28
Mean	1.04	2.48	2.73	5.02	2.31	4.34	2.42	2.76	.97	1.15	.34	1.92
Standard deviation	.72	2.49	3.31	6.31	2.52	6.85	2.03	2.92	.92	1.36	.36	1.80
Standard error	.23	.48	.65	1.63	.49	2.17	.68	.45	.31	.38	.18	.34
Range	.10-	.12-	.18-	.36-	.01-	.51-	.25-	.12-	.06-	.06-	.05-	.19-
	2.39	10.83	12.67	19.80	8.26	23.23	5.58	10.30	2.50	4.62	.85	6.38
Stem (dry grams)												
Number of observations	10	27	26	15	27	10	9	43	9	13	4	28
Mean	69.62	87.99	78.69	43.38	89.39	76.30	48.18	88.23	19.49	19.24	1.32	75.50
Standard deviation	73.00	87.81	77.76	63.38	84.50	81.78	50.14	102.96	19.69	22.14	1.00	73.51
Standard error	23.09	16.90	15.25	16.36	16.26	25.86	16.71	15.70	6.56	6.14	.50	13.89
Range	.81-	.34-	.34-	.36-	.48-	.51-	1.30-	.57-	.27-	.20-	.20-	.42-
	207.11	345.90	268.00	171.40	258.72	247.69	131.57	380.75	52.52	65.20	2.62	234.69
Total woody (dry grams)												
Number of observations	10	27	26	15	27	10	9	43	9	13	4	28
Mean	70.66	90.47	81.41	48.39	91.70	80.65	50.60	91.00	20.46	20.39	1.66	77.42
Standard deviation	73.50	89.10	78.74	68.93	86.07	83.38	52.12	105.66	20.54	23.02	1.13	74.79
Standard error	23.24	17.15	15.44	17.80	16.56	26.37	17.37	16.11	6.85	6.38	.56	14.13
Range	.94-	.66-	1.05-	.72-	.96-	1.02-	1.55-	.60-	.54-	.27-	.25-	.84-
	208.19	351.90	271.00	181.37	259.83	251.62	137.15	391.05	54.86	66.98	2.94	239.04

Table 2.--Regressions for estimation of biomass of 23 shrubs from

Regression factors on	<i>Acer</i>	<i>Acer</i>	<i>Alnus</i>	<i>Alnus</i>	<i>Amelanchier</i>	<i>Betula</i>	<i>Cornus</i>	<i>Corylus</i>	<i>Lonicera</i>	<i>Lonicera</i>	<i>Lonicera</i>	<i>Populus</i>
stem diameter class (cm) (X)	<i>nigrum</i>	<i>spicatum</i>	<i>crispa</i>	<i>rugosa</i>	spp.	<i>papyrifera</i>	<i>rugosa</i>	<i>cornuta</i>	<i>canadensis</i>	<i>hirsuta</i>	<i>oblongifolia</i>	<i>tremula</i>
Leaf biomass (dry grams) (Y)												
a	---	2.869	5.650	3.123	5.432	3.421	8.616	4.808	6.592	3.926	6.009	2.227
b	---	3.669	2.222	3.071	2.008	1.838	2.541	3.571	2.681	1.163	3.115	4.258
R <sup>2</sup>	---	.608	.748	.718	.875	.990	.940	.991	.901	.242	.749	.963
S y.x	---	7.788	4.706	4.815	2.486	.524	2.701	1.568	.586	2.644	.217	1.956
Probability of larger	---	.005	.005	.005	.005	.010	.005	.005	.100	NS	NS	.005
F value	---											
Current annual woody twig biomass (dry g)												
a	0.7676	2.174	2.349	4.122	1.510	.588	1.813	1.196	1.325	1.974	3.756	.894
b	1.4903	1.008	1.119	2.318	1.870	4.802	2.478	3.070	1.866	1.571	3.116	2.501
R <sup>2</sup>	.724	.260	.196	.815	.426	.282	.946	.774	.801	.281	.284	.430
S y.x	.4021	2.186	3.028	2.815	1.947	6.155	.506	1.406	.440	1.208	.373	1.381
Probability of larger	.005	.010	.025	.005	.005	NS	.005	.005	.005	.100	NS	.001
F value												
Stem biomass (dry grams)												
a	21.0780	41.074	37.137	27.452	36.439	22.260	30.648	36.858	26.766	44.156	14.338	33.711
b	3.6892	2.706	2.772	3.122	2.995	3.452	3.184	3.272	2.180	3.578	3.082	2.871
R <sup>2</sup>	.908	.779	.845	.985	.922	.717	.955	.845	.930	.684	.563	.911
S y.x	23.5026	42.107	31.242	8.069	24.132	46.119	11.340	41.061	5.566	13.006	.806	22.021
Probability of larger	.005	.005	.005	.005	.005	.005	.005	.005	.005	.010	NS	.001
F value												
Total woody biomass (dry grams)												
a	21.7300	43.660	39.684	31.328	37.909	22.865	32.421	38.031	28.090	46.002	18.093	34.501
b	3.6535	2.630	2.696	3.050	2.963	3.502	3.152	3.267	2.166	3.402	3.089	2.871
R <sup>2</sup>	.911	.778	.855	.991	.924	.764	.957	.847	.931	.678	.695	.911
S y.x	23.3043	42.776	30.571	6.824	24.158	42.982	11.590	41.881	5.759	13.635	.762	22.311
Probability of larger	.005	.005	.005	.005	.005	.005	.005	.005	.005	.005	NS	.001
F value												

<sup>1</sup>All the woody growth was determined to be current annual. Stem and total woody biomass regressions were identical to current twig.

In this study, then, the data for each species-variable combination were subjected to an iterative nonlinear, least squares regression analysis using the allometric relation

$$Y = aX^b \quad (1)$$

where Y is the biomass in grams dry weight of the dependent variable and X is the shrub diameter in cm at 15 cm above ground. The

regressions were tested for significance using a standard F-test.

The resultant equation elements are presented in table 2.

r 23 shrub species from northeastern Minnesota

<i>nicera</i>	<i>Populus</i>	<i>Prunus</i>	<i>Prunus</i>	<i>Rhamnus</i>	<i>Ribes</i>	<i>Rosa</i>	<i>Rosa</i>	<i>Rubus</i>	<i>Rubus</i>	<i>Salix</i>	<i>Sorbus</i>	<i>Viburnum</i>
<i>longifolia</i>	<i>tremuloides</i>	<i>pennsylvanica</i>	<i>virginiana</i>	<i>alnifolia</i>	spp.	<i>acicularis</i>	<i>blanda</i>	<i>parviflorus</i>	<i>strigosus</i>	spp.	<i>americana</i>	<i>rafinosquianum</i>
4	28	9	9	6	9	8	9	9	11	9	9	9
.44	1.14	1.17	.75	1.08	.50	.53	.50	.50	.48	1.17	1.17	1.00
.13	.53	.54	.43	.49	.22	.21	.22	.22	.21	.54	.54	.43
.06	.10	.18	.14	.20	.07	.07	.07	.07	.06	.18	.18	.14
.25-	.50-	.50-	.25-	.50-	.25-	.25-	.25-	.25-	.25-	.50-	.50-	.50-
.50	1.75	1.75	1.25	1.75	.75	.75	.75	.75	.75	1.75	1.75	1.50
4	14	3	4	6	9	8	9	9	11	9	3	8
.54	6.77	.68	2.33	5.66	.30	1.06	1.07	1.17	.87	15.57	.26	11.08
.35	9.81	.43	1.52	6.21	.34	.73	1.10	1.04	1.09	16.49	.40	12.27
.18	2.62	.25	.76	2.54	.11	.26	.37	.35	.33	5.50	.23	4.34
.08-	.16-	.26-	.16-	.32-	.08-	.24-	.14-	.10-	.10-	.37-	.01-	.46-
.89	25.37	1.11	3.68	17.24	1.14	2.19	3.69	3.17	3.64	43.28	.72	33.96
4	28	9	9	6	9	8	9	9	11	9	9	9
.34	1.92	2.57	.97	1.91	.11	.48	.64	1.41	.15	2.08	3.41	2.60
.36	1.80	1.80	.92	2.06	.09	.44	.61	1.40	.13	1.83	4.16	3.30
.18	.34	.60	.31	.84	.03	.16	.20	.47	.04	.61	1.39	1.10
.05-	.19-	.72-	.06-	.11-	.01-	.12-	.09-	.07-	.03-	.46-	.01-	.20-
.85	6.38	6.13	2.50	5.23	.23	1.28	1.78	3.62	.48	5.72	11.73	9.48
4	28	9	9	6	9	8	9	9	11	9	9	9
1.32	75.50	97.74	75.67	51.91	2.59	2.54	3.32	---	1.17	109.34	83.60	82.89
1.00	73.51	89.27	68.67	54.01	3.64	2.36	4.13	---	1.61	131.46	99.72	95.10
.50	13.89	29.76	22.89	22.05	1.21	.83	1.38	---	.49	43.82	33.24	31.70
.20-	.42-	1.19-	.70-	1.10-	.09-	.24-	.18-	---	.08-	.64-	.30-	1.45-
2.62	234.69	247.12	185.79	139.82	10.69	6.56	10.42	---	5.39	351.67	269.03	222.19
4	28	9	9	6	9	8	9	9	11	9	9	9
1.66	77.42	100.31	78.78	53.82	2.70	3.03	3.96	2.81	1.32	111.42	87.01	85.49
1.13	74.79	90.42	71.49	56.06	3.71	2.74	4.69	2.80	1.73	133.21	103.07	98.01
.56	14.13	30.14	23.83	22.89	1.24	.97	1.56	.93	.52	44.40	34.36	32.67
.25-	.84-	2.38-	.71-	1.21-	.16-	.36-	.36-	.14-	.16-	1.28-	.31-	1.85-
2.94	239.04	251.57	193.49	145.05	10.92	7.84	12.20	7.24	5.87	357.39	278.12	231.67

23 shrubs from northeastern Minnesota (allometric relation  $Y = aX^b$ )

<i>nicera</i>	<i>Populus</i>	<i>Prunus</i>	<i>Prunus</i>	<i>Rhamnus</i>	<i>Ribes</i>	<i>Rosa</i>	<i>Rosa</i>	<i>Rubus</i>	<i>Rubus</i>	<i>Salix</i>	<i>Sorbus</i>	<i>Viburnum</i>
<i>longifolia</i>	<i>tremuloides</i>	<i>pennsylvanica</i>	<i>virginiana</i>	<i>alnifolia</i>	spp.	<i>acicularis</i>	<i>blanda</i>	<i>parviflorus</i>	<i>strigosus</i>	spp.	<i>americana</i>	<i>rafinosquianum</i>
6.009	2.227	4.947	7.953	2.009	1.513	3.286	4.160	4.595	7.081	4.514	2.885	8.526
3.115	4.258	2.836	1.954	3.835	3.023	2.004	2.302	2.376	3.871	3.692	3.454	3.007
.749	.963	.000	.900	.881	.549	.879	.597	.720	.748	.894	.000	.951
.217	1.956	.601	3.522	2.394	.245	.275	.745	.586	.577	5.735	.560	2.925
NS	.005	NS	.005	.010	.025	.005	.025	.005	.005	.005	NS	.005
3.756	.894	2.095	1.516	1.040	.321	1.671	2.230	8.214	.491	.545	.841	1.096
3.116	2.501	1.037	2.605	2.885	1.719	2.312	2.078	3.283	1.498	3.690	4.023	4.362
.284	.430	.257	.837	.808	.604	.513	.568	.957	.332	.787	.717	.766
.373	1.382	1.654	1.317	1.011	.059	.333	.429	.312	.110	.902	2.366	1.707
NS	.005	NS	.005	.025	.025	.050	.025	.005	.100	.005	.005	.005
14.338	33.718	48.927	34.049	29.929	32.958	12.890	30.900	---	10.782	17.344	13.177	38.820
3.082	2.876	2.545	2.708	2.760	5.458	3.162	4.519	---	4.111	4.939	4.393	4.125
.563	.914	.950	.856	.874	.776	.908	.873	---	.745	.923	.917	.981
.806	22.024	21.366	27.863	21.452	1.843	.773	1.570	---	.857	39.009	30.800	14.172
NS	.005	.005	.005	.010	.005	.005	.005	---	.005	.005	.005	.005
18.093	34.502	49.916	35.575	30.971	32.001	14.527	31.182	---	11.519	17.815	13.982	39.921
3.089	2.874	2.547	2.704	2.764	5.256	3.042	4.074	---	4.032	4.919	4.900	4.132
.695	.914	.952	.861	.872	.780	.878	.844	---	.734	.923	.922	.981
.762	22.316	21.233	28.520	22.456	1.861	1.035	1.979	---	.938	39.551	30.738	14.504
NS	.005	.005	.005	.010	.005	.005	.005	---	.005	.005	.005	.005