

Utah State University

DigitalCommons@USU

Foundations of Wave Phenomena

Open Textbooks

8-2014

Problem Set 10

Charles G. Torre
charles.torre@usu.edu

Follow this and additional works at: https://digitalcommons.usu.edu/foundation_wave

 Part of the [Physics Commons](#)

To read user comments about this document and to leave your own comment, go to https://digitalcommons.usu.edu/foundation_wave/35

Recommended Citation

Torre, Charles G., "Problem Set 10" (2014). *Foundations of Wave Phenomena*. 35.
https://digitalcommons.usu.edu/foundation_wave/35

This Book is brought to you for free and open access by the Open Textbooks at DigitalCommons@USU. It has been accepted for inclusion in Foundations of Wave Phenomena by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.



PROBLEM SET 10**Problem 10.1**

Show that if ϕ is any solution of the sine-Gordon equation then $\tilde{\phi} = -\phi$ and $\tilde{\phi} = \phi + 2n\pi$, where n is an integer, are also solutions to the sine-Gordon equation. (These are *discrete symmetries* of the sine-Gordon equation.)

Problem 10.2

Verify the continuity equation (21.17) for the sine-Gordon equation.

Problem 10.3

Plot the soliton solution (21.12) and its energy density as a function of x , verifying that the soliton is a localized “lump” of energy. How do these graphs change as you vary λ and m ? (Here is a good place to use a computer.)

Problem 10.4

Verify the identities (21.14) and (21.15).

Problem 10.5

The results of this problem will be used in the next problem. Define

$$u = \frac{m}{2}(t - x), \quad v = \frac{m}{2}(t + x), \quad \tilde{\phi}(u, v) = \frac{\sqrt{\lambda}}{m}\phi(t(u, v), x(u, v)).$$

Show that the sine-Gordon equation can be written as

$$\frac{\partial^2 \tilde{\phi}}{\partial u \partial v} + \sin(\tilde{\phi}) = 0. \quad (21.30)$$

Problem 10.6

Using the notation of the previous problem, show that the quantities

$$\rho = -\frac{1}{2} \left(\frac{\partial^2 \tilde{\phi}}{\partial u^2} \right)^2 + \frac{1}{8} \left(\frac{\partial \tilde{\phi}}{\partial u} \right)^4,$$

and

$$j = \frac{\partial^2 \tilde{\phi}}{\partial u^2} \frac{\partial^2 \tilde{\phi}}{\partial u \partial v} \sin(\tilde{\phi}) + \frac{\partial^2 \tilde{u}}{\partial u^2} \sin(\tilde{\phi}) - \frac{1}{2} \left(\frac{\partial \tilde{\phi}}{\partial u} \right)^2 \cos(\tilde{\phi}).$$

satisfy a continuity equation

$$\frac{\partial \rho}{\partial v} + \frac{\partial j}{\partial u} = 0,$$

when $\tilde{\phi}$ satisfies (21.30). The resulting conservation law is but one of an infinite hierarchy of conserved quantities that are intimately related to the stability of the sine-Gordon solitons.

The End

Congratulations. You started off trying to understand trig functions and the harmonic oscillator equation. A relatively short time later you were contemplating non-linear wave equations, solitons, and infinitely many conservation laws. And we're just getting started...