

Math. J. Okayama Univ. **57** (2015), 85–98

## STEENROD-ČECH HOMOLOGY-COHOMOLOGY THEORIES ASSOCIATED WITH BIVARIANT FUNCTORS

KOHEI YOSHIDA

ABSTRACT. Let  $\mathbf{NG}_0$  denote the category of all pointed numerically generated spaces and continuous maps preserving base-points. In [SYH], we described a passage from bivariant functors  $\mathbf{NG}_0^{\text{op}} \times \mathbf{NG}_0 \rightarrow \mathbf{NG}_0$  to generalized homology and cohomology theories. In this paper, we construct a bivariant functor such that the associated cohomology is the Čech cohomology and the homology is the Steenrod homology (at least for compact metric spaces).

### 1. INTRODUCTION

According to [Du], a topological space  $X$  is said to be  $\Delta$ -generated if it has the final topology with respect to its singular simplexes. CW-complexes are typical examples of such  $\Delta$ -generated spaces. In [SYH], we showed that the category of  $\Delta$ -generated spaces is equivalent to the subcategory of the category  $\mathbf{Diff}$  of diffeological spaces consisting of those special type of objects which we call numerically generated spaces. Throughout this paper, we use term “numerically generated” instead of “ $\Delta$ -generated”. Let  $\mathbf{NG}_0$  be the category of pointed numerically generated spaces and pointed continuous maps. In [SYH], we showed that  $\mathbf{NG}_0$  is a symmetric monoidal closed category with respect to the smash product, and that every bilinear enriched functor  $F : \mathbf{NG}_0^{\text{op}} \times \mathbf{NG}_0 \rightarrow \mathbf{NG}_0$  gives rise to a pair of generalized homology and cohomology theories, denoted by  $h_\bullet(-, F)$  and  $h^\bullet(-, F)$  respectively, such that

$$h_n(X, F) \cong \pi_0 F(S^{n+k}, \Sigma^k X), \quad h^n(X, F) \cong \pi_0 F(\Sigma^k X, S^{n+k})$$

hold whenever  $k$  and  $n + k$  are non-negative.

As an example, consider the bilinear enriched functor  $F$  which assigns to  $(X, Y)$  the mapping space from  $X$  to the topological free abelian group  $AG(Y)$  generated by the points of  $Y$  modulo the relation  $* \sim 0$ . The Dold-Thom theorem says that if  $X$  is a CW-complex then the groups  $h_n(X, F)$  and  $h^n(X, F)$  are, respectively, isomorphic to the singular homology and cohomology groups of  $X$ . But this is not the case for general  $X$ ; there exists a space  $X$  such that  $h_n(X, F)$  (resp.  $h^n(X, F)$ ) is not isomorphic to the singular homology (resp. cohomology) group of  $X$ .

---

*Mathematics Subject Classification.* 55N20, 55N05, 55N07.

*Key words and phrases.* Čech cohomologies, Steenrod homologies, bivariant functors.

The aim of this paper is to construct a bilinear enriched functor such that for any space  $X$  the associated cohomology groups are isomorphic to the Čech cohomology groups of  $X$ . Interestingly, it turns out that the corresponding homology groups are isomorphic to the Steenrod homology groups for any compact metrizable space  $X$ . Thus we obtain a bibariant theory which ties together the Čech cohomology and the Steenrod homology theories.

Let  $\mathbf{NGC}_0$  be the full subcategory of  $\mathbf{NG}_0$  consisting of compact metric spaces. For given a linear enriched functor  $T : \mathbf{NG}_0 \rightarrow \mathbf{NG}_0$ , let

$$\check{F} : \mathbf{NG}_0^{\text{op}} \times \mathbf{NGC}_0 \rightarrow \mathbf{NG}_0$$

be a bifunctor which maps  $(X, Y)$  to the space  $\varinjlim_{\lambda} \text{map}_0(X_{\lambda}, \varprojlim_{\mu_i} T(Y_{\mu_i}^{\check{C}}))$ . Here  $\lambda$  runs through coverings of  $X$ , and  $X_{\lambda}$  is the Vietoris nerve corresponding to  $\lambda$  ([P]). The main results of the paper can be stated as follows.

**Theorem 1.1.** *The functor  $\check{F}$  is a bilinear enriched functor.*

**Theorem 1.2.** *Let  $X$  be a compact metrizable space. Then  $h_n(X, \check{F}) = H_n^{\text{st}}(X, \mathbb{S})$  is the Steenrod homology group with coefficients in the spectrum  $\mathbb{S} = \{T(S^k)\}$ .*

In particular, let  $T$  be the functor which assigns to every  $X$  the topological abelian group  $AG(X)$ , and let

$$\check{C} : \mathbf{NG}_0^{\text{op}} \times \mathbf{NGC}_0 \rightarrow \mathbf{NG}_0$$

be the corresponding bifunctor.

**Theorem 1.3.** *For any pointed space  $X$ ,  $h^n(X, \check{C})$  is the Čech cohomology group of  $X$ , and  $h_n(X, \check{C})$  is the Steenrod homology group of  $X$  if  $X$  is a compact metrizable space.*

Recall that the Steenrod homology group is related to the Čech homology group of  $X$  by the exact sequence

$$0 \longrightarrow \varprojlim_{\lambda_i}^1 \tilde{H}_{n+1}(X_{\lambda_i}^{\check{C}}) \longrightarrow H_n^{\text{st}}(X) \longrightarrow \tilde{H}_n(X) \longrightarrow 0.$$

According to [KKS], if  $X$  is a movable compactum then we have  $\varprojlim_{\lambda_i}^1 \tilde{H}_{n+1}(X_{\lambda_i}^{\check{C}}) = 0$ , and hence the following corollary follows.

**Corollary 1.4.** *Let  $X$  be a movable compactum. Then  $h_n(X, \check{C})$  is the Čech homology group of  $X$ .*

The paper is organized as follows. In Section 2 we recall from [SYH] the category  $\mathbf{NG}_0$  and the passage from bilinear enriched functors to generalized homology and cohomology theories. We also recall the definition of Čech

cohomology and Steenrod homology group, and Vietoris and Čech nerves; In Section 3 we prove Theorem 1.1; Finally, in Section 4 we prove Theorems 1.2 and 1.3.

## 2. PRELIMINARIES

**2.1. Homology and cohomology theories via bifunctors.** Let  $\mathbf{NG}_0$  be the category of pointed numerically generated topological spaces and pointed continuous maps. In [SYH] we showed that  $\mathbf{NG}_0$  satisfies the following properties:

- (1) It contains pointed CW-complexes;
- (2) It is complete and cocomplete;
- (3) It is monoidally closed in the sense that there is an internal hom  $Z^Y$  satisfying a natural bijection  $\text{hom}_{\mathbf{NG}_0}(X \wedge Y, Z) \cong \text{hom}_{\mathbf{NG}_0}(X, Z^Y)$ ;
- (4) There is a coreflector  $\nu: \text{Top}_0 \rightarrow \mathbf{NG}_0$  such that the coreflection arrow  $\nu X \rightarrow X$  is a weak equivalence;
- (5) The internal hom  $Z^Y$  is weakly equivalent to the space of pointed maps from  $Y$  to  $Z$  equipped with the compact-open topology.

Throughout the paper, we write  $\text{map}_0(Y, Z) = Z^Y$  for any  $Y, Z \in \mathbf{NG}_0$ .

A map  $f: X \rightarrow Y$  between topological spaces is said to be numerically continuous if the composite  $f \circ \sigma: \Delta^n \rightarrow Y$  is continuous for every singular simplex  $\sigma: \Delta^n \rightarrow X$ . We have the following.

**Proposition 2.1.** ([SYH]) *Let  $f: X \rightarrow Y$  be a map between numerically generated spaces. Then  $f$  is numerically continuous if and only if  $f$  is continuous.*

From now on, we assume that  $\mathbf{C}_0$  satisfies the following conditions: (i)  $\mathbf{C}_0$  contains all finite CW-complexes. (ii)  $\mathbf{C}_0$  is closed under finite wedge sum. (iii) If  $A \subset X$  is an inclusion of objects in  $\mathbf{C}_0$  then its cofiber  $X \cup CA$  belongs to  $\mathbf{C}_0$ ; in particular,  $\mathbf{C}_0$  is closed under the suspension functor  $X \mapsto \Sigma X$ .

**Definition 2.2.** Let  $\mathbf{C}_0$  be a full subcategory of  $\mathbf{NG}_0$ . A functor  $T: \mathbf{C}_0 \rightarrow \mathbf{NG}_0$  is called *enriched (or continuous)* if the map

$$T: \text{map}_0(X, X') \rightarrow \text{map}_0(T(X), T(X')),$$

which assigns  $T(f)$  to every  $f$ , is a pointed continuous map.

Note that if  $f$  is constant, then so is  $T(f)$ .

**Definition 2.3.** An enriched functor  $T$  is called *linear* if for any pair of a pointed space  $X$ , a sequence

$$T(A) \rightarrow T(X) \rightarrow T(X \cup CA)$$

induced by the cofibration sequence  $A \rightarrow X \rightarrow X \cup CA$ , is a homotopy fibration sequence.

**Example 2.4.** Let  $AG : CW_0 \rightarrow \mathbf{NG}_0$  be the functor which assigns to a pointed CW-complex  $(X, x_0)$  the topological abelian group  $AG(X)$  generated by the points of  $X$  modulo the relation  $x_0 \sim 0$ . Then  $AG$  is a linear enriched functor. (see [SYH])

**Theorem 2.5.** ([SYH, Th 6.4]) *A linear enriched functor  $T$  defines a generalized homology  $\{h_n(X, T)\}$  satisfying*

$$h_n(X, T) = \begin{cases} \pi_n T(X), & n \geq 0 \\ \pi_0 T(\Sigma^{-n} X), & n < 0. \end{cases}$$

Next we introduce the notion of a bilinear enriched functor, and describe a passage from a bilinear enriched functor to generalized cohomology and generalized homology theories. We assume that  $\mathbf{C}'_0$  satisfies the same conditions of  $\mathbf{C}_0$ .

**Definition 2.6.** Let  $\mathbf{C}_0$  and  $\mathbf{C}'_0$  be full subcategories of  $\mathbf{NG}_0$ . A bifunctor  $F : \mathbf{C}_0^{\text{op}} \times \mathbf{C}'_0 \rightarrow \mathbf{NG}_0$  is a function which

- (1) to each objects  $X \in \mathbf{C}_0$  and  $Y \in \mathbf{C}'_0$  assigns an object  $F(X, Y) \in \mathbf{NG}_0$ ;
- (2) to each  $f \in \text{map}_0(X, X')$ ,  $g \in \text{map}_0(Y, Y')$  assigns a continuous map  $F(f, g) \in \text{map}_0(F(X', Y), F(X, Y'))$ .

$F$  is required to satisfy the following equalities:

- (a)  $F(1_X, 1_Y) = 1_{F(X, Y)}$ ;
- (b)  $F(f, g) = F(1_X, g) \circ F(f, 1_Y) = F(f, 1_{Y'}) \circ F(1_{X'}, g)$ ;
- (c)  $F(f' \circ f, 1_Y) = F(f, 1_Y) \circ F(f', 1_Y)$ ,  $F(1_X, g' \circ g) = F(1_X, g') \circ F(1_X, g)$ .

**Definition 2.7.** A bifunctor  $F : \mathbf{C}_0^{\text{op}} \times \mathbf{C}_0 \rightarrow \mathbf{NG}_0$  is called *enriched* if the map

$$F : \text{map}_0(X, X') \times \text{map}_0(Y, Y') \rightarrow \text{map}_0(F(X', Y), F(X, Y')),$$

which assigns  $F(f, g)$  to every pair  $(f, g)$ , is a pointed continuous map.

Note that if either  $f$  or  $g$  is constant, then so is  $F(f, g)$ .

**Definition 2.8.** For any pairs of pointed spaces  $(X, A)$  and  $(Y, B)$ ,  $F$  is *bilinear* if the sequences

- (1)  $F(X \cup CA, Y) \rightarrow F(X, Y) \rightarrow F(A, Y)$
- (2)  $F(X, B) \rightarrow F(X, Y) \rightarrow F(X, Y \cup CB)$ ,

induced by the cofibration sequences  $A \rightarrow X \rightarrow X \cup CA$  and  $B \rightarrow Y \rightarrow Y \cup CB$ , are homotopy fibration sequences.

**Example 2.9.** Let  $T : \mathbf{NG}_0 \rightarrow \mathbf{NG}_0$  be a linear enriched functor, and let  $F(X, Y) = \text{map}_0(X, T(Y))$  for  $X, Y \in \mathbf{NG}_0$ . Then  $F : \mathbf{NG}_0^{\text{op}} \times \mathbf{NG}_0 \rightarrow \mathbf{NG}_0$  is a bilinear enriched functor.

**Theorem 2.10.** ([SYH, Th 7.4]) *A bilinear enriched functor  $F$  defines a generalized cohomology  $\{h^n(-, F)\}$  and a generalized homology  $\{h_n(-, F)\}$  such that*

$$h_n(Y, F) = \begin{cases} \pi_0 F(S^n, Y) & n \geq 0 \\ \pi_0 F(S^0, \Sigma^{-n}Y) & n < 0, \end{cases} \quad h^n(X, F) = \begin{cases} \pi_0 F(X, S^n) & n \geq 0 \\ \pi_{-n} F(X, S^0) & n < 0, \end{cases}$$

hold for any  $X \in \mathbf{C}_0$  and  $Y \in \mathbf{C}'_0$ .

**Proposition 2.11.** ([SYH]) *If  $X$  is a CW-complex, we have  $h_n(X, F) = H_n(X, \mathbb{S})$  and  $h^n(X, F) = H^n(X, \mathbb{S})$ , the generalized homology and cohomology groups with coefficients in the spectrum  $\mathbb{S} = \{F(S^0, S^n) \mid n \geq 0\}$ .*

**2.2. Čech cohomology and Steenrod homology groups.** We recall that the Čech cohomology group of  $X$  with coefficients group  $G$  is defined to be the colimit of the singular cohomology groups

$$\check{H}^n(X, G) = \varinjlim_{\lambda} H^n(X_{\lambda}^{\check{C}}, G),$$

where  $\lambda$  runs through coverings of  $X$  and  $X_{\lambda}^{\check{C}}$  is the Čech nerve corresponding to  $\lambda$ , i.e.  $v \in X_{\lambda}^{\check{C}}$  is a vertex of  $X_{\lambda}^{\check{C}}$  corresponding to an open set  $V \in \lambda$ . On the other hand, the Steenrod homology group of a compact metric space  $X$  is defined as follows. As  $X$  is a compact metric space, there is a sequence  $\{\lambda_i\}_{i \geq 0}$  of finite open covers of  $X$  such that  $\lambda_0 = \{X\}$ ,  $\lambda_i$  is a refinement of  $\lambda_{i-1}$ , and  $X$  is the inverse limit  $\varprojlim_i X_{\lambda_i}^{\check{C}}$ . According to [F], the Steenrod homology group of  $X$  with coefficients in the spectrum  $\mathbb{S}$  is defined to be the group

$$H_n^{st}(X, \mathbb{S}) = \pi_n \underline{\text{holim}}_{\lambda_i} (X_{\lambda_i}^{\check{C}} \wedge \mathbb{S})$$

where  $\underline{\text{holim}}$  denotes the homotopy inverse limit. (See also [KKS] for the definition without using subdivisions.)

**2.3. Vietoris and Čech nerves.** For each  $X \in \mathbf{NG}_0$ , let  $\lambda$  be an open covering of  $X$ . According to [P], the Vietoris nerve of  $\lambda$  is a simplicial set in which an  $n$ -simplex is an ordered  $(n+1)$ -tuple  $(x_0, x_1, \dots, x_n)$  of points contained in an open set  $U \in \lambda$ . Face and degeneracy operators are respectively given by

$$d_i(x_0, \dots, x_n) = (x_0, x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$$

and

$$s_i(x_0, x_1, \dots, x_n) = (x_0, x_1, \dots, x_{i-1}, x_i, x_i, x_{i+1}, \dots, x_n), \quad 0 \leq i \leq n.$$

We denote the realization of the Vietoris nerve of  $\lambda$  by  $X_{\lambda}$ . If  $\lambda$  is a refinement of  $\mu$ , then there is a canonical map  $\pi_{\mu}^{\lambda} : X_{\lambda} \rightarrow X_{\mu}$  induced by the identity map of  $X$ .

The relation between the Vietoris and the Čech nerves is given by the following Proposition due to Dowker.

**Proposition 2.12.** ([Do]) *The Čech nerve  $X_\lambda^{\check{C}}$  and the Vietoris nerve  $X_\lambda$  have the same homotopy type.*

According to [Do], for arbitrary topological space, the Vietoris and Čech homology groups are isomorphic and the Alexander-Spanier and Čech cohomology groups are isomorphic.

### 3. PROOF OF THEOREM 1.1

Let  $T$  be a linear enriched functor. We define a bifunctor  $\check{F} : \mathbf{NG}_0^{\text{op}} \times \mathbf{NGC}_0 \rightarrow \mathbf{NG}_0$  as follows. For  $X \in \mathbf{NG}_0$  and  $Y \in \mathbf{NGC}_0$ , we put

$$\check{F}(X, Y) = \lim_{\rightarrow \lambda} \text{map}_0(X_\lambda, \overleftarrow{\text{holim}}_{\mu_i} T(Y_{\mu_i}^{\check{C}})),$$

where  $\lambda$  is an open covering of  $X$  and  $\{\mu_i\}_{i \geq 0}$  is a set of finite open covers of  $Y$  such that  $\mu_0 = \{Y\}$ ,  $\mu_i$  is a refinement of  $\mu_{i-1}$ , and  $Y$  is the inverse limit  $\overleftarrow{\lim}_i Y_{\mu_i}^{\check{C}}$ .

Given based maps  $f : X \rightarrow X'$  and  $g : Y \rightarrow Y'$ , we define a map

$$\check{F}(f, g) \in \text{map}_0(\check{F}(X', Y), \check{F}(X, Y'))$$

as follows. Let  $\nu$  and  $\gamma$  be open covering of  $X'$  and  $Y'$  respectively, and let  $f^\#\nu = \{f^{-1}(U) \mid U \in \nu\}$  and  $g^\#\gamma = \{g^{-1}(V) \mid V \in \gamma\}$ . Then  $f^\#\nu$  and  $g^\#\gamma$  are open coverings of  $X$  and  $Y$  respectively. By the definition of the nerve, there are natural maps  $f_\nu : X_{f^\#\nu} \rightarrow X'_\nu$  and  $g_\gamma : Y_{g^\#\gamma}^{\check{C}} \rightarrow (Y')_\gamma^{\check{C}}$ . Hence we have the map

$$T(g_\gamma)^{f_\nu} : T(Y_{g^\#\gamma}^{\check{C}})^{X'_\nu} \rightarrow T((Y')_\gamma^{\check{C}})^{X_{f^\#\nu}}$$

induced by  $f_\nu$  and  $g_\gamma$ . Thus we can define

$$\check{F}(f, g) = \lim_{\rightarrow \nu} \overleftarrow{\text{holim}}_{\gamma} T(g_\gamma)^{f_\nu} : \check{F}(X', Y) \rightarrow \check{F}(X, Y').$$

**Theorem 1.1.** The functor  $\check{F}$  is a bilinear enriched functor.

First we prove that the sequence

$$\check{F}(X \cup CA, Z) \rightarrow \check{F}(X, Z) \rightarrow \check{F}(A, Z)$$

induced by the sequence  $A \rightarrow X \rightarrow X \cup CA$ , is a homotopy fibration sequence. Let  $\lambda$  be an open covering of  $X \cup CA$ , and let  $\lambda_X$ ,  $\lambda_{CA}$  and  $\lambda_A$  be the coverings of  $X$ ,  $CA$  and  $A$  consisting of those  $U \in \lambda$  such that  $U$  intersects with  $X$ ,  $CA$ , and  $A$ , respectively. We need the following lemma.

**Lemma 3.1.** *We have a homotopy equivalence*

$$(X \cup CA)_\lambda^{\check{C}} \simeq X_{\lambda_X}^{\check{C}} \cup C(A_{\lambda_A}^{\check{C}}).$$

*Proof.* By the definition of the Čech nerve, we have  $(X \cup CA)_\lambda^{\check{C}} = X_{\lambda_X}^{\check{C}} \cup (CA)_{\lambda_{CA}}^{\check{C}}$ . Since

$$X_{\lambda_X}^{\check{C}} \cup (CA)_{\lambda_{CA}}^{\check{C}} \simeq X_{\lambda_X}^{\check{C}} \cup A_{\lambda_A}^{\check{C}} \times I \cup (CA)_{\lambda_{CA}}^{\check{C}},$$

and since  $(CA)_{\lambda_{CA}}^{\check{C}} \simeq *$ , we have

$$X_{\lambda_X}^{\check{C}} \cup (CA)_{\lambda_{CA}}^{\check{C}} \simeq X_{\lambda_X}^{\check{C}} \cup C(A_{\lambda_A}^{\check{C}}).$$

Hence we have  $(X \cup CA)_\lambda \simeq X_{\lambda_X}^{\check{C}} \cup C(A_{\lambda_A}^{\check{C}})$ .  $\square$

By Proposition 2.12 and Lemma 3.1, the sequence

$$A_{\lambda_A} \rightarrow X_{\lambda_X} \rightarrow (X \cup CA)_\lambda$$

is a homotopy cofibration sequence. Hence the sequence

$$[(X \cup CA)_\lambda, Z] \rightarrow [X_{\lambda_X}, Z] \rightarrow [A_{\lambda_A}, Z]$$

is an exact sequence for any  $\lambda$ . Since the nerves of the form  $\lambda_X$  (resp.  $\lambda_A$ ) are cofinal in the set of nerves of  $X$  (resp.  $A$ ), we conclude that the sequence

$$\check{F}(X \cup CA, Z) \rightarrow \check{F}(X, Z) \rightarrow \check{F}(A, Z)$$

is a homotopy fibration sequence.

Now we show that the sequence  $\check{F}(Z, A) \rightarrow \check{F}(Z, X) \rightarrow \check{F}(Z, X \cup CA)$  is a homotopy fibration sequence. By the linearity of  $T$ , the sequence

$$T(A_{\lambda_A}^{\check{C}}) \rightarrow T(X_{\lambda_X}^{\check{C}}) \rightarrow T((X \cup CA)_\lambda^{\check{C}})$$

is a homotopy fibration sequence. Since the fibre  $T(A_{\lambda_A}^{\check{C}})$  is homeomorphic to the inverse limit

$$\varprojlim (* \rightarrow T((X \cup CA)_\lambda^{\check{C}}) \leftarrow T(X_{\lambda_X}^{\check{C}})),$$

we have

$$\begin{aligned} & \varprojlim (* \rightarrow \varprojlim_\lambda T((X \cup CA)_\lambda^{\check{C}}) \leftarrow \varprojlim_{\lambda_X} T(X_{\lambda_X}^{\check{C}})) \\ & \simeq \varprojlim \varprojlim_\lambda (* \rightarrow T((X \cup CA)_\lambda^{\check{C}}) \leftarrow T(X_{\lambda_X}^{\check{C}})) \\ & \simeq \varprojlim_\lambda \varprojlim (* \rightarrow T((X \cup CA)_\lambda^{\check{C}}) \leftarrow T(X_{\lambda_X}^{\check{C}})) \\ & \simeq \varprojlim_\lambda T(A_{\lambda_A}^{\check{C}}). \end{aligned}$$

This implies that the sequence

$$\varprojlim_{\lambda_A} T(A_{\lambda_A}^{\check{C}}) \rightarrow \varprojlim_{\lambda_X} T(X_{\lambda_X}^{\check{C}}) \rightarrow \varprojlim_\lambda T((X \cup CA)_\lambda^{\check{C}})$$

is a homotopy fibration sequence, hence so is  $\check{F}(Z, A) \rightarrow \check{F}(Z, X) \rightarrow \check{F}(Z, X \cup CA)$ .

Next we prove the continuity of  $\check{F}$ . Let  $F(X, Y) = \text{map}_0(X, \varprojlim_{\mu_i} T(Y_{\mu_i}^{\check{C}}))$ , so that we have  $\check{F}(X, Y) = \varinjlim_{\lambda} F(X_\lambda, Y)$ . We need the following lemma.

**Lemma 3.2.** *The functor  $F$  is an enriched bifunctor.*

*Proof.* Let  $F_1(Y) = \varprojlim_{\mu_i} T(Y_{\mu_i}^{\check{C}})$  and  $F_2(X, Z) = \text{map}_0(X, Z)$ , so that we have  $F(X, Y) = F_2(X, F_1(Y))$ . Clearly  $F_2$  is continuous.

Let  $G_1$  be the functor which maps  $Y$  to  $\varprojlim_{\mu_i} Y_{\mu_i}^{\check{C}}$ . Since  $T$  is enriched,  $F_1$  is continuous if so is  $G_1$ . It suffices to show that the map  $G'_1: \text{map}_0(Y, Y') \times \varprojlim_{\mu_i} Y_{\mu_i}^{\check{C}} \rightarrow \varprojlim_{\lambda_j} (Y')_{\lambda_j}^{\check{C}}$ , adjoint to  $G_1$ , is continuous for any  $Y$  and  $Y'$ . Given an open covering  $\lambda$  of  $Y'$ , let  $p_\lambda^n$  be the natural map  $\varprojlim_{\lambda} (Y')_{\lambda}^{\check{C}} \rightarrow \text{map}_0(\Delta^n, (Y')_{\lambda}^{\check{C}})$ . Then  $G'_1$  is continuous if so is the composite

$$p_\lambda^n \circ G'_1: \text{map}_0(Y, Y') \times \varprojlim_{\mu_i} Y_{\mu_i}^{\check{C}} \rightarrow \text{map}_0(\Delta^n, (Y')_{\lambda}^{\check{C}})$$

for every  $\lambda \in \text{Cov}(Y')$  and every  $n$ . Here we may assume by [SYH, Proposition 4.3] that  $\text{map}_0(\Delta^n, (Y')_{\lambda}^{\check{C}})$  is equipped with the compact open topology. Let  $(g, \alpha) \in \text{map}_0(Y, Y') \times \varprojlim_{\mu_i} Y_{\mu_i}^{\check{C}}$ , and let  $W_{K,U} \subset \text{map}_0(\Delta^n, (Y')_{\lambda}^{\check{C}})$  be an open neighborhood of  $p_\lambda^n(G'_1(g, \alpha))$ , where  $K$  is a compact set of  $\Delta^n$  and  $U$  is an open set of  $(Y')_{\lambda}^{\check{C}}$ .

Let us choose simplices  $\sigma$  of  $Y_{g^\# \lambda}^{\check{C}}$  with vertices  $g^{-1}(U(\sigma, k))$ , where  $U(\sigma, k) \in \lambda$  for  $0 \leq k \leq \dim \sigma$ . Let

$$O(\sigma) = \bigcap_{0 \leq k \leq \dim \sigma} U(\sigma, k) \subset Y'.$$

Let us choose a point  $y_\sigma \in \bigcap_{0 \leq k \leq \dim \sigma} g^{-1}(U(\sigma, k))$ , then  $g(y_\sigma) \in O(\sigma)$ . Let  $W_1$  be the intersection of all  $\overline{W}_{y_\sigma, O(\sigma)}$ .

There is an integer  $l$  such that

$$\mu_l > \overline{\mu}_l > g^\# \lambda$$

where  $\overline{\mu}_l$  is a closed covering  $\{\overline{V} | V \in \mu_l\}$  of  $Y$ . Thus for any  $U \in \mu_l$ , there is an open set  $V_U \in g^\# \lambda$  such that  $\overline{U} \subset g^{-1}(V_U)$ . Since  $Y$  is a compact set,  $\overline{U}$  is compact. Let  $W_2$  be the intersection of  $W_{\overline{U}, V_U}$ , and let  $W = W_1 \cap W_2$ .

Since  $\mu_l > g^\# \lambda$ , we have

$$p_\lambda^n(G'_1(g, \alpha)) = (g_\lambda)_*(\pi_{g^\# \lambda}^{\mu_l})_* p_{\mu_l}^n \alpha.$$

where  $(g_\lambda)_*$  and  $(\pi_{g^\# \lambda}^{\mu_l})_*$  are induced by  $g_\lambda: Y_{g^\# \lambda}^{\check{C}} \rightarrow (Y')_{\lambda}^{\check{C}}$  and  $\pi_{g^\# \lambda}^{\mu_l}: Y_{\mu_l}^{\check{C}} \rightarrow Y_{g^\# \lambda}^{\check{C}}$ , respectively. Let

$$W' = (p_{\mu_l}^n)^{-1}(W_{K, (\pi_{g^\# \lambda}^{\mu_l})^{-1}(g_\lambda)^{-1}(U)}).$$



Then  $W \times W'$  is a neighborhood of  $(g, \alpha)$  in  $\text{map}_0(Y, Y') \times \varprojlim_{\mu_i} Y_{\mu_i}$ . To see that  $p_\lambda \circ G'_1$  is continuous at  $(g, \alpha)$ , we need only show that  $W \times W'$  is contained in  $(p_\lambda \circ G'_1)^{-1}(U)$ . Suppose  $(h, \beta)$  belongs to  $W \times W'$ . Since  $W$  is contained in  $W_1$ , we have

$$y_\sigma \in h^{-1}(O(\sigma)) \subset \bigcap_{0 \leq k \leq \dim \sigma} h^{-1}(U(\sigma, k)).$$

This means that the vertices  $h^{-1}(U(\sigma, k)) \in h^\# \lambda$ ,  $0 \leq k \leq \dim \sigma$ , determine simplices  $\sigma'$  of  $Y_{h^\# \lambda}$  each corresponding to each  $\sigma \subset Y_{g^\# \lambda}$ . Thus we have an isomorphism

$$s : Y_{h^\# \lambda}^{\check{C}} \rightarrow Y_{g^\# \lambda}^{\check{C}},$$

$$h^{-1}(U(\sigma, k)) \mapsto g^{-1}(U(\sigma, k)).$$

Moreover since  $W$  is contained in  $W_2$ , we have  $\bar{\mu}_l > h^\# \lambda$ .

Since the commutative diagram

$$\begin{array}{ccccc} Y_{\mu_l}^{\check{C}} & \longrightarrow & Y_{g^\# \lambda}^{\check{C}} & \xrightarrow{g_\lambda} & (Y')_{\lambda}^{\check{C}} \\ & \searrow & \uparrow s & \nearrow h_\lambda & \\ & & Y_{h^\# \lambda}^{\check{C}} & & \end{array}$$

is commutative, we have the equation

$$p_\lambda^n \circ G'_1(h, \beta)(K) = h_\lambda \pi_{h^\# \lambda}^{\mu_l}(\beta)(K) = g_\lambda \pi_{g^\# \lambda}^{\mu_l}(\beta)(K)$$

Since  $g_\lambda \pi_{g^\# \lambda}^{\mu_l}(\beta)(K)$  is contained in  $U$ , so is  $p_\lambda^n \circ G'_1(h, \beta)(K)$ .

Thus  $p_\lambda^n \circ G'_1$  is continuous for all  $\lambda \in \text{Cov}(Y')$ , and hence so is

$$G'_1 : \text{map}_0(Y, Y') \times \varprojlim_{\mu_i} Y_{\mu_i}^{\check{C}} \rightarrow \varprojlim_{\lambda_j} (Y')_{\lambda_j}^{\check{C}}.$$

□

We are now ready to prove Theorem 1.1. For given pointed spaces  $X$ ,  $Y$  and a covering  $\mu$  of  $X$ , let  $i_\mu$  denote the natural map  $F(X_\mu, Y) \rightarrow \varinjlim_{\mu} F(X_\mu, Y)$ . To prove the theorem, it suffices to show that the map

$$\begin{aligned} \check{F}' \circ (1 \times i_\lambda) : \text{map}_0(X, X') \times F(X'_\lambda, Y) &\rightarrow \text{map}_0(X, X') \times \varinjlim_{\lambda} F(X'_\lambda, Y) \\ &\rightarrow \varinjlim_{\mu} F(X_\mu, Y) \end{aligned}$$

which maps  $(f, \alpha)$  to  $i_{f^\# \lambda}(F(f_\lambda, 1_Y)(\alpha))$ , is continuous for every covering  $\lambda$  of  $X$ .

Let

$$R_\lambda : \text{map}_0(X, X') \rightarrow \varinjlim_{\mu} \text{map}_0(X_\mu, X'_\lambda)$$

be the map which assigns to  $f : X \rightarrow X'$  the image of  $\text{map}_0(X, X')$ ,  $f_\lambda \in \text{map}_0(X_{f^\#\lambda}, X'_\lambda)$  in  $\varinjlim_{\mu} \text{map}_0(X_\mu, X'_\lambda)$ , and let  $Q_\lambda$  be the map

$$\begin{aligned} \varinjlim_{\mu} \text{map}_0(X_\mu, X'_\lambda) \times F(X'_\lambda, Y) &\rightarrow \varinjlim_{\mu} F(X_\mu, Y), \\ [f, \alpha] &\mapsto i_{f^\#\lambda} f_\lambda \circ \alpha = i_{f^\#\lambda}(F(f_\lambda, 1_Y)(\alpha)). \end{aligned}$$

Since we have  $\check{F}' \circ (1 \times i_\lambda) = Q_\lambda \circ (R_\lambda \times 1)$ , we need only show the continuity of  $Q_\lambda$  and  $R_\lambda$ . Since  $Q_\lambda$  is induced by the maps  $\text{map}_0(X_\mu, X'_\lambda) \times F(X'_\lambda, Y) \rightarrow F(X_\mu, Y)$ ,  $Q_\lambda$  is continuous.

To see that  $R_\lambda$  is continuous, let  $W_{K^f, U}$  be a neighborhood of  $f_\lambda$  in  $\text{map}_0(X_{f^\#\lambda}, X'_\lambda)$ , where  $K^f$  is a compact subset of  $X_{f^\#\lambda}$  and  $U$  is an open subset of  $X'_\lambda$ . Since  $K^f$  is compact, there is a finite subcomplex  $S^f$  of  $X_{f^\#\lambda}$  such that  $K^f \subset S^f$ . Let  $\tau_i^f$ ,  $0 \leq i \leq m$ , be simplexes of  $S^f$ . By taking a suitable subdivision of  $X_{f^\#\lambda}$ , we may assume that there is a simplicial neighborhood  $N_{\tau_i^f}$  of each  $\tau_i^f$ ,  $1 \leq i \leq m$ , such that  $K^f \subset S^f \subset \cup_i N_{\tau_i^f} \subset f_\lambda^{-1}(U)$ .

Let  $\{x_k^i\}$  be the set of vertices of  $\tau_i^f$  and let  $W$  be the intersection of all  $W_{\{x_k^i\}, U_{(\tau_i^f)'}}$  where  $U_{(\tau_i^f)'}$  is an open set of  $X'_\lambda$  containing the set  $\{f(x_k^i)\}$ . Then  $W$  is a neighborhood of  $f$ . We need only show that  $R_\lambda(W) \subset i_{f^\#\lambda}(W_{K^f, U})$ . Suppose that  $g$  belongs to  $W$ . Since  $\{x_k^i\}$  is contained in  $g^{-1}(U_{(\tau_i^f)'})$  for any  $i$ , a simplex  $\tau_i^g$  spanned by the vertices is contained in  $X_{g^\#\lambda}$ . Let  $S^g$  be the finite subcomplex of  $X_{g^\#\lambda}$  consists of simplexes  $\tau_i^g$ . By the construction,  $S^f$  and  $S^g$  are isomorphic. Moreover there is a compact subset  $K^g$  of  $X_{g^\#\lambda}$  such that  $K^g$  and  $K^f$  are homeomorphic. On the other hand, since  $g(\{x_k^i\}) \subset U_{(\tau_i^f)'}$ , there is a simplex of  $X'_\lambda$  having  $g_\lambda(\tau_i^g)$  and  $(\tau_i^f)'$  as its faces. This means that  $g_\lambda(\tau_i^g) \subset f_\lambda(\cup_i N_{\tau_i^f})$ . Thus we have  $g_\lambda(K^g) = \cup_i g_\lambda(\tau_i^g) \subset f_\lambda(\cup_i N_{\tau_i^f})$ .

Let  $f^\#\lambda \cap g^\#\lambda$  be an open covering

$$\{f^{-1}(U) \cap g^{-1}(V) \mid U, V \in \lambda\}$$

of  $X$ . We regard  $X_{f^\#\lambda}$  and  $X_{g^\#\lambda}$  as a subcomplex of  $X_{f^\#\lambda \cap g^\#\lambda}$ . Since  $g_\lambda|_{X_{f^\#\lambda \cap g^\#\lambda}}$  is contiguous to  $f_\lambda|_{X_{f^\#\lambda \cap g^\#\lambda}}$ , we have a homotopy equivalence  $g_\lambda|_{X_{f^\#\lambda \cap g^\#\lambda}} \simeq f_\lambda|_{X_{f^\#\lambda \cap g^\#\lambda}}$ . By the homotopy extension property of  $g_\lambda|_{X_{f^\#\lambda \cap g^\#\lambda}} : X_{f^\#\lambda \cap g^\#\lambda} \rightarrow X'_\lambda$  and  $f_\lambda : X_{f^\#\lambda} \rightarrow X'_\lambda$ ,  $g_\lambda|_{X_{f^\#\lambda \cap g^\#\lambda}}$  extends to map  $G : X_{f^\#\lambda} \rightarrow X'_\lambda$ .

We have the relation  $G \sim \pi_{f^\#\lambda}^{f^\#\lambda \cap g^\#\lambda} G = g_\lambda|_{X_{f^\#\lambda \cap g^\#\lambda}} = \pi_{g^\#\lambda}^{f^\#\lambda \cap g^\#\lambda} g_\lambda \sim g_\lambda$ , where  $\sim$  is the relation of the direct limit. Moreover by  $G(K^f) \subset$

$f_\lambda(\cup_i N_{\tau_i^f}) \subset U$ , we have  $[g_\lambda] = [G] \in i_{f\#\lambda}(W_{K^f, U})$ . Hence  $R_\lambda$  is continuous, and so is  $\check{F}'$ .

#### 4. PROOFS OF THEOREMS 1.2 AND 1.3

To prove Theorems 1.2 and 1.3, we need several lemmas.

**Lemma 4.1.** *There exists a sequence  $\lambda_1^n < \lambda_2^n < \dots < \lambda_m^n < \dots$  of open coverings of  $S^n$  such that :*

- (1) *For each open covering  $\mu$  of  $S^n$ , there is an  $m \in \mathbb{N}$  such that  $\lambda_m^n$  is a refinement of  $\mu$ :*
- (2) *For any  $m$ ,  $S_{\lambda_m^n}^n$  is homotopy equivalent to  $S^n$ .*

*Proof.* We prove by induction on  $n$ . For  $n = 1$ , we define an open covering  $\lambda_m^1$  of  $S^1$  as follows. For any  $i$  with  $0 \leq i < 4m$ , we put

$$U(i, m) = \left\{ (\cos \theta, \sin \theta) \mid \frac{(4i-3)\pi}{8m} < \theta < \frac{(4i+5)\pi}{8m} \right\}.$$

Let  $\lambda_m^1 = \{U(i, m) \mid 0 \leq i < 4m\}$ . Then the set  $\lambda_m^1$  is an open covering of  $S^1$  and is a refinement of  $\lambda_{m-1}^1$ . Clearly  $(S^1)_{\lambda_m^1}^{\check{C}}$  is homeomorphic to  $S^1$ , hence  $S_{\lambda_m^1}^1$  is homotopy equivalent to  $S^1$ . Moreover for any open covering  $\mu$  of  $S^1$ , there exists an  $m$  such that  $\lambda_m^1$  is a refinement of  $\mu$ . Hence the lemma is true for  $n = 1$ . Assume now that the lemma is true for  $1 \leq k \leq n-1$ . Let  $\lambda_m^n$  be the open covering  $\lambda_m^{n-1} \times \lambda_m^1$  of  $S^{n-1} \times S^1$  and let  $\lambda_m^n$  be the open covering of  $S^n$  induced by the natural map  $p : S^{n-1} \times S^1 \rightarrow S^{n-1} \times S^1 / S^{n-1} \vee S^1$ . Since  $S_{\lambda_m^{n-1}}^{n-1}$  is a homotopy equivalence of  $S^{n-1}$ , we have

$$S_{\lambda_m^n}^n \approx (S^{n-1} \times S^1 / S^{n-1} \vee S^1)_{\lambda_m^n} \approx (S_{\lambda_m^{n-1}}^{n-1} \times S_{\lambda_m^1}^1) / (S_{\lambda_m^{n-1}}^{n-1} \vee S_{\lambda_m^1}^1) \approx S^n.$$

Thus the sequence  $\lambda_1^n < \lambda_2^n < \dots < \lambda_m^n < \dots$  satisfies the required conditions.  $\square$

**Lemma 4.2.**  $h_n(X, \check{F}) \cong \pi_n \mathop{\text{holim}}_{\leftarrow \mu} T(X_\mu^{\check{C}})$  for  $n \geq 0$ .

*Proof.* By Lemma 4.1, we have an isomorphism

$$\lim_{\rightarrow \lambda} [S_\lambda^n, \mathop{\text{holim}}_{\leftarrow \mu} T(X_\mu^{\check{C}})] \cong [S^n, \mathop{\text{holim}}_{\leftarrow \mu} T(X_\mu^{\check{C}})].$$

Thus we have

$$\begin{aligned}
h_n(X, \check{F}) &= \pi_0 \check{F}(S^n, X) \\
&= \pi_0 \varinjlim_\lambda \text{map}_0(S_\lambda^n, \varprojlim_\mu T(X_\mu^{\check{C}})) \\
&\cong \varinjlim_\lambda [S^0, \text{map}_0(S_\lambda^n, \varprojlim_\mu T(X_\mu^{\check{C}}))] \\
&\cong \varinjlim_\lambda [S_\lambda^n, \varprojlim_\mu T(X_\mu^{\check{C}})] \\
&\cong [S^n, \varprojlim_\mu T(X_\mu^{\check{C}})] \\
&\cong \pi_n \varprojlim_\mu T(X_\mu^{\check{C}}).
\end{aligned}$$

□

Now we are ready to prove Theorem 1.2. Let  $X$  be a compact metric space and let  $\mathbb{S} = \{T(S^k) \mid k \geq 0\}$ . Since  $X$  is a compact metric space, there is a sequence  $\{\mu_i\}_{i \geq 0}$  of finite open covers of  $X$  with  $\mu_0 = X$  and  $\mu_i$  refining  $\mu_{i-1}$  such that  $X = \varprojlim_i X_{\mu_i}^{\check{C}}$  holds. Let us denote  $X_{\mu_i}^{\check{C}} = X_i^{\check{C}}$  and  $X_{\mu_i} = X_i$  if there is no possibility of confusion. According to [F], there is a short exact sequence

$$0 \longrightarrow \varprojlim_i^1 H_{n+1}(X_i^{\check{C}}, \mathbb{S}) \longrightarrow H_n^{st}(X, \mathbb{S}) \longrightarrow \varprojlim_i H_n(X_i^{\check{C}}, \mathbb{S}) \longrightarrow 0$$

where  $H_n(X, \mathbb{S})$  is the homology group of  $X$  with coefficients in the spectrum  $\mathbb{S}$ . (This is a special case of the Milnor exact sequence [MI].) On the other hand, by [BK], we have the following.

**Lemma 4.3.** ([BK]) *There is a natural short exact sequence*

$$0 \longrightarrow \varprojlim_i^1 \pi_{n+1} T(X_i^{\check{C}}) \longrightarrow \pi_n \varprojlim_i T(X_i) \longrightarrow \varprojlim_i \pi_n T(X_i^{\check{C}}) \longrightarrow 0.$$

By Proposition 2.11, we have a diagram

$$\begin{array}{ccccccc}
(4.1) & 0 & \longrightarrow & \varprojlim_i^1 H_{n+1}(X_i^{\check{C}}, \mathbb{S}) & \longrightarrow & H_n^{st}(X, \mathbb{S}) & \longrightarrow & \varprojlim_i H_n(X_i^{\check{C}}, \mathbb{S}) & \longrightarrow & 0 \\
& & & \downarrow \cong & & & & \downarrow \cong & & \\
& 0 & \longrightarrow & \varprojlim_i^1 \pi_{n+1}(T(X_i^{\check{C}})) & \longrightarrow & \pi_n(\varprojlim_i T(X_i^{\check{C}})) & \longrightarrow & \varprojlim_i \pi_n(T(X_i^{\check{C}})) & \longrightarrow & 0.
\end{array}$$

Hence it suffices to construct a natural homomorphism

$$H_n^{st}(X, \mathbb{S}) \rightarrow \pi_n(\varprojlim_i T(X_i^{\check{C}}))$$

making the diagram (4.1) commutative.

Since  $T$  is continuous, the identity map  $X \wedge S^k \rightarrow X \wedge S^k$  induces a continuous map  $i' : X \wedge T(S^k) \rightarrow T(X \wedge S^k)$ . Hence we have the composite homomorphism

$$\begin{aligned} H_n^{st}(X, \mathbb{S}) &= \pi_n \mathop{\text{holim}}\limits_i (X_i^{\check{C}} \wedge \mathbb{S}) \\ &\cong \varinjlim_k \pi_{n+k} (\mathop{\text{holim}}\limits_i (X_i^{\check{C}} \wedge T(S^k))) \\ &\xrightarrow{I} \varinjlim_k \pi_{n+k} (\mathop{\text{holim}}\limits_i T(X_i^{\check{C}} \wedge S^k)) \\ &\cong \pi_n (\mathop{\text{holim}}\limits_i T(X_i^{\check{C}})) \end{aligned}$$

in which  $I = \varinjlim_k i'_*$  is induced by the homomorphisms

$$i'_* : \pi_{n+k} (\mathop{\text{holim}}\limits_i (X_i^{\check{C}} \wedge T(S^k))) \rightarrow \pi_{n+k} (\mathop{\text{holim}}\limits_i T(X_i^{\check{C}} \wedge S^k)).$$

Clearly resulting the homomorphism  $H_n^{st}(X, \mathbb{S}) \rightarrow \pi_n (\mathop{\text{holim}}\limits_i T(X_i^{\check{C}}))$  makes the diagram (4.1) commutative. Thus  $h_n(X, \check{F})$  is isomorphic to the Steenrod homology group coefficients in the spectrum  $\mathbb{S}$ .

Finally, to prove Theorem 1.3 it suffices to show that  $h^n(X, \check{C})$  is isomorphic to the Čech cohomology group of  $X$ .

By Lemma 4.1, we have a homotopy commutative diagram

$$\begin{array}{ccccccc} \cdots & \xrightarrow{=} & AG(S^n) & \xrightarrow{=} & AG(S^n) & \xrightarrow{=} & \cdots \\ & & \downarrow \simeq & & \downarrow \simeq & & \\ \cdots & \longrightarrow & AG(S_{\lambda_{m-1}}^n) & \xrightarrow{\simeq} & AG(S_{\lambda_m}^n) & \longrightarrow & \cdots \end{array}$$

Hence we have  $AG(S^n) \simeq \mathop{\text{holim}}\limits_i AG(S_{\lambda_i}^n)$ .

Thus we have

$$\begin{aligned} h^n(X, \check{C}) &= \pi_0 \check{C}(X, S^n) \\ &= \pi_0 \varinjlim_{\lambda} \text{map}_0(X_{\lambda}, \mathop{\text{holim}}\limits_{\mu} AG((S^n)_{\mu}^{\check{C}})) \\ &\cong [S^0, \varinjlim_{\lambda} \text{map}_0(X_{\lambda}, AG(S^n))] \\ &\cong \varinjlim_{\lambda} [S^0, \text{map}_0(X_{\lambda}, AG(S^n))] \\ &\cong \varinjlim_{\lambda} [S^0 \wedge X_{\lambda}, AG(S^n)] \\ &\cong \varinjlim_{\lambda} [X_{\lambda}, AG(S^n)]. \end{aligned}$$

Hence  $h^n(X, \check{C})$  is isomorphic to the Čech cohomology group of  $X$ .

#### ACKNOWLEDGEMENT

The author expresses hearty thanks to Professor K. Shimakawa for calling his attention to the subject and for many helpful suggestions through the preparation of manuscript during the preparation period for manuscript. We also thank Claude Schochet for pointing their work [KKS].

## REFERENCES

- [BK] A.K.Bousfield and D.M.Kan, *Homotopy limits, completions and localization*, Springer Lecture Notes in Math. 304 Springer-Verlag, Berlin, 1972.
- [Do] C. Dowker, *Homology Groups of Relations*, Ann. of Math., 56, No. 1, 1952, 84-95.
- [Du] D.Dugger, *Notes on Delta-Generated Spaces*, available at <http://www.uoregon.edu/ddugger/delta.html>.
- [F] S.C. Ferry, *Remarks on Steenrod homology*, in: Proc. 1993 Oberwolfach Conf. on the Novikov Conjectures, Index Theorems and Rigidity, Vol. 2, London Math. Soc. Lecture Notes Ser., Vol. 227, Cambridge, UK, 1995, 148-166.
- [KKS] Daniel S. Kahn, Jerome Kaminker, and Claude Schochet *Generalized homology theories on compact metric spaces*, Michigan Math. J., 24, 1977, 203-224.
- [MI] J. Milnor, *On the Steenrod homology theory*, in: Proc. 1993 Oberwolfach Conf. on the Novikov Conjectures, Index Theorems and Rigidity, Vol. 1, London Math. Soc. Lecture Notes Ser., Vol. 226, Cambridge, UK, 1995, 79-96.
- [P] T.Porter, *Čech Homotopy I*, J.London Math. Soc. (2), 6, 1973, 429-436
- [SYH] Shimakawa, K., Yoshida, K., and Haraguchi, T *Homology and cohomology associated with a bivariant functor*. arXiv:1010.3336

KYOTO RAKUHOKU HIGH SCHOOL  
59 SHIMOGAMO UMENOKICHO, SAKYO-KU, KYOTO 606-0851, JAPAN  
*e-mail address:* sc09121@s.okadai.jp