

RESEARCH ARTICLE

Open Access



Health-related quality of life after laparoscopic repair of giant paraesophageal hernia: how does recurrence in CT scan compare to clinical success?

Henriikka Hietaniemi^{1,2*} , Ilkka Ilonen^{1,2}, Tommi Järvinen^{1,2}, Juha Kauppi^{1,2}, Saana Andersson^{1,2}, Harri Sintonen³ and Jari Räsänen^{1,2}

Abstract

Background: Computed tomography (CT) is widely used in the diagnosis of giant paraesophageal hernias (GPEH) but has not been utilised systematically for follow-up. We performed a cross-sectional observational study to assess mid-term outcomes of elective laparoscopic GPEH repair. The primary objective of the study was to evaluate the radiological hernia recurrence rate by CT and to determine its association with current symptoms and quality of life.

Methods: All non-emergent laparoscopic GPEH repairs between 2010 to 2015 were identified from hospital medical records. Each patient was offered non-contrast CT and sent questionnaires for disease-specific symptoms and health-related quality of life.

Results: The inclusion criteria were met by 165 patients (74% female, mean age 67 years). Total recurrence rate was 29.3%. Major recurrent hernia (> 5 cm) was revealed by CT in 4 patients (4.3%). Radiological findings did not correlate with symptom-related quality of life. Perioperative mortality occurred in 1 patient (0.6%). Complications were reported in 27 patients (16.4%).

Conclusions: Successful laparoscopic repair of GPEH requires both expertise and experience. It appears to lead to effective symptom relief with high patient satisfaction. However, small radiological recurrences are common but do not affect postoperative symptom-related patient wellbeing.

Keywords: Paraesophageal hernia, Laparoscopy, Computerized tomography, Quality of life

Background

The term giant paraesophageal hernia (GPEH) is used when at least one third of the stomach is situated above the diaphragm. GPEHs constitute approximately 2 to 5% of all paraesophageal hernias [1]. Symptoms caused by GPEH include chest and stomach pain, reflux, heartburn,

regurgitation, vomiting, bloating, and shortness of breath. Symptoms can vary from mild or temporary to acute and severe. Emergency surgery is necessary in cases of incarceration, volvulus, and perforation of the stomach [2, 3]. In rare cases, a GPEH can be asymptomatic and discovered incidentally in chest X-rays or computed tomography (CT) scans [4].

The risk of a major medical emergency associated with GPEH is approximately 1% annually and leads to a lifetime risk of approximately 18% at 65 years [5]. The mortality rate after hospitalization for symptomatic GPEH

* Correspondence: henriikka.hietaniemi@helsinki.fi

¹Department of General Thoracic and Esophageal Surgery, Helsinki University Hospital, Haartmaninkatu 4, 00290 Helsinki, Finland

²Department of Surgery, Cliniicum, University of Helsinki, Helsinki, Finland

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

has been reported to be 16.4% for conservative treatment [6], while mortality after emergency operations is higher than after elective surgery [7]. Due to the increased risk and eventual likelihood for emergency surgery, GPEH are commonly treated with elective surgery [6, 8, 9].

The long-term outcomes and durability of laparoscopic repair are debated [10–12]. Complex laparoscopic repair of GPEH has been reported to be safe and successful [13–15] and the postoperative length of stay is shorter and postoperative pain is less severe than after open surgery [11, 16]. On the other hand, radiological evidence of recurrence after laparoscopic surgery is often higher than after open repair [17].

In previous studies, the imaging of choice has usually been barium esophagogram [17, 18]. However, the barium swallow has become so rare in our clinical practice that few radiologists have experience of executing and interpreting it. Additionally, a barium esophagogram does not reliably show the size of a paraesophageal hernia preoperatively [19] and a sensitivity of only 30% before obesity surgery has been reported [20]. Computed tomography (CT) is currently in common use for both elective and emergency situations and has been used to measure the size of the hiatal orifice with high sensitivity [21].

Health-related quality of life (HRQoL) has been measured individually in previous studies with generic or disease-specific instruments. Minimal correlation between radiological findings and HRQoL has been shown [14, 18, 22].

The objective of this study was to evaluate the treatment outcomes of elective laparoscopic GPEH repair in a single tertiary care centre. The primary outcome was radiological recurrence in CT and its association with current symptoms and HRQoL. We also evaluated possible pre- and perioperative factors that could be linked to unsatisfactory outcomes or GPEH recurrence.

Methods

This was a consecutive case-series study on laparoscopic GPEH repair with follow up for radiological surveillance of hernia recurrence and patient postoperative HRQoL.

Patient cohort

The patient cohort included patients diagnosed with GPEH and subsequently operated electively with laparoscopy for GPEH in a single tertiary care hospital between 2010 and 2015. Patient information was collected retrospectively from electronic medical records (EMR) and included date of operation, preoperative symptoms, comorbidities, smoking, body mass index (BMI), preoperative proton pump inhibitor (PPI) use, pre- and postoperative laboratory and radiological tests, length of hospital stay, complications, reoperations, mortality, readmissions, intensive care unit stay, duration of

operation, urgency of operation, possible additional procedures, referring hospital, operating surgeon, and follow-up data. The size of the hernia was based primarily on the original radiologist's report of preoperative CT, or if unavailable, the assessment in the operation report or preoperative gastroscopy.

Surgical technique

The operations were performed laparoscopically using five ports with a Nathanson liver retractor. During the operation, the hernia sac is dissected circumferentially and reduced from the mediastinum. The esophagus is mobilised to obtain > 2 cm of tension-free esophagus beneath the diaphragm. A hiatoplasty is performed with sutures and the diaphragmatic crura are reinforced with mesh if needed. A fundoplication is commonly performed, predominantly a floppy Nissen fundoplication with a thick nasogastric tube as a bougie. In patients not suitable for fundoplication, a simple gastropexy is performed. The operation reports were reviewed to obtain more detailed information about the surgery. We examined the reports for possible fundoplication, use of mesh reinforcement, lengthening of esophagus, and other features.

Quality of life

The identified patients were sent an invitation to participate in the study with a letter of information, study consent, and questionnaires for generic 15D HRQoL and disease-specific Gastroesophageal Reflux Disease-Health Related Quality of Life (GERD-HRQL) instruments. The 15D is a generic, 15-dimensional self-administered quality of life instrument [23]. The 15D score was obtained from the patients that reported scores for each of the 15 dimensions. These scores were compared to the scores from the general population from the Terveys 2011 health study [24]. The GERD-HRQL was developed to measure symptom severity related to gastroesophageal reflux disease but was used here as similar symptoms are reported with GPEH [25, 26]. The questionnaires were administered only postoperatively. The patients were divided into groups of symptom severity by their GERD-HRQL scores. A score of 0–5 was considered excellent, 6–10 good, 11–15 fair, and 16 and over, poor.

Radiological recurrence

The patients who consented to participate were offered a non-contrast CT scan of the chest and upper abdomen. A radiologist's report was received to determine the recurrence of the hernia. A recurrent hernia was defined as a hernia with > 2 cm of the stomach above the diaphragm. The hernia was considered small, or minor recurrence, if it was 2 to 5 cm and a large recurrent

hernia, or major recurrence, was > 5 cm, as defined in prior studies [18, 22, 27].

Statistical analysis

We analysed results with statistical tests using IBM SPSS Statistics Version 24. Chi-squared or Fisher's exact test were used between two categorical variables. The Mann-Whitney test was used for analyses with interval independent variables and categorical dependent variables. The Kruskal-Wallis test was used for analyses with interval independent variables and ordinal dependent variables. Spearman correlation was used for analyses with two interval variables. A simple logistic regression was used for the analyses with categorical independent variables and interval dependent variables. Results were considered statistically significant when the two-tailed *p*-value was < 0.05.

Ethics approval

The study was submitted to and approved by the Research Ethics Committee of the Faculty of Medicine of Helsinki University (code 419/13/02/02/2015) and by the Institutional Review Board (IRB) of the Helsinki University Hospital Heart and Lung Centre (decision 8/2016).

Results

Patient demographics are presented in Table 1 and the patient flow in Fig. 1. A total of 227 patients underwent laparoscopic repair of paraesophageal hernia between 2010 and 2015. We excluded open procedures, both planned and converted, patients who had been previously operated on (*n* = 29), emergency operations (*n* = 20), patients with non-giant paraesophageal hernia (*n* = 9) and 4 patients who had been operated at a regional hospital. We included 165 patients in our study. The total number available for QoL analysis was 123 and a total of 93 for radiological analysis. The median follow-up time from the operation to receiving the questionnaires was 33 months (range 10 to 72 months).

Surgical features

Most operations (*n* = 134, 81.2%) were performed by one surgeon (JR) and altogether five different surgeons performed these operations. Mesh reinforcement was used in 8 patients (4.2%) and absorbable mesh was used in all except one of them. Esophageal lengthening was considered necessary after mobilization in none of the patients. A fundoplication was performed in 149 patients (90.3%). The mean duration of operation was 125 min (SD ± 51, range 51–348 min). Robot-assisted surgery was used for 9 patients (5.5%).

Table 1 Patient demographics

	n (%)
Gender	
Male	43 (26.1)
Female	122 (73.9)
Age (years)	
< 50	6 (3.6)
51–60	35 (21.2)
61–70	64 (38.8)
≥ 70	60 (36.4)
Age-adjusted CCI ^a	
0–2	71 (43.0)
≥ 3	94 (57.0)
Pulmonary disease ^b	36 (21.8)
History of ever smoking	42 (25.5)
Preoperative BMI > 35 kg/m ² ^c	12 (9.0)
Anemia	111 (67.3)
Symptomatic	158 (95.8)
Hernia size	
30–49%	26 (16.6)
50–74%	77 (46.7)
75–99%	22 (13.3)
100%	32 (19.4)
Hernia class	
III	148 (89.7)
IV	17 (10.3)
Fundoplication	149 (90.3)
Crural mesh	8 (4.8)
Length of stay (days)	
1–2	66 (40.2)
3–4	51 (30.9)
5–7	26 (15.8)
≥ 7	21 (13.1)


^aCCI/ Charlson comorbidity index

^bAsthma or chronic obstructive pulmonary disease (COPD)

^cBody Mass Index (BMI), data was available for 133 patients

Adverse events

Complications after laparoscopic operation were reported in 27 patients (16.4%); 4 patients had more than one complication. The complications were classified according to the Clavien-Dindo classification [28, 29]. There were 18 patients (10.9%) with grade-II complications with a median Charlson comorbidity index (CCI) of 1 [30]. A grade-III complication was reported in 7 patients (4.2%) with a median CCI of also 1. One patient (0.6%) had a grade-IV complication and one patient (0.6%) had a grade-V complication with CCIs of 2 and 4, respectively. Complications are summarised in Table 2.



Nine deaths occurred during follow up. There was one postoperative death within 30 days. This patient was considered high risk preoperatively, with an age-adjusted CCI of 6. According to autopsy, death was due to cryoglobulinemic vasculitis which caused intestinal perforation. The other eight deaths were not directly related to GPEH and occurred a mean of 22 months ($SD \pm 14.6$) after operation.

In total, 16 patients (9.7%) required reoperation. Of these, 10 (6.1%) occurred within 30 days of the primary operation, with reasons including recurrent hernia ($n = 3$), gastric paralysis ($n = 2$), small intestine strangulation ($n = 1$), suspected bleeding ($n = 1$), gastric perforation ($n = 1$), small intestine perforation ($n = 1$), and leakage at the GE junction ($n = 1$). The reoperations were carried out mainly using open technique, either laparotomy or thoracotomy. The patient with suspected bleeding was reoperated laparoscopically and for two patients endoscopic intervention with PEG was sufficient. The causes for a later reoperation were hernia recurrence and in one case gastric strangulation.

The median hospital stay postoperatively was 3 days (range 1 to 34 days).

Recurrence and patient reported outcomes

Of the 165 operated patients, 158 (95.8%) were symptomatic preoperatively. Disease-specific pre- and postoperative symptoms are presented in Table 3. The scores derived from the GERD-HRQL questionnaire were mainly excellent (66%) or good (12%). A fair score was achieved by 12 patients (10%) and a poor score by 15 patients (13%). The median GERD-HRQL score was 2 (range 0 to 56).

A total of 118 patients (71.5%) answered the question regarding current overall satisfaction. Seven (5.9%) patients reported overall dissatisfaction for symptom control postoperatively. The GERD score correlated with satisfaction ($p = 0.001$). The unsatisfied patients had a median score of 19 (range 6 to 54) and the satisfied patients had a median score of 1 (range 0 to 19).

The 15D quality of life survey was returned by 121 patients (73.3%). The results are presented in Fig. 2. The overall 15D HRQoL score was close to the age-adjusted average. The mean 15D score was 0.85 for patients and 0.90 for the general population ($p = 0.001$). The mean 15D total score was slightly better in the group with radiological recurrence (0.89) than in patients without

Table 2 Complications by Clavien-Dindo classification

Complications	n (%)
Grade II	18 (10.9)
Wound infection	6 (3.6)
Other infection	3 (1.8)
Lung embolism	3 (1.8)
Exacerbation of pulmonary disease	2 (1.2)
Urinary retention	2 (1.2)
Atrial fibrillation	1 (0.6)
Partial infarction of the spleen	1 (0.6)
Grade III	7 (4.2)
Chylothorax	1 (0.6)
Esophageal stricture	1 (0.6)
GE-junction perforation	1 (0.6)
Small intestine perforation	1 (0.6)
Small intestine strangulation	1 (0.6)
Gastric paralysis	1 (0.6)
Gastric strangulation	1 (0.6)
Grade IV	1 (0.6)
Gastric perforation	1 (0.6)
Grade V	1 (0.6)
Perforation of the duodenum	1 (0.6)

Table 3 Patient-reported symptoms pre- and postoperatively based on electronic medical records and current information obtained

Symptom	n = 162	preoperative n (%)	postoperative n (%)
Pain		94 (57.0)	13 (8.0)
Heartburn		40 (24.2)	3 (1.9)
Regurgitation		31 (18.8)	2 (1.2)
Vomiting		37 (22.4)	0 (0)
Dysphagia		49 (29.7)	13 (8.0)
Difficulty swallowing			
solid		35 (21.2)	0 (0)
soft		2 (1.2)	0 (0)
liquid		2 (1.2)	0 (0)
Dyspnea		25 (15.2)	2 (1.2)
Bloating		1 (0.6)	9 (5.6)
Early satiety		30 (18.2)	0 (0)
Aspiration		9 (5.5)	0 (0)
Cough		10 (6.1)	0 (0)
PPI ^a		97 (58.8)	16 (9.9)

^aDaily use of proton pump inhibitors

recurrence (0.84) ($p = 0.03$). The only dimension with a statistically significant difference between patients with and without recurrence was in the bladder and bowel functions (excretion) ($p = 0.012$).

A current CT scan was obtained from 92 patients (59% of operated patients). The median follow-up period from operation to CT was 39 months (range 12–79). Major recurrence, as defined in the Methods section, was revealed by CT in 4 patients (4.3%). The total recurrence rate in our patients, including the patients reoperated for recurrence preceding our follow up, was 29.3%. The symptoms evaluated by overall GERD-HRQL scores did not correlate with radiological findings ($p = 0.124$) (Fig. 3).

Risk factors


Known preoperative pulmonary disease was associated with postoperative complications ($p = 0.036$), a poorer quality of life in the 15D score ($p = 0.011$) and greater symptoms in GERD-HRQL ($p = 0.002$). Older age and longer operating time correlated with a prolonged length of stay (both $p = 0.001$). We did not identify any factors from patient or operation characteristics that could predict hernia recurrence.

Discussion

The most important finding of this study is that laparoscopic repair of GPEH in a specialised tertiary referral centre is safe, has a low recurrence rate for large hernia, and has good long-term patient satisfaction as measured by generic HRQoL and disease-specific symptom questionnaires. However, there is a moderate risk for grade I or II complications and overall hernia recurrence.

We observed a considerable rate of radiological recurrence after laparoscopic GPEH repair. Recurrent hernia occurred in 29% of the patients. However, only 4 patients (4.3%) had major recurrent hernia, which may still be susceptible to the risks associated with giant paraesophageal hernia, most notably incarceration, volvulus and perforation of the stomach. Recurrent hernia does not seem to affect patient HRQoL, which may explain the minimal reported need for reoperations in the literature for recurrent hernia (approximately 0.5 to 4.4%) even though radiological recurrence is common [18, 31, 32].

In initial studies, the radiological recurrence rate for GPEH was often higher in laparoscopic than in open procedures, with laparoscopic recurrence rates from 13% up to 42% [17]. Despite this early outcome report, laparoscopic operations have become the predominant operative approach [33]. As surgeons have improved laparoscopic operation techniques, results have improved over time and experience. The recurrence rates of laparoscopic repair of GPEH in retrospective series ranged from 12 to 21% [11, 34]. However, in a recent



*=Dimension with statistically significant difference (p<0.05)

Fig. 2 15D Quality of life score of patients with laparoscopically repaired giant paraesophageal hernia compared to the general population. The graph displays the scores for the 15 different dimensions and the total score for quality of life in our patients and the general population. GPEH = Giant paraesophageal hernia




Fig. 3 Symptom-related quality of life and recurrence. The graph shows the percentage of patients with recurrent paraesophageal hernia, diagnosed by computerized tomography, in different groups of symptom severity

prospective series the rate was as high as 32.7% at 1 year. [19] The total recurrence rate of 29.3% in our series is similar.

Our relatively high recurrence rate may depend on several factors. First, we performed follow up with CT, which may be more sensitive to detect recurrent hernias than a barium swallow test, especially true paraesophageal and sliding hernias. Second, we did not perform Collis gastroplasty, which may increase the number of radiological recurrences. We avoided lengthening procedures and mesh reinforcement to reduce additional risks. These include leakage and later stricture formation at the site of gastroplasty and erosion, stenosis, and dysphagia caused by mesh. The value of recurrence prevention by the most commonly currently used biological meshes in long-term follow up has been questioned [35]. As our patients had good HRQoL even after minor radiological hernia, it is unclear if better results would have been achieved with additional procedures [36].

In our data, the HRQoL as defined by the 15D score was close to the population average. A slight inferiority was detected in breathing, and bladder and bowel functions. The reasons for this result remain unclear and further examination would require a much larger series. The 15D score was slightly higher in the group of patients with recurrent hernia. While this group had fewer problems in bladder and bowel functions, the association with GPEH is uncertain. The preoperative quality of life was not measured here, but overall HRQoL improves with a laparoscopic GPEH repair [22].

Symptom relief (as measured by the GERD-HRQL score) after laparoscopic GPEH repair was good in our series. This has also been shown in previous studies, with good to excellent results reported by 77 to 90% of patients [17, 18, 32]. We also observed that as in many previous studies [17, 18] the radiological findings did not correlate with clinical symptoms.

Perioperative mortality was low (0.6%), similar to the 0 to 1.6% previously reported. The 30-day reoperation rate was 6.1%. The complication rate (16.4%) was comparable to the rates reported in previous studies (9 to 19%). Most complications were minor (Clavien-Dindo grade II), with wound infections being the most common. There were some cases of leakage and perforation, which is comparable to other series [17, 18, 31].

The strength of our study was that we reached a high percentage of our patients for follow-up surveys and CT scans. This study also has some limitations. This was a retrospective cross-sectional study with a single timepoint surveillance for imaging joined with HRQoL questionnaires. In addition, this study was limited to laparoscopic operations that were successfully completed; conversions to open repair were excluded. Thus, we did not use an intention-to-treat analysis, but rather concentrated on the

end results of laparoscopy. We also acknowledge that there were no structured questionnaires for both HRQoL and symptoms preoperatively, which could cause recall bias in the presentation of symptoms. Despite a significant portion of patients willing to participate in this study, some patients declined an additional follow-up CT scan, which could cause selection bias in the results.

Conclusions

In conclusion, although laparoscopic repair of GPEH is a complex procedure, when executed by an experienced surgeon it appears to lead to effective symptom relief with high patient satisfaction. Minor radiological recurrence is common but does not affect HRQoL or satisfaction. These findings suggest that the laparoscopic approach is a feasible first-line surgical strategy for GPEH repair. Further research with longer follow-up times and wider, population-based studies are required to determine the optimal approach to repair GPEH.

Abbreviations

BMI: Body Mass Index; CCI: Charlson Comorbidity Index; COPD: Chronic obstructive pulmonary disease; CT: Computed tomography; EMR: Electronic medical records; GERD-HRQL: Gastroesophageal reflux disease –health related quality of life; GPEH: Giant paraesophageal hernia; HRQoL: Health-related quality of life; IRB: Institutional review board; PPI: Proton pump inhibitor

Acknowledgements

Not applicable.

Authors' contributions

Conception and design of the work: H.H., I.I., J.R. Acquisition of data: H.H., I.I. Analysis and interpretation of data: H.H., I.I., T.J. Analysis and interpretation of data concerning the 15D quality of life instrument: H.S. Drafting of the manuscript: H.H. Review and editing the manuscript: H.H., I.I., T.J., J.K., S.A., J.R. All authors have read and approved the manuscript.

Funding

The research was carried out with state funding.

Availability of data and materials

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The study was submitted to and approved by the Research Ethics Committee of the Faculty of Medicine of Helsinki University (code 419/13/02/02/2015) and by the Institutional Review Board (IRB) of the Helsinki University Hospital Heart and Lung Centre (decision 8/2016). All procedures performed were in accordance with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all participants.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of General Thoracic and Esophageal Surgery, Helsinki University Hospital, Haartmaninkatu 4, 00290 Helsinki, Finland. ²Department of Surgery, Clinicum, University of Helsinki, Helsinki, Finland. ³Department of Public Health, University of Helsinki, Helsinki, Finland.

Received: 5 February 2020 Accepted: 11 May 2020

Published online: 20 May 2020

References

- Mitiek MO, Andrade RS. Giant hiatal hernia. *Ann Thorac Surg*. 2010;89(6):S2168–73.
- Maziak DE, Todd TR, Pearson FG. Massive hiatus hernia: evaluation and surgical management. *J Thorac Cardiovasc Surg*. 1998;115(1):53–60 discussion 1–2.
- Allen MS, Trastek VF, Deschamps C, Pairolero PC. Intrathoracic stomach. Presentation and results of operation. *J Thorac Cardiovasc Surg*. 1993;105(2):253–8 discussion 8–9.
- Awais O, Luketich JD. Management of giant paraesophageal hernia. *Minerva Chir*. 2009;64(2):159–68.
- Stylopoulos N, Gazelle GS, Rattner DW. Paraesophageal hernias: operation or observation? *Ann Surg*. 2002;236(4):492–500 discussion –1.
- Sihvo E, Salo JA, Rasanen JV, Rantanen TK. Fatal complications of adult paraesophageal hernia: a population-based study. *J Thorac Cardiovasc Surg*. 2009;137(2):419–24.
- Tam V, Luketich JD, Winger DG, Sarkaria IS, Levy RM, Christie NA, et al. Non-elective Paraesophageal hernia repair portends worse outcomes in comparable patients: a propensity-adjusted analysis. *J Gastrointest Surg*. 2017;21(1):137–45.
- Polomsky M, Jones CE, Sepesi B, O'Connor M, Matousek A, Hu R, et al. Should elective repair of intrathoracic stomach be encouraged? *J Gastrointest Surg*. 2010;14(2):203–10.
- Stylopoulos N, Rattner DW. Paraesophageal hernia: when to operate? *Adv Surg*. 2003;37:213–29.
- Schauer PR, Ikramuddin S, McLaughlin RH, Graham TO, Slivka A, Lee KK, et al. Comparison of laparoscopic versus open repair of paraesophageal hernia. *Am J Surg*. 1998;176(6):659–65.
- Dallemagne B, Quero G, Lapergola A, Guerriero L, Fiorillo C, Perretta S. Treatment of giant paraesophageal hernia: pro laparoscopic approach. *Hernia*. 2017.
- Lebenthal A, Waterford SD, Fischella PM. Treatment and controversies in paraesophageal hernia repair. *Front Surg*. 2015;2:13.
- Luketich JD, Raja S, Fernando HC, Campbell W, Christie NA, Buenaventura PO, et al. Laparoscopic repair of giant paraesophageal hernia: 100 consecutive cases. *Ann Surg*. 2000;232(4):608–18.
- Pierre AF, Luketich JD, Fernando HC, Christie NA, Buenaventura PO, Litle VR, et al. Results of laparoscopic repair of giant paraesophageal hernias: 200 consecutive patients. *Ann Thorac Surg*. 2002;74(6):1909–15 discussion 15–6.
- Pitcher DE, Curet MJ, Martin DT, Vogt DM, Mason J, Zucker KA. Successful laparoscopic repair of paraesophageal hernia. *Arch Surg*. 1995;130(6):590–6.
- Draaisma WA, Gooszen HG, Tournioij E, Broeders IA. Controversies in paraesophageal hernia repair: a review of literature. *Surg Endosc*. 2005;19(10):1300–8.
- Hashemi M, Peters JH, DeMeester TR, Huprich JE, Quek M, Hagen JA, et al. Laparoscopic repair of large type III hiatal hernia: objective followup reveals high recurrence rate. *J Am Coll Surg*. 2000;190(5):553–60 discussion 60–1.
- Luketich JD, Nason KS, Christie NA, Pennathur A, Jobe BA, Landreneau RJ, et al. Outcomes after a decade of laparoscopic giant paraesophageal hernia repair. *J Thorac Cardiovasc Surg*. 2010;139(2):395–404 e1.
- Koch OO, Schurich M, Antoniou SA, Spaun G, Kaindstorfer A, Pointner R, et al. Predictability of hiatal hernia/defect size: is there a correlation between pre- and intraoperative findings? *Hernia*. 2014;18(6):883–8.
- Goitein D, Sakran N, Rayman S, Szold A, Goitein O, Raziel A. Barium swallow for hiatal hernia detection is unnecessary prior to primary sleeve gastrectomy. *Surg Obes Relat Dis*. 2017;13(2):138–42.
- Ouyang W, Dass C, Zhao H, Kim C, Criner G, Investigators CO. Multiplanar MDCT measurement of esophageal hiatus surface area: association with hiatal hernia and GERD. *Surg Endosc*. 2016;30(6):2465–72.
- Stringham JR, Phillips JV, McMurry TL, Lambert DL, Jones DR, Isbell JM, et al. Prospective study of giant paraesophageal hernia repair with 1-year follow-up. *J Thorac Cardiovasc Surg*. 2017;154(2):743–51.
- Sintonen H. An approach to measuring and valuing health states. *Soc Sci Med Med Econ*. 1981;15(2):55–65.
- Koskinen S, Lundqvist A, Ristiluoma N. Terveys, toimintakyky ja hyvinvointi Suomessa 2011. Helsinki: Terveystieteiden tutkimuskeskus; 2012.
- Velanovich V, Vallance SR, Gusz JR, Tapia FV, Harkabus MA. Quality of life scale for gastroesophageal reflux disease. *J Am Coll Surg*. 1996;183(3):217–24.
- Velanovich V. The development of the GERD-HRQL symptom severity instrument. *Dis Esophagus*. 2007;20(2):130–4.
- Jones R, Simorov A, Lomelin D, Tadaki C, Oleynikov D. Long-term outcomes of radiologic recurrence after paraesophageal hernia repair with mesh. *Surg Endosc*. 2015;29(2):425–30.
- Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004;240(2):205–13.
- Clavien PA, Barkun J, de Oliveira ML, Vauthey JN, Dindo D, Schulick RD, et al. The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg*. 2009;250(2):187–96.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373–83.
- Dallemagne B, Kohlen L, Perretta S, Weerts J, Markiewicz S, Jehaes C. Laparoscopic repair of paraesophageal hernia. Long-term follow-up reveals good clinical outcome despite high radiological recurrence rate. *Ann Surg*. 2011;253(2):291–6.
- Nason KS, Luketich JD, Qureshi I, Keeley S, Trainor S, Awais O, et al. Laparoscopic repair of giant paraesophageal hernia results in long-term patient satisfaction and a durable repair. *J Gastrointest Surg*. 2008;12(12):2066–75 discussion 75–7.
- McLaren PJ, Hart KD, Hunter JG, Dolan JP. Paraesophageal hernia repair outcomes using minimally invasive approaches. *JAMA Surg*. 2017;152(12):1176–8.
- Zehetner J, Demeester SR, Ayazi S, Kilday P, Augustin F, Hagen JA, et al. Laparoscopic versus open repair of paraesophageal hernia: the second decade. *J Am Coll Surg*. 2011;212(5):813–20.
- Oelschlager BK, Pellegrini CA, Hunter JG, Brunt ML, Soper NJ, Sheppard BC, et al. Biologic prosthesis to prevent recurrence after laparoscopic paraesophageal hernia repair: long-term follow-up from a multicenter, prospective, randomized trial. *J Am Coll Surg*. 2011;213(4):461–8.
- Nason KS, Luketich JD, Awais O, Abbas G, Pennathur A, Landreneau RJ, et al. Quality of life after collis gastroplasty for short esophagus in patients with paraesophageal hernia. *Ann Thorac Surg*. 2011;92(5):1854–60 discussion 60–1.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

