

ANTAGONISM AND MODES OF ACTION OF *CHAETOMIUM GLOBOSUM* SPECIES GROUP, POTENTIAL BIOCONTROL AGENT OF BARLEY FOLIAR DISEASES

PAULINA MOYA^{1,3*}, DEBORA PEDEMONTE¹, SUSANA AMENGUAL¹, MARIO E. E. FRANCO^{1,4}, MARINA N. SISTERNA^{1,2}

Resumen: Antagonismo y mecanismos de acción de hongos del grupo de especies *Chaetomium globosum*, potenciales biocontroladores de enfermedades fúngicas foliares de la cebada. La "mancha en red" y la "mancha borrosa" son enfermedades foliares de cebada causadas por *Drechslera teres* y *Bipolaris sorokiniana*, respectivamente. Una alternativa para su manejo es el control biológico. El objetivo de este trabajo fue identificar dos aislamientos del género *Chaetomium* (C2 y C5), endófitos de plántulas de cebada y estudiar sus interacciones *in vitro* con *D. teres* y *B. sorokiniana*, aisladas de semillas del mismo hospedante. Todos los microorganismos se caracterizaron cultural y morfológicamente. Los aislamientos de *Chaetomium* además se caracterizaron molecularmente. Se realizaron pruebas de cultivos duales y se registraron los efectos del antagonista a nivel microscópico en la morfología de los patógenos. Los resultados confirmaron la identidad de los patógenos y de los aislamientos de *Chaetomium* spp. como *Chaetomium globosum* grupo de especies. *Bipolaris sorokiniana* fue inhibida en un 30% por C2 y en un 31.2 % por C5 respecto al control. *D.teres* fue inhibida en un 40% por C2 y en un 36% por C5 en referencia al control. Los mecanismos de acción frente a *B. sorokiniana* fueron antibiosis y competencia. Microscópicamente se observaron conidios aberrantes. Frente a *D. teres* se registró competencia y micoparasitismo. Microscópicamente se evidenció plasmólisis, enrollamiento y pigmentación anaranjada.

Palabras clave: *Bipolaris sorokiniana*, *Chaetomium globosum*, *Drechslera teres*, antibiosis, endófito, micoparasitismo.

Summary: "Net blotch" (*Drechslera teres*) and "Bipolaris spot blotch" (*Bipolaris sorokiniana*) are foliar diseases of barley. Biological control is currently considered as an efficient alternative to chemical management of these plant diseases. The aim of the present study was to identify 2 isolates of *Chaetomium* (C2 and C5), endophytes on barley seedlings and to study the *in vitro* interactions with *D. teres* and *B. sorokiniana*, isolated from seeds of the same host. Cultural and morphological characterization of all microorganisms was done. In addition, molecular characterization of *Chaetomium* spp. was conducted and dual culture tests were carried out to find, by microscopic observations, the effects of the antagonist on the morphology of the pathogens. The results confirm the identity of the pathogens and the isolates of *Chaetomium* spp. as *Chaetomium globosum* species group. Inhibition of *B. sorokiniana* and *D. teres* by C2 and C5 accounted for 30% and 31.2 %, and 40% and 36% respectively, compared with the control. The mechanisms of action against *B. sorokiniana* and *D. teres* were antibiosis and competition and mycoparasitism, respectively. Microscopic observation revealed deformed conidia in *B. sorokiniana* and plasmolysis, coiling and orange pigmentation in *D. teres*.

Key words: *Bipolaris sorokiniana*, *Chaetomium globosum*, *Drechslera teres*, antibiosis, endophyte, mycoparasitism.

¹ Centro de Investigaciones de Fitopatología (CIDEFI) - Facultad de Ciencias Agrarias y Forestales, Universidad Nacional de La Plata. 60 y 119, (1900) La Plata, Buenos Aires, Argentina.

² Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CICPBA)

³ Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

⁴ Agencia Nacional de Promoción Científica y Tecnológica

Autor para correspondencia : p_moya_@hotmail.com

INTRODUCTION

Malting barley crop (*Hordeum distichum* L.) plays a key role in the cereal global production. In Argentina, malting barley production covers an area of approximately 1.000.000 hectares (MAGyP, 2014). As a result of minimum tillage and the use of susceptible cultivars, diseases have become an increasing problem (Carmona 2009). "Net blotch" (*Drechslera teres* (Sacc.) Shoem. (teleomorph *Pyrenophora teres* Drechsler), is an endemic disease, it means that remains in a particular area, with a prevalence of 76 % and an incidence of 100% , being responsible for yield losses of 20% (Carmona *et al.*, 1999). The most affected components include grain weight and the number of grains per square meter. Also, the malt extract decreases, affecting malt quality for beer production (Carmona 2009). "Bipolaris spot blotch" (*Bipolaris sorokiniana*) (Sacc.) Shoemaker causes foliar lesions and this pathogen also affects other plant parts producing discoloration grain, seedling blight and root rot (Fernández Valiela, 1978; Wiese, 1987). In Argentina, frequency and incidence of *Bipolaris* spot blotch has increased in regions with more moderate temperatures. In addition, the barley growing region extends to the central-north provinces (Santa Fe, Entre Ríos) resulting in a highest risk of disease spread due to the fact that temperate humid regions are the most favorable areas for the disease occurrence (Carmona, 2009). All these causes have established *Bipolaris* spot blotch as a re-emerging disease (Sisterna, 2014).

Biocontrol is an environmental-friendly and efficient alternative to chemical pesticide management of plant diseases. In this sense, endophyte organisms, particularly fungi, have received major attention in recent years. Endophytes are capable to reduce in the host the effect of fungi diseases, through secondary metabolites production as alkaloids (Abello & Kelemu 2006). The plant-endophyte relationship is so strong that could be involved in the systemic induced resistance (Waller *et al.*, 2005). All these mechanisms lead to a higher competitive ability compared to plant endophyte free (Vila-Aiub *et al.*, 2003).

There are many studies with promising results on using endophyte *Chaetomium* spp. as a biocontrol agent. Antagonistic mechanisms of this fungus include competition for space and nutrients (Vannaci

& Harman 1987), mycoparasitism (Mandal *et al.*, 1999) and metabolite production (antibiosis) such as chaetomin, chaetoglobosin, cochliodinol, chaetosin and prenisatin (Brewer *et al.*, 1970; Brewer *et al.*, 1972; Brewer & Taylor, 1978). Other microscopic studies analyzed deformed conidia with distorted walls, lysis and formation of holes in the host pathogen mycelium, inhibition in conidial germination and hyphal elongation (Mandal *et al.*, 1999; Biswas *et al.*, 2000).

Chaetomium spp. is reported as antagonist of soil and foliar pathogens (Soytong *et al.*, 2001) in rice (Soytong, 1992), maize (Soytong, 1991) and wheat (Biswas *et al.*, 2000; Aggarwal *et al.*, 2004; Istifadah & McGee, 2006; Istifadah *et al.*, 2006).

The aim of the present study was to analyze *in vitro* interactions between two isolates of *Chaetomium globosum* species group, previously morphocultural and molecular characterized, with *Bipolaris sorokiniana* and *Drechslera teres*, fungi phytopathogens of barley.

MATERIALS AND METHODS

Isolation of microorganisms

Drechslera teres and *B. sorokiniana* were isolated from samples of naturally infected barley seeds from cv. Scarlet, cultivated in Tres Arroyos locality (Buenos Aires Province). The agar method was used following the rules imparted by ISTA (Neegaard, 1979). On the other hand, 12 isolates of *Chaetomium* spp. were obtained from previous biocontrol greenhouse tests. In them, from barley seedlings, small pieces of leaves were disinfected in 70% ethanol (1½ min), 5% sodium hypochlorite (1½ min), rinsed in sterile distilled water (3 min) and placed in Petri dishes with potato dextrose agar (PDA) medium 2%. The dishes were incubated for 5-6 days at 25°C, until colony growth was observed. Pure cultures used in subsequent studies were stored on slants of PDA.

Cultural and morphological characterization of isolates of Chaetomium spp. Drechslera teres and Bipolaris sorokiniana

Among the 12 *Chaetomium* isolates, two groups with cultural and morphological differences were *a priori* distinguished. From them, one isolate from each one (C2 and C5) were chosen as

representatives.

Vegetative and reproductive structures of both isolates were studied. Also their cultural characteristics were analyzed on different media: potato - dextrose agar (PDA), oatmeal agar (OA) and malt extract agar (MEA). The isolates were grown in darkness at temperature 23° and five replications from each treatment were done. The parameters evaluated at 7, 14 and 21 days were: average diameter and colour (according to Rayner 1970).

Ascomatas descriptions and spores microscopic measurements were taken from colonies grown on OA. This was the medium where the isolates gave the best sporulation. Domsch *et al.* (2007) keys were followed to the systematic classification. Other specific literature (Dreyfuss *et al.*, 1986; Ames 1963; Von Arx *et al.*, 1986) were consulted for the morfo-cultural characterization.

Drechslera teres and *B. sorokiniana* were characterized based on cultural and morphological characteristics, following routine phytopathologic methods and specific literature (Ellis, 1971, 1976; Sivanesan, 1987)

Molecular characterization of Chaetomium isolates

From the monosporic cultures of the two isolates of *Chaetomium* spp. C2 and C5, DNA was isolated using the CTAB method of Doyle & Doyle (1990), modified by Bornet and Branchard (2001). Amplification was performed using ribosomal DNA fragment that conforms the ITS (Internal Transcribed Spacer ITS1 - ITS4), with primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White *et al.*, 1990). The amplification reaction was performed under the following conditions: one cycle of initial denaturation at 94 °C for 4 minutes; 33 cycles at 94 °C for 1 minute, 45 seconds at 56 °C and at 72 minutes and finally an elongation of 5 minutes at 72 °C (Curtis *et al.*, 1994; Goodwin *et al.*, 2001). The reaction mixture consisted of 1X reaction buffer (500 mM KCl; 100 mM Tris-HCl, pH 9.0 at 25; 1% Triton X-100 (no magnesium)); 1.5 mM magnesium chloride; 0.2 mM of each dNTP; 0.3 mM of each primer; 1 U of T-Plus DNA polymerase (Molecular Biology-Highway INBIO-UNICEN) and 50 ng of template DNA per reaction, in a final volume of 15 ul. The amplified fragments were purified by adding one volume of

isopropanol and 0.1 volume of 3M Sodium Acetate. The mixture was kept at -18 ° C for 12 hours and then centrifuged at 15,000 g for 15 minutes. The DNA was washed with 70% ethanol, then dried and dissolved in sterile distilled water (Maniatis *et al.*, 1982). The two strands of the DNA fragment amplified by PCR were sequenced in Macrogen Inc. Seoul, Korea. The sequences were deposited in databases DDBJ / EMBL / GenBank under the access numbers KX011856 for isolate C2 and KX011855 for isolate C5.

In vitro antagonistic test

The inhibitory action of C2 and C5 against the two pathogens was tested in dual culture (Dal Bello *et al.*, 1994). An antagonist disc (5 mm. diameter) was placed in the Petri dish containing 9 ml. PDA. After three days, a pathogen disc was placed leaving 4 cm. distance from the first disc. Three replicates of each pathogen-antagonist interaction were done. A culture of each pathogen alone served as control. The Petri dishes with the dual cultures were incubated for 7 days at 25°C. Two diameters perpendicular to each pathogenic colony were measured for all treatments and average values were determined. These values were analyzed by the Kruscal Wallis test and then Tukey test was conducted to find the differences among treatments.

To reveal the effects of the antagonist on the pathogen morphology, microscopic observations were performed.

RESULTS

Cultural and morphological characterization of isolates of Chaetomium spp., D. teres and B. sorokiniana

In OA the isolates showed olivaceous-yellow colonies according to the color of (Rayner, 1970). Isolate C5 showed globose or oval ostiolar ascomata with straight or undulate hairs (Fig. 1A) unlike isolate 2, which showed irregular hairs with no undulations (Fig. 1B). Both isolates showed terminal hairs all unbranched and displayed one-celled, lemon-shaped ascospores, with an apical germ pore and regular in shape with sizes: C2: 8.4µm x 6µm; C5: 9.6µm x 7.68µm (Fig. 1C). In PDA isolate C5 was slightly yellower than C2 and showed margins wavy or irregular, while C2 developed full edges. In

MEA both isolates were transparent. The growth on the several media did not evidence major differences. It was slightly higher on OA, followed by APG and MEA. At 7 days, on the three media, C2 grew faster than C5. Then the growth stabilized. The morphocultural characteristics of C2 and C5 isolates were very close to *C. globosum*.

The isolates of *B. sorokiniana* showed grey olivaceous to black velvet-like colonies with wavy or irregular margins, abundantly sporulating (Fig. 1D). *B. sorokiniana* displayed curved conidia, though in culture they are frequently straight, fusoid or elipsoid, dark brown in color, smooth, 4 to 12 pseudosepta, 40-120 (80) x 17-28(22.5) µm. (Figure 1E).

D. teres showed typical cultural characteristics of the species. The mycelium was initially medium grey (Rayner, 1970) with white tuft-like formations (Fig. 1F) near the edge of the colony. The conidia were straight, cylindrical, rounded at the ends, subhyaline, yellowish brown to olivaceous brown, 4 to 6 pseudosepta, 72-105 (88.5) x 12-21 (14.4) µm L/A (Fig. 1G).

Molecular identification of Chaetomium isolates

ITS amplicons for strains C2 and C5 had sizes of 557 and 556 bp. respectively. A search of both sequences in the GenBank database using BLAST tool found that the sequence KX011855 was 99% identical to its counterpart *Chaetomium globosum* strain CGMCC of 3.14295 (JN209864.1), isolated in China, while KX011856 what the sequence was 100% for *Chaetomium globosum* voucher CIAT563 (KR012922.1) of Colombian origin. However, it also were found high levels of sequence identity with other species of the same genus so this controversial result made the identification impossible only based on the nucleotide sequence of the ITS.

In vitro culture test

The results of the *C. globosum* – *B. sorokiniana* interaction were showed in Table 1.

Seven days after incubation significant differences were observed among treatments (K: 23.49 p= 0.000008). According to the Tukey test strains C2 and C5 did not show significant differences in their inhibitory capacity between them. However,




Fig. 1. Monoespecific cultures: (A) Ascomata from *Chaetomium globosum* strain 5, globose or oval ostiolar with straight or undulate hairs. OM 40X scar = 200 µm. (B) Ascomata from *Chaetomium globosum* strain 2, with irregular hairs without undulations. OM 40X scar = 200 µm. (C) Unicellular, lemon-shaped ascospores of *C. globosum* strain 2; OM 40X scar = 10 µm. (D) Colonies of *Bipolaris sorokiniana* and (E) conidia. OM 40X scar = 50 µm. (F) Mycelium of *Drechslera teres* with white tuft-like formations. (G) Cylindrical and rounded ends conidia of *D. teres* OM 40X scar = 50 µm.

Table 1. Average values of the mycelial growth of *Bipolaris sorokiniana* in the presence of *C. globosum* C2, C5 and controls. Within the average column, means followed by the same letter are not significantly different ($P < 0.001$) by the Tukey test.

Treatments	N	Average	S.E.
Control	12	7.15 b	± 0.04
C2	12	5.02 a	± 0.58
C5	12	4.92 a	± 0.34

Table 2. Average values of the mycelial growth of *Drechslera teres* in the presence of *C. globosum* C2, C5 and controls. Within the average column, means followed by the same letter are not significantly different ($P < 0.001$) by the Tukey test.

Treatment	N	Average	S.E.
Control	12	8.72 b	± 0.05
C2	12	5.29 a	± 0.28
C5	12	5.56 a	± 0.12

significant differences were observed with respect to the control. The inhibitory effect of C2 and C5 were 30 % and 31%, respectively.

Similar conditions regarding *C. globosum* - *D. teres* interaction were observed in Table 2 (K: 23.76 $p = 0.000007$) with significant differences with respect to the control and no significant differences between the isolates. C2 inhibited 40% and C5 36% with respect to the control (Fig. 2).

The predominant mechanism of action of *C. globosum* species group against *B. sorokiniana* was antibiosis (Fig. 3A). Other mechanisms,

such as competition were also observed (Fig. 3B). Microscopic observation of *B. sorokiniana* revealed abnormal conidia as a result of the contact with *C. globosum* species group. (Fig. 3C).

The most important mechanisms displayed by *C. globosum* - *D. teres* included competition and mycoparasitism, with an orange pigmentation in the interaction zone (Fig. 3D y 3E). Microscopic observations of *C. globosum* - *D. teres* interaction revealed plasmolysis (Fig. 4A), hyphal coiling (Fig. 4B) and orange pigmentation (Fig. 4C).




Fig. 2. Percent of mycelial growth inhibition of *Bipolaris sorokiniana* and *Drechslera teres* caused by *Chaetomium globosum* strains C2 and C5.




Fig. 3. Antagonistic interactions of dual cultures of *C. globosum* against *B. sorokiniana* : (A) Antibiosis; (B) Competition. (C) *B. sorokiniana* abnormal conidia. OM 40X scar = 50 μ m. (D) Orange pigmentation in the interaction area between *D. teres*- C2 ; (E) *D. teres*-C5.

DISCUSSION

Following biocontrol steps, *in vitro* assays are necessary to carry out a second stage of *in vivo* assays. This investigation gives a first approach on the knowledge for *C. globosum* application *in vivo*, as a biocontrol agent of spot blotch and net blotch. For that, the identification to a specific level is of great importance, as well as its characterization. The study of each species allows deeper knowledge to evaluate antagonistic behavior.

Chaetomium globosum is by far the commonest and most cosmopolitan *Chaetomium* species especially on plant remains, seeds, compost paper and other cellulosic substrates. (Domsch *et al.*, 2007).

Several authors have tried unsuccessfully to define sections of *Chaetomium* based on morphological characters, which later proved to be heterogeneous. Ames (1963) distinguished 10

groups based on the shape of ascumatal hairs. Sörgel (1960) suggested a subdivision of *Chaetomium* into 13 species groups based on ascospore features. Seth (1970) used ascospore shape and ascumatal hairs separately to distinguish eight species groups. A comprehensive classification system of *Chaetomium* was proposed by Dreyfuss (1976). Based on morphological characters (ascogonia, asci, ascospores and anamorphs) together with some physiological traits (temperature and nutritional requirements, growth and fruiting rate, compatibility, etc.), he divided the genus into four species groups, namely *C. aureum*, *C. bostrychodes*, *C. globosum* and *C. murorum*. Von Arx *et al.* (1986) provided a more comprehensive revision, which largely followed the concepts of Sörgel (1960) and Dreyfuss (1976), recognizing only 134 species.

Cultural and morphological characterization of C2 and C5 isolates agrees with the description




Fig. 4. *Chaetomium globosum* - *Drechslera teres* interaction area (A) plasmolysis; (B) hyphal coiling; (C) orange pigmentation. OM 40X scar = 50 μ m.

of Dreyfus *et al.* (1976) for the *Chaetomium globosum* species group. Its main characteristics are the presence of limoniform spores and peridial hairs, ranging from wavy to straight. A peridium exclusively of *textura intricata* or in combinations with *textura epidermoidea*, ascospores ranging from limoniform with apiculate ends to ellipsoidal or nearly spherical without apiculate ends, and with one (rarely two) germ pores (Asgari & Zare 2011). In the present study, colour and measurements of colonies, ascospores size and type and shape of peridial hairs agree with that reported by Domsch *et al.* (2007).

Scarce investigations have been performed on molecular taxonomy. *Chaetomium* spp. and other members of the Chaetomiaceae were studied with 18S and 28S rDNA sequences individually (Lee & Hanlin 1999, Untereiner *et al.* 2001) or in combination with *tef* and *rpb2* genes (Zhang *et al.*, 2006). These studies have supported the monophyly of the Chaetomiaceae, occupying a sister relationship to other families of Sordariales, especially Lasiosphaeriaceae.

Asgari & Zare (2011) analyzed 21 species of *Chaetomium* taxa, on the basis of morphological and molecular characters, using a combined sequence dataset of the ITS region, partial LSU rDNA, and b-tubulin gene.

In this investigation high levels of C2 and C5 identity with Gene Bank sequences of *C. globosum* were found but also high levels with other species of the genera. This result difficulties the identification, based exclusively in the ITS nucleotidic sequence. Rodríguez Morejón (2003) found similar difficulties. This researcher used

the highly conserved region D1 and D2 from 288 ADNr and demonstrated that a unique gen cannot be used in the molecular study of *C. globosum*. To confirm the identity of both isolates through molecular tools it will be necessary to combine sequence dataset of the ITS region, partial LSU rDNA, and b-tubulin (Asgari & Zare 2011). So we can conclude that, in this study, isolates C2 and C5 are classified inside the *C. globosum* species group, not being possible to reach to the species level with the sequences analyzed.

The present study reveals that in *in vitro* dual tests, both *Chaetomium* isolates, C2 and C5, inhibited the two pathogens. Aggarwal *et al.*, (2004), Biswas *et al.*, (2000) and Mandal *et al.*, (1999) found similar results for *B. sorokiniana*. Also, Tathan *et al.*, (2012) working *in vitro* with *B. oryzae* demonstrated antagonist activity of several species of *Chaetomium*.

No studies are available for *D. teres* on this regard. However, Istifadah & McGee (2006); Istifadah *et al.* (2006) worked on the interaction between isolates of *Chaetomium globosum* and *D. tritici-repentis*, another species of this genus, which is responsible for the wheat tan spot.

The orange pigmentation observed in the interaction area *C. globosum*-*D. teres*, both at microscopic and macroscopic level was also described in other interactions with *D. teres* and the antagonist *Trichoderma* spp. (Moya & Sisterna 2012; Moya *et al.*, 2015). Related studies referred this coloration to the presence of anthraquinones, particularly catenarin. These metabolites could play an antifungal and antibacterial role (Engström *et al.*, 1993; Wakuliński *et al.*, 2003).

Antibiosis was the most prevalent mechanism of action between *C. globosum*-*B. sorokiniana* in agreement with studies by Aggarwal *et al.* (2004). Biswas *et al.* (2000) and Mandal *et al.* (1999) also observed this mechanism in their experiments. Istifadah *et al.* (2006) found the same effect against *D. tritici-repentis*. The role of this mechanism in the antagonist activity of *C. globosum* against *Venturia inaequalis*, *Pythium ultimum*, *Fusarium nivale*, *Sclerotinia sclerotiorum* and *Helminthosporium* spp. has also been recorded by several researchers (Tveit & Wood, 1955; Hubbard *et al.*, 1982; Cullen & Andrews, 1984; Di Pietro, 1992; Nakashima *et al.*, 1991). *Chaetomium* spp. produces a diverse amount of bioactive metabolites such as chaetoglobosin A, chaetomin, cochliodinol and prenisatin that inhibit the growth of the pathogens *in vitro* (Brewer *et al.*, 1972; Breinholt *et al.*, 1996). Other authors suggest that the antibiosis must not be the most important mechanism of action of an antagonist, since there is a risk of occurrence of pathogenic strains resistant to antibiotics (Infante *et al.*, 2009).

Competition constitutes a major antagonistic mechanism, which is favoured by the biocontrol agent characteristics such as ecological plasticity, growth velocity and development and by external factors such as soil type, pH, temperature, humidity, among others (Infante *et al.*, 2009). In the present study, antagonism by competition, with reduction of the growth of the pathogen mycelium was also observed both in *B. sorokiniana* and *D. teres* interactions. Few studies on this mechanism of action were found in *Chaetomium* spp. (Vanacci & Harman, 1987). Mycoparasitism was observed in the interaction between *C. globosum* and *B. sorokiniana*. This mode of action produced conidia deformation. Similar results were reported by Biswas *et al.* (2000). Also, Mandal *et al.* (1999) observed, between these organisms, hyphal interaction with lysis and perforation of the pathogen wall caused by the antagonist.

In the interaction between *C. globosum* and *D. teres*, hyphal coiling was observed. The same was found by Vannacci & Harman (1987) who studied the interaction *C. globosum* - *Alternaria brassicicola* and *C. globosum* - *A. raphani* and between this antagonist and *Rhizoctonia solani*. These interactions also revealed intracellular growth of *C. globosum* in the pathogen hyphae (Gao *et al.*, 2005).

In Argentina, no studies *in vitro* on the interaction between the endophyte *Chaetomium globosum* species group and phytopathogenic agents have been reported. Therefore, the present study provides information regarding the relationship between this antagonist and phytopathogenic fungi of barley. Future research should consider the use of this endophyte as a useful control tool in an integrated strategy for disease treatments.

ACKNOWLEDGEMENTS

We want to especially thank Dr. Pedro Balatti from UNLP (Universidad Nacional de La Plata) for their contribution to molecular characterization of *Chaetomium* spp. This paper was performed with financial support from CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas), CICPBA (Comisión de Investigaciones Científicas de la Provincia de Bs. As.) and UNLP.

BIBLIOGRAPHY

- ABELLO, J. F & S. KELEMU. 2006. Hongos endófitos: ventajas adaptativas que habitan en el interior de las plantas. *Revista CORPOICA* 7: 55-57. Disponible en: http://www.corpoica.org.co/SitioWeb/Archivos/Revista/6_HongosEndfitos_RevCorpoica_V7N2_2006_CorregidoDEF.pdf
- AGGARWAL, R., A. K. TEWARI, K. D. SRIVASTAVA & D.V. SINGH. 2004. Role of antibiosis in the biological control of spot blotch (*Cochliobolus sativus*) of wheat by *Chaetomium globosum*. *Mycopathologia* 157: 369-377.
- AMES, L. M. 1963. A monograph of the Chaetomiaceae. Ser. 2: 1-125 *Army Res Dev.*, Washington DC.
- ATSCHUL, S. F, W. GISH, W. MILLER, E.W. MYERS & D. J. LIPMAN. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215: 403-410.
- ASGARI, B. & R. ZARE. 2011. The genus *Chaetomium* in Iran, a phylogenetic study including six new species. *Mycologia* 103: 863-882.
- BISWAS, S. K., K. D. SRIVASTAVA, R. AGGARWAL, P. DUREJA & D. V. SINGH. 2000. Antagonism of *Chaetomium globosum* to *Drechslera sorokiniana*, the spot blotch pathogen of wheat. *Indian Phytopath* 53: 436-440.
- BORNET, B. & M. BRANCHARD. 2001. Nonanchored Inter Simple Sequence Repeat (ISSR) markers: reproducible and specific tools for genome fingerprinting. *Plant Mol Biol Rep* 19: 209-215.

- BREINHOLT, J., H. DEMUTH, M. HEIDE, G. W. JENSEN, M. MØLLER, R. I. NIELSEN, C. E. OLSEN & C. N. ROSENDAHL. 1996. Prenisatin (5-(3-methyl-2-butenyl)-indole-2,3-dione): an antifungal isatin derivative from *Chaetomium globosum*. *Acta Chem Scand* 50: 443-445.
- BREWER, D., J. M. DUNCAN, W. A. JERRAM, C. K. LEACH, S. SAFE S & A. TAYLOR. 1972. Ovine ill-thrift in Nova Scotia 5. The production and toxicology of chaetomium, metabolites of *Chaetomium* spp. *Can J Microbiol* 18: 1129-1137.
- BREWER, D., W. A. JERRAM, D. MEILER & A. TAYLOR. 1970. The toxicity of cochliodinol, and antibiotic metabolite of *Chaetomium* spp. *Can J Microbiol* 16: 433-440.
- BREWER, D. & A. TAYLOR. 1978. The production of toxic metabolites by *Chaetomium* spp. isolated from soil of permanent pasture. *Can J Microbiol* 24: 1082-1086.
- CARMONA, M. A. 2009. Enfermedades causadas por hongos necrotrofos. Manual de Identificación y manejo de las enfermedades del cultivo de la Cebada. 14-18.
- CARMONA, M. A., D. E. BARRETO & E. M. REIS. 1999. Detection, transmission and control of *Drechslera teres* in barley seed. *Seed Sci Technol* 27:761-769.
- CHENNA, R., H. SUGAWARA, T. KOIKE, R. LOPEZ, T. J. GIBSON, D. G. HIGGINS & J. D. THOMPSON. 2003. Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res* 31:3497-3500.
- CULLEN, D. & J. H. ANDREWS. 1984. Evidence for the role of antibiosis in the antagonism of *Chaetomium globosum* to the apple scab pathogen, *Venturia inaequalis*. *Can J Bot* 62: 1819-1823.
- CURTIS, M. D., J. GORE & R. P. OLIVER. 1994. The phylogeny of the tomato leaf mold fungus *Cladosporium fulvum* syn. *Fulvia fulva* by analysis of rDNA sequences. *Current Genet* 25: 318-322.
- DAL BELLO, G. M., C. I. MÓNACO & M. N. SISTERNA. 1994. Efecto de *Trichoderma* spp. sobre el control del tizón de la plántula en trigo ocasionada por *Bipolaris sorokiniana*. *Fitopatol Bras* 19: 394-400.
- DI PIETRO, A., M. GUT-RELLA, J. P. PACHLATKE & F. J. SCHWIN. 1992. Role of antibiotics by *Chaetomium globosum* in biological control of *Pythium ultimum*, a causal agent of damping off. *Phytopathology* 82: 131-135.
- DOMSCH, K. H., W. GAMS & T-H. ANDRESON. 2007. Compendium of soil fungi. vols 1 y 2. Academic Press, London.
- DOYLE, J. J. & J. L. DOYLE. 1990. Isolation of plant DNA from fresh tissue. *Focus* 12: 13-15.
- DREYFUSS, M. 1976. Taxonomische Untersuchungen innerhalb der Gattung *Chaetomium*. *Sydowia* 28:50-133.
- ELLIS, M. B. 1971. Dematiaceous Hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey.
- ELLIS, M. B. 1976. More Dematiaceous Hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey.
- ENGSTROM, K., S. BRISHAMMAR, C. H. SVENSSON, M. BENGTTSSON & R. ANDERSSON. 1993. Anthraquinones from some *Drechslera* species and *Bipolaris sorokiniana*. *Mycol. Res.* 97: 381-384.
- FERNANDEZ VALIELA, M. V. 1978. Introducción a la Fitopatología. Vol. III: Hongos. 3ª. Ed. Colección Científica del INTA. Buenos Aires.
- GAO, K., X. LIU, Z. KANG & K. MENDGEN. 2005. Mycoparasitism of *Rhizoctonia solani* by Endophytic *Chaetomium spirale* ND35: Ultrastructure and Cytochemistry of the Interaction. *J Phytopathol* 153: 280-290.
- GOODWIN, S. B., L. D. DUNKLE & V. L. ZISMANN. 2001. Phylogenetic analysis of *Cercospora* and *Mycosphaerella* based on the internal transcribed spacer region of ribosomal DNA. *Phytopathology* 91: 648- 658.
- HUBBARD, J. P., G. E. HARMAN & C. J. ECKENRODE. 1982. Interaction of a biological control agent *Chaetomium globosum* with seed coat mycoflora. *Canadian J Microbiol* 28: 431-437.
- INFANTE, D., B. MARTÍNEZ, N. GONZALEZ & Y. REYES. 2009. Mecanismos de acción de *Trichoderma* frente a hongos fitopatógenos. *Rev. Protección Veg.* 24: 14-21.
- ISTIFADAH, N. & P. A. MCGEE. 2006. Endophytic *Chaetomium globosum* reduces development of tan spot in wheat caused by *Pyrenophora tritici-repentis*. *Australas Plant Path*35: 411-418.
- ISTIFADAH, N., J. SALEEBA & P. MCGEE. 2006. Isolates of endophytic *Chaetomium* spp. inhibit the fungal pathogen *Pyrenophora tritici-repentis* in vitro. *Can J Bot* 84: 1148- 1155.
- LEE, S. J. & R. T. HANLIN. 1999. Phylogenetic relationships of *Chaetomium* and similar genera based on ribosomal DNA sequences. *Mycologia* 91: 434-442.
- MANDAL, S, K. D. SRIVASTAVA, R. AGGARWAL & D. V. SINGH. 1999. Mycoparasitic action of some fungi on spot blotch pathogen *Drechslera sorokiniana* of wheat. *Indian Phytopath* 52: 39-43
- MAGyP. Manual de estimaciones agrícolas. Diciembre de 2014. Disponible en http://www.minagri.gov.ar/dimeagro/indicadores/ofere-dem-int/2014/01_ofere_dem_int_jul2014.pdf. [Acceso: 12 de abril de 2015].
- MANIATIS, T., E. FRITSCH & J. SAMBROOK. 1982. Molecular Cloning. A Laboratory Manual. New York: Cold Spring Harbor Laboratory.
- MOYA, P. & M. N. SISTERNA. 2012. Estudios preliminares de Biocontrol de “Mancha en red” de la cebada en semilla, con cepas de *Trichoderma* spp. Ensayos “in vitro” *Revista Sanidad de Semillas* 2: 49-49.

- MOYA, P., C. CARRIÓN & M. N. SISTERNA. 2015. Caracterización de pigmentos formados en la interacción fúngica *Drechslera teres* - *Trichoderma* spp. XXXV Jornadas Argentinas de Botánica, 23 al 26 de septiembre de 2015. Salta, Argentina.
- NEERGAARD, P. 1979. Seed Pathology. vol. I and II. Revised Edition, Mac Millan Press, London.
- NAKASHIMA, N., Z. MOROMIZATO & N. MATSUYAMA. 1991. *Chaetomium* spp., antagonistic microorganisms phytopathogenic fungi. *Journal of the Faculty of Agriculture*, Kyushu University 36: 109-115.
- RAYNER, R. W. 1970. *A mycological color chart*. Commonwealth Mycological Institute and British Mycological Society. Kew.
- RODRÍGUEZ MOREJÓN, K. C. 2003. Estudio taxonómico (morfológico y molecular) de especies del género *Chaetomium* y géneros afines. Tesis doctoral. Facultad de Medicina y Ciencias de la Salud, Universidad Rovira y Virgili, Cataluña.
- SETH, H. K. 1970. A monograph of the genus *Chaetomium*. *Beih. Nova Hedwig* 37:1-133.
- SISTERNA, M. N. 2014. Mancha borrosa del trigo. En: Cordo, C. & M. Sистерна (eds.), *Enfermedades del trigo: avances científicos en la Argentina*, pp. 173-186. Editorial de la Universidad de La Plata. La Plata.
- SIVANESAN, A. 1987. Graminicolous species of *Bipolaris*, *Curvularia*, *Drechslera*, *Exserohilum* and their teleomorphs. *Mycological Papers* 158:1-261.
- SÖRGE, G. 1960. Zum Problem der Trennung von Arten bei Pilzen, dargestellt am Beispiel der Ascomycetengattung *Chaetomium*. *Arch Mikrobiol* 36:51-66.
- SOYTONG, K. 1991. Antagonism of *Chaetomium globosum* to leaf spot disease of corn (*Zea mays*). Proceedings of the 29th Kasetsart University pp. 269-275. Plant Science, Kasetsart Univ., Bangkok (Thailand)- Bangkok.
- SOYTONG, K. 1992. Biological Control of rice blast disease by seed coating with antagonistic fungi. *Songklanakarin Journal of Science and Technology* 14: 59-65.
- SOYTONG, K., S. KANOKMEDHAKUL, V. KUKONGVIRIYAPA & M. ISOBE. 2001. Application of *Chaetomium* species (*Ketomium*®) as a new broad spectrum biological fungicide for plant disease control: A review article. *Fungal Divers*. 7: 1-15.
- TAMURA, K., D. PETERSON, N. PETERSON, M. NEI & S. KUMAR. 2011. MEGA5: Molecular Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol. Biol. Evol.* 28: 2731-2739.
- TATHAN, S., P. SIBOUNNAVONG, K. SOYTONG & C. TO-ANUN. 2012. Biological metabolites from *Chaetomium* spp. to inhibit *Drechslera oryzae* causing leaf spot of rice. *Int J Agric Tech* 8: 1691-1701.
- THOMPSON, J. D., T. J. GIBSON, F. PLEWNIAK, F. JEANMOUGIN & D. G. HIGGINS. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids. Res.* 25: 4876-4882.
- TVEIT, M. & R. K. S. WOOD. 1955. The control of *Fusarium* blight in oat seedling with antagonistic species of *Chaetomium*. *Ann Appl Biol* 43: 538-552.
- UNTEREINER, W. A., V. DÉBOIS & F. A. NAVEAU. 2001. Molecular systematics of the ascomycete genus *Farrowia* (Chaetomiaceae). *Can J Bot* 79:321-333.
- VANNACCI, G. & G. E. HARMAN. 1987. Biocontrol of seed-borne *Alternaria raphani* and *A. brassicicola*. *Can J Microbiol* 33: 850-856.
- VILA-AIUB, M. M., M. MARTÍNEZ-GHERSA & C. GHERSA. 2003. Evolution of herbicide resistance in weeds: vertically transmitted fungal endophytes as genetic entities. *Evol Ecol* 17: 441-456.
- VON ARX, J. A., J. GUARRO, M. J. FIGUERAS. 1986. The ascomycete genus *Chaetomium*. *Beih Nova Hedwig* 84:1-162.
- WAKULIŃSKI, W., P. KACHLICKI, P. SOBICZEWSKI, M. SCHOLLENBERGER, C. Z. ZAMORSKI, B. LOTOCKA & J. SAROVA. 2003. Catenarin production by isolates of *Pyrenophora tritici-repentis* (Died.) Drechsler and its antimicrobial activity. *J Phytopathol* 151: 74-79.
- WALLER, F., B. ACHATZ, H. BALTRUSCHAT, J. FODOR, K. BECKER, M. FISCHER, T. HEIER, R. HÜCKELHOVEN, C. NEUMANN, D. WETTSTEIN, P. FRANKEN & K. KOGEL. 2005. The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance, and higher yield. *Proc. Natl. Acad. Sci.* 102: 13386-13391.
- WHITE, T. J., T. BRUNS, S. LEE & J. TAYLOR. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis, M. A., Gelfand, D. H., Sninsky, J. J. & T. J. White (eds.), *PCR Protocols: A Guide to Methods and Applications*, pp. 315-322. Academic Press, Inc., New York.
- WIESE, M. V. 1987. *Compendium of wheat diseases*. American Phytopathological Society: St Paul.

Recibido el 16 de diciembre de 2015, aceptado el 19 de mayo de 2016.