



## Mid Atlantic Regional Chapter of the American College of Sports Medicine

Annual Scientific Meeting, November 1<sup>st</sup> – 2<sup>nd</sup>, 2019  
Conference Proceedings

International Journal of Exercise Science, Volume 9, Issue 8



### Development of a Time Efficient Protocol for Cross-Limb Comparisons of Muscle Mitochondrial Capacity Using NIRS

Rewais Hanna<sup>1</sup>, Jigar Gosalia<sup>1</sup>, Zachary Hobson<sup>1</sup>, Jocelyn Delgado<sup>1</sup>, Alaina Demalis<sup>1</sup>, Kevin McCully<sup>2</sup>, Brian Irving<sup>3</sup>, Swapan Mookerjee<sup>4</sup>, Giampietro Vairo<sup>1</sup> and David Proctor<sup>1</sup>. <sup>1</sup>Penn State University, University Park, PA, <sup>2</sup>University of Georgia, Athens, GA, <sup>3</sup>Louisiana State University, Baton Rouge, LA, <sup>4</sup>Bloomsburg University, Bloomsburg, PA

The non-invasive determination of muscle oxidative capacity via Near Infrared Spectroscopy (NIRS) typically involves voluntary contraction of a single limb and requires as many as 22 brief ischemic occlusions per measurement. This limits the number of oxidative capacity measurements that can be completed in a given test session and also makes cross-limb muscle comparisons challenging. **PURPOSE:** To establish the efficacy of a recently developed protocol that utilizes fewer (i.e. 6) ischemic occlusions combined with surface electrical stimulation (E-stim) in both limbs simultaneously. **METHODS:** The test employs 2 upper thigh cuffs and 2 NIRS sensors placed directly over the vastus lateralis (VL) muscles (supine position) or the semi-tendinosis (ST) muscles (prone position). Metabolic rate is temporarily increased via E-stim pads placed above and below each NIRS sensor. A standard 6Hz frequency is employed using a pre-modulation setting, with the intensity (mV) increased sufficient to raise metabolic rate ( $\geq 3$  fold), but within the tolerance of each participant. The mitochondrial capacity protocol involves 4 separate sets of 30 sec of E-stim followed by 6 x 5 sec cuff inflation/5 sec cuff deflation cycles. Analysis consists of calculating oxygenation recovery rate constants (Tc) for each muscle (i.e., 4 repeated measurements per muscle, per limb) using a customized software program. **RESULTS:** In preliminary tests of moderately active younger adults, Tc ranged from 24 to 44 sec in the VL and 32 to 53 sec in the ST. Variability of repeated tests (CV%) averaged <10% (range 6.2-17.2%) in both muscles. Metabolic rate increased from pre- (slope = -0.011) to post- (slope = -0.018) stimulation. **CONCLUSION:** This bilateral E-stim protocol is time efficient and should facilitate cross-limb comparisons of muscle mitochondrial capacity.