



Mid Atlantic Regional Chapter of the American College of Sports Medicine

Annual Scientific Meeting, November 1st – 2nd, 2019
Conference Proceedings

International Journal of Exercise Science, Volume 9, Issue 8



Lower Body Kinematics Do Not Differ Between Flat Ground and Mound Baseball Throwing

Joshua P. Perez¹, Maryellen M. Crain¹, Andrew C. Venezia¹, Bryon C. Applequist²

¹The University of Scranton, Scranton, PA, ²The University of Hawaii at Hilo, Hilo, HI

It is well known that a throwing program is essential for baseball pitchers to build up their arm strength and stamina. Many research studies have been conducted to determine the best way to perform a throwing program, but no definitive answer for how to properly conduct such a program has been determined. Many programs are designed and implemented on flat ground, however, a training program utilizing a mound enables a pitcher to push off and generate downward force prior to throwing the ball and provides a better transfer to real game situations. Long tossing from a mound has the potential to simulate in-game throwing and could better prepare a pitcher to pitch during a game. **PURPOSE:** The purpose of the study was to determine if lower body kinematics differed between throwing off flat ground and throwing off a mound. **METHODS:** Six healthy individuals ($20Y \pm 1.26$) with previous pitching experience were recruited for this study. Subjects had 38 retro-reflective markers placed in various anatomical locations to quantify lower extremity kinematics during the throwing motion using a motion capture system. Subjects completed 10 total throws, five from flat ground, and five from a mound at a throwing distance of 67 meters. Pelvic rotation angular velocity, trunk rotation angular velocity, and stride length were calculated and analyzed with PitchTrak software. A dependent t-test was used to compare the flat ground and mound conditions for each dependent variable. **RESULTS:** There were no differences between the flat ground and mound throwing conditions for pelvic rotation angular velocity (Flat Ground 715.17 ± 306.88 , Mound 640.97 ± 155.80 , $p > .05$), trunk rotation angular velocity (Flat Ground 888.63 ± 84.57 , Mound 857.33 ± 120.45 , $p > .05$), and stride length (Flat Ground 73.47 ± 9.50 , Mound 73.83 ± 10.32 , $p > .05$). **CONCLUSION:** Lower body kinematics did not differ between long tossing off a mound or flat ground. Due to this, coaches may implement flat ground throwing programs without any negative consequences to pitching mechanics. Consequently, they may also implement throwing programs from a mound to simulate pitching in a game without any increased risk of potential injury.

Supported by Research as a High Impact Practice Grant 841024