brought to you by 🎚 CORE

133

MAURI ORA, 1982, 10; 133-137

ADDITIONAL RECORDS OF MARINE FISHES FROM KAIKOURA

CLIVE D. ROBERTS and J. VAN BERKEL 2

- Department of Zoology, Victoria University of Wellington Wellington, New Zealand
 - 2. Edward Percival Field Station, Department of Zoology University of Canterbury, Kaikoura, New Zealand

ABSTRACT

Twenty-nine marine fish species are listed with information on reference specimens, details of capture, habitat and biological observations. Ten species are recorded for the first time from the Kaikoura area and the occurrence of two species, previously reported as unverified records, is confirmed.

KEYWORDS: fish, marine, new records, distribution, Kaikoura.

INTRODUCTION

In 1979, Francis published a check-list of the marine fishes of Kaikoura recording 179 species. This paper presents additional records from Kaikoura compiled since 1979.

Specimens listed here were all obtained onboard commercial fishing vessels or were brought to the Edward Percival Field Station by local fishermen. They were identified and either preserved and placed in a curated fish collection, or photographed and discarded if badly damaged or very large. Nineteen of these species have been recorded by Francis (1979) and are included here because of the valuable data accompanying each record.

Classification used in the species list follows Greenwood et al. (1966) for the teleosts and format follows that used by Francis (1979). Unless stated otherwise, fish were captured by commercial set net. In addition, information is presented on depth, location and date of capture, body size, sex and reproductive state where available.

SPECIES LIST

Abbreviations used in this species list are as follows:

- species recorded for the first time from the Kaikoura area - Edward Percival Field Station, Kaikoura EPFS - National Museum of New Zealand, Wellington NMNZ - identification confirmed by; D.A. Robertson (DAR), id. P.J. McMillan (PJM), C.D. Paulin (CDP), J. Moreland (JM) or J.A.F. Garrick (JAFG). - specimen(s) spec. FT. - Fork Length - Total Length TLF - Female - Male the "Hole" - sea area about 5 km southwest of South Bay, Kaikoura

the "Front" - sea area off the seaward end of the Kaikoura Peninsula the "Island" - the northern Conway Ridge, 15 km southeast of the Haumuri Bluffs.

Galeorhinus australis (Macleay, 1881) Triakidae school or grey shark EPFS KA 509B 1 juvenile TL 52 cm, 180 m, Goose Bay 2.7.82.

Isurus oxyrinchus Rafinesque, 1810 Lamnidae mako shark 1 spec. 160 m, the Hole

1.7.80.

*Echinorhinus cookei Pietschmann, 1928 Echinorhinidae

Cook's bramble shark EPFS KA503M photo id.

JAFG 1 spec. 120 m, Haumuri Bluffs

11.12.79.

Narkidae Typhlonarke aysoni (Hamilton, 1902)

blind electric ray EPFS KA508R id. JAFG 1

spec. Kaikoura 1982.

*Harriota raleighana Goode & Bean, 1895 Rhinochimaeridae

long-nosed ghost shark EPFS KA508K id. JAFG 1 spec. 300m, off South Bay 3.11.81.

Chauliodontidae *Chauliodus sloanei Bloch & Schneider, 1801

viperfish EPFS KA503G id. JM l spec. in "bonita" stomach, off Kaikoura 17.7.79.

Alepisauridae Alepisaurus brevirostris Gibbs, 1960

lancet fish EPFS KA508A l damaged spec. intertidal, Goose Bay 5.10.81; 1 spec., Kaikoura 12.10.81.

Myctophidae Lampanyctodes hectoris (Gunther, 1876)

lantern fish EPFS KA508U id. DAR, 1 spec.,

2 spec. washed up dead, South Bay 24 to

31.5.82.

Moridae

Pseudophycis barbata Gunther, 1863 Southern bastard red cod NMNZ 8031 id. CDP 1 spec. 110-150m trawl, off Kaikoura Peninsula 11.7.77. (= P. breviusculus in Francis. 1979: 66).

Macrouridae

*Coelorinchus bollonsi McCann & McKnight 1980 rat-tail EPFS KA508Z, KA509A id. PJM 2 spec. 220m, Goose Bay, 3.7.82; several spec. id. PJM 180-220m, Goose Bay 30.6.82 and 2.7.82.

*Coelorinchus biclinozonalis Arai & McMillan, 1982 rat-tail EPFS FA509C id. PJM 1 spec. 150-250m, the Front 24.10.80; 2 spec. id. PJM 150m, Goose Bay 30.6.82.

*Coelorinchus oliverianus Phillipps, 1927 rat-tail EPFS KA509J id. PJM l spec. in hoki stomach 150m, Goose Bay 3.7.81.

Lepidorhynchus denticulatus (Richardson, 1846) javelin fish EPFS KA504J id. PJM l spec., off South Bay 22.11.80.

Oreosomatidae

*Neocyttus rhomboidalis Gilchrist, 1906 spikey oreo dory EPFS KA505R id. PJM 1 spec. 360-430 m, the Island 29.7.81.

Trachipteridae

*Desmodema jacksonensis Ramsay, 1881 ribbon fish EPFS KA505N photo id. CDP 1 spec. FL. 171 cm F. 110-165 m, the Hole 19.6.81.

Apogonidae

Epigonus telescopus (Risso, 1810) black cardinal fish EPFS KA508L and photo id. CDP l spec. 370 m, off Kaikoura 29.12.81.

Carangidae

Trachurus declivis (Jenyns, 1842) jack mackerel EPFS KA509D id. DAR 1 spec. 180 m, Goose Bay 2.7.81.

Naucrates ductor (Linnaeus, 1758) pilot fish EPFS KA508Q id. CDP l spec. 60-100 m, off Haumuri Bluffs -1.82.

Bramidae

Brama brama (Bonnaterre, 1788)
Ray's bream NMNZ 9660 id. CDP 2 spec. FL.
56, 57 cm 100 m, the Hole 19.7.80; 2 spec.,
off South Bay 22.12.80; 2 spec. FL. 63 cm
M., FL. 55 cm F. 150 m, the Hole 25.6.81;
EPFS KA505P and photo 1 spec. FL. 58cm
100 m, the Hole 26.6.81.

Emmelichthyidae

Emmelichthys nitidus Richardson, 1845 red baitfish EPFS KA503U 1 spec. 55 m, Goose Bay 8.1.80; EPFS KA508S photo 1 spec., the Hole 15.2.82.

Pentacerotidae

*Paristiopterus labiosus (Gunther, 1871) giant boarfish EPFS KA505Q l spec. FL. 41 cm 30 m, Goose Bay 15.5.81; EPFS KA508X and photo l spec., Kaikoura 16.6.82.

Latridae

Mendosoma lineatum Guichenot, 1849 telescope fish NMNZ 9659 id. CDP 1 spec. 18 m, off South Bay 28.10.80.

Percophididae

Hemerocoetes monopterygius (Bloch & Schneider, 1801) opal fish NMNZ 9658 id. CDP 1 spec. in stargazer stomach 110-130 m, the Hole 19.7.80.

Scombridae

Thunnus maccoyi Castelnau, 1872 southern bluefin tuna 1 spec. 98 kg 130 m, the Hole 1.7.80; 1 spec., the Hole 30.6.81.

Gasterochisma melampus Richardson, 1845 butterfly tuna EPFS KA508P id. CDP 1 spec. of several in school of kawhai seine net, South Bay 19.1.82.

Centrolophidae

Seriolella brama (Gunther, 1860) common or blue warehou NMNZ 12214 3 juveniles FL 2.9, 5.8, 5.9 cm in school of 1 dozen under floating rope, Goose Bay 28.4.80; EPFS KA509E photo 1 spec., FL 64 m F. (ovary 9.3% body weight) 120 m, the Hole 30.6.80.

Seriolella punctata (Bloch & Schneider, 1801) silver warehou EPFS KA509F photo 1 spec. FL 57 cm F., (ovary 14.6% body weight) 150-250 m, the Front 25.10.80.

*Seriolella caerulea Guichenot, 1848 white warehou EPFS KA509G photo 1 spec. FL 55 cm M. (undeveloped) 180 m, the Front 25.10.80; several spec. 150-250 m, the Front 20-30.10.80.

Hyperoglyphe antarctica (Carmichael, 1818) stoney-eye EPFS KA509H photo 1 spec. FL 50.5 cm 150-250 m, the Front 25.10.80; 1 spec. FL 96 cm F. (spent) 200 m, Goose Bay 28.6.81.

DISCUSSION

Records of 29 species of marine fish are listed. Ten species are recorded for the first time and bring the total number of species recorded from the Kaikoura area to 189. None of the ten new records are of fish considered to be rare. The occurrence of Isurus oxyrinchus and Echinorhinus sp., previously listed as unverified by Francis, is confirmed. On-going taxonomic revisions (e.g. Macrouridae) and increased sampling effort, particularly in water of over 100 m, account for most of the new species listed.

ACKNOWLEDGEMENTS

We thank the following for help with identification:
D.A. Robertson and P.J. McMillan (Fisheries Research Division,
MAF, Wellington), C.D. Paulin and J. Moreland (National Museum
of New Zealand, Wellington) and J.A.F. Garrick (Victoria
University of Wellington). We are most grateful for the help
and co-operation given to us by local fishermen and in particular
we thank B. Fearnley, W. Jones, T. Forbes, R. Huggins and
R. Reinke. During the course of this work one of us (CDR) was in
receipt of a Victoria University Grants Committee travel grant.

LITERATURE CITED

FRANCIS, M.P. 1979. Checklist of the marine fishes of Kaikoura, New Zealand.

Mauri Ora, 7: 63-71.

GREENWOOD, P.H., D.E. ROSEN, S.H. WEITZMAN and G.S. MYERS, 1966.

Phyletic studies of teleostean fishes, with a provisional classification of living forms. Bulletin of the American Museum of Natural History, 131: 339-456.