

MODELS OF ROSETTA/OSIRIS 67P DUST COMA PHASE FUNCTION

F. Moreno

*Instituto de Astrofísica de Andalucía, CSIC, c/ Glorieta de la Astronomía s/n, 18008
Granada, Spain*

fernando@iaa.es

D. Guirado

*Instituto de Astrofísica de Andalucía, CSIC, c/ Glorieta de la Astronomía s/n, 18008
Granada, Spain*

O. Muñoz

*Instituto de Astrofísica de Andalucía, CSIC, c/ Glorieta de la Astronomía s/n, 18008
Granada, Spain*

I. Bertini

*Department of Physics and Astronomy 'G. Galilei', University of Padova, Vicolo dell'
Osservatorio 3, I-35122 Padova, Italy*

C. Tubiana

*Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg, 3, D-37077
Göttingen, Germany*

C. Güttler

*Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg, 3, D-37077
Göttingen, Germany*

M. Fulle

Osservatorio Astronomico, Via Tiepolo 11, I-34143 Trieste, Italy

A. Rotundi

*INAF - Istituto di Astrofisica e Planetologia Spaziali, Via Fosso del Cavaliere, 100, I-00133
Rome, Italy*

*Università degli Studi di Napoli Parthenope, Dipdi Scienze e Tecnologie, CDN IC4,
I-80143 Naples, Italy*

V. Della Corte

*INAF - Istituto di Astrofisica e Planetologia Spaziali, Via Fosso del Cavaliere, 100, I-00133
Rome, Italy*

*Universit' a degli Studi di Napoli Parthenope, Dipdi Scienze e Tecnologie, CDN IC4,
I-80143 Naples, Italy*

S.L. Ivanovski

*INAF - Istituto di Astrofisica e Planetologia Spaziali, Via Fosso delCavaliere, 100, I-00133
Rome, Italy*

G. Rinaldi

*INAF - Istituto di Astrofisica e Planetologia Spaziali, Via Fosso delCavaliere, 100, I-00133
Rome, Italy*

D. Bockel' ee-Morvan

*LESIA, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universit' es,
UPMC Univ. Paris 06, Univ. Paris-Diderot, Sorbonne Paris Cit' e, 5 place Jules Janssen,
F-92195 Meudon, France*

V.V. Zakharov

*Laboratoire de M' et' eorologie Dynamique, UPMC, Sorbonne Universit' es, Paris, France
INAF - Istituto di Astrofisica e Planetologia Spaziali, Via Fosso delCavaliere, 100, I-00133
Rome, Italy*

J. Agarwal

*Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg, 3, D-37077
G' ottingen, Germany*

S. Mottola

*Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Planetenforschung,
Rutherfordstraße 23, 12489 Berlin, Germany*

I. Toth

*MTA CSFK Konkoly Observatory of the Hungarian Academy of Sciences, Konkoly Thege
M. ut 15-17. H1121 Budapest, Hungary*

E. Frattin

*Department of Physics and Astronomy 'G. Galilei', University of Padova, Vicolo dell'
Osservatorio 3, I-35122 Padova, Italy*

L.M. Lara

*Instituto de Astrof' isica de Andaluc' ia, CSIC, c/ Glorieta de la Astronom' ia s/n, 18008
Granada, Spain*

P.J. Gutiérrez

*Instituto de Astrofísica de Andalucía, CSIC, c/ Glorieta de la Astronomía s/n, 18008
Granada, Spain*

Z.Y. Lin

*National Central University No.300 Jhongda Rd., Jhongli city, Taoyuan County, 320
Taiwan*

L. Kolokolova

*Planetary Data System Group, Department of Astronomy, Rm1207D, Atlantic Bldg.,
University of Maryland, College Park, MD, 20742, USA*

H. Sierks

*Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077
Göttingen, Germany*

G. Naletto

*University of Padova, Department of Physics and Astronomy “Galileo Galilei”, Via
Marzolo 8, 35131 Padova, Italy*

*University of Padova, Center of Studies and Activities for Space (CISAS) “G. Colombo”,
Via Venezia 15, 35131 Padova, Italy*

CNR-IFN UOS Padova LUXOR, Via Trasea 7, 35131 Padova, Italy

P. L. Lamy

*Laboratoire Atmosphères, Milieux et Observations Spatiales, CNRS & Université de
Versailles Saint-Quentin-en-Yvelines, 11 Boulevard d'Alembert, 78280 Guyancourt, France*

R. Rodrigo

*Centro de Astrobiología, CSIC-INTA, 28850 Torrejón de Ardoz, Madrid, Spain
International Space Science Institute, Hallerstrasse 6, 3012 Bern, Switzerland*

D. Koschny

*Science Support Office, European Space Research and Technology Centre/ESA, Keplerlaan
1, Postbus 299, 2201 AZ Noordwijk ZH, The Netherland*

B. Davidsson

*Jet Propulsion Laboratory, M/S 183-401, 4800 Oak Grove Drive, Pasadena, CA 91109,
USA*

M. A. Barucci

*LESIA, Observatoire de Paris, PSL Research University, CNRS, Univ.Paris Diderot,
Sorbonne Paris Cité, UPMC Univ.Paris 06, Sorbonne Universités, 5 place Jules Janssen,
92195 Meudon, France*

J.-L. Bertaux

*Laboratoire Atmosphères, Milieux et Observations Spatiales, CNRS & Université de
Versailles Saint-Quentin-en-Yvelines, 11 Boulevard d'Alembert, 78280 Guyancourt, France*

D. Bodewits

Department of Astronomy, University of Maryland, College Park, MD 20742-2421, USA

G. Cremonese

*INAF, Astronomical Observatory of Padova, Vicolo dell'Osservatorio 5, 35122 Padova,
Italy*

V. Da Deppo

CNR-IFN UOS Padova LUXOR, Via Trasea 7, 35131 Padova, Italy

S. Debei

*University of Padova, Department of Industrial Engineering, Via Venezia 1, 35131 Padova,
Italy*

M. De Cecco

University of Trento, Faculty of Engineering, Via Mesiano 77, 38121 Trento, Italy

J. Deller

*Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077
Göttingen, Germany*

S. Fornasier

*LESIA, Observatoire de Paris, PSL Research University, CNRS, Univ.Paris Diderot,
Sorbonne Paris Cité, UPMC Univ.Paris 06, Sorbonne Universités, 5 place Jules Janssen,
92195 Meudon, France*

W.-H. Ip

*Graduate Institute of Astronomy, National Central University, 300 Chung-Da Rd,
Chung-Li 32054, Taiwan*

*Space Science Institute, Macau University of Science and Technology, Avenida Wai Long,
Taipa, Macau*

H. U. Keller

*Institut für Geophysik und extraterrestrische Physik, Technische Universität Braunschweig,
Mendelssohnstr.3, 38106 Braunschweig, Germany*

*Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Planetenforschung,
Rutherfordstraße 23, 12489 Berlin, Germany*

M. Lazzarin

*University of Padova, Department of Physics and Astronomy “Galileo Galilei”, Vicolo
dell’Osservatorio 3, 35122 Padova, Italy*

J. J. López-Moreno

*Instituto de Astrofísica de Andalucía (CSIC), c/ Glorieta de la Astronomía s/n, 18008
Granada, Spain*

F. Marzari

*University of Padova, Department of Physics and Astronomy “Galileo Galilei”, Via
Marzolo 8, 35131 Padova, Italy*

and

X. Shi

*Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077
Göttingen, Germany*

ABSTRACT

The phase function of the dust coma of comet 67P has been determined from Rosetta/OSIRIS images (Bertini et al. 2017). This function shows a deep minimum at phase angles near 100° and a strong backscattering enhancement. These two properties cannot be reproduced by regular models of cometary dust, most of them based on wavelength-sized and randomly-oriented aggregate particles. We show, however, that an ensemble of oriented elongated particles of a wide variety of aspect ratios, with radii $r \gtrsim 10 \mu\text{m}$, and whose long axes are perpendicular to the direction of the solar radiation, are capable of reproducing the observed

phase function. These particles must be absorbing, with an imaginary part of the refractive index of about 0.1 to match the expected geometric albedo, and with porosity in the 60-70% range.

Subject headings: Minor planets, asteroids: individual (67P/Churyumov-Gerasimenko)
— Methods: numerical

1. INTRODUCTION

One of the goals of the Rosetta mission was the characterization of the dust environment of comet 67P, with several instruments being devoted to this task. In particular, the dust phase function has been measured in the optical range from images taken by the OSIRIS cameras (Bertini et al. 2017). The phase function was retrieved in a wide range of phase angles in a time interval of about two hours. As these conditions cannot be met from the ground, the OSIRIS measurements are unique. So far, only for a handful of short-period comets observed from Earth the phase function is available, albeit being most of the times restricted to the backscattering domain (see Bertini et al. 2017, and references therein). In addition, for Earth based observations the phase angle dependence is always inherently mixed with temporal and heliocentric distance variability in the coma, so that it is often difficult to disentangle the intrinsic activity from the phase effect. Notwithstanding this, an overall agreement of the OSIRIS phase functions with those derived from the ground is found (Bertini et al. 2017).

2. THE OSIRIS PHASE FUNCTION

Bertini et al. (2017) have retrieved phase functions from OSIRIS WAC and NAC images, at varying heliocentric and nucleocentric distances, with the Orange F22 filter (effective wavelength $\lambda_{eff} = 642.2$ nm), and the Green F21 filter ($\lambda_{eff} = 537.2$ nm) (see their figure 4). During the phase function measurements, the phase angle Sun-comet-spacecraft (i.e. the nucleus elongation) remained close to 90°. The geometry of the observations can be seen in Figure 1b of Bertini et al. (2017).

For the purpose of comparison with models, we only consider the MTP020/071 phase function (obtained near perihelion), which is representative of most phase curves, although we also display in most figures the MTP025/092 curve ($r = 2.18$ au post-perihelion) for completeness. That phase curve is the one that shows the shallowest slope in the backscattering regime (see Figure 1). The corresponding spacecraft (S/C) ranges were 420 km and 80

km, respectively, indicating that the particles along the line of sight of the OSIRIS cameras were well outside the gas acceleration region (at ~ 12 km, see Gerig et al. 2018). The large nucleocentric distances of the S/C (particularly during the perihelion measurements) and the fact that the pointing is directly away from the nucleus position would suggest that the phase function can only be minimally affected by different optical depth along distinct lines of sight. In any case, the true 3D nature of the dust distribution, which would be needed to obtain the optical depth along any line of sight, is unknown, and will only be unveiled by complex coupled gas and dust dynamics modeling (e.g. Crifo 2006; Zakharov et al. 2018, and references therein).

3. MODELING THE PHASE FUNCTION

3.1. MODELS ASSUMING SIZE DISTRIBUTIONS OF RANDOMLY-ORIENTED PARTICLES

All previous models of cometary dust (e.g. Kimura et al. 2003; Bertini et al. 2007; Moreno et al. 2007; Lasue et al. 2009; Kolokolova et al. 2015; Zubko et al. 2016) are built under the hypothesis that the coma is a cloud of particles in random orientations. However, none of those models is able to reproduce the phase function curve derived from the OSIRIS images. In addition, the models based on fluffy, wavelength-sized aggregate particles (e.g. Kimura et al. 2003) typically display a maximum in the degree of linear polarization that is too high in comparison with observations. Kimura et al. (2006) and Kolokolova & Mackowski (2012) explain this fact as a computer limitation associated to the limited amount of monomers, showing that the maximum of polarization decreases as the amount of monomers increase. A rough spheroid model has been introduced by Kolokolova et al. (2015), which consists of a wide size distribution of such spheroids. That model matches much better the observed degree of linear polarization and the color properties but, as it has been stated, the modeled phase functions show only a modest backscattering enhancement and have minima at phase angles much smaller than 100° , otherwise in agreement with the composite phase function curve obtained from a compilation of ground-based observations from various comets by D. Schleicher (<http://asteroid.lowell.edu/comet/dustphase.html>). On the other hand, to explain accurately the observations of linear polarization versus phase angle, models of mixtures of compact and aggregate particles have been introduced by (e.g. Lasue et al. 2009; Das et al. 2011), as well as mixtures of weakly and highly absorbing agglomerated debris particles (Zubko et al. 2016).

All of those previous models assume either monodisperse distributions of wavelength-sized particles, or polydisperse distributions peaking in the submicrometer range. However, there are indications that the dominant scatterers in 67P might be larger, as indicated by the

small amount of submicron and micron-sized particles detected by Rosetta/MIDAS, much less than initially expected (Mannel et al 2017; Güttler et al. 2018). Also, the analysis of thermal spectra of the quiescent coma performed with Rosetta/VIRTIS-H implies a minimum radius of compact particles of $10 \mu\text{m}$ (in combination with 25% of fluffy aggregate particles in number, whose contribution to the scattered light is minimal) (Bockel'ee-Morvan et al. 2017a,b). The study of the evolution of the 67P dust size distribution by Fulle et al. (2016b) confirms that the coma brightness at perihelion and later is in fact dominated by particles $<100 \mu\text{m}$. In addition, the analysis of a large ground-based image dataset by Monte Carlo dust tail modeling (Moreno et al. 2017) agrees with that constraint. This model replicated tail images from the current apparition, from about 4.5 au pre-perihelion to 3 au post-perihelion, as well as trail data from the current and previous orbits. The minimum model particle size was $10 \mu\text{m}$ in all cases.

Given these facts, we started by trying models having randomly-oriented particles of sizes larger than the nominal incident wavelength, assumed at $\lambda=0.6 \mu\text{m}$. For this task, we used available light scattering codes such as the T-matrix code of spheroids (Mishchenko et al. 1996), the Multi-Sphere T-matrix (MSTM) code (Mackowski & Mishchenko 2011), and the geometric optics code by Macke & Mishchenko (1996) for ellipsoids. Except for the geometric optics code, the computing time and memory requirements are a rapidly increasing functions of the size parameter, $X=2\pi r/\lambda$. This imposes strong limits to the simulations, for which we should keep the number of possible combinations of input parameters to a minimum.

One important constraint to the model is the observed geometric albedo. The nucleus geometric albedo at 649 nm is 0.0677 ± 0.0039 (Fornasier et al. 2015), and it is reasonable to assume that large particles in the coma will have the same geometric albedo, provided they do not experience any change in their physical properties after ejection, such as sublimation of volatiles or fragmentation. These processes have been found to be negligible in the Rosetta studies of the 67P's coma (Fulle et al. 2015).

To meet the geometric albedo constraint we must assume absorbing particles. The precise value of the mean refractive index of 67P dust coma particles is unknown. We used a refractive index of $m=1.6+0.1i$, a value comprised between low-absorbing silicates and strongly absorbing organic and carbonaceous materials at red wavelengths (see e.g. Jenniskens 1993). To calculate the scattering pattern of relatively large particles, we used the MSTM code for an array of spherules. Since we are searching for scatterers giving a high backscattering enhancement, we started by following a procedure similar to that devised by Mishchenko et al. (2007). To demonstrate the onset and development of the coherent backscattering mechanism, which is the responsible for the backscattering enhancement,

Mishchenko et al. (2007) considered a certain scattering volume filled with an increasing number of small scatterers. We performed a similar calculation by considering a spherical volume of $r=2 \mu\text{m}$ containing 500, 1000, and 2500 randomly placed $0.1\text{-}\mu\text{m}$ radius spherules, giving a porosity, P , of 93.7%, 87.5%, and 69.0%, respectively (see Figure 1, top panels). To obtain the scattering functions, we assumed that the array of scatterers is illuminated from a given direction, and then compute the azimuthally-averaged scattering matrix, i.e., the average scattering matrix for all the scattering planes about the direction of the incoming beam. Then, for a given number of spherules, we repeated the simulation for a large number of such randomly-generated targets. This procedure is much less CPU time consuming than generating the scattering matrix for a single randomly-oriented scatterer.

The resulting phase functions are displayed in Figure 1. Also shown are the results corresponding to a single homogeneous $r=2 \mu\text{m}$ sphere calculated from Mie theory, just to show the large discrepancies of that simple model with the MSTM models in the phase function. Concerning the MSTM models, we see that as the number of spherules in the volume increases, the backscattering enhancement also increases. In order to check whether this trend is maintained for still lower porosities, we first increase the number of spheres up to ~ 3100 , corresponding to $P = 61\%$, but we obtained very similar results to that of $P = 69\%$. This $P = 61\%$ is essentially the lower limit reachable with our simple random packing procedure. To decrease P we had to use a more complex technique than random packing. Among the techniques available we used a “falling down” algorithm (Fulle & Blum 2017). With this code, we were able to generate spherical volumes having P in the range 55%-45%, i.e., containing some 3600 to 4400 spherules. At these porosity levels, we noticed a reversal in the trend, i.e., a decrease in backscattering enhancement as compared with that found in the $P \sim 60\text{-}70\%$ range. We then conclude that this porosity range is the one that gives the highest backscattering enhancement. We underline that this conclusion is based on the specific spherical volume described above (see Figure 1, top panels). Larger volumes would have been desirable to test but, as stated above, we are strongly constrained by computer memory and CPU time limitations.

It is interesting to note that the porosity range for which the highest backscattering enhancement is observed agrees with the nucleus porosity estimated at $71 \pm 8\%$ by Fulle et al. (2016a), and is also consistent with the value found specifically for the upper layers of the surface, which is below 74% from Philae SESAME/CASSE and MUPUS (Knapmeyer et al. 2018). It is also not far from the Rosetta/MIDAS and Rosetta/GIADA results for relatively compact particles ($\sim 50\%$, see Fulle & Blum 2017).

The following step was to explore other configurations of arrays of spherules of total size larger than the wavelength to test whether an improvement in the fits could be achieved.

We built larger particles by increasing the number of spherical monomers, but owing to the limitations in computer time and memory, we had to combine larger spherical monomers with smaller ones, although we could not find a satisfactory fit in any case. An example of those configurations is shown in Figure 2, in which a cluster of four 2- μm spheres is randomly peppered by a total of 2000 0.1 to 0.15 μm radius spherules having a total equivalent spherical radius (radius of a sphere of equal volume), $r_{eq} \sim 3.4 \mu\text{m}$. This particular arrangement gives an overall better fit to the phase function, mainly for the forward lobe, but fails at reproducing the observed backscattering enhancement.

To test the performance of still larger particles, we had to resort to the geometric optics code. As mentioned, we used the code for ellipsoidal particles by Macke & Mishchenko (1996). We have tested a variety of ellipsoids of different sizes and axes ratios, but the computed phase functions are all flat at backscattering, similarly to large homogeneous spheres. In conclusion, none of the considered models of randomly-oriented particles at a wide range of sizes can reproduce reliably the OSIRIS phase function.

3.2. MODELS ASSUMING SIZE DISTRIBUTIONS OF PARTICLES ALIGNED TO THE SOLAR RADIATION DIRECTION

The presence of aligned particles in the cometary environment has been subject of research for many years, since the early work of Dolginov & Mytrophanov (1976). In the nucleus acceleration region comet gravity, and aerodynamic forces should dominate (Fulle et al. 2015; Ivanovski et al. 2017a). Outside that region, radiative torques, although ignored for many years, might play an important role (see e.g. Draine & Weingartner 1996; Lazarian 2003; Rosenbush et al. 2007). Interestingly, alignment of the particles with respect to the solar radiation will place the long axes of the particles perpendicular to radiation (Rosenbush et al. 2007).

For the radiative torques to be efficient as the alignment mechanism, the characteristic time of alignment should be shorter than the flight time of the particles along the line of sight of the OSIRIS camera Kolokolova et al. (2016), based on a study by Hoang & Lazarian (2014), estimate that for the 67P environment, a 10- μm oblate spheroid would be aligned in $\sim 3 \times 10^5$ s. Assuming a radial trajectory and a constant speed of 10 m s^{-1} (Lin et al. 2016), that particle would become aligned at a distance of 3000 km from the nucleus. Thus, it appears that this mechanism alone would not be sufficient to provide the required alignment, at least for those particles in the vicinity of the S/C. However, in addition to radiative torques, there is another mechanism that could be playing a role, which is the mechanical alignment by gas-particle relative motion (Gold 1952) and aerodynamic force (e.g. Ivanovski

et al. 2017b). The Gold alignment mechanism, which is efficient for supersonic gas flow (as in a cometary coma, see e.g. Zakharov et al. 2018) would be characterized by particles having their longer axes directed preferentially along the gas flow due to the tendency of the particle to minimize its cross section in the gas flow (Rosenbush et al. 2007). The gas flow is radial at distances greater than about 20 km from the nucleus (Marschall et al. 2016). The particles experience this radial gas flow until the gas decoupling distance, at some 20 nuclear radii (i.e., about 30 km). After that, there is no other mechanism that can affect their orientation except radiative torque which produce alignment in the same direction - long axis perpendicular to the radiation, thus, keeping the original gas alignment. This direction is nearly perpendicular to the direction of the solar radiation for particles in the vicinity of the S/C during the perihelion phase function measurements (see Figure 1b by Bertini et al. 2017). Thus, the radiative torque and the Gold mechanism working together might explain the alignment of the particles. However, although a quantitative evaluation of the efficiency of those mechanisms in terms of the physical properties of the particles should be certainly performed, this is beyond the scope of this paper.

To model aligned scatterers, we consider that the particles have their long axes perpendicular to radiation, and calculate the azimuthally averaged scattering matrix. For these simulations, we assumed spheroidal particles, because in this way we can encompass a wide range of sizes with same geometry. The spheroids have axes (a, b, c) , where $a=b>c$ for oblate, and $a>b=c$ for prolate spheroids, respectively. The axis ratio is defined as a/c . Thus, oblate spheroids are set with the c -axis parallel to the solar radiation, and prolate spheroids are set with the a -axis perpendicular to the solar radiation.

Although we know that the dominant scatterers in 67P must be larger than the wavelength, for completeness, we performed simulations for a wider range of sizes, starting from the submicron domain. This will show that the dominant scatterers must be, in fact, much larger. Thus, for distributions of small sized particles, we used the T-matrix method for spheroids by Mishchenko et al. (1996). We start by considering oblate spheroids only, because they provide a higher backscattering enhancement than prolate spheroids for a given r_{eq} . In addition, oblate spheroids might indeed provide a good description of cometary dust particles as has been found from laboratory experiments by Stephens & Gustafson (1991), and recently by Bischoff et al. (2018). Both studies dealt with mixtures of volatile and refractory materials and showed that during the sublimation of volatile materials, the refractory materials form a thin mantle that eventually cracks, being lifted from the surface. Bischoff et al. show that these cracked pieces have a diameter of up to 5 to 10 times larger than the thickness of the mantle, i.e. they are flake-like particles resembling oblate spheroids of large axis ratio.

We generated scattering matrices for size distributions of such spheroidal particles, for which we assumed $q=2$. Figure 3 shows two simulations with such spheroids being distributed according to a differential power law of index -3.5 (a typical value for 67P dust coma), and with three different values of the minimum size, 0.1 , 0.3 , and $1.5 \mu\text{m}$, the maximum size being $r_{\text{max}}=2.5 \mu\text{m}$ in all cases. From Figure 3, it is clear that the larger the sizes the better the agreement with the OSIRIS phase function. On the other end, submicron-dominated size distribution functions fail at fitting the phase function.

Sizes substantially larger than the maximum radius assumed in the previous model of spheroids ($r_{\text{max}}=2.5 \mu\text{m}$ with $q=2$) cannot be efficiently handled with the T-matrix code. To build larger particles, we fill oblate spheroidal volumes with spherules of the order of the nominal wavelength or smaller, as input to the MSTM code, with an average porosity of 65-70%, i.e., near the optimal values found for the $2 \mu\text{m}$ spherical volume described in the previous section. We built oblate spheroidal volumes with axial ratios in the range $q=2-3$, and with equivalent spherical radius of $3.1 \mu\text{m}$, $4.6 \mu\text{m}$, and $7.0 \mu\text{m}$, filled with approximately 1000, 1500, and 4000 spherules, respectively, where the radii of the spherules range from 0.1 to $0.6 \mu\text{m}$. Owing to computing time limitations, the number of realizations for each size was limited to 256, 36, and 20, respectively. Figure 4 shows the resulting phase functions for the three sizes, where we can notice again how the agreement with the typical OSIRIS phase function tends to improve as the particle size increases, in both forward and backward lobes.

So far, only oblate spheroids up to a size of $r_{\text{eq}}=7 \mu\text{m}$ have been considered as model particles. We have to extend the upper size limit in the model to larger sizes, as measured by several instruments on board Rosetta, and inferred from the ground. In addition, we have to extend the particle shape distribution by taking into account a mixture of oblate and prolate particles, as would be required to simulate any natural dust sample. Thus, for a given equivalent spherical radius, we assumed a uniform distribution of axes ratios, i.e., the same amount of prolate and oblate spheroids.

To generate the shape-averaged scattering matrix, we used the individual matrices calculated using the geometric optics code by Macke & Mishchenko (1996). Figure 5 shows a comparison of the results obtained for three equivalent spherical radii ($r_{\text{eq}}=10 \mu\text{m}$, $r_{\text{eq}}=100 \mu\text{m}$, and $r_{\text{eq}}=1000 \mu\text{m}$) with the OSIRIS phase function, for a shape distribution having axes ratios in the interval $q \in [0.25, 4]$. This wide axes ratio distribution is not surprising, in view of the findings by Fulle et al. (2017) who inferred aspect ratios of 5 or more in order to explain the bulk densities of the particles measured by GIADA during the whole mission.

As it is seen, the synthetic phase functions are close at backscattering, but the one for the smaller radius gives a better fit at large phase angles. In addition, the resulting geometric albedo is 0.06, very close to that obtained at the nucleus surface (Fornasier et al.

2015). The inferred particle size is consistent with the ground-based tail modeling, where differential power-law size distributions of index -3 and minimum particle radius of $10 \mu\text{m}$ were found to be consistent with the observations (Moreno et al. 2017). This is also consistent with MIDAS results, where a very small amount of micron-sized particles or smaller were detected (Mannel et al 2017; Giller et al. 2018), and with VIRTIS-H thermal spectra modeling where minimum sizes of relatively compact $10 \mu\text{m}$ particles (combined with 25% fractals in number) following a size distribution of power index -3 or smaller are found (Bockelée-Morvan et al. 2017a,b). It is also important to note that the phase function shape is maintained for up to mm-sized particles in the backscattering lobe, explaining in that way the results obtained by Fulle et al. (2018), in that the OSIRIS phase function should be applied to dust diameters up to 2.5 mm at least for the conversion of dust brightness into dust sizes.

4. CONCLUSIONS

We have shown that regular models of cometary dust based on wavelength-sized and randomly-oriented aggregate particles cannot reproduce the dust phase function determined from Rosetta/OSIRIS observations. Such phase function can, however, be adequately fitted by assuming that the main scatterers in 67P coma are large compared to the visual wavelengths ($>10 \mu\text{m}$), show a wide distribution of aspect ratios, and are aligned with their long axes perpendicular to the solar radiation direction. Porosity levels for the particles of 60-70% would be particularly favoured, since those particles would give the largest backscattering enhancement. For a refractive index of $m=1.6+0.1i$, the geometric albedo for such particles would be ~ 0.06 , i.e., close to that measured for the nucleus surface.

OSIRIS was built by a consortium of the Max-Planck-Institut für Sonnensystemforschung, in Göttingen, Germany, CISAS-University of Padova, Italy, the Laboratoire d'Astrophysique de Marseille, France, the Instituto de Astrofísica de Andalucía, CSIC, Granada, Spain, the Research and Scientific Support Department of the European Space Agency, Noordwijk, The Netherlands, the Instituto Nacional de Técnica Aeroespacial Madrid, Spain, the Universidad Politécnica de Madrid, Spain, the Department of Physics and Astronomy of Uppsala University, Sweden, and the Institut für Datentechnik und Kommunikationsnetze der Technischen Universität Braunschweig, Germany. The support of the national funding agencies of Germany (DLR), France (CNES), Italy (ASI), Spain (MEC), Sweden (SNSB; Grant No. 74/10:2), and the ESA Technical Directorate is gratefully acknowledged. H. Rickman was also supported by Grant No. 2011/01/B/ST9/05442 of the Polish National Science Center. We thank the ESA teams at ESAC, ESOC and ESTEC for their work in support of the

Rosetta mission We thank the Rosetta Science Ground Segment at ESAC, the Rosetta Mission Operations Centre at ESOC, and the Rosetta Project at ESTEC for their outstanding work enabling the science return of the Rosetta Mission.

We thank Evgenij Zubko for his constructive comments on the paper, and to Daniel Mackowski, Michael Mishchenko, and Andreas Macke for making their light scattering codes available.

This work was supported by contracts AYA2015-67152-R and AYA2015-71975-REDT from the Spanish Ministerio de Economía y Competitividad.

REFERENCES

- Bertini, I., Thomas, N., & Barbieri, C., 2007, *A&A*, 461, 351
- Bertini, I., La Forgia, F., Tubiana, C. et al., 2017, *MNRAS*, 469, 404
- Bischoff, D., Gundlach, B., Neuhaus, M. et al., 2018, *MNRAS*, submitted
- Bockel´ee-Morvan, D., Rinaldi, G., Erard, S., et al., 2017a, *MNRAS*, 469, 443
- Bockel´ee-Morvan, D., Rinaldi, G., Erard, S., et al., 2017b, *MNRAS*, 469, 842
- Crifo, J.F., 2006, *Adv. Space Res.*, 38, 1911
- Das, H.S., Paul, D., Suklabaidya, A., et al., 2011, *MNRAS*, 416, 94
- Dolginov, A.Z., & Mytrophanov, I.G., 1976, *Astrophys. Space Sci.*, 43, 291
- Draine, B.T., & Weingartner, J.C., 1996, *ApJ*, 470, 551
- Fornasier, S., Hasselmann, P.H., Barucci, M.A., et al., 2015, *A&A*, 583, 30
- Fulle, M., Ivanovski, S.L., Bertini, I., et al., 2015, *A&A*, 583, A14
- Fulle, M., Della Corte, V., Rotundi, A., et al., 2016a, *MNRAS*, 462, 132
- Fulle, M., Marzari, F., Della Corte, V., et al., 2016b, *ApJ*, 821, 19
- Fulle, M., & Blum, J., 2017, *MNRAS*, 469, 39
- Fulle, M., Della Corte, V., Rotundi, A., et al., 2017, *MNRAS*, 469, 45
- Fulle, M., Bertini, I., Della Corte, V., et al., 2018, *MNRAS*, 476, 2835

- Gerig, S.-B., Marschall, R., Thomas, N., et al., 2018, *Icarus*, 311, 1
- Gold, T., 1952, *MNRAS*, 112, 215
- Güttler, C., et al., 2018, in preparation
- Hoang, T., & Lazarian, A., 2014, *MNRAS*, 438, 680
- Ivanovski, S.L., Della Corte, V., Rotundi, A., et al., 2017a, *MNRAS*, 469, 774
- Ivanovski, S.L., Zakharov, V.V., Della Corte, V., et al., 2017b, *Icarus*, 282, 333
- Jenniskens, P., 1993, *A&A*, 274, 653
- Kimura, H., Kolokolova, L., & Mann, I., 2003, *A&A*, 407, L5
- Kimura, H., Kolokolova, L., & Mann, I., 2006, *A&A*, 449, 1243
- Knapmeyer, M., Fischer, H.-H., Knollenberg, J., et al., 2018, *Icarus*, 310, 165
- Kolokolova, L., & Mackowski, D., 2012, *J. Quant. Spectrosc. Rad. Transfer*, 113, 2567
- Kolokolova, L., Das, H.S., Dubovik, O., et al., 2015, *Planet. Space Sci.*, 116, 30
- Kolokolova, L., Koenders, C., Goetz, C., et al., 2016, *MNRAS*, 462, 422
- Lasue, J., Levasseur-Regourd, A.C., Hadamcik, E., et al. 2009, *Icarus*, 199, 129
- Lazarian, A., 2003, *J. Quant. Spectrosc. Rad. Transfer*, 79, 881
- Levasseur-Regourd, A.C., Hadamcik, E., & Renard, J.B., 1996, *A&A*, 313, 327
- Lin, Z.-Y., Lai, I.L., Su, C.C., et al., 2016, *A&A*, 588, L3
- Macke, A., & Mishchenko, M.I., 1996, *Appl. Optics*, 35, 429
- Mackowski, D.M., & Mishchenko, M.I., 2011, *J. Quant. Spectrosc. Rad. Transfer*, 112, 2182
- Mannel, T., Bentley, M.S., Boakes, P., et al., 2017, *European Planetary Science Congress*,
id. EPSC2017-258
- Marschall, R., Su, C.C., Liao, Y., et al., 2016, *A&A*, 589, A90
- Mishchenko, M.I., Travis, L.D., & Mackowski, D.W., 1996, *J. Quant. Spectrosc. Rad. Transfer*, 55, 535
- Mishchenko, M.I., Liu, L., Mackowski, D.W., et al., 2007, *Opt. Express*, 15, 2822

- Moreno, F., Muñoz, O., Guirado, D., et al., 2007, *J. Quant. Spectrosc. Rad. Transfer*, 106, 348
- Moreno, F., Muñoz, O., Gutiérrez, P.J., et al., 2017, *MNRAS*, 469, 186
- Muñoz, O., Moreno, F., Vargas-Martín, F., et al., 2017, *ApJ*, 846, 85
- Rosenbush, V., Kolokolova, L., Lazarian, A., et al., 2007, *Icarus*, 186, 317
- Stephens, J.R., & Gustafson, B.A.S., 1991, *Icarus*, 94, 209
- Zakharov, V.V., Ivanovski, S.L., Crifo, J.F., et al., 2018, *Icarus*, 312, 121
- Zubko, E., Videen, G., Hines, D.C., et al., 2016, *Planet. Space. Sci.*, 123, 63




Fig. 1.— Backscattering enhancement test. Top three panels: 2- μm radius spherical volumes filled with (left to right) 500, 1000, and 2500 randomly located 0.1- μm radius spherules, leading to porosities of 93.7%, 87.5%, and 69.0%, respectively. XYZ labels are in μm . Bottom panel: OSIRIS phase functions (red solid circles for MTP020/071 and red open circles for MTP025/092) compared with MSTM calculations for the above clusters. Blue, red, and black thin lines correspond to porosities of 93.7%, 87.5%, and 69.0%, respectively. The thick green line corresponds to the Mie theory phase function for a homogeneous and compact sphere of 2- μm radius. All phase functions are normalized to unity at a phase angle of 100°.




Fig. 2.— OSIRIS phase functions (red solid circles for MTP020/071 and red open circles for MTP025/092) compared with MSTM calculations (solid black line) for peppered sphere clusters such as that shown on the inset. XYZ labels on the inset are given in μm . All phase functions are normalized to unity at a phase angle of 100.




Fig. 3.— OSIRIS phase functions (red solid circles for MTP020/071 and red open circles for MTP025/092) compared with T-matrix calculations for size distributions of oriented, compact, and oblate ($q=2$, as that shown) spheroids: the black, blue, and green lines are for size distributions having $r_{min} = 0.1, 0.3, \text{ and } 1.5 \mu\text{m}$, respectively, with $r_{max} = 2.5 \mu\text{m}$. All phase functions are normalized to unity at a phase angle of 100




Fig. 4.— OSIRIS phase functions (red solid circles for MTP020/071 and red open circles for MTP025/092) compared with MSTM simulations of oriented oblate spheroidal volumes filled with 0.1 to 0.6- μm spherules. Black, blue, and green lines are for equivalent spherical radii of 3.1, 4.6, and 7.0 μm . The inset illustrates the array of spherules for the largest particle, where the XYZ labels are in μm . All phase functions are normalized to unity at a phase angle of 100




Fig. 5.— OSIRIS phase function measurements for MTP020/071 (solid red circles) compared to geometric optics simulations for three distributions of spheroids with uniform axes ratio in the $\epsilon = [0.25, 4.0]$ interval, and having equivalent spherical radii of $10 \mu\text{m}$ (black solid line), $100 \mu\text{m}$ (black dashed line), and $1000 \mu\text{m}$ (black dotted line). All phase functions are normalized to unity at 100° of phase angle. As in all the simulations in this work, the refractive index is set to $m = 1.6 + 0.1i$.