

Highlights on Spanish Astrophysics X, Proceedings of the XIII Scientific Meeting of the Spanish Astronomical Society held on July 16–20, 2018, in Salamanca, Spain. B. Montesinos, A. Asensio Ramos, F. Buitrago, R. Schödel, E. Villaver, S. Pérez-Hoyos, I. Ordóñez-Etxeberria (eds.), 2019

Dust-to-gas ratio in a complete sample of type-1 AGN.

I. Ordovás-Pascual¹, S. Mateos¹, F. J. Carrera¹, A. Caccianiga², R. Della Ceca², P. Severgnini², A. Moretti², L. Ballo³ and A. Corral¹

¹ Instituto de Física de Cantabria (CSIC-UC), E-39005, Santander, Spain

² INAF-Osservatorio Astronomico di Brera, Via Brera 28, I-20121, Milan, Italy

³ XMM-Newton Science Operations Centre, ESAC/ESA, PO Box 78, E-28692 Villanueva de la Cañada, Madrid, Spain

Abstract

According to the Unified Model of Active Galactic Nuclei (AGN), unobscured AGN based on its optical spectrum (detection of rest-frame UV-optical broad emission lines, type-1 AGN) should appear as X-ray unabsorbed AGN. However, there is an important fraction (10-30%) of AGN whose optical and X-ray classifications do not match, and the origin of the discrepancy is not clear. To provide insight into this topic, we have conducted a statistical analysis of the optical obscuration and X-ray absorption properties of the optically type-1 AGN from the Bright Ultra-hard XMM-Newton Survey (BUXS) with $L_{2-10\text{keV}} > 10^{42}$ erg s⁻¹ and $z=0.05-1$. We have high-quality spectra from XMM-Newton and either SDSS spectra or proprietary observations for the selected sample. In order to provide the most complete sample as possible, we have conducted a detailed analysis of the emission lines to provide a reliable classification of the AGNs. We derive the X-ray absorption by fitting their XMM-Newton spectra and the optical extinction using UV/optical spectral continuum fits. As BUXS is a flux limited X-ray selected sample at hard energies ($f_{4.5-10\text{keV}} \leq 6 \times 10^{-14}$ erg s⁻¹ cm⁻²), it is complete for N_{H} column densities up to the Compton-thick limit ($\sim 10^{24}$ cm⁻²). Our preliminary results show that most type-1 AGN in our sample show consistent optical and X-ray classification, but there is a large fraction (20%) of objects with large N_{H} column densities ($N_{\text{H}} > 4 \times 10^{21}$ cm⁻²). (See poster).