



Indian Journal of Traditional Knowledge
Vol 19 (2), April 2020, pp 268-276



Traditional processing associated changes in chemical parameters of wild Yam (*Dioscorea*) tubers from Koraput, Odisha, India

Debabrata Panda^{*1,+}, Meghali Biswas¹, Bandana Padhan¹ & Sangram K Lenka²

¹Department of Biodiversity and Conservation of Natural Resources, Central University of Orissa, Koraput 764 021, Odisha, India

²The Energy and Resources Institute, Gurugram 122 001, Haryana, India

E-mail: ⁺dpanda80@gmail.com

Received 08 May 2019; revised 20 January 2020

Wild yam tuber considered as famine food and played a prime role in the food habit of tribal people of Koraput district of Odisha, India. The tribal people employed a range of processing of these yams such as boiled, soaked and sun dried for detoxification of antinutrients in accordance with their needs. There is a scarcity of documented information on their nutrient composition/retention by traditional processing methods. The current study assessed the traditional processing (boiled, soaked and sun dried) associated changes in chemical composition and physico-functional characteristics of 6 wild and 1 cultivated *Dioscorea* tubers collected from Koraput, India. Different processing led to a significant reduction of proximate compositions and nutrient content compared to the raw tuber, whereas physico-functional parameters increased significantly ($p < 0.05$). In addition, there was significant decrease in the antinutrients, minerals and vitamin content by different processing in studied yam species. Results suggested that wild *Dioscorea* tubers as safe food sources for mass consumption and should be used in boiling form, as it retains higher nutrients coupled with significant removal of antinutritional compositions.

Keywords: *Dioscorea* tubers, Nutritional composition, Proximate composition, Traditional processing

IPC Code: Int. Cl.²⁰: A23L 33/20, A61K 36/00

Roots and tubers form a significant component of food in the tropics and sub-tropics of the world and it has assumed importance in the dietary habits of indigenous communities¹. Yam (*Dioscorea*) being a high value crop, it contributes about 10% of the total root and tubers production around the world²⁻³. In the developing world, while food security is concerned, wild yams with high calorie and rich carbohydrate content occupy a prominent position⁴. *Dioscorea* are annual or perennial climber species with underground edible tuber in the family Dioscoreaceae and worldwide more than 600 species are represented under the genus⁵⁻⁶. *Dioscorea* has a great diversity of species that are used as indigenous food by small and marginal rural families and forest-dwelling communities^{4,7,8}. Further, because of presence various bioactive constituents, these species are regarded unique in terms of their food, medicinal and economic value⁹. In different parts of the world varieties of edible wild yam species are found to differ in their chemical composition and nutritional values¹⁰. Though, there are

some reports on chemical compositions of the *Dioscorea* species of Nepalese¹⁰, Indian¹¹ and Ghanaian¹² species but there is a dearth of information on wild edible *Dioscorea* species of Koraput.

Koraput districts in the Indian state of Odisha is one of the agro-biodiversity hotspots and a storehouse of great diversity of food crops and plant species¹³. Several cultivated and wild *Dioscorea* tubers have been used as food and medicine by the forest-dwelling communities of Koraput^{9,14}. These yams were used in different forms by employing different processing, mainly fried, baked or boiled. However, their wider utilisation has been limited because of presence of many antinutritional compositions^{4,8}. Most of the wild yams are acrid in nature and are caused irritation, inflammation leading to gastrointestinal disturbances¹⁰. The tribal/rural people collected wild tubers and have practised different traditional processing such as boiled, soaked and sun dried for detoxification of antinutrients in accordance to their needs⁹. Limited scientific information is available on the chemical parameters in the wild *Dioscorea* species and also scientifically their

*Corresponding author

indigenous detoxification or food processing practice is not yet been studied. The knowledge on wild tuber and their uses by tribal people has the potential for addressing food security and physico-functional properties have become important for bio-prospecting of flour out of tuber in food industries¹⁵. The present work aims to ascertain the effect of traditional processing practices on their nutrient and antinutrient retentions in wild *Dioscorea* species of Koraput. The study aims to suggest a method to remove the antinutrient/toxins from the wild *Dioscorea* species and make these edible tubers as safe food sources for mass consumption.

Materials and methods

Collection and processing of yam tuber

Six edible wild yam species namely *Dioscorea bulbifera*, *D. glabra*, *D. hamiltonii*, *D. hispida*, *D. pentaphylla* and *D. pubera*, including *D. alata*, a cultivated species was used in the study. These are consumed frequently as food as well as vegetables by tribals of Koraput. The details of yam tubers and their traditional consumption practice are presented in Table 1. The matured tubers of each species were collected from the garden of Central University of Orissa, Koraput, which were grown under same agro-climatic conditions¹⁶. The freshly collected tubers were sliced into pieces of ~2.5 mm diameter, divided into 4 sections and subjected to different traditional processing methods, such as raw, boiled, water soaked and sun dried. The first section which was peeled, was used as raw yam; the second portion was made into small pieces followed by boiling for 30 min at 100°C. The third portion of the raw yam which were peeled and sliced were made to soak for 24 h in distilled water and used as soaked sample. The peeled, washed and sliced raw yam in the fourth section was subjected to sun drying for 4 days. All the

processed samples were dried in shade for 4-6 days. After processing the samples were grounded into powder mechanically and used for analysis.

Measurement of proximate composition

For measurement of moisture content, samples were oven dried at 105°C for 24 h to a constant weight¹⁷. The determination of crude fat was done for 3 h in a lipid extractor and the solvent used petroleum ether¹⁷. To determine the ash content, the samples were ignited at 600°C for a period of 3 h in muffle furnace and weight of the residue was taken¹⁷. Using standard methods crude fibre content were determined¹⁷.

Measurement of physico-functional parameters

water solubility index (WSI) of *Dioscorea* tubers were carried out following the method of Anderson *et al.*¹⁹. Water absorption capacity (WAC) was measured following the method of Phillips *et al.*¹⁸. Foam capacity (FC) was determined as per the method of Coffman and Garcia²⁰. Paste clarity (PC) of flour was measured following the method of Craig *et al.*²¹

Determination of nutritional Compositions

Starch, reducing sugar and amylose content of yam tuber sample were determined following the method of Sadasivam and Manickam²². Protein content of tubers was estimated as per the procedure of Lowry *et al.* (1951)²³. Ascorbic acid (Vitamin C) content was measured following the method of Omaye *et al.*²⁴ and the value was expressed as mg/100 g on dry weight basis. The α -Tocopherol (vitamin E) was measured following the method of Baker *et al.*²⁵ by using α -tocopherol as standard.

Determination of anti-nutritional compositions

Diosgenin content was determined followed by the method of Uematsu *et al.*²⁶ by using diosgenin as

Table 1 — Details of yam species with their tribal consumption practice from Koraput, India.

Accession No	Yam species	Tribal consumption practice
CUO 001	<i>Dioscorea bulbifera</i> L.	Tubers are sliced, boiled and kept overnight to remove bitterness and cooked as curry.
CUO 002	<i>Dioscorea hamiltonii</i> Hook. f.	Tuber are boiled and used to make curry.
CUO 003	<i>Dioscorea pubera</i> Blume	Tubers are boiled and cooked as vegetable.
CUO 004	<i>Dioscorea hispida</i> Dennst.	Tubers are sliced and soaked in water for 24 h and consumed. The processed tuber slices are also sundried and used as chips.
CUO 005	<i>Dioscorea pentaphylla</i> L.	Tubers are boiled and used to make curry.
CUO 006	<i>Dioscorea glabra</i> Roxb.	Tubers are boiled and cooked to make curry.
CUO 007	<i>Dioscorea alata</i> L.	Tubers are boiled and cooked as curry.

standard. Oxalate was determined by following the method of Meena *et al.*²⁷ using oxalic acid as standard. Phytate content of yam sample was determined spectrophotometrically as per the method described by Wheeler and Ferrel²⁸. Total phenol content was determined as per the method of Sadasivam and Manickam²² using gallic acid as standard. Tannin content of yam flour was determined by the Folin-Dennis spectrophotometric method of Sadasivam and Manickam²² by using tannic acid as standard. The α -amylase inhibitor was measured by following the method of Alonso *et al.*²⁹

Determination of mineral composition

The mineral compositions were estimated by digestion of tuber sample (1 g each) in nitric, sulphuric and perchloric acid (6:1:2) at 100°C following the method AOAC¹⁷. The mineral compositions such as

calcium, copper, iron, magnesium, manganese and zinc contents were measured by using Atomic Absorption Spectrophotometer¹⁷. The digested sample was also used for the measurement of sodium and potassium content in a Flame Photometer (Systronics-105)¹⁷.

Data analysis

All the parameters were statistically analysed by two-way analysis of variance using the software CROPSTAT (IRRI, Philippines). The mean value of all the parameters were compared by performing the Fisher's LSD test.

Results

Test of significance

The test of significance of different parameters in yam tubers were subjected to different processing methods is presented in Table 2. In different parameters,

Table 2 — Sum square is the absolute value and percentage of total (in bracket) of main effect resulting from analysis of variance of studied parameters in yam tubers subjected to different processing methods.

Parameters	Source of variation		
	Species (S, <i>df</i> =6)	Processing (P, <i>df</i> =3)	Species x Processing (SxP, <i>df</i> =18)
Moisture	2300** (87.9)	210** (8.0)	107* *(4.1)
Ash	32.00** (46.9)	15.14** (22.2)	20.22** (29.6)
Crude fat	10.12** (71.0)	3.59* (25.0)	0.43 ns (3.0)
Crude fibre	11.73** (63.0)	4.70** (25.0)	2.09* (11.0)
Water absorption capacity	242344** (78.1)	2490** (12.2)	1881 ** (8.7)
Water solubility index	4022** (73.3)	1127** (20.5)	303** (5.2)
Foam capacity	2754** (77.5)	684** (19.2)	65** (1.8)
Paste clarity	12572** (92.9)	866** (6.4)	63* (2.4)
Protein	1327** (51.2)	641** (24.8)	634** (24.5)
Sugar	15388** (26.9)	24538** (42.9)	16082** (28.1)
Starch	76457** (49.7)	2919* (1.9)	59020** (38.3)
Amylose	622.11** (88.0)	69.85** (9.0)	11.94* (1.0)
Ascorbic acid	2396** (94.0)	89.81* (3.0)	53.15* (2.0)
Vitamin-E	8.65** (83.0)	1.16* (11.0)	0.52* (5.0)
Oxalate	116.79** (41.0)	110.88** (38.0)	55.25** (19.0)
Phytate	192.51** (62.0)	70.79** (23.0)	41.48** (13.0)
Diosgenin	4.65** (25.0)	11.71** (64.0)	1.88* (10.0)
Phenol	517.13** (33.0)	332.22** (21.2)	711.62** (45.4)
Tannin	74.81** (68.5)	15.10 ** (13.8)	19.01** (17.4)
Mg	719494** (55.2)	410898** (29.3)	164476** (13.0)
Ca	111580** (22.2)	390139** (61.2)	157193** (13.6)
Cu	563.70** (50.2)	426.27** (38.0)	132.62** (11.8)
Fe	55886* (8.6)	46671* (7.2)	212676** (32.6)
Mn	12467* (9.7)	9458* (7.3)	41371** (32.1)
Zn	59.90** (45.5)	52.77** (40.1)	18.86* *(14.3)
Na	254522** (69.8)	43409** (11.9)	66504** (18.2)
K	470529** (32.2)	24564** (6.5)	660824** (62.3)

df: degrees of freedom, Total *df*=83, The P of overall ANOVA for species, processing and Species x Processing interaction for each parameter **p*<0.05, ***p*<0.01, ns: not significant

among the replications the differences were non-significant ($p > 0.05$), but in the case of different species, different traditional processing method and their interaction were statistically significant ($p < 0.05$).

Proximate Composition

The proximate compositions in studied yam tubers are shown in Table 3. There was significant ($p < 0.05$) differences of moisture content was observed among the studied yam species and the value ranged from 66.07 to 78.05% (Table 3). The cause of variance for moisture content was the species (87.9%), followed by processing (8.0%) and interactions of species×processing (4.1%) (Table 2). There were significant ($p < 0.05$) differences of ash content, crude fibre and crude fat observed among the studied yam tubers (Table 3). The different processing methods led to a significant ($p < 0.05$) decrease in ash and crude fat content compared to raw yams. Similarly, boiling and soaking led to significant decrease of crude fibre content, while in sun dried samples no significant changes was observed.

Physico-functional parameters

Different physico-functional parameters in studied yam tubers were shown in Table 3. There were

significant ($p < 0.05$) difference of WAC observed among the *Dioscorea* species (Table 3). Boiling of yam caused a significant ($p < 0.05$) increase of WAC only in *D. bulbifera* and *D. pentaphylla* but the WAC was not significantly changed by soaking and sun drying in the studied *Dioscorea* species. In addition, foam capacity was significantly increased by boiling and soaking in comparison to the raw tuber, while FC was not significantly changed by sun drying. Paste clarity (PC) of tuber flours of all the *Dioscorea* species was significantly improved by boiling, soaking and sun drying compared to the raw tuber. Water solubility index was significantly increased by boiling compared to other processing methods.

Nutritional parameters

Different nutritional parameters in studied yam tubers are presented in Table 4. Sugar, starch and amylose content of raw tubers ranged from 54.50 to 88.00 (mg/g dry weight), 101.6 to 173.2 (mg/g dry weight) and 5.46 to 15.55 (mg/g dry weight), respectively. Significant decrease of sugar content was observed by soaking compared to the raw tuber, while sun drying significantly improved the level of sugar only in *D. glabra*, *D. hispida* and *D. alata*. The starch content was higher in raw sample and was

Table 3 — Proximate analysis and physico-functional parameters in different wild and cultivated species of *Dioscorea* subjected to different processing.

Species	Moisture content (%)				Ash content (%)				Crude fat (g/100 g dwt.)				Crude fibre (g/100 g dwt.)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D. bulbifera</i>	74.84 ^b	78.57 ^{a*}	76.59 ^{a*}	71.25 ^{b*}	2.31 ^b	1.52 ^{b*}	1.99 ^{b*}	1.52 ^{b*}	2.29 ^a	1.27 ^{b*}	1.78 ^{a*}	2.05 ^a	1.78 ^a	0.79 ^{b*}	1.21 ^{a*}	1.64 ^a
<i>D. hamiltoni</i>	70.89 ^c	74.18 ^{b*}	72.34 ^{b*}	68.36 ^{b*}	2.52 ^b	1.16 ^{b*}	1.53 ^{b*}	1.58 ^{b*}	2.40 ^a	1.12 ^{b*}	1.16 ^{b*}	2.07 ^{a*}	1.91 ^a	0.99 ^{a*}	1.34 ^{a*}	1.61 ^a
<i>D. pubera</i>	66.07 ^d	69.35 ^{c*}	68.45 ^{c*}	63.14 ^{c*}	2.96 ^a	1.53 ^{b*}	1.05 ^{b*}	1.49 ^{b*}	2.47 ^a	1.06 ^{b*}	1.58 ^{a*}	2.31 ^a	1.65 ^a	0.94 ^{b*}	1.13 ^{a*}	1.44 ^a
<i>D. hispida</i>	72.46 ^c	73.4 ^b	74.74 ^{b*}	68.24 ^{b*}	2.58 ^b	1.53 ^{b*}	1.52 ^{b*}	1.06 ^{b*}	2.06 ^b	1.47 ^{b*}	1.64 ^{a*}	1.76 ^{b*}	1.56 ^a	0.86 ^{b*}	1.33 ^{a*}	1.43 ^a
<i>D. pentaphylla</i>	70.53 ^c	74.80 ^{b*}	72.83 ^{b*}	66.09 ^{c*}	2.30 ^b	2.20 ^{a*}	2.68 ^{a*}	2.33 ^{a*}	2.42 ^a	1.11 ^{b*}	1.61 ^{b*}	2.05 ^{a*}	1.75 ^a	1.08 ^{a*}	1.36 ^{a*}	1.58 ^a
<i>D. glabra</i>	78.05 ^a	78.36 ^a	79.06 ^a	74.47 ^{a*}	2.96 ^b	2.02 ^{a*}	2.56 ^a	2.58 ^a	1.87 ^b	1.65 ^a	1.86 ^a	1.74 ^b	1.04 ^b	0.41 ^{c*}	0.71 ^{b*}	0.86 ^b
<i>D. alata</i>	70.40 ^c	73.49 ^{b*}	73.86 ^{b*}	66.64 ^{c*}	3.19 ^a	2.41 ^{a*}	2.58 ^{a*}	2.98 ^a	1.38 ^c	1.07 ^{b*}	1.16 ^c	1.07 ^{c*}	0.74 ^b	0.22 ^{c*}	0.47 ^{b*}	0.64 ^b
LSD(P<0.05)	2.34				0.56				0.31				0.29			
Species	Water absorption capacity (%)				Foam capacity (%)				Paste clarity (%)				Water solubility index (%)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D. bulbifera</i>	353.6 ^a	383.0 ^{a*}	340.5 ^a	352.7 ^a	31.8 ^b	40.2 ^{b*}	35.5 ^{b*}	32.9 ^b	48.5 ^b	60.0 ^{b*}	54.5 ^{b*}	51.5 ^{b*}	15.85 ^c	30.35 ^{d*}	25.75 ^c	19.55 ^d
<i>D. hamiltoni</i>	261.1 ^c	270.6 ^c	265.9 ^c	263.9 ^c	22.5 ^d	32.3 ^{d*}	29.2 ^{c*}	24.0 ^d	33.0 ^f	38.0 ^{e*}	36.0 ^{e*}	32.5 ^c	34.50 ^a	39.50 ^{b*}	38.00 ^b	36.25 ^b
<i>D. pubera</i>	216.5 ^d	227.5 ^d	220.5 ^d	213.7 ^d	37.2 ^a	44.6 ^{a*}	40.3 ^{a*}	38.3 ^a	41.0 ^d	51.5 ^{d*}	49.0 ^{d*}	44.0 ^{d*}	18.15 ^c	29.60 ^{d*}	22.50 ^f	19.60 ^d
<i>D. hispida</i>	144.6 ^f	161.8 ^f	150.7 ^c	146.1 ^c	16.9 ^e	26.7 ^{e*}	23.4 ^{d*}	18.3 ^c	14.5 ^e	28.5 ^{f*}	22.5 ^{f*}	18.5 ^{f*}	35.93 ^a	60.20 ^{a*}	46.50 ^a	42.75 ^a
<i>D. pentaphylla</i>	306.2 ^b	337.7 ^{b*}	314.1 ^b	311.5 ^b	25.8 ^c	35.3 ^{c*}	29.4 ^{c*}	26.4 ^c	39.5 ^d	52.0 ^{d*}	49.0 ^{d*}	44.0 ^{d*}	26.4 ^b	33.15 ^{c*}	30.15 ^d	28.55 ^c
<i>D. glabra</i>	255.6 ^c	268.4 ^c	265.7 ^c	262.5 ^c	16.6 ^e	22.3 ^{f*}	19.4 ^{e*}	18.5 ^c	69.7 ^a	78.0 ^{a*}	75.0 ^{a*}	70.5 ^{a*}	17.70 ^c	26.45 ^{e*}	21.10 ^f	20.35 ^d
<i>D. alata</i>	197.2 ^e	211.3 ^e	206.8 ^d	201.5 ^d	26.5 ^c	40.1 ^{b*}	36.4 ^{b*}	32.4 ^{b*}	45.5 ^c	56.0 ^{c*}	52.0 ^{c*}	48.5 ^{c*}	24.5b	39.10 ^{b*}	35.10 ^{c*}	29.85 ^c
LSD(P<0.05)	15.7				2.68				1.9				2.26			

Means followed by a common letter in the same column are not significantly different at $p < 0.05$ level by Fisher's least significance difference (LSD) test. *: represents the significantly different from Raw at $p < 0.05$.

Table 4 — Nutritional compositions of *Dioscorea* species subjected to different processing. Data are the mean of three replication (n=3).

Species	Sugar (mg/g dwt.)				Starch (mg/g dwt.)				Amylose (mg/g dwt.)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D.bulbifera</i>	87.6 ^a	78.0 ^a	60.9 ^{a*}	96.6 ^a	172.1 ^a	107.9 ^{c*}	130.2 ^{b*}	174.6 ^a	11.06 ^b	9.35 ^{a*}	9.91 ^b	10.67 ^b
<i>D.hamiltoni</i>	89.6 ^a	79.9 ^a	67.6 ^{a*}	94.4 ^a	171.3 ^a	156.5 ^a	150.8 ^{a*}	191.1 ^a	15.55 ^a	12.38 ^{a*}	14.59 ^a	16.25 ^a
<i>D.pubera</i>	88.0 ^a	83.5 ^a	68.6 ^{a*}	90.8 ^a	173.2 ^a	125.1 ^{b*}	107.2 ^{c*}	189.9 ^a	15.34 ^a	11.49 ^{a*}	13.49 ^a	15.11 ^a
<i>D.hispida</i>	61.4 ^c	56.6 ^b	46.5 ^{b*}	71.5 ^{b*}	132.6 ^c	104.8 ^{c*}	126.1 ^b	138.3 ^c	8.84 ^c	6.65 ^{b*}	9.60 ^b	10.22 ^b
<i>D.pentaphylla</i>	54.5 ^c	46.4 ^b	41.3 ^{b*}	75.8 ^{b*}	101.6 ^d	96.2 ^c	98.6 ^c	118.5 ^d	8.57 ^c	6.55 ^{b*}	7.24 ^c	9.11 ^b
<i>D.glabra</i>	48.8 ^d	39.6 ^b	37.9 ^{c*}	64.0 ^{c*}	121.5 ^c	88.4 ^{d*}	86.8 ^{d*}	128.7 ^c	5.46 ^d	3.03 ^{c*}	4.59 ^d	5.87 ^c
<i>D.alata</i>	78.0 ^b	75.0 ^a	68.0 ^{a*}	81.9 ^b	151.3 ^b	147.3 ^a	145.9 ^a	156.0 ^b	10.24 ^b	6.32 ^{b*}	8.60 ^b	10.12 ^b
LSD(P<0.05)	10.3				18.5				2.1			
Species	Protein (mg/g dwt.)				Ascorbic acid (mg/100 g dwt.)				Vitamin-E (mg/100 g dwt.)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D.bulbifera</i>	21.60 ^a	12.90 ^{a*}	22.65 ^a	16.75 ^{a*}	6.77 ^b	5.50 ^{a*}	5.35 ^{b*}	3.59 ^{b*}	0.450 ^c	0.350 ^{b*}	0.230 ^{b*}	0.165 ^{c*}
<i>D.hamiltoni</i>	17.55 ^b	9.65 ^{b*}	17.75 ^b	12.45 ^{b*}	5.48 ^c	3.21 ^{b*}	3.69 ^{c*}	4.02 ^{b*}	0.760 ^a	0.320 ^{b*}	0.435 ^{a*}	0.550 ^{a*}
<i>D.pubera</i>	14.10 ^c	8.30 ^{b*}	13.75 ^c	12.70 ^{b*}	9.85 ^a	6.63 ^{a*}	7.32 ^{a*}	8.48 ^{a*}	0.660 ^b	0.370 ^{b*}	0.515 ^{a*}	0.530 ^{a*}
<i>D.hispida</i>	9.40 ^d	6.25 ^{c*}	10.90 ^d	4.55 ^{d*}	1.96 ^e	0.55 ^{d*}	0.57 ^{e*}	1.60 ^c	0.740 ^a	0.540 ^{a*}	0.270 ^{b*}	0.465 ^{a*}
<i>D.pentaphylla</i>	8.80 ^d	4.85 ^{d*}	9.70 ^d	7.80 ^c	3.28 ^d	2.07 ^{c*}	2.48 ^d	2.60 ^c	0.430 ^c	0.255 ^{c*}	0.315 ^{b*}	0.395 ^{b*}
<i>D.glabra</i>	8.30 ^d	6.70 ^{c*}	8.50 ^e	7.20 ^c	3.61 ^d	2.21 ^{c*}	3.27 ^c	2.76 ^c	0.310 ^d	0.175 ^{c*}	0.215 ^{b*}	0.285 ^b
<i>D.alata</i>	12.95 ^c	6.10 ^{d*}	12.45 ^c	8.10 ^{c*}	4.90 ^c	3.62 ^{b*}	4.03 ^c	4.39 ^b	0.310 ^d	0.130 ^{d*}	0.220 ^{c*}	0.275 ^b
LSD(P<0.05)	1.57				1.2				0.09			

Means followed by a common letter in the same column are not significantly different at $p < 0.05$ level by Fisher's least significance difference (LSD) test. *: represents the significantly different from Raw at $p < 0.05$

significantly decreased by boiling and soaking in most of the yam tubers. Whereas, amylose content of yam tubers was significantly declined under boiling compared to raw, soaking and sundry samples. Raw yam tuber was high in protein content, ascorbic acid and vitamin-E content but significantly reduced in different processing. Significant reduction of protein content was observed in boiled sample compared to raw tuber, while ascorbic acid and vitamin-E were declined in boiled, soaked and sun-dried sample.

Antinutritional compositions

Different anti-nutritional parameters, such as diosgenin, oxalate, phytate, phenols, tannins and α -amylase inhibitor contents of studied raw and processed yam tubers were shown in Table 5. There was significant ($p < 0.05$) higher levels of anti-nutrients were observed in wild yam species compared to the cultivated one. In addition, the raw yam tubers showed a very high anti-nutrient content compared to the processed sample and all the processing methods such as soaked, boiled and sun-dried resulted in reduction of anti-nutrients in the tubers.

Mineral compositions

Different mineral compositions in studied yam tubers were presented in Table 6. In raw yam tubers

the range of Na was 16.3 to 83.16 mg/100 g, K was 456 to 934 mg/100 g, Mg was 21.8 to 67.5 mg/100 g, Cu was 9.73 to 14.62 mg/100 g, Mn was 2.35 to 23.7 mg/100 g, Ca was 22.11 to 46.3 mg/100 g, Fe was 8.66 to 23.55 mg/100 g and Zn was 1.06 to 3.04 mg/100 g. Different processing of yam led to significant decrease of mineral contents. Based on the results, K was the abundant mineral, ranging from 465 to 934 mg/100 g dry weight, while Zn was the least abundant mineral among the studied yam tubers. Some wild yam species (*D. hamiltonii*, *D. pubera* and *D. bulbifera*) exhibited remarkably higher mineral compositions in comparison to the cultivated species.

Discussion

Consumption of indigenous foods is being encouraged worldwide for meeting dietary needs of people. For the studied wild edible *Dioscorea* species, there is no previous report on chemical compositions/retention in relation to different traditional processing methods. Proximate compositions such as moisture, ash, crude fat and crude fibre are the major components of food³⁰⁻³¹. Boiled and soaked yam showed significantly ($p < 0.05$) higher moisture content compared to the raw tuber and it may be the result of presence of water from boiling and soaking, while it is decreased by sun

Table 5 — Anti-nutritional compositions in indifferent species of *Dioscorea* subjected to different processing (mg/100 g). Data are the mean of three replication (n=3).

Species	Diosgenin (mg/100 g dwt.)				Phytate (mg/100 g dwt.)				Oxalase (mg/100 g dwt.)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D.bulbifera</i>	4.87 ^a	3.66 ^{a*}	2.36 ^{c*}	3.52 ^{a*}	5.30 ^e	4.44 ^{c*}	3.74 ^{b*}	4.60 ^{b*}	3.29 ^c	2.40 ^{a*}	0.54 ^{c*}	2.33 ^{b*}
<i>D.hispida</i>	5.05 ^a	3.67 ^{a*}	2.64 ^{c*}	3.96 ^{a*}	9.72 ^a	5.36 ^{b*}	3.47 ^{b*}	4.84 ^{b*}	4.26 ^b	2.81 ^{a*}	2.17 ^{b*}	0.95 ^{d*}
<i>D.pubera</i>	4.17 ^b	3.15 ^{a*}	3.18 ^{b*}	3.66 ^a	4.49 ^f	2.65 ^{d*}	3.94 ^b	3.52 ^{b*}	3.62 ^c	0.55 ^{c*}	0.49 ^{c*}	0.79 ^{d*}
<i>D.hamiltoni</i>	4.14 ^b	3.00 ^{b*}	3.17 ^{b*}	3.19 ^{b*}	5.89 ^d	2.79 ^{d*}	3.58 ^{b*}	4.64 ^{b*}	3.50 ^c	0.23 ^{c*}	0.27 ^{c*}	0.69 ^{d*}
<i>D.pentaphylla</i>	4.96 ^a	3.13 ^{b*}	4.24 ^{a*}	3.76 ^{a*}	7.64 ^c	6.28 ^{a*}	6.61 ^{a*}	5.27 ^{a*}	4.39 ^b	0.29 ^{c*}	0.55 ^{c*}	0.72 ^{d*}
<i>D.glabra</i>	4.93 ^a	3.63 ^{a*}	3.77 ^{b*}	3.33 ^{a*}	8.82 ^b	4.50 ^{c*}	6.64 ^{a*}	5.66 ^{a*}	7.54 ^a	1.57 ^{b*}	3.54 ^{a*}	5.82 ^{a*}
<i>D.alata</i>	3.18 ^c	2.36 ^{b*}	2.57 ^c	2.87 ^b	3.40 ^g	2.22 ^{d*}	2.41 ^{c*}	2.94 ^d	2.78 ^d	0.19 ^{c*}	0.59 ^{c*}	1.70 ^{c*}
LSD(P<0.05)	0.65				0.56				0.45			
Species	α Amylase Inhibitor (Units/ g dwt.)				Phenol (mg/ g dwt.)				Tanin (mg/100 g dwt.)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D.bulbifera</i>	26.04 ^a	5.34 ^{b*}	4.59 ^{b*}	11.55 ^{b*}	4.12 ^d	2.98 ^{a*}	1.58 ^{b*}	3.39 ^a	2.96 ^a	0.59 ^{d*}	0.97 ^{d*}	1.25 ^{c*}
<i>D.hamiltoni</i>	22.80 ^b	6.26 ^{b*}	8.95 ^{a*}	6.27 ^{d*}	4.33 ^d	2.27 ^{a*}	3.46 ^a	2.79 ^{a*}	2.59 ^b	1.62 ^{b*}	1.28 ^{c*}	1.73 ^{a*}
<i>D.pubera</i>	24.72 ^a	9.46 ^{a*}	9.27 ^{a*}	13.95 ^{a*}	8.87 ^a	2.35 ^{a*}	2.25 ^{b*}	2.89 ^{a*}	3.13 ^a	2.63 ^{a*}	2.45 ^{a*}	1.50 ^{b*}
<i>D.hispida</i>	22.45 ^b	5.26 ^{b*}	1.38 ^{c*}	6.82 ^{c*}	5.39 ^c	2.62 ^{a*}	1.15 ^{c*}	2.93 ^{a*}	2.65 ^b	0.66 ^{d*}	0.21 ^{e*}	0.23 ^{d*}
<i>D.pentaphylla</i>	20.13 ^c	4.78 ^{b*}	3.92 ^{b*}	5.23 ^{d*}	6.57 ^b	2.35 ^{a*}	1.68 ^{b*}	3.02 ^{a*}	2.94 ^a	1.53 ^{c*}	0.96 ^{d*}	1.89 ^{a*}
<i>D.glabra</i>	18.48 ^d	3.73 ^{c*}	7.42 ^{a*}	15.51 ^{a*}	1.96 ^e	0.67 ^{b*}	0.92 ^{c*}	1.33 ^b	0.96 ^c	0.16 ^{e*}	0.27 ^{e*}	0.36 ^{d*}
<i>D.alata</i>	20.71 ^c	5.29 ^{b*}	5.81 ^{b*}	8.59 ^{c*}	3.64 ^d	2.87 ^a	2.16 ^{b*}	2.58 ^{a*}	2.55 ^b	1.81 ^{b*}	1.97 ^{b*}	1.82 ^{a*}
LSD(P<0.05)	1.99				0.88				0.21			

Means followed by a common letter in the same column are not significantly different at $p < 0.05$ level by Fisher's least significance difference (LSD) test. *: represents the significantly different from Raw at $p < 0.05$.

drying (Table 3). The ash content, crude fibre and fat contents in the studied raw yam tubers were higher than earlier reported value in different tropical yam tubers¹⁰⁻¹¹. The variation of proximate compositions among the yam species may be due to their origin and genetic makeup. Various traditional processing method resulted a remarkable decrease in ash content, crude fat and crude fibre content compared to raw yams.

The physico-functional properties such as WAC, FC, PC and WSI are necessary for testing the quality tuber flour for using in food industries¹⁵. Boiling of yam led to significant increase of WAC and WSI, whereas soaking significantly improved the FC and PC (Table 3). Based on the results of physico-functional properties, it suggested that these wild yam tuber flours were capable of holding higher content of water because of the starch gelatinization³². Higher WAC of the tuber flour is useful for making various food products¹⁵. Similar findings also observed in other tuber flours like *C. esculenta* and potato¹⁵. Paste clarity is known to influence by the amylose content of the flour and some wild yam flours showed higher paste clarity and better useful in food industries³³.

The value of nutritional parameters such as sugar, starch, amylose and protein content in raw tubers

were in the range of earlier reported value in yam species of Nepal¹⁰, Ethiopia³⁴ and from Sri Lanka³⁵. The processing of yam tuber by boiling and soaking led to the significant decrease of sugar, starch and amylose content compared to raw tuber. Adepoju *et al.*⁸ and Adepoju³⁶ were reported that the cooking and soaking of yam tubers caused leaching of nutrient and results in decrease of carbohydrate content in processed tuber flour and its products. In contrast, significant increase of sugar content of tuber flour under sun-dried sample was believed to be due to less moisture retention in the tubers. Presence of higher protein and vitamin content in yam species such as *D. hamiltonii*, *D. pubera* and *D. oppositifolia* over the other species indicated its nutritional superiority. Different processing led to significant decline of ascorbic acid and vitamin E compared to the raw tuber.

Different anti-nutritional parameter such as diosgenin, oxalate and phytate were associated with the health implications and affect the digestion of food^{30,37}. The levels of antinutrients in studied raw yam tubers were lower than the earlier reported value in tropical tubers¹². Based on the results levels of anti-nutrients were decreased by processing of yams before consumption. Generally wild yams are consumed by employing different traditional

Table 6 — Mineral compositions of different species of *Dioscorea* subjected to different processing (mg/100 g). Data are the mean of three replication (n=3).

Species	Na (mg/100 g dwt.)				K (mg/100 g dwt.)				Mg (mg/100 g dwt.)				Cu (mg/100 g dwt.)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D. bulbifera</i>	62.21 ^b	26.50 ^{b*}	22.63 ^{c*}	60.13 ^b	845.15 ^a	456.88 ^{a*}	431.00 ^{b*}	600.55 ^{b*}	48.15 ^d	38.55 ^{a*}	32.00 ^{c*}	45.60 ^d	12.25 ^a	3.13 ^{a*}	4.23 ^{b*}	11.15 ^b
<i>D. hamiltoni</i>	83.17 ^a	29.62 ^{b*}	47.48 ^{a*}	72.95 ^a	934.15 ^a	341.53 ^{a*}	589.85 ^{a*}	629.38 ^{b*}	52.60 ^c	23.75 ^{c*}	35.45 ^{b*}	51.01 ^c	12.78 ^a	3.94 ^{a*}	6.85 ^{a*}	12.46 ^a
<i>D. pubera</i>	64.35 ^b	58.50 ^a	21.77 ^{c*}	64.27 ^b	611.57 ^b	452.12 ^{a*}	356.75 ^{b*}	602.33 ^b	67.55 ^a	32.75 ^{b*}	45.51 ^{a*}	66.30 ^a	14.62 ^a	3.21 ^{a*}	6.46 ^{a*}	14.50 ^a
<i>D. hispida</i>	16.30 ^f	9.55 ^{d*}	10.25 ^d	15.80 ^c	465.14 ^c	315.85 ^{b*}	367.22 ^{b*}	439.84 ^c	21.85 ^e	15.25 ^{d*}	18.35 ^{d*}	20.15 ^f	12.03 ^a	4.46 ^{a*}	4.11 ^{b*}	11.94 ^b
<i>D. pentaphylla</i>	28.74 ^e	21.55 ^{c*}	22.15 ^{c*}	28.27 ^d	483.34 ^b	251.74 ^{b*}	223.30 ^{c*}	380.85 ^c	43.28 ^e	32.66 ^{b*}	38.73 ^{b*}	42.15 ^d	12.88 ^a	3.73 ^{a*}	6.22 ^{a*}	9.26 ^{d*}
<i>D. glabra</i>	33.17 ^d	29.27 ^b	23.47 ^{b*}	26.63 ^d	612.22 ^b	321.42 ^{a*}	456.77 ^{a*}	521.32 ^b	27.68 ^f	10.71 ^{d*}	19.40 ^{d*}	27.17 ^e	12.06 ^b	4.24 ^{a*}	6.02 ^{a*}	10.18 ^b
<i>D. alata</i>	45.00 ^c	31.04 ^{b*}	29.55 ^{b*}	43.98 ^c	645.31 ^b	342.15 ^{a*}	531.60 ^{a*}	609.39 ^a	58.66 ^b	23.99 ^{c*}	29.55 ^{c*}	58.09 ^b	9.73 ^b	3.27 ^{a*}	5.37 ^{a*}	9.23 ^d
LSD(P<0.05)	6.12				136.5				3.89				2.41			
Species	Mn (mg/100 g dwt.)				Ca (mg/100 g dwt.)				Fe (mg/100 g dwt.)				Zn (mg/100 g dwt.)			
	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried	Raw	Boiled	Soaked	Sun dried
<i>D. bulbifera</i>	13.73 ^c	4.25 ^{b*}	7.12 ^{c*}	13.42 ^{c*}	32.64 ^c	22.43 ^{b*}	25.02 ^{b*}	31.53 ^b	23.55 ^c	13.15 ^{b*}	16.88 ^{b*}	23.07 ^c	2.83 ^b	0.90 ^{b*}	1.43 ^{b*}	2.21 ^{c*}
<i>D. hamiltoni</i>	16.68 ^b	3.57 ^{b*}	10.56 ^{b*}	16.32 ^{b*}	46.31 ^a	11.22 ^{d*}	32.61 ^{a*}	44.21 ^a	28.95 ^b	12.55 ^{b*}	19.44 ^{b*}	28.61 ^b	2.76 ^b	0.63 ^{c*}	1.15 ^{c*}	2.63 ^{b*}
<i>D. pubera</i>	23.77 ^a	9.37 ^{a*}	13.89 ^{a*}	23.43 ^a	34.25 ^c	28.59 ^{a*}	24.32 ^{b*}	28.43 ^{c*}	65.07 ^a	24.38 ^{a*}	43.07 ^{a*}	64.34 ^a	2.76 ^b	0.84 ^{b*}	1.79 ^{a*}	1.65 ^{d*}
<i>D. hispida</i>	2.13 ^f	0.50 ^{d*}	0.80 ^{e*}	1.79 ^g	25.63 ^d	13.11 ^{d*}	18.91 ^{c*}	14.37 ^{e*}	8.66 ^e	4.23 ^{c*}	5.15 ^d	8.41 ^e	1.64 ^d	0.09 ^{e*}	0.22 ^{e*}	1.55 ^e
<i>D. pentaphylla</i>	3.74 ^e	0.74 ^{d*}	1.63 ^{e*}	3.21 ^e	39.00 ^b	23.00 ^{b*}	26.12 ^{b*}	29.34 ^{b*}	12.63 ^d	11.84 ^b	8.70 ^d	4.22 ^{f*}	1.86 ^c	0.84 ^{b*}	1.32 ^{b*}	1.75 ^d
<i>D. glabra</i>	2.35 ^f	0.44 ^{d*}	1.03 ^{e*}	2.79 ^f	26.06 ^d	17.33 ^{c*}	10.45 ^{d*}	20.26 ^{d*}	10.34 ^e	3.24 ^{c*}	5.34 ^{d*}	9.66 ^e	1.06 ^e	0.33 ^{d*}	0.55 ^{d*}	0.86 ^{f*}
<i>D. alata</i>	7.80 ^d	2.45 ^{c*}	4.65 ^{d*}	7.37 ^d	22.11 ^e	15.12 ^{c*}	10.32 ^{d*}	20.34 ^d	16.47 ^d	7.35 ^{c*}	13.20 ^c	16.31 ^d	3.04 ^a	1.45 ^{a*}	1.31 ^{b*}	2.88 ^{a*}
LSD(P<0.05)	0.93				2.41				4.39				0.13			

Means followed by a common letter in the same column are not significantly different at $p < 0.05$ level by Fisher's least significance difference (LSD) test. *: represents the significantly different from Raw at $p < 0.05$.

processing methods which led to removal of anti-nutrients^{8,36}. The levels of antinutrients in yam tuber after processing were negligible and does not affect the bioavailability of nutritional parameters⁸.

The results of mineral compositions showed that raw yam tubers were very rich in minerals, such as Na, K, Ca, Mg, Fe and Zn. The range of minerals in the studied wild yam tubers were higher than earlier reported value of cultivated species³. Different processing led to decrease in mineral content in studied yam tuber, However, the decrease was more pronounced by boiling and soaking due to leaching of the minerals. In the studied yam tubers potassium was the abundant mineral and the level of potassium was higher than the sodium content. This suggests that studied yam tubers are safe food and better for human health³⁸. Different traditional processing methods led to significant reduction of nutrient and anti-nutrient compositions with improvement of tuber flour physico-functional parameter. Though processing of tuber by sun drying retained more nutrients and minerals compositions compared to boiling and soaking, but it is not suitable for detoxification/removal of antinutrients from the wild *Dioscorea* tubers.

Conclusion

Some wild yam species, such as *D. hamiltonii*, *D. pubera* and *D. bulbifera* showed higher nutritional and mineral compositions compared to the cultivated species. Processing of yam tuber by boiling led to greater retention of nutrients and moderate loss of minerals as well as with successive removal of anti-nutrients. Though, sun drying is also a competitive practice for retaining the maximum nutrients after processing but because of presence of anti-nutrients, it is not suitable for consumption. Hence, it is suggested that these wild yams should be consumed in boiled form. Processing of these yam tubers interestingly improves its physico-functional properties, having bioprospecting potential for food industries.

Acknowledgements: Authors are thankful to Prof S K Palita, HOD, DBCNR for valuable suggestions and University Grant Commission (UGC) is acknowledged for providing Non-NET Fellowship.

Conflicts of Interest: The authors declare that there are no conflicts of interest.

References

- 1 Kumar S, Parida AK & Jena PK, Ethno-Medico-Biology of Bān-Aālu (*Dioscorea* species): A neglected tuber crops of Odisha, India. *Int J Pharmacy Life Sci*, 4 (2013) 3143.
- 2 IITA, Yam. Research review. Ibadan, Nigeria: International Institute of Tropical Agriculture, (2006) 1–4.
- 3 FAO, Food and Agriculture Organization Statistics database agriculture. FAOSTAT, Rome (2002).
- 4 Behera KK, Maharana T, Sahoo S & Prusti A, Biochemical quantification of protein, fat, starch, crude fibre, ash and dry matter content in different collection of greater yam (*Dioscorea alata* L.) found in Orissa. *Nature Science*, 7 (2009) 24.
- 5 Kouakou MD, Dabonne S, Guehi T & Kuoame LP, Effects of post-harvest storage on some biochemical parameters of different parts of two yams species (*Dioscorea* spp.). *Afr J Food Sci Technol*, 1 (2010) 1.
- 6 Amanze NJ, Agbo NJ, Eke-Okoro ON & Njoku DN, Selection of yam seeds from open pollination for adoption in yam (*Dioscorea rotundata* Poir) production zones in Nigeria. *J Plant Breed Crop Sci*, 3 (2011) 68.
- 7 Dansi A, Mignouna HD, Zoundjihékpon J, Sangare A, Asiedu R & Quin FM, Morphological diversity, cultivar groups and possible descent in the cultivated yams (*Dioscorea cayenensis*-*D. rotundata* complex) in Benin. *Genet Resour Crop Evol*, 46 (1999) 371.
- 8 Adepoju OT, Boyejo O & Adeniji PO, Effects of processing methods on nutrient and antinutrient composition of yellow yam (*Dioscorea cayenensis*) products. *Food Chem*, 11 (2018) 428.
- 9 Padhan B & Panda D, Wild tuber species diversity and its ethno-medicinal use by tribal people of Koraput district of Odisha. *Ind J Natural Products Resour*, 2 (2016) 33.
- 10 Bhandari MR, Kasai T & Kawabata J, Nutritional evaluation of wild yam (*Dioscorea* spp.) tubers of Nepal. *Food Chem*, 82 (2003) 619.
- 11 Shajeela PS, Mohan VR, Jesudas LL & Soris PT, Nutritional and antinutritional evaluation of wild yam (*Dioscorea* spp.). *Trop Subtrop Agroecosyst*, 14 (2011) 723.
- 12 Polycarp D, Afoakwa EO, Budu AS & Otoo E, Characterization of chemical composition and anti-nutritional factors in seven species within the Ghanaian yam (*Dioscorea*) germplasm. *Int Food Res J*, 19 (2012) 985.
- 13 Mishra S & Chaudhury SS, Ethnobotanical flora used by four major tribes of Koraput, Odisha, India. *Genet Resour Crop Evol*, 59 (2012) 793.
- 14 Mishra S, Swain S, Chaudhary S & Ray S, Wild edible tubers (*Dioscorea* spp.) and their contribution to the food security of tribes of Jeypore tract, Orissa, India. *PGR Newsletter*, 56 (2011) 63.
- 15 Kaur M, Kaushal P & Sandhu KS, Studies on physicochemical and pasting properties of taro (*Colocasia esculenta* L.) flour in comparison with a cereal, tuber and legume flour. *J Food Sci Technol*, 50 (2011) 94.
- 16 Padhan B & Panda D, Variation of photosynthetic characteristics and yield in wild and cultivated species of yams (*Dioscorea* spp.) from Koraput, India. *Photosynthetica* 56 (2018) 1010.
- 17 AOAC, Official methods of analysis of the Association of Official Analytical Chemists, (15th ed.). Washington DC, 1990.
- 18 Phillips RD, Chinnan MS, Branch AL, Miller J & Mcwatters KH, Effects of pre-treatment on functional and nutritional properties of cowpea meal. *J Food Sci*, 53 (1988) 805.
- 19 Anderson RA, Conway HF, Pfeifer VF & Griffin EL, Roll and extrusion-cooking of grain sorghum grits. *Cereal Sci Today*, 14 (1969) 372.
- 20 Coffman CW & Garcia W, Functional properties and amino acid content of protein isolate from mung bean flour. *J Food Technol*, 12 (1977) 473.
- 21 Craig SAS, Maningat CC, Seib PA & Hoseney RC, Starch paste clarity. *Cereal Chem*, 66 (1989) 173.
- 22 Sadasivam S & Manickam A, Biochemical Methods. New Age International (P) Limited, Biochemistry, 2007, 284.
- 23 Lowry OH, Rosebrough NJ, Farr AL & Randall RJ, Protein measurement with Folin-Phenol reagent. *J Biol Chem*, 193 (1951) 263.
- 24 Omaye ST, Tuenbull TD & Sauberlich HE, Selected method for the determination of ascorbic acid in animal cells, tissues and fluid. In: Mc Cormic, D. B., wright, D.L. (Eds.), *Methods enzymol*, Vol. 62. Academic press New York, 1979, 3.
- 25 Baker H, Frank O, Angelis B & Feingold S, Plasma tocopherol in man at various times after ingesting free or acetylated tocopherol. *Nutr Rep Int*, 21 (1980) 531.
- 26 Uematsu Y, Hirata K, Saito K & Kudo I, Spectrophotometric determination of saponin in Yucca extract used as food additive. *JAOAC Int*, 83 (2000) 1451.
- 27 Meena BA, Umapathy KP, Pankaja N & Prakash J, Soluble and insoluble oxalates in selected foods. *J Food Sci Technol*, 24(1987) 43.
- 28 Wheeler EL & Ferrel RE, A method for Phytic Acid Determination in Wheat and Wheat Fractions. *Cereal Chem*, 48 (1971) 312.
- 29 Alonso R, Orúe E & Marzo F, Effects of extrusion and conventional processing methods on protein and antinutritional factor contents in pea seeds. *Food Chem*, 63 (1998) 505.
- 30 Fekadu H, Beyene F & Desse G, Effect of traditional processing methods on nutritional composition and anti-nutritional factors of anchote (*Coccinia Abyssinica* (lam.) Cogn) tubers grown in Western Ethiopia. *J Food Process Technol*, 4 (2013) 249.
- 31 Aruah BC, Uguru MI & Oyiga BC, Genetic variability and inter-relationship among some Nigerian pumpkin accessions (*Cucurbita* spp.). *Int J Plant Breed*, 6 (2012) 34.
- 32 Bhatt A. & Patel V, Antioxidant potential of banana: Study using simulated gastrointestinal model and conventional extraction. *Ind J of Exp Biol*, 53(2015) 367
- 33 Amon AS, Soro RY, Assemmand EF, Dué EA & Kouamé LP, Effect of boiling time on chemical composition and physico-functional properties of flours from taro (*Colocasia esculenta* cv *foué*) corm grown in Côte d'Ivoire. *J Food Sci Technol*, 51 (2014) 855.
- 34 Bradbury JH & Holloway WD, Chemistry of tropical root crops: Significance for nutrition and agriculture in the Pacific. Australian Centre for International Agricultural Research. Canberra, 1988, 89.

- 35 Wanasundera JPD & Ravindran G, Nutritional assessment of yam (*Dioscorea alata*) tubers. *Plant Foods Hum Nutr*, 46(1994) 33.
- 36 Adepoju OT, Effects of processing methods on nutrient retention and contribution of white yam (*Dioscorea rotundata*) products to nutritional intake of Nigerians. *Afr J Food Sci*, 6 (2012) 163.
- 37 Phillippy BQ, Lin M & Rasco B, Analysis of phytate in raw and cooked potatoes. *J Food Compos Anal*, 17 (2004) 217.
- 38 Aradhana, Rao AR, & Kale RK, Diosgenin-a growth stimulator of mammary gland of ovariectomized mouse. *Ind J of Exp Biol*, 30(1992) 367.