

Endocrine

Antineoplastic activity of Artemisin in Adrenocortical Carcinoma

--Manuscript Draft--

Manuscript Number:	ENDO-D-19-00433R2
Full Title:	Antineoplastic activity of Artemisin in Adrenocortical Carcinoma
Article Type:	Research Letter
Corresponding Author:	Luigi Iorini, M.D. Azienda Socio Sanitaria Territoriale degli Spedali Civili di Brescia ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Azienda Socio Sanitaria Territoriale degli Spedali Civili di Brescia
Corresponding Author's Secondary Institution:	
First Author:	Luigi Lorini, M.D.
First Author Secondary Information:	
Order of Authors:	Luigi Lorini, M.D. Salvatore Grisanti Roberta Ambrosini Deborah Cosentini Marta Laganà Luigi Grazioli Guido Alberto Massimo Tiberio Sandra Sigala Alfredo Berruti
Order of Authors Secondary Information:	
Funding Information:	
Response to Reviewers:	<p>Professor Sebastiano Filetti Editor in Chief Endocrine</p> <p>Dear professor Filetti</p> <p>Here attached please find the revised version of the paper entitled "Antineoplastic activity of Artemisin in Adrenocortical Carcinoma." By Luigi Lorini et al that we wish to resubmit as Research Letter to Endocrine. As itemized below, in the new version all the minor comments/chriticisms raised by the referee 2 were addressed. Looking forward to the decision of the Editorial Office we remain</p> <p>Yours sincerely</p> <p>Luigi Lorini on behalf of all co-authors</p> <p>Replies to the Reviewer #2 comments</p> <p>I read with attention the manuscript entitled "antineoplastic activity of artemisin in adrenocortical carcinoma". It addresses an interesting case of a possible</p>

antineoplastic activity of artemisin in a patient with advanced adrenocortical carcinoma. Despite, as mentioned by the authors, is difficult to establish if artemisin has had a role in slowing the disease progression, this is a promising observation that could warrant the implementation of preclinical and clinical studies in the following years, considering the limited availability of anticancer drugs in the treatment of adrenal carcinoma, especially in advanced patients.

We thank the reviewer for the appreciation of our paper

I suggest to report how was PET (or CT or RM) after 33 months of EDP-M scheme
To address this issue the following sentence was added: "In November 2011 an abdomen MRI showed a hypervascularized nodule attached to the left diaphragmatic side, confirmed by a subsequent PET with fdg ." (page 3 lines 11-12)

What do you mean by indolent progression? Please specify the radiologic findings during the subsequent chemotherapy lines and if chemotherapy was going on when there was disease progression

We have better clarified the point and the following paragraph was added in the text:
"Second line chemotherapy with carboplatin + paclitaxel was introduced leading to a radiological response lasting 10 months of both diaphragmatic (15mm vs 23mm) and mesenteric (5 mm vs 10mm) lesions. At disease progression, Gemcitabine plus capecitabine was administered but this regimen was interrupted after 2 months due to inefficacy. Mitotane therapy was never interrupted.

In December 2013, a further disease progression was observed: at TC scan the 2 abdominal lesions converged in a large left adrenal lesion (51x12mm).

During the subsequent months, the disease showed only limited progression.

Therefore, due to the rather indolent behavior of the disease, in March 2015 the patient underwent surgery with radical intent. The histological examination confirmed the ACC recurrence" (page 3, lines 13-20).

[Click here to view linked References](#)

Antineoplastic activity of Artemisin in Adrenocortical Carcinoma.

¹Luigi Lorini,¹Salvatore Grisanti,²Roberta Ambrosini,¹Deborah Cosentini,¹Marta Laganà,²Luigi Grazioli,
⁴Guido AM Tiberio,³Sandra Sigala,¹Alfredo Berruti

¹Medical Oncology Unit, Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, University of Brescia. ASST Spedali Civili, Brescia, Italy

²Department of Radiology, ASST Spedali Civili, Brescia, Italy

³Section of Pharmacology, Department of Molecular and Translational Medicine, University of Brescia, Brescia, Italy

⁴Surgical Clinic, Department of Clinical and Experimental Sciences, University of Brescia and ASST Spedali Civili di Brescia, Brescia, Italy.

Correspondence to:

Alfredo Berruti

Oncologia Medica

ASST-Spedali Civili di Brescia

Piazzale Spedali Civili 1

25123 Brescia

Italy

Tel +39 0303995410

Email: alfredo.berruti@gmail.com

INTRODUCTION

Artemisia annua is a medicinal plant that is used in China to treat various diseases for over two millennia. Preclinical studies revealed that the compound has a cytotoxic activity against various types of cancer cells(1)

Data on anticancer activity of artemisin in humans are limited, but case reports and small case series have shown encouraging results in patients affected by metastatic breast cancer, colorectal cancer and uveal melanoma (2).

The main mechanism of artemisin antineoplastic activity is not fully elucidated. This drug and relevant derivatives display a multi-target activity, as frequently observed in most natural products (3). Indeed, artemisin compounds could block the activation of intracellular pathways such as Wnt/ β -catenin, BCR/ABL, or by inducing cell cycle arrest, alteration of DNA repair mechanisms and inducing death via apoptosis and non-apoptotic pathways.

Concerning the artemisin safety profile, reported adverse effects, such as renal, hepatic and neurological toxicity were unfrequent and mild. Only one case of toxic brainstem encephalopathy was observed in a patient after only 2 week treatment of artemisin (400mg)(4).

The good safety profile and the multi-target mechanism of action makes this drug potentially useful in association with current available therapies, in order to potentiate their efficacy. In particular, the observed activity of artemisin in reversing resistance to chemotherapy is suggestive for the potential synergism of action between artemisin and cytotoxic drugs (5). Indeed, the drug administered in association with gemcitabine or cyclophosphamide was found to improve the response rate in patients with pancreatic cancer and lung cancer, respectively (6) (7). Further synergistic effects were observed with others antineoplastic compounds, such as cisplatin, carboplatin, doxorubicin and temozolomide(8).

AdrenoCortical Carcinoma (ACC) is a rare and aggressive neoplasm. The prognosis of metastatic patients is poor(9)(10)

Mitotane or mitotane plus cytotoxic chemotherapy (etoposide, doxorubicin and cisplatin–EDP-M scheme) are the standard systemic treatments in the management of metastatic ACC patients (11). No effective second-line therapies are available (12)(13)(14). Modern molecular target therapies and immunotherapies failed to demonstrate a significant activity (15)(16). On these basis, new treatment strategies are needed. p53 mutation and wnt- β catenin amplification are frequently involved in the ACC carcinogenesis. Since both pathways were reported to be targeted by artemisin and its derivatives, this observation makes artemisin potentially active in the management of ACC(17).

1 We present in this paper the case of a male patient, bearing ACC with liver and peritoneal metastases, who
2 obtained a durable disease response after oral assumption of artemisin, taken after failure of several
3 antineoplastic therapies.

4 5 6 CASEPRESENTATION

7
8 A 51 years old patient underwent left adrenalectomy for ACC in May 2008. Surgery was complicated by
9 capsular rupture. The histological examination showed: adrenal lesion of 7 x 5.5 x 4.5 cm. No adjuvant
10 mitotane therapy was prescribed.

11
12
13 Disease progression occurred 18 months later, when a FDG PET, performed in November 2009, showed
14 local recurrence (45mm) together with multiple metastatic lesions located between spleen and
15 diaphragma. The patient was submitted to EDP-M scheme followed by maintenance of mitotane therapy,
16 leading to a partial response lasting 33 months. In November 2011 a MRI showed hypervascularized
17 nodules on left diaphragma and mesenterial tissue near the spleen, confirmed by a subsequent FDG-PET.
18
19
20

21
22
23 Second line chemotherapy with carboplatin + paclitaxel was introduced leading to a radiological response
24 lasting 10 months of both diaphragmatic (15mm vs 23mm) and mesenteric (5 mm vs 10mm) lesions. At
25 disease progression, Gemcitabine plus capecitabine was administered but this regimen was interrupted
26 after 2 months due to inefficacy. Mitotane therapy was never interrupted.
27
28
29

30
31
32 In December 2013, a further disease progression was observed: at TC scan the 2 abdominal lesions
33 converged in a large left adrenal lesion (51x12mm). During the subsequent months, the disease showed
34 only limited progression. Therefore, due to the rather indolent behavior of the disease, in March 2015 the
35 patient underwent surgery with radical intent. The histological examination confirmed the ACC recurrence.
36
37
38

39
40 A newer disease progression was observed in September 2015 when a diaphragmatic mass (42 mm) was
41 detected. This lesion underwent a local regional therapy with gamma knife attaining a 20% size reduction.
42

43
44 The disease remained stable till April 2017 when a MRI showed in T2 weighted images (T2W) two slightly
45 hyperintense metastatic lesions in liver segment 8 (S8) and S7 (8 mm and 7 mm) with restricted diffusion in
46 diffusion weighted images (DWI b800). The lesions were hypointense in ADC (apparent diffusion coefficient)
47 map and hepatobiliary phase post Gadoteric acid injection (figure 1a)
48
49
50

51 In May 2017, after a consultation to an herbalist, the patient started taking Artemisia Annuata tabs
52
53 (Artemisin 99%) (600mg die for five days followed by 5 days off). The treatment was well tolerated, the
54 patient did not suffer from any symptoms and no clinical sign of toxicity was observed at clinical
55
56
57 examination.

58
59 In July 2017, a MRI showed a marked size reduction of the 2 metastases in S8 and S7 as confirmed at
60 DWI b800 and ADC map (5 mm vs 8 mm and 5 mm vs 7 mm) (figure 1b).
61

1 The response obtained was maintained at the MRI performed in January 2018 showing stable disease of
2 the lesion in S7 and S8 at T2W and DWI/ ADC (figure 1c).

3
4 Artemisin assumption was prudentially interrupted in February 2018, as recommended by the herbalist,
5
6 due to the occurrence of an Herpes Zoster episode on the left side of abdomen.

7
8 In May 2018 a MRI showed an increase in size and number of the hepatic metastases in S8-S7 as compared
9
10 with previous MRI (8,2mm vs 5 mm and 6mm vs 5mm) (figure 1d).

11
12 In June 2018 the patient resumed the assumption of Artemisia Annua at the same dosage and schedule.
13
14 However a MRI performed on August 2018 showed moderate progression of disease on the two liver
15
16 lesions that showed an increased in size at S8 (11vs7mm), S7 (20 vs 8mm). In September 2018, the two
17
18 lesions at S7 and S8 underwent thermoablation leading to size reduction.

19
20 The last MRI performed on March 2019 showed further limited disease progression on the known liver
21
22 lesions in S7 and S8 (11mm vs 7 mm and 8mm vs 6mm).

23
24 At the last follow-up examination on May 2019, the patient was in good condition, he is still continuing
25
26 taking the assumption of Artemisia Annua at the same dosage without significant side effects without
27
28 taking any further specific antineoplastic treatment.

29 30 DISCUSSION

31
32 The present case describes for the first time an antineoplastic activity of Artemisin in an advanced ACC
33
34 patient. As observed in other rclinical eports, the drug was very well tolerated and the disease response
35
36 duration (about 12 months) is noteworthy, considering the clinical setting (heavily pre-treated metastatic
37
38 ACC) in which the drug was taken. It should be underlined that the disease response to artemisin in our
39
40 patient was longer to that observed with second and third line therapies, previously administered, namely
41
42 cisplatin + taxol and gemcitabine +capecitabine.

43
44 The patient continued taking artemisin after disease progression as the only systemic therapy and the
45
46 disease displayed a rather indolent behavior, without televant side effects. It is difficult to establish
47
48 whether this compound has had a role in slowing the disease progression

49
50 Due to the very limited availability of anticancer drugs in the treatment of ACC, the activity of artemisin
51
52 observed in the case presented in this paper warrants the implementation of preclinical and clinical
53
54 studies, in order to understand the molecular mechanisms of the artemisin cytotoxicity in ACC, as well as its
55
56 clinicalefficacy.

57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

COMPLIANCE WITH ETHICAL STANDARDS:

This study was supported with the contribution of FIRM Foundation (Cremona, Italy).

Conflict of Interest: Dr Luigi Lorini declares that he has no conflict of interest

Informed consent was obtained from the patient included in the study.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES

1. Zhang Y, Xu G, Zhang S, Wang D, Saravana Prabha P, Zuo Z. Antitumor Research on Artemisinin and Its Bioactive Derivatives. *Nat Prod Bioprospect*. August 2018;8(4):303–19.
2. Li Z, Li Q, Wu J, Wang M, Yu J. Artemisinin and Its Derivatives as a Repurposing Anticancer Agent: What Else Do We Need to Do? *Molecules*. 7 October 2016;21(10):1331.
3. Efferth T, Koch E. Complex Interactions between Phytochemicals. The Multi-Target Therapeutic Concept of Phytotherapy. *CDT*. 1 January 2011;12(1):122–32.
4. Panossian LA, Garga NI, Pelletier D. Toxic brainstem encephalopathy after artemisinin treatment for breast cancer. *Ann Neurol*. November 2005;58(5):812–3.
5. Liu L, Zuo LF, Guo JW. Reversal of multidrug resistance by the anti-malaria drug artesunate in the esophageal cancer Eca109/ABCG2 cell line. *Oncology Letters*. November 2013;6(5):1475–81.
6. Zhang J-L, Wang Z, Hu W, Chen S-S, Lou X-E, Zhou H-J. DHA regulates angiogenesis and improves the efficiency of CDDP for the treatment of lung carcinoma. *Microvascular Research*. May 2013;87:14–24.
7. Wang S-J, Gao Y, Chen H, Kong R, Jiang H-C, Pan S-H, et al. Dihydroartemisinin inactivates NF- κ B and potentiates the anti-tumor effect of gemcitabine on pancreatic cancer both in vitro and in vivo. *Cancer Letters*. July 2010;293(1):99–108.
8. Efferth T. Cancer combination therapies with artemisinin-type drugs. *Biochemical Pharmacology*. September 2017;139:56–70.
9. Libé R, Borget I, Ronchi CL, Zaggia B, Kroiss M, Kerkhofs T, et al. Prognostic factors in stage III–IV adrenocortical carcinomas (ACC): an European Network for the Study of Adrenal Tumor (ENSAT) study. *Ann Oncol*. October 2015;26(10):2119–25.
10. Berruti A, Libè R, Laganà M, Eddaieb H, Sukkari MA, Bertherat J, et al. Morbidity and mortality of bone metastases in advanced adrenocortical carcinoma: a multicenter retrospective study. *European Journal of Endocrinology*. May 2019;180(5):311–20.
11. Fassnacht M, Dekkers OM, Else T, Baudin E, Berruti A, de Krijger RR, et al. European Society of Endocrinology Clinical Practice Guidelines on the management of adrenocortical carcinoma in adults, in collaboration with the European Network for the Study of Adrenal Tumors. *European Journal of Endocrinology*. October 2018;179(4):G1–46.
12. Ferrari L, Claps M, Grisanti S, Berruti A. Systemic Therapy in Locally Advanced or Metastatic Adrenal Cancers: A Critical Appraisal and Clinical Trial Update. *European Urology Focus*. February 2016;1(3):298–300.
13. Sperone P, Ferrero A, Daffara F, Priola A, Zaggia B, Volante M, et al. Gemcitabine plus metronomic 5-fluorouracil or capecitabine as a second-/third-line chemotherapy in advanced adrenocortical carcinoma: a multicenter phase II study. *Endocrine-Related Cancer*. June 2010;17(2):445–53.
14. Henning JEK, Deutschbein T, Altieri B, Steinhauer S, Kircher S, Sbierra S, et al. Gemcitabine-Based

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Chemotherapy in Adrenocortical Carcinoma: A Multicenter Study of Efficacy and Predictive Factors. *The Journal of Clinical Endocrinology & Metabolism*. 1 November 2017;102(11):4323–32.

15. Terzolo M, Daffara F, Ardito A, Zaggia B, Basile V, Ferrari L, et al. Management of adrenal cancer: a 2013 update. *J Endocrinol Invest*. March 2014;37(3):207–17.

16. Cosentini D, Grisanti S, Dalla Volta A, Laganà M, Fiorentini C, Perotti P, et al. Immunotherapy failure in adrenocortical cancer: where next? *Endocrine Connections*. December 2018;E5–8.

17. Assié G, Letouzé E, Fassnacht M, Jouinot A, Luscap W, Barreau O, et al. Integrated genomic characterization of adrenocortical carcinoma. *Nat Genet*. June 2014;46(6):607–12.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Legends to figures

Figure 1: Magnetic Resonance Imaging (T2 weighted sequences) of liver metastases in a patient with adrenocortical carcinoma submitted to artemisin: 1a) baseline condition April 2017, 1b) July 2017 after 3 month treatment, 1c) January 2018, after 9 month treatment, 1d) May 2018 at disease progression after 12 month treatment.




Figure 1a: MRI April 2017, T2w sequence

Figure 1b: MRI July 2017 T2W sequence

Figure 1c: MRI January 2018 T2W sequence

Figure 1d: MRI May 2018 T2W sequence

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45