

ISSN: 2224-0616

Int. J. Agril. Res. Innov. & Tech. 3 (2): 66-71, December, 2013

Available online at <http://www.ijarit.webs.com>

EFFECT OF INTEGRATED APPROACH OF PLANT NUTRIENTS ON YIELD AND YIELD ATTRIBUTES OF DIFFERENT CROPS IN WHEAT-SESAME-T. AMAN CROPPING PATTERN

M.A. Islam^{1*}, Mst. A. Begum² and M.M. Jahangir³

Received 27 September 2013, Revised 13 December 2013, Accepted 25 December 2013, Published online 31 December 2013

Abstract

The experiment was carried out at FSRD site, Pushpopara, Pabna, during November, 2010 to December, 2011 to observe the comparative performance of integrated plant nutrients management (IPNS) system through the use of organic (cowdung, cowdung slurry) manure and inorganic fertilizer on wheat, sesame and T. Aman crops under wheat-sesame-T. Aman cropping pattern. The experiment was consisted with four treatments viz. T₁: Soil test based inorganic fertilizer dose for high yield goal, T₂: Cowdung @ 5 t ha⁻¹ + IPNS basis inorganic fertilizer dose for high yield goal, T₃: Cowdung slurry @ 5 t ha⁻¹ + IPNS basis inorganic fertilizer dose for high yield goal and T₄: Fertilizer dose usually practiced by the farmers. In case of wheat, the highest grain yield (3.80 t ha⁻¹) was obtained from bio-slurry treated plot that means T₃ treatment followed by T₂ and the lowest (3.31 t ha⁻¹) from T₄. Higher seed yield (1.31 t ha⁻¹) of sesame was obtained from T₃ that was statistically identical to T₂ and T₁ and the lower (1.01 t ha⁻¹) from T₄. For T. Aman rice, the highest grain yield (4.89 t ha⁻¹) was obtained from T₃ which was statistically indistinguishable from T₁ where as the lowest grain yield (4.1 t ha⁻¹) was recorded from T₄. Considering the whole pattern, it is observed that the highest gross return (271100 Tk ha⁻¹) was obtained from T₃ followed by T₂ and the lowest (225650 Tk ha⁻¹) from T₁ treatment. Total variable cost was recorded as the highest (100368 Tk ha⁻¹) in T₂ followed by T₃ and the lowest (86775 Tk ha⁻¹) in T₄ treatment. The highest marginal value of product (45450 Tk ha⁻¹) was recorded in T₃ followed by T₂ where as the minimum (28710 Tk ha⁻¹) was found in T₁ over the T₄ treatment. Marginal variable cost was observed as the highest (13593 Tk ha⁻¹) in T₂ treatment followed by T₃ and the minimum (8899 Tk ha⁻¹) was recorded in T₁ treatment. The highest MBCR (4.15) was recorded from T₃ followed by T₂ and the minimum (2.31) from T₄ treatment.

Keywords: IPNS System, Organic Manure, Rice Equivalent Yield, Cropping Pattern

¹On Farm Research Division, Regional Agricultural Research Station, BARI, Pabna, Bangladesh

²Department of Agronomy, Bangladesh Agricultural University, Mymensingh, Bangladesh

³Department of Soil Science, Bangladesh Agricultural University, Mymensingh, Bangladesh

*Corresponding author's email: amin_bau@yahoo.com (M.A. Islam)

Introduction

The basic concept of Integrated Plant Nutrition System (IPNS) is the management of all available plant nutrient sources, organic and inorganic, to provide optimum and sustainable crop production conditions within the prevailing farming system. Therefore, in IPNS an appropriate combination of mineral fertilizers, organic manures, crop residues, compost, N-fixing crops and bio fertilizer is used according to the local ecological conditions, land use systems and the individual farmer's social and economic conditions.

The main aim of integrated plant nutrition system is to increase and sustain soil fertility to provide a sound basis for flexible food production systems that, within the constraints of soil and climate, can grow a wide range of

crops to meet changing needs (FAO, 2001). Therefore, it is necessary to use inorganic and organic fertilizers in an integrated way so as to obtain economically profitable crop yield, without incurring loss to soil fertility (Haque *et al.*, 2001). IPNS can produce comparable or higher crop yield compared to sole fertilizer use (BARC, 2005).

Soil fertility deterioration is a major constraint for higher crop production in Bangladesh. The increasing land use intensity without adequate and balanced use of chemical fertilizers and with little or no use of organic manure have caused severe fertility deterioration of our soils resulting in stagnating or even declining of crop productivity. The farmers of this country use on an average, 215 kg nutrients/ha annually (149 kg N + 37 kg P₂O₅ + 22 kg K₂O + 7 kg S, Zn, B and

others), while the crop removal is about 280-350 kg ha⁻¹ (Islam, 2008). Since fertile soil is the fundamental resource for higher crop production, its maintenance is a prerequisite for long-term sustainable crop productivity. Soil organic matter is a key factor for sustainable soil fertility and crop productivity. Organic matter undergoes mineralization with the release of substantial quantities of N, P, and S and smaller amount of micronutrients. In Bangladesh, most of the cultivated soils have less than 1.5% organic matter and some soils even less than 1%, while a good agricultural soil should contain at least 2.5% organic matter (BARC, 2005). Moreover, this important component of soils is declining with time due to intensive cropping and use of higher doses of nitrogenous fertilizers with little or no addition of organic manure.

In Bangladesh, major food crops remove about 2.98 million tons of nutrients annually against a total addition of 0.72 million ton (Rahman *et al.*, 2008). According to an appraisal report of Bangladesh soil resources, soils of about 6.10 m ha contain very low (less than 1%) organic matter, 2.15 m ha contain low (1-2%) organic matter and the remaining 0.90 ha contain more than 2 % organic matter. Ali (1997) reported that during the years 1967-1995, the highest depletion of organic matter occurred in soils of Meghna River Floodplain (35%) followed by Madhupur Tract (29%), Brahmaputra Floodplain (21%), Old Himalayan Piedmont Plains (18%) and Gangetic Floodplain (15%).

The average organic matter content of top soils has decline by 20-46% over past 20 years due to intensive cropping without inclusion of legume crops, imbalance use of fertilizer, use of modern varieties and scanty use of organic manure. It is agreed that decrease in soil fertility is a major constraint for higher crop production in Bangladesh. The beneficial effect of organic manure in crops production has been demonstrated by many workers (Joshi *et al.*, 1994; Batsai *et al.*, 1979; Singh *et al.*, 1970; Subhan, 1991).

A suitable combination of organic and inorganic source of nutrients is necessary for sustainable agriculture that can ensure food production with high quality (Reganold *et al.*, 1990). Nambiar (1991) viewed that integrated use of organic manure and chemical fertilizers would be quite promising not only in providing greater stability in production, but also in maintaining better soil fertility. The long-term research of BRRRI revealed that the application of cowdung 5 t/ha/yr improved rice productivity as well as prevented the soil resources from degradation (Bhuiyan, 1991). Thus, it is necessary to use fertilizer and manure in an integrated way in order to obtain sustainable crop yield without

affecting soil fertility. Based on the soil fertility problem as discussed above, the present study was undertaken to investigate the effect of combined use of chemical fertilizers and organic manures in wheat-sesame-T. Aman rice cropping pattern regarding yield and economic return.

Material and Methods

The experiment was carried out at FSRD site, Pushpopara, Pabna, during November, 2010 to December, 2011 to observe the comparative performance of integrated plant nutrients management (IPNS) system through the use of organic (cowdung, cowdung slurry) and inorganic fertilizer on wheat, sesame and T. Aman crops under wheat-sesame-T. Aman cropping pattern. The trial was conducted in 6 different locations at the farmer's field. Before conducting the experiment, initial composite soil samples (0-15 cm depth) were collected from the experimental plots and were analyzed. Nutrient packages was calculated on the basis of soil test value according to the instructions outlined by BARC fertilizer recommendation guide 2005.

$$F_r = U_f - X \frac{C_i}{C_s} (S_t - L_s)$$

Where

F_r = Fertilizer nutrient required for given soil test value

U_f = Upper limit of the recommended fertilizer nutrient for the respective STVI class

C_i = Units of class intervals used for fertilizer nutrient recommendation

C_s = Units of class intervals used for STVI class

S_t = Soil test value

L_s = Lower limit of the soil test value within STVI class

The experiment was laid out in Randomized Complete Block Design with a unit plot size of 5m×6m. Four fertilizer treatments viz. T₁: Soil test based inorganic basis fertilizer dose for high yield goal, T₂: Cowdung @ 5 t ha⁻¹ + IPNS basis inorganic fertilizer dose for high yield goal, T₃: Cowdung slurry @ 5 t ha⁻¹ + IPNS basis inorganic fertilizer dose for high yield goal and T₄: Farmers practice. Variety BARI Gom 24, BARI Til 4 and Swarna was selected for wheat, sesame and T. Aman respectively by the cooperator farmers and used in the trial. Wheat seeds were sown in the field on December 15-25, 2010 maintaining spacing of 20 cm × 5 cm. Fifty percent urea and entire amount of TSP, MoP, Gypsum, Zinc sulphate, Boric acid and organic manure were applied as basal as per treatment specification. Rest 50% urea was applied at 18-21 days after sowing (DAS). The seeds of sesame were broadcasted @ 7 kg ha⁻¹ on April 6, 2011. All fertilizers with full amount except urea were applied during final land preparation. Fifty percent urea was applied as basal and the rest amount was applied as top dress during first irrigation. In case of T. Aman rice, thirty day's old seedling was transplanted on 16 and 17

August, 2011. Entire amount of all fertilizers including cowdung manure and cowdung slurry except urea were applied as basal. Urea was applied in two equal splits at 15 and 55 DAT as top dress. Intercultural operations viz. weeding, thinning, irrigation and spraying of pesticides were done as and when required in order to support normal plant growth and development. The crops were harvested after full maturity. Wheat, sesame and T. Aman were harvested on

March 12-17, July 7 and 25-29 November, 2011 respectively.

The recorded data were statistically analyzed following Gomez and Gomez (1984). All types of variable production cost are recorded to find out the marginal benefit cost ratio (MBCR). Economic analysis with respect to net return was carried out to evaluate the profitability of different treatments.

Table 1. Soil test value (Nutrient status) of the initial soil of the experimental field at FSRD site, Pushpopara, Pabna

pH	OM (%)	Total N (%)	P	S	B	K (me/100g soil)
			(µg/g soil)			
8.23	1.91	0.11	17	21	0.39	0.23
Alkaline	M	L	M	M	M	O

Table 2. Analytical value of N, P and K nutrient from different manure (Cowdung and bioslurry)

Kind of Manure	Nutrient supply (kg) from 1 ton material		
	N	P	K
Cowdung	3	1	3
Cowdung slurry	4.6	1.6	5

Table 3. Calculated amount of different nutrients as per treatment specification applied for different crops under wheat – sesame – T. Aman cropping pattern (Soil test based)

Treatment	Manure	Wheat						Sesame						T. aman					
		N	P	K	S	B	Zn	N	P	K	S	B	Zn	N	P	K	S	B	Zn
T ₁	-	112	18	47	6	1	1.5	73	16	25	9	1	1.5	84	9	27	5	1	1
T ₂	CD @ 5 t ha ⁻¹	97	13	32	6	1	1.5	58	11	10	9	1	1.5	69	4.0	12	5	1	1
T ₃	CDS @ 5 t ha ⁻¹	89	10.5	22	6	1	1.5	50	8.5	-	9	1	1.5	61	1.5	2	5	1	1
T ₄	-	69	22	19	-	-	-	35	12	18	-	-	-	69	20	37	10	-	-

Results and Discussion

Yield and yield attributes of wheat

Significantly higher plant height (98.84 cm) was obtained from T₃ that was statistically similar to T₂ and lower from T₄ treatment. The spike m⁻² (333.55) and grain spike⁻¹ (44.05) were observed as the highest from T₃ followed by T₂ and the lowest from T₄ treatment. Significantly the highest 1000 grain weight (42.61 g) was obtained from T₃ and the lowest (3.31 g) from T₄. The highest grain yield (3.80 t ha⁻¹) was obtained from bio-slurry treated plot that means T₃ treatment followed by T₂ and the lowest (3.31 t ha⁻¹) from T₄ treatment. The highest straw yield (5.92 t ha⁻¹) was also obtained from bio-slurry

treated plot i.e. from T₃ and the lowest (5.38 t ha⁻¹) from T₄. Significant variation of grain yield indicated that nutrient management packages had significant influence on wheat production. It was observed that IPNS systems with organic manure (cowdung, bio-slurry) based nutrient packages showed better performance over T₁ and T₄ packages as those dealt with chemical fertilizers only. This might be due to the positive effect of organic manure (bio-slurry, cowdung) on yield and yield contributing characters of wheat. The result of yield increment of wheat was supported by the findings of Bodruzzaman *et al.* (2002), Karki *et al.* (1996), Kologi *et al.* (1993), Maskey (1978).

Table 4. Yield and yield contributing characters of wheat at FSRD site pushpapara, Pabna

Treatment	Plant height (cm)	Spikem ⁻²	Filled grainspike ⁻¹	1000 grain weight (g)	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)
T ₁	95.63 b	312.27 c	38.58 c	42.21 b	3.51 c	5.50 c
T ₂	98.38 a	320.05 b	39.75 b	42.28 b	3.60 b	5.80 b
T ₃	98.84 a	333.55 a	44.05 a	42.61 a	3.80 a	5.92 a
T ₄	95.91 b	308.00 c	38.00 c	42.10 c	3.31 d	5.38 d
CV (%)	8.49	11.77	5.90	2.24	6.22	4.77

Within column values followed by same letter(s) did not differ significantly by DMRT

Yield and yield attributes of sesame

Data presented in the Table 5 revealed that plant height, plant population, nos. of siliqua plant⁻¹ and nos. of seeds siliqua⁻¹ of sesame were statistically non significant but varied numerically among the treatments. The higher plant height (136.8 cm) was recorded from T₃ followed by T₂ and the lower (130.1 cm) from T₄ treatment. Similar trend was found in case of plant populations. The maximum nos. of siliqua plant⁻¹ (36.16) was obtained from T₃ followed by T₂ and the minimum (33.42) from T₄ treatment. Numerically higher nos. of seeds siliqua⁻¹ (47.00) was found in T₃ followed by T₂ and the lower (44.47) from T₄ treatment. Higher seed yield (1.31 t ha⁻¹) of sesame was obtained from T₃ that was statistically identical to T₂ and T₁ and the lower (1.01 t ha⁻¹) from T₄. The cumulative

positive effect of nos. of siliqua plant⁻¹, nos. of seeds siliqua⁻¹ and weight of 1000 seeds might be contributed to higher seed yield in T₃. Lower seed yield was attained from farmers practice treatment probably the poor performance of yield contributing characters. Maximum stover yield (3.53 t ha⁻¹) was also recorded in T₃ was statistically indistinguishable from T₂ and T₁ and the minimum (3.32 t ha⁻¹) from T₄. The availability of nutrients and balanced uptake of nutrient might be enhanced optimum plant growth and finally maximized grain yield in T₃ treatment as compared to other treatments. Haruna and Abimiku, (2011) reported about higher yield of sesame from application of organic manure.

Table 5. Yield and yield contributing characters of sesame at FSRD site, Pushpapara, Pabna

Treatment	Plant height (cm)	Plant population	Nos. of siliqua plant ⁻¹	Nos. of seeds siliqua ⁻¹	1000 seeds weight (g)	Seed yield (t ha ⁻¹)	Stover yield (t ha ⁻¹)
T ₁	134.0	34.00	34.45	44.52	2.62 b	1.21 ab	3.41 ab
T ₂	135.9	35.16	35.42	45.00	2.64 ab	1.25 ab	3.44 ab
T ₃	136.8	35.42	36.16	47.00	2.68 a	1.31 a	3.53 a
T ₄	130.1	33.42	33.42	44.47	2.62 b	1.01 b	3.32 b
CV (%)	8.04	14.71	11.49	8.52	4.32	12.07	12.42

Within column values followed by same letter(s) did not differ significantly by DMRT.

Yield and yield attributes of T. Aman rice

The yield and yield contributing characters were statistically significant except panicle length (Table 6). Yield attributes like filled grain panicle⁻¹, 1000 grain weight were found as the highest in T₃ and the minimum in T₄. The results revealed that the yield contributing characters exhibited better performance due to IPNS with CDS based fertilizer management. The highest grain yield (5.05 t ha⁻¹) was obtained from T₃ which was statistically indistinguishable from T₂ where as the lowest grain yield (4.26 t ha⁻¹) was recorded from T₄ treatment. The maximum yield in T₃ treatment might be due to the cumulative positive effect of yield contributing parameters. Probably integrated fertilizer management using

both inorganic and organic sources improved the availability of nutrients and their balanced uptake facilitated optimum growth and development of the crop which ultimately increased grain yield. The poor performance of yield attributes might be attributed to lower grain yield in T₄. The highest straw yield (7.90 t ha⁻¹) was obtained from T₃ followed by T₂ and the lowest (6.6 t ha⁻¹) from T₄ treatment. Similar trend of yield increment in rice with the application of bioslurry was also found by Bodruzzaman *et al.* (2002), Kanthaswamy (1993), Tripathi (1993), Gupta (1991), Singh *et al.* (1995), Maskey (1978).

Table 6. Yield and yield contributing characters of T. aman rice at FSRD site, Pushpapara, Pabna

Treatment	Plant height (cm)	Nos. of tiller hill ⁻¹	Nos. of panicle m ⁻²	Nos. of filled grain panicle ⁻¹	1000 grain wt (g)	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)
T ₁	108.4 b	8.96 b	276.1 b	117.8 c	20.28ab	4.81 b	7.06 c
T ₂	103.1 d	8.57 c	285.7 ab	123.4 b	20.17bc	4.91 ab	7.35 b
T ₃	105.8 c	9.38 a	293.7 a	128.7 a	20.35a	5.05 a	7.90 a
T ₄	111.0 a	9.31 a	297.3 a	108.4 d	20.14c	4.26 c	6.6 d
CV (%)	6.54	8.07	10.17	7.35	4.66	5.59	6.06

Within column values followed by same letter(s) did not differ significantly by DMRT

Economics

Gross return and total variable cost for wheat, sesame and T. Aman are presented against different treatments in the Table 7. It is observed that the highest gross return (276130 Tk ha⁻¹) was obtained from T₃ followed by T₂ and the lowest (230080 Tk ha⁻¹) from T₁ treatment. Many research work related to integrated use of fertilizer have found profitable (Haque *et al.*, 2001; Gupta, 1991). Total variable cost for purchasing different inputs i.e. fertilizers, pesticides and carrying out various intercultural operations recorded as the highest (97674 Tk ha⁻¹) in T₁ followed by T₂ and the lowest (86775 Tk ha⁻¹) in T₄ treatment. Comparatively high variable cost in T₂ than T₃ treatment might be due to higher labour cost for intercultural operations i.e. weed management. Fresh cowdung contains viable weed seeds that compete with the crops and requires farmers to put extra labours for weeding where as bioslurry is also reported to be free from weed seeds (Tripathi, 1993; Van Brake, 1980). The lowest variable cost found in T₄ treatment due to less

input cost for purchasing fertilizers as farmers do not use optimum dose of fertilizers for crop production. This statement agrees to the findings reported by Jahiruddin *et al.* (2009). This has created unbalanced use of fertilizers which produces negative impact on soil fertility and crop yield and there by ultimately on economic return. The highest marginal value of product (46050 Tk ha⁻¹) was recorded in T₃ followed by T₂ where as the minimum (27440 Tk ha⁻¹) was found in T₁ over the T₄ treatment. The highest marginal value of product in T₄ treatment is mainly of higher yield and economic return facilitated by IPNS with cowdung slurry. Marginal variable cost was observed as the highest (10899 Tk ha⁻¹) in T₂ treatment followed by T₃ and the minimum (7949 Tk ha⁻¹) was recorded in T₁ treatment. Comparatively low marginal variable cost in T₂ and T₃ over T₁ treatment probably due to less input cost for purchasing chemical fertilizers. The highest MBCR (5.79) was recorded from T₃ followed by T₂ and the minimum (2.52) from T₁ treatment.

Table 7. Profitability of the IPNS system of fertilizer application over the farmers' practice obtained from wheat-sesame-T. Aman rice cropping pattern

Treatment	Rice equivalent yield (t ha ⁻¹)	Gross return from Wheat- Sesame- T. Aman Pattern (Tk ha ⁻¹)	Total variable cost for Wheat- Sesame-T. Aman Pattern (Tk ha ⁻¹)	Marginal value of product (Tk ha ⁻¹)	Marginal variable cost (Tk ha ⁻¹)	MBCR
T ₁	12.88	257520	97674	27440	10899	2.52
T ₂	13.22	264365	95425	34285	8650	3.96
T ₃	13.81	276130	94724	46050	7949	5.79
T ₄	11.50	230080	86775	-	-	-

Item of Input	Unit price (Tk kg ⁻¹)	Item of Output	Unit price (Tk kg ⁻¹)
Cowdung / Cowdung slurry	1.0	Wheat grain	20.0
Urea	20.0	Wheat straw	1.00
TSP	24.0	Rice grain	20
MoP	15.0	Rice straw	2.5
Gypsum	10.0	Sesame seed	55
ZnSO ₄	130.0	Sesame stover	1.0
Boric acid	130.0		

Conclusion

Considering yield and return it can be concluded that treatment T₃ that means IPNS with cowdung slurry is the most profitable as compared to other treatments. Many researchers claim that IPNS based fertilizer management systems have been found promising for maintaining soil health. In order to maintain sustainable agriculture we should give priority for creating awareness and using IPNS based fertilizer package for successful crop production keeping sustenance of soil health.

References

- Ali, M.M., Saheed, S.M. and Kubota, D. 1997. Soil degradation during the period 1967-1995 in Bangladesh. II. Selected chemical characters. *Soil Sci. Plant Nutr.* 43 (4): 879-890.
- BARC. 2005. Fertilizer Recommendation Guide. Bangladesh Agricultural Research Council, Farmgate, Dhaka-1215. 260 p.
- Batsai, S.T., Polyakev, A.A. and Nedbal, R.F. 1979. Effect of organic and mineral fertilizers on the yield and quality of

- irrigated late white cabbage in the steppe region of the Crimea. *Hort. Abst.* 49 (11): 730.
- Bhuiyan, N.I. 1991. Issues concerning declining/stagnating productivity in Bangladesh Agriculture. A paper presented at the National Workshop on Risk Management in Bangladesh Agriculture, held at BARC, Dhaka. August, 1991. pp. 24-27.
- Bodruzzaman, M., Sadat, M.A., Meisner, C.A., Hossain, A.B.S. and Khan, H.H. 2002. Direct and residual effects of applied organic manures on yield in a wheat-rice cropping pattern. Paper presented in 17th WCSS Symposium held in Thailand. 14-21 August, 2002. pp. 7-8.
- Brady, N.C. 1974. Organic matter of mineral soils. *In: The nature and properties of soils.* (Buckman, H.O. and Brady N.C. ed.) Macmillan Publishing Co., New York, pp. 137-163.
- FAO. 2001. Report of the 2nd Research Co-ordination Meeting of the FAO/IAEA Co-ordinated Research Project on The Use of Nuclear Techniques for developing Integrated Nutrient and Water Management Practices for Agroforestry Systems. 7-11 May, Kuala Lumpur. IAEA - 311-D1-RC-735.2.
- Gomez, K.A. and Gomez, A.A. 1984. Statistical Procedure for Agricultural Research (2nd edition), Jhon Willey and Sons, New York. pp. 202-215.
- Gupta, D.R. 1991. Bio-Fertilizer from Biogas Plants/ In Changing Villages, Vol. 10, No. 1. Jan- Mar., 1991. pp. 25-26.
- Haque, M.Q., Rahman, M.H., Islam, F., Rijpma, J. and Kadir, M.M. 2001. Integrated nutrient management in relation to soil fertility and yield sustainability under Wheat-Mung-T.Aman cropping pattern. *J. Biol. Sci.* 1 (8): 731-734.
- Haruna, I.M. and Abimiku, M.S. 2011. Yield of Sesame (*Sesamum indicum* L.) as Influenced by Organic Fertilizers in the Southern Guinea Savanna of Nigeria. *African J. Biotech.* 10 (66): 14881-14887.
- Islam, M.S. 2008. Soil fertility history, present status and future scenario in Bangladesh. Paper presented at the IPI-BFA-BRRI International workshop on Balanced Fertilization for Increasing and Sustaining Crop Productivity held at Hotel Rajmoni Ishakha, Dhaka, Bangladesh. 30 Mar.-01 Apr. 2008. 78 p.
- Jahiruddin, M., Islam, M.R. and Miah, M.A.M. 2009. Constraints of farmers' access to fertilizer for food production. Final Report. National Food Policy Capacity Strengthening Programme. FAO. Dhaka. 52 p.
- Joshi, J.R., Moncrief, J.F., Swan, J.B. and Malzer, G.L. 1994. Long-term conservation tillage and liquid dairy manure effects on corn. II. Nitrate concentration in soil water. *Soil & Till. Res.* 31 (2-3): 225-233.
- Kanthaswamy, V. 1993. Effect of bio-digested slurry in Rice. In Biogas Slurry Utilisation. New Deihi: CORT. 46 p.
- Karki, Krishna, B. and Gurung, B. 1996. Evaluation of Biogas Slurry Extension Pilot Programme. Kathmandu: BSP, SNV-Nepal. 2 p.
- Kologi, S.D., Nagalika, S.S. and Hirevendkanagoudar, L.V. 1993. Effect of Biogas Slurry in Crop Yield/ In Biogas Slurry Utilisation. New Delhi:CORT. pp. 20-22.
- Maskey, S.L. 1978. Manurial Value of Biogas Slurry: Some Observations. (Paper presented at International Workshop on Microbiological aspects of Biogas production. May 31- June 3, Kathmandu). 2 p.
- Nambiar, K.K.M. 1991. Long-term fertility effects on wheat productivity. Wheat for the nontraditional warm areas. Proceedings International Conference, Mexico, DF (Mexico) CIMMYT. pp. 516-521.
- Rahman, M.A., Ullah, M.M., Sen, R., Hasan, M.K., Islam, M.B. and Khan, M.S. 2008. Bio-Slurry management and its Effect on Soil Fertility and Crop Production. Project Report, Sept. 2008. 5 p.
- Reganold, J.P., Robert, I.P. and Parr, J.F. 1990. Sustainability of agriculture in the United States- An overview. *Sustainable Agric. Sci. Am.* 262: 112-120.
- Singh, K., Gill, J.S. and Verma, O.P. 1970. Studies on poultry manure in relation to vegetable production. *Indian. J. Hort.* 27: 42-47.
- Singh, S.P., Verma, H.N., Vatsa, D.K. and Kalia, A.K. 1995. Effect of Biogas Digested Slurry on Pea, Okra, Soybean and Maize/ In Biogas Forum Vol. IV, No. 63. 14 p.
- Subhan, 1991. Effect of organic materials on growth and production of cabbage (*Barssica oleracca* L.) *Soils & Fert.* 54 (4): 587.
- Tripathi, A.K. 1993. Biogas Slurry - A Boon for Agriculture Crops/ In Biogas Slurry Utilisation. New Delhi: CORT. pp. 11-14.
- Van Brake. 1980. The Ignis Fatuus of Biogas. Small Scale Anaerobic digesters (biogas plants): a critical review of the pre-1970 literature Delft, the Netherlands. Delft University Press. 106 p.