



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

Complete mitochondrial genome of the gray reef shark, *Carcharhinus amblyrhynchos* (Carcharhiniformes: Carcharhinidae)

Nicholas Dunn, Shaili Johri, David Curnick, Chris Carbone, Elizabeth A. Dinsdale, Taylor K. Chapple, Barbara A. Block & Vincent Savolainen

To cite this article: Nicholas Dunn, Shaili Johri, David Curnick, Chris Carbone, Elizabeth A. Dinsdale, Taylor K. Chapple, Barbara A. Block & Vincent Savolainen (2020) Complete mitochondrial genome of the gray reef shark, *Carcharhinus amblyrhynchos* (Carcharhiniformes: Carcharhinidae), *Mitochondrial DNA Part B*, 5:3, 2080-2082, DOI: [10.1080/23802359.2020.1765208](https://doi.org/10.1080/23802359.2020.1765208)

To link to this article: <https://doi.org/10.1080/23802359.2020.1765208>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 14 May 2020.



Submit your article to this journal [↗](#)



Article views: 228



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

Complete mitochondrial genome of the gray reef shark, *Carcharhinus amblyrhynchos* (Carcharhiniformes: Carcharhinidae)

Nicholas Dunn^{a,b}, Shailli Johri^{c,d}, David Curnick^a, Chris Carbone^a, Elizabeth A. Dinsdale^d, Taylor K. Chapple^e, Barbara A. Block^c and Vincent Savolainen^b

^aInstitute of Zoology, Zoological Society of London, London, UK; ^bDepartment of Life Sciences, Imperial College London, Ascot, UK; ^cHopkins Marine Station, Stanford University, Pacific Grove, CA, USA; ^dDepartment of Biology, San Diego State University, San Diego, CA, USA; ^eCoastal Oregon Marine Experiment Station, Oregon State University, Newport, OR, USA

ABSTRACT

We report the first mitochondrial genome sequences for the gray reef shark, *Carcharhinus amblyrhynchos*. Two specimens from the British Indian Ocean Territory were sequenced independently using two different next generation sequencing methods, namely short read sequencing on the Illumina HiSeq and long read sequencing on the Oxford Nanopore Technologies' MinION sequencer. The two sequences are 99.9% identical and are 16,705 base pairs (bp) and 16,706 bp in length. The mitogenome contains 22 tRNA genes, two rRNA genes, 13 protein-coding genes and two non-coding regions; the control region and the origin of light-strand replication (OL).

ARTICLE HISTORY

Received 28 March 2020
Accepted 25 April 2020

KEYWORDS

Shark; British Indian Ocean Territory; *Carcharhinus amblyrhynchos*; mitochondrial genome

Main text



The gray reef shark *Carcharhinus amblyrhynchos*, is a highly-social, reef-dependent species distributed widely in the tropical Indo-Pacific and currently listed as 'Near Threatened' in the IUCN Red List (Smale 2009). Populations have declined due to illegal fishing activities (Osgood and Baum 2015; Ferretti et al. 2018). Whilst there have been genetic studies conducted on the species (Holmes et al. 2009; Momigliano et al. 2015, 2017), its mitogenome has not yet been described.

We describe the complete mitochondrial genome of *C. amblyrhynchos*. Tissue was sampled as fin clips from two specimens in the British Indian Ocean Territory in March 2018. Specimen 1 (GenBank MT093205) was a female tagged at location -5.46386° 71.77841° and specimen 2 (GenBank MT104515) was a male tagged at -5.24956° 71.79906° . Samples were stored at Hopkins Marine Station before specimen 1's tissue was transferred to Silwood Park, Imperial College London. The samples were then analyzed independently in the two laboratories. The DNA from specimen 1 is available at Silwood Park DNA & Tissue Bank (CITES GB038) under accession VS8956-20002085971; DNA from specimen 2 is available at Hopkins Marine Station, Stanford University under accession 020002232485. For specimen 1, genomic DNA was extracted using Qiagen's Blood & Tissue Kit and was sequenced using an Illumina HiSeq. The mitochondrial genome sequence was assembled using ABySS v2.0.2 (Jackman et al. 2017) and GapCloser v1.12 (Luo et al. 2012).

For specimen 2, the DNA was extracted and sequenced using the Oxford Nanopore Technologies' MinION sequencer following Johri et al. (Johri et al. 2019). The MitoFish mitoannotator (Iwasaki et al. 2013) was used to annotate the sequences, and these were aligned against one another and mitogenomes from other Carcharhinid species using MUSCLE (Edgar 2004) within Geneious Prime (v2019.0.4). A phylogenetic tree was produced in Geneious Prime using MrBayes (Huelsenbeck and Ronquist 2001; Ronquist et al. 2012) plugin (v.3.2.6, substitution model: HKY85, burn-in length: 100,000) using the gray bamboo shark (*Chiloscyllium griseum*; NC_017882) and scalloped hammerhead shark (*Sphyrna lewini*; NC_022679) as outgroups.

The complete mitochondrial genomes are 16,705 bp (specimen 1) and 16,706 bp (specimen 2) in length. Each contains two rRNAs, 22 tRNAs, 13 protein-coding genes and a non-coding control region. The nucleotide base composition is identical with 31.5% A, 25.2% C, 13.2% G and 30.1% T, the overall GC content is 38.4%. The two sequences have 16,682 identical sites (99.9% pairwise identity). The differences include one base addition and 23 substitutions. Four substitutions result in a change to the amino acid sequence of COI. These differences could be due to the sequencing methods or represent evidence of population structure within the species in BIOT despite high spatial connectivity across the territory (Carlisle et al. 2019).

Whilst the fine-scale phylogenetic relationships within Carcharhinidae remain unresolved (Naylor et al. 2012), the tree (Figure 1) supports the placement of *C. amblyrhynchos* in

CONTACT Vincent Savolainen  v.savolainen@imperial.ac.uk  Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst Road, Ascot, SL5 7PY, UK

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Cladogram showing the phylogenetic relationship of species with complete mitogenome sequences in the genus *Carcharhinus* including *Prionace glauca*, with the scalloped hammerhead shark (*Sphyrna lewini*) and gray bamboo shark (*Chiloscyllium griseum*) as outgroups. The new sequences for the gray reef shark (*Carcharhinus amblyrhynchos*) are in bold. Families are indicated by vertical lines and represented by silhouettes accessed from PhyloPic (phylopic.org). Values at each node represent the Bayesian posterior probability at each node, GenBank accession numbers for each sequence are in brackets.

a clade with *C. albimarginatus*, *C. falciformis*, and *Prionace glauca*. The low posterior probability that supports the placement of *P. glauca* with *C. albimarginatus* and *C. falciformis* suggests that further work is required to fully resolve the tree. However, the high support for deeper clades within *Carcharhinus* adds to calls for a taxonomic revision of *P. glauca* (Naylor et al. 2012; Johri et al. 2019). The new mitochondrial genomes presented here will aid in conservation genetics, environmental DNA and population studies as researchers move toward assessing populations using genome sequences.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Author contributions

ND and SJ contributed to concept, bioinformatics and analysis, ND wrote the manuscript with comments from all authors, SJ contributed to sequencing and EAD provided laboratory support. BAB, VS and DC contributed to the concept of the manuscript and financial support of the project. BAB, TKC and DC contributed the samples.

Funding

We thank the Bertarelli Foundation and the UK Natural Environment Research Council for funding, Robert Edwards and Adrian Cantu for computing support, Danny Coffey, Tim White, Nathan Truelove, Robert Schallert, and David Jacoby for collecting samples, Theodore Reimer for logistics and CD-genomics for assistance with Illumina sequencing.

Data availability statement

The data that support the findings of this study are openly available in GenBank at <https://www.ncbi.nlm.nih.gov/nuccore/MT093205> reference number MT093205.1 and <https://www.ncbi.nlm.nih.gov/nuccore/MT104515> reference number MT104515.1.

References

- Carlisle AB, Tickler D, Dale JJ, Ferretti F, Curnick DJ, Chapple TK, Schallert RJ, Castleton M, Block BA. 2019. Estimating space use of mobile fishes in a large marine protected area with methodological considerations in acoustic array design. *Front Mar Sci.* 6:256. Available from: <https://www.frontiersin.org/article/10.3389/fmars.2019.00256/full>.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32(5):1792–1797.
- Ferretti F, Curnick D, Liu K, Romanov EV, Block BA. 2018. Shark baselines and the conservation role of remote coral reef ecosystems. *Sci Adv.* 4(3):eaag0333.

- Holmes BH, Steinke D, Ward RD. 2009. Identification of shark and ray fins using DNA barcoding. *Fish Res.* 5(2-3):280–288.
- Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics.* 17(8):754–755.
- Iwasaki W, Fukunaga T, Isagozawa R, Yamada K, Maeda Y, Satoh TP, Sado T, Mabuchi K, Takeshima H, Miya M, et al. 2013. Mitofish and mitoannotator: a mitochondrial genome database of fish with an accurate and automatic annotation pipeline. *Mol Biol Evol.* 30(11):2531–2540.
- Jackman SD, Vandervalk BP, Mohamadi H, Chu J, Yeo S, Hammond SA, Jahesh G, Khan H, Coombe L, Warren RL, et al. 2017. ABySS 2.0: resource-efficient assembly of large genomes using a Bloom filter. *Genome Res.* 27(5):768–777.
- Johri S, Solanki J, Cantu VA, Fellows SR, Edwards RA, Moreno I, Vyas A, Dinsdale EA. 2019. ‘Genome skimming’ with the MinION hand-held sequencer identifies CITES-listed shark species in India’s exports market. *Sci Rep.* 9(1):1–13
- Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, et al. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *GigaSci.* 1(1):2047–217X.
- Momigliano P, Harcourt R, Robbins WD, Jaiteh V, Mahardika GN, Sembiring A, Stow A. 2017. Genetic structure and signatures of selection in grey reef sharks (*Carcharhinus amblyrhynchos*). *Heredity (Edinb).* 119(3):142–153.
- Momigliano P, Harcourt R, Robbins WD, Stow A. 2015. Connectivity in grey reef sharks (*Carcharhinus amblyrhynchos*) determined using empirical and simulated genetic data. *Sci Rep.* 5(1):13229.
- Naylor GJ, Caira JN, Jensen K, Rosana K. a, Straube N, Lakner C. 2012. Elasmobranch phylogeny: a mitochondrial estimate based on 595 species. In: *Biology of sharks and their relatives*. 2nd (Carrier, J. C., Musick, J. A. & Heithaus, M. R.), 31–56. Boca Raton, FL: CRC Press
- Osgood GJ, Baum JK. 2015. Reef sharks: recent advances in ecological understanding to inform conservation. *J Fish Biol.* 87(6):1489–1523.
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol.* 61(3):539–42
- Smale M. 2009. *Carcharhinus amblyrhynchos*, Grey Reef Shark. IUCN Red List Threat Species. 8235:e.T39365A10216946.