



UNIVERSITY OF LEEDS

This is a repository copy of *The (theta, wheel)-free graphs Part IV: Induced paths and cycles*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/161693/>

Version: Accepted Version

Article:

Radovanović, M, Trotignon, N and Vuskovic, K (2021) The (theta, wheel)-free graphs Part IV: Induced paths and cycles. *Journal of Combinatorial Theory: Series B*, 146. pp. 495-531. ISSN 0095-8956

<https://doi.org/10.1016/j.jctb.2020.06.002>

© 2020, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

The (theta, wheel)-free graphs

Part IV: induced paths and cycles

Marko Radovanović*, Nicolas Trotignon†, Kristina Vušković‡

May 27, 2020

Abstract

A hole in a graph is a chordless cycle of length at least 4. A theta is a graph formed by three internally vertex-disjoint paths of length at least 2 between the same pair of distinct vertices. A wheel is a graph formed by a hole and a node that has at least 3 neighbors in the hole. In this series of papers we study the class of graphs that do not contain as an induced subgraph a theta nor a wheel. In Part II of the series we prove a decomposition theorem for this class, that uses clique cutsets and 2-joins. In this paper we use this decomposition theorem to solve several problems related to finding induced paths and cycles in our class.

1 Introduction

In this paper all graphs are finite and simple. We say that a graph G contains a graph H if H is isomorphic to an induced subgraph of G , and that G is H -free if it does not contain H . For a family of graphs \mathcal{H} , G is \mathcal{H} -free if for every $H \in \mathcal{H}$, G is H -free.

A hole in a graph is a chordless cycle of length at least 4. A theta is a graph formed by three paths between the same pair of distinct vertices so that the union of any two of the paths induces a hole. This implies that each of the three paths has length at least 2 (a path of length 1 would form a chord of the cycle induced by the two other paths). A wheel is a graph formed by a hole and a node that has at least 3 neighbors in the hole.

In this series of papers we study the class of (theta, wheel)-free graphs, that we denote by \mathcal{C} throughout the paper. This project is motivated and explained in more detail in Part I

*University of Belgrade, Faculty of Mathematics, Belgrade, Serbia. Partially supported by Serbian Ministry of Education, Science and Technological Development project 174033. E-mail: markor@matf.bg.ac.rs

†Univ Lyon, EnsL, UCBL, CNRS, LIP, F-69342, LYON Cedex 07, France. Partially supported by the LABEX MILYON ANR-10-LABX-0070. E-mail: nicolas.trotignon@ens-lyon.fr

‡School of Computing, University of Leeds, Leeds, UK and Faculty of Computer Science (RAF), Union University, Belgrade, Serbia. Partially supported by EPSRC grant EP/N019660/1, and Serbian Ministry of Education, Science and Technological Development projects 174033 and III44006. E-mail: k.vuskovic@leeds.ac.uk

of the series [7], where two subclasses of \mathcal{C} are studied. In Part II of the series [20], we prove a decomposition theorem for graphs in \mathcal{C} that uses clique cutsets and 2-joins, and use it to obtain a polynomial time recognition algorithm for the class. In Part III of the series [21] we use the decomposition theorem from [20] to obtain further properties of graphs in the class and to construct polynomial time algorithms for maximum weight clique, maximum weight stable set, and coloring problems. In this part we use the decomposition theorem from [20] to obtain polynomial time algorithms for several problems related to finding induced paths and cycles.

The DISJOINT PATHS problem is to test whether a graph G with k pairs of specified vertices $(s_1, t_1), \dots, (s_k, t_k)$ contains vertex-disjoint paths P_1, \dots, P_k such that, for $i = 1, \dots, k$, P_i is a path from s_i to t_i . When k is part of the input, this problem is NP-complete [15]. If k is any fixed integer (i.e. not part of the input) then the problem is called k -DISJOINT PATHS and can be solved in $\mathcal{O}(n^3)$ time as shown by the linkage algorithm of Robertson and Seymour [22]. (Throughout the paper n will denote the number of vertices and m the number of edges of an input graph.) In this paper we consider the induced variant of this problem. We note that there are slight differences in the definition of this problem (for example in [13]; see also Subsection 4.4).

INDUCED DISJOINT PATHS problem (G, \mathcal{W})

Instance: A graph G and a set $\mathcal{W} = \{(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)\}$ of pairs of vertices of G such that all $2k$ vertices are distinct and the only edges between these vertices are of the form $s_i t_i$, for some $1 \leq i \leq k$.

Question: Does G contain k vertex-disjoint paths $P_i = s_i \dots t_i$, $i \in \{1, \dots, k\}$, such that no vertex of P_i is adjacent to a vertex of P_j , for every $i \neq j$?

For chordal graphs [1] this problem can be solved in time $\mathcal{O}(kn^3)$. When k is fixed (i.e. not part of the input) then the problem is called k -INDUCED DISJOINT PATHS. The k -INDUCED DISJOINT PATHS is NP-complete whenever $k \geq 2$ as proved by Bienstock [2], even when restricted to several classes of graphs (see [18]). It has been studied in several classes of graphs (but not many if compared to other problems such as graph coloring or maximum stable set for instance). It can be solved for line graphs as a simple corollary of the linkage algorithm mentioned above. For any fixed k , it can be solved in linear time for planar graphs [16] and circular-arc graphs [14], and in polynomial time for AT-free graphs [12] and claw-free graphs [9].

Golovach, Paulusma and van Leeuwen [13] proved that for claw-free graphs the k -INDUCED DISJOINT PATHS is fixed-parameter tractable when parameterized by k , meaning that it can be solved for any fixed k in time $f(k)n^c$, where c is a constant that does not depend on k , and f is a computable function that depends only on k .

Here, we prove that the INDUCED DISJOINT PATHS problem remains NP-complete on \mathcal{C} , and we give a polynomial time algorithm for the k -INDUCED DISJOINT PATHS problem on \mathcal{C} . We also consider a number of related problems, namely k -IN-A-PATH, k -IN-A-TREE

and H -INDUCED-TOPOLOGICAL MINOR, to be defined in Section 4. These problems were already studied in different settings and classes of graphs, see [5, 6, 8, 19].

Outline of the paper

In Section 2 we state several results for the class \mathcal{C} that are proved in previous parts of this series and that will be needed here. Most importantly, we give the decomposition theorem for \mathcal{C} from [20] and recall some results that will help us manage the cutsets that appear in this decomposition theorem. In fact, fundamental for our algorithms are the 2-join decomposition techniques developed in [25], which we also describe here.

In Section 3 we prove that for a graph $G \in \mathcal{C}$ the only obstruction for the existence of an induced cycle through the given two non-adjacent vertices is a clique cutset of G that separates these vertices. Note that a similar result holds for claw-free graphs as proved by Bruhn and Saito [3]. Using our result we derive an $\mathcal{O}(nm)$ -time algorithm for the problem of finding a chordless cycle through two prescribed vertices of a graph in \mathcal{C} . Also, we show that k -IN-A-CYCLE problem is fixed-parameter tractable (when parameterized by k) for graphs in \mathcal{C} .

In Section 4 we give an $\mathcal{O}(n^{2k+6})$ -time algorithm for the k -INDUCED DISJOINT PATHS for graphs in \mathcal{C} . As an intermediate step we show that the INDUCED DISJOINT PATHS problem is fixed-parameter tractable (when parameterized by k) for the class of graphs of \mathcal{C} that do not have clique cutsets. We also show that if \mathcal{G} is any hereditary class of graphs such that there exists an $\mathcal{O}(n^c)$ -time algorithm (where c is a constant that does not depend on k) for the k -INDUCED DISJOINT PATHS on the graphs of \mathcal{G} that do not have clique cutsets, then the k -INDUCED DISJOINT PATHS problem can be solved on \mathcal{G} in $\mathcal{O}(n^{2k+c})$ -time.

In Section 4 we show that the generalization of the k -INDUCED DISJOINT PATHS problem where the terminals (i.e. vertices $s_1, \dots, s_k, t_1, \dots, t_k$) are not necessarily distinct and where there could possibly be edges between the terminal vertices, can be solved in $\mathcal{O}(n^{4k+6})$ -time. Consequently we obtain polynomial-time algorithms for a number of related problems (on \mathcal{C}) such as k -IN-A-PATH, k -IN-A-TREE and H -INDUCED TOPOLOGICAL MINOR.

Finally, in Section 5 we give several NP-completeness results.

Terminology and notation

A *clique* in a graph is a (possibly empty) set of pairwise adjacent vertices. A *stable set* in a graph is a (possibly empty) set of pairwise nonadjacent vertices. A *diamond* is a graph obtained from a complete graph on 4 vertices by deleting an edge. A *claw* is a graph induced by nodes u, v_1, v_2, v_3 and edges uv_1, uv_2, uv_3 .

A *path* P is a sequence of distinct vertices $p_1 p_2 \dots p_k$, $k \geq 1$, such that $p_i p_{i+1}$ is an edge for all $1 \leq i < k$. Edges $p_i p_{i+1}$, for $1 \leq i < k$, are called the *edges of* P . Vertices p_1 and

p_k are the *endnodes* of P . A *cycle* C is a sequence of vertices $p_1p_2 \dots p_kp_1$, $k \geq 3$, such that $p_1 \dots p_k$ is a path and p_1p_k is an edge. Edges $p_i p_{i+1}$, for $1 \leq i < k$, and edge p_1p_k are called the *edges of C* . Let Q be a path or a cycle. The vertex set of Q is denoted by $V(Q)$. The *length* of Q is the number of its edges. An edge $e = uv$ is a *chord* of Q if $u, v \in V(Q)$, but uv is not an edge of Q . A path or a cycle Q in a graph G is *chordless* if no edge of G is a chord of Q .

Let G be a graph. We denote the vertex set of G by $V(G)$. For $v \in V(G)$ and $T \subseteq V(G)$, $N_T(v)$ is the set of neighbors of v in T . Also, if $T = V(G)$, then we use $N(v)$ to denote $N_T(v)$. For $S \subseteq V(G)$, $N_G(S)$ (or simply $N(S)$ when clear from context) denotes the set of vertices $u \in V(G) \setminus S$ such that for some $v \in S$, uv is an edge of G . Also, for $S \subseteq V(G)$, $G[S]$ denotes the subgraph of G induced by S and $G \setminus S$ the subgraph of G induced by $V(G) \setminus S$. For disjoint subsets A and B of $V(G)$, we say that A is *complete* (resp. *anticomplete*) to B if every vertex of A is adjacent (resp. nonadjacent) to every vertex of B .

When clear from the context, we will sometimes write G instead of $V(G)$.

2 Decomposition of (theta, wheel)-free graphs

To state the decomposition theorem for graphs in \mathcal{C} we first define the basic classes involved and then the cutsets used.

Basic classes

If R is a graph, then the *line graph* of R , denoted by $L(R)$, is the graph whose vertices are the edges of R , and such that two vertices of $L(R)$ are adjacent if and only if the corresponding edges are adjacent in R . A graph G is *chordless* if no cycle of G has a chord.

In the decomposition theorem for (theta, wheel)-free graphs obtained in [20] we have two types of *basic graphs*: line graphs of triangle-free chordless graphs and P-graphs. Their union is denoted by \mathcal{B} . The definition of a P-graph is a bit technical (see [20]), and in this paper we do not need it in full. We therefore just list the properties that we need. Fortunately, these properties are common to both types of basic graphs. Here they are:

- every graph $G \in \mathcal{B}$ is induced by $V(L(R)) \cup K$, where $V(L(R))$ and K are disjoint;
- R is triangle-free and chordless and K is a clique (it is possible that $K = \emptyset$);
- each vertex of $L(R)$, that corresponds to an edge of R incident with a degree 1 vertex (in R), is adjacent to at most one vertex of K , and these are the only edges between $L(R)$ and K ;
- if a vertex v of $L(R)$, that is of degree at least 2 in $L(R)$, is adjacent to a vertex w of K , then w has no other neighbors in $L(R)$ (note that this implies: if a vertex v of

$L(R)$ is adjacent to a vertex w of K that has some other neighbor in $L(R)$, then the degree of v in G is 2).

We say that R is the skeleton and K the special clique of G . Clearly every line graph of a triangle-free chordless graph satisfies the above properties. The fact that P-graphs satisfy them follows from conditions (i), (vi) and (viii) in the definition of P-graphs in [20]. We also observe that all induced subgraphs of graphs in \mathcal{B} satisfy these properties. Note that each $G \in \mathcal{B}$ is diamond-free (since $L(R)$ does not contain a diamond, each vertex of $L(R)$ has at most one neighbor in K and no vertex of K is adjacent to more than 1 vertex that is of degree at least 2 in $L(R)$), and center of every claw of G is contained in K .

Cutsets

In a graph G , a subset S of nodes and/or edges is a *cutset* if its removal yields a disconnected graph. A node cutset S is a *clique cutset* if S is a clique. Note that every disconnected graph has a clique cutset: the empty set.

An *almost 2-join* in a graph G is a pair (X_1, X_2) that is a partition of $V(G)$, and such that:

- For $i \in \{1, 2\}$, X_i contains disjoint nonempty sets A_i and B_i , such that A_1 is complete to A_2 , B_1 is complete to B_2 , and there are no other adjacencies between X_1 and X_2 .
- For $i \in \{1, 2\}$, $|X_i| \geq 3$.

An almost 2-join (X_1, X_2) is a *2-join* when for $i \in \{1, 2\}$, X_i contains at least one path from A_i to B_i , and if $|A_i| = |B_i| = 1$ then $G[X_i]$ is not a chordless path.

We say that $(X_1, X_2, A_1, A_2, B_1, B_2)$ is a *split* of this (almost) 2-join, and the sets A_1, A_2, B_1, B_2 are the *special sets* of this (almost) 2-join. We often use the following notation: $C_i = X_i \setminus (A_i \cup B_i)$ (possibly, $C_i = \emptyset$).

We are ready to state the decomposition theorem from [20].

Theorem 2.1 ([20]) *If G is (theta, wheel)-free, then G is a line graph of a triangle-free chordless graph or a P-graph, or G has a clique cutset or a 2-join.*

It trivially follows that if $G \in \mathcal{C}$ then $G \in \mathcal{B}$ or G has a clique cutset or a 2-join. We now describe how we decompose a graph from \mathcal{C} into basic graphs using the cutsets in the above theorem.

Decomposing with clique cutsets

If a graph G has a clique cutset K , then its node set can be partitioned into sets (A, K, B) , where A and B are nonempty and anticomplete. We say that (A, K, B) is a *split* for the clique cutset K . When (A, K, B) is a split for a clique cutset of a graph G , the *blocks*

of decomposition of G with respect to (A, K, B) are the graphs $G_A = G[A \cup K]$ and $G_B = G[K \cup B]$.

A *clique cutset decomposition tree of depth p* for a graph G is a rooted tree T defined as follows.

- (i) The root of T is $G_0 = G$.
- (ii) The non-leaf nodes of T are G_0, G_1, \dots, G_{p-1} . Each non-leaf node G_i has two children: one is G_{i+1} and the other one is G_{i+1}^B .
The leaf-nodes of T are graphs $G_1^B, G_2^B, \dots, G_p^B$ and G_p . Graphs $G_1^B, G_2^B, \dots, G_p^B, G_p$ have no clique cutset.
- (iii) For $i \in \{0, 1, \dots, p-1\}$, G_i has a clique cutset split (A_i, K_i, B_i) and graphs $G_{i+1} = G[A_i \cup K_i]$ and $G_{i+1}^B = G[B_i \cup K_i]$ are blocks of decomposition of G_i with respect to this clique cutset split.

Also, we set $B_p = A_{p-1}$, $K_p = K_{p-1}$ and $G_{p+1}^B = G_p$. So, the leaves of T are the graphs $G_{i+1}^B = G[K_i \cup B_i]$, for $i \in \{0, 1, \dots, p\}$.

Theorem 2.2 ([24]) *A clique cutset decomposition tree of an input graph G can be computed in time $\mathcal{O}(nm)$ and has $\mathcal{O}(n)$ nodes.*

Note that for a non-leaf node G_i of T the corresponding clique cutset K_i is also a clique cutset of G . The following lemmas proved in [7] will also be needed.

Lemma 2.3 (Lemma 3.2 in [7]) *If G is a wheel-free graph that contains a diamond, then G has a clique cutset.*

A *star cutset* in a graph is a node cutset S that contains a node (called a *center*) adjacent to all other nodes of S . Note that a nonempty clique cutset is a star cutset.

Lemma 2.4 (Lemma 3.3 in [7]) *If $G \in \mathcal{C}$ has a star cutset, then G has a clique cutset.*

Decomposing with 2-joins

We first state some properties of 2-joins in graphs with no clique cutset. Let \mathcal{D} be the class of all graphs from \mathcal{C} that do not have a clique cutset. By Lemma 2.4 no graph from \mathcal{D} has a star cutset and by Lemma 2.3 no graph from \mathcal{D} contains a diamond.

An almost 2-join with a split $(X_1, X_2, A_1, A_2, B_1, B_2)$ in a graph G is *consistent* if the following statements hold for $i = 1, 2$:

- (i) Every component of $G[X_i]$ meets both A_i, B_i .
- (ii) Every node of A_i has a non-neighbor in B_i .

- (iii) Every node of B_i has a non-neighbor in A_i .
- (iv) Either both A_1, A_2 are cliques, or one of A_1 or A_2 is a single node, and the other one is a disjoint union of cliques.
- (v) Either both B_1, B_2 are cliques, or one of B_1, B_2 is a single node, and the other one is a disjoint union of cliques.
- (vi) $G[X_i]$ is connected.
- (vii) For every node v in X_i , there exists a path in $G[X_i]$ from v to some node of B_i with no internal node in A_i .
- (viii) For every node v in X_i , there exists a path in $G[X_i]$ from v to some node of A_i with no internal node in B_i .

Note that the definition contains redundant statements (for instance, (vi) implies (i)), but it is convenient to list properties separately as above.

Lemma 2.5 (Lemma 6.1 in [7]) *If $G \in \mathcal{D}$, then every almost 2-join of G is consistent.*

We now define the blocks of decomposition of a graph with respect to a 2-join. Let G be a graph and $(X_1, X_2, A_1, A_2, B_1, B_2)$ a split of a 2-join of G . Let k_1 and k_2 be positive integers. The *blocks of decomposition* of G with respect to (X_1, X_2) are the two graphs $G_1^{k_1}$ and $G_2^{k_2}$ that we describe now. We obtain $G_1^{k_1}$ from G by replacing X_2 by a *marker path* $P^2 = a_2 \dots b_2$ of length k_1 , where a_2 is a node complete to A_1 , b_2 is a node complete to B_1 , and $V(P_2) \setminus \{a_2, b_2\}$ is anticomplete to X_1 . The block $G_2^{k_2}$ is obtained similarly by replacing X_1 by a marker path $P^1 = a_1 \dots b_1$ of length k_2 .

In [20] the blocks of decomposition with respect to a 2-join that we used in construction of a recognition algorithm had marker paths of length 2. In this paper we will use blocks whose marker paths are of length 5. So, unless otherwise stated, when we say that G_1 and G_2 are blocks of decomposition with respect to a 2-join we will mean that their marker paths are of length 5. This will be discussed in more detail in Section 4.

Lemma 2.6 (Lemma 2.10 in [21]) *Let G be a graph from \mathcal{D} . Let (X_1, X_2) be a 2-join of G , and G_1, G_2 the blocks of decomposition with respect to this 2-join whose marker paths are of length at least 2. Then G_1 and G_2 are in \mathcal{D} and they do not have star cutsets.*

A 2-join (X_1, X_2) of G is a *minimally-sided 2-join* if for some $i \in \{1, 2\}$ the following holds: for every 2-join (X'_1, X'_2) of G , neither $X'_1 \subsetneq X_i$ nor $X'_2 \subsetneq X_i$. In this case X_i is a *minimal side* of this minimally-sided 2-join.

A 2-join (X_1, X_2) of G is an *extreme 2-join* if for some $i \in \{1, 2\}$ and all $k \geq 3$ the block of decomposition G_i^k has no 2-join. In this case X_i is an *extreme side* of such a 2-join.

Graphs in general do not necessarily have extreme 2-joins (an example is given in [25]), but it is shown in [25] that graphs with no star cutset do. It is also shown in [25] that if G has no star cutset then the blocks of decomposition with respect to a 2-join whose marker paths are of length at least 3, also have no star cutset. This is then used to show that in a graph with no star cutset, a minimally-sided 2-join is extreme. We summarize these results in the following lemma.

A *flat path* of G is any path of G of length at least 3, whose interior vertices are of degree 2 (this definition is slightly different than in [25], but this does not affect the proof of part (iii) of the following lemma). The following statement can be extracted from Lemmas 3.2, 4.2, 4.3 and 4.4 of [25].

Lemma 2.7 ([25]) *Let G be a graph with no star cutset. Let $(X_1, X_2, A_1, A_2, B_1, B_2)$ be a split of a minimally-sided 2-join of G with X_1 being a minimal side, and let G_1 and G_2 be the corresponding blocks of decomposition whose marker paths are of length at least 3. Then the following hold:*

- (i) $|A_1| \geq 2$, $|B_1| \geq 2$, and in particular all the vertices of $A_2 \cup B_2$ are of degree at least 3.
- (ii) If G_1 and G_2 do not have star cutsets, then (X_1, X_2) is an extreme 2-join, with X_1 being an extreme side (in particular, G_1 has no 2-join).
- (iii) If P is a flat path of G , such that $P \cap X_1 \neq \emptyset$ and $P \cap X_2 \neq \emptyset$, then one of the following holds:
 - (a) for an endnode u of P , $P \setminus u \subseteq X_1$ and $u \in A_2 \cup B_2$;
 - (b) for endnodes u and v of P , $u \in A_2$, $v \in B_2$, $P \setminus \{u, v\} \subseteq X_1$, the length of P is at least 3 and $G[X_1]$ has exactly two connected components that are both a path with one end in A_1 , one end in B_1 and interior in C_1 .

Remark 2.8 *We will be applying Lemma 2.7 only to graphs $G \in \mathcal{D}$, and so by Lemmas 2.4 and 2.6, all of G, G_1, G_2 have no star cutsets. Furthermore, for $G \in \mathcal{D}$, by Lemma 2.5, every 2-join is consistent, so outcome (b) of (iii) of Lemma 2.7 is not possible (i.e. (a) is the only possible outcome of Lemma 2.7(iii)).*

In [25] it is shown that one can decompose a graph with no star cutset using a sequence of ‘non-crossing’ 2-joins into graphs with no star cutset and no 2-join (which will in our case be basic). In this paper we will use minimally-sided 2-joins (as opposed to [25] and [21], where minimally-sided 2-joins were ‘moved’ to allow marker paths to be disjoint). This will be particularly important when solving the induced paths problem. We now describe this 2-join decomposition.

2-Join decomposition tree T_G of depth $p \geq 1$ of a graph G that has no star cutset and has a 2-join

- (i) The root of T_G is $G^0 = G$.
- (ii) The non-leaf nodes of T_G are G^0, G^1, \dots, G^{p-1} . Each non-leaf node G^i has two children: one is G^{i+1} and the other one is G_B^{i+1} .

The leaf-nodes of T_G are graphs $G_B^1, G_B^2, \dots, G_B^p$ and G^p . Graphs $G_B^1, G_B^2, \dots, G_B^p, G^p$ have no star cutset nor 2-join.

- (iii) For $i \in \{0, 1, \dots, p-1\}$, G^i has a 2-join (X_1^i, X_2^i) that is minimally-sided with minimal side X_1^i . We denote the split of this 2-join by $(X_1^i, X_2^i, A_1^i, A_2^i, B_1^i, B_2^i)$. Graphs G^{i+1} and G_B^{i+1} are blocks of decomposition of G^i with respect to (X_1^i, X_2^i) whose marker paths are of length 5. The block G_B^{i+1} corresponds to the minimal side X_1^i , i.e. $X_1^i \subseteq V(G_B^{i+1})$. We denote with P^{i+1} the marker path used to build G^{i+1} .

Lemma 2.9 *For $G \in \mathcal{D}$ the depth of any decomposition tree T_G is at most n .*

PROOF — To prove this lemma we can use almost the identical proof as for Lemma 8.1 from [25]; the only difference is that in the last paragraph of that proof, in order to conclude that $|A_1^i|, |B_1^i| \geq 2$ and that G^i has no star cutset now we use Lemmas 2.7 and 2.6. \square

Lemma 2.10 *There is an algorithm with the following specification.*

Input: *A graph $G \in \mathcal{D}$ that has a 2-join.*

Output: *A 2-join decomposition tree T_G of depth at most n , such that all graphs that correspond to the nodes of T_G belong to \mathcal{D} , and all graphs that correspond to the leaves of T_G (i.e. G_B^1, \dots, G_B^p, G^p) belong to \mathcal{B} .*

Running time: $\mathcal{O}(n^4m)$.

PROOF — Let $G^0 = G$. Suppose that a decomposition tree of depth $i \geq 0$ of G has been constructed. By Lemmas 2.4 and 2.6 graph G^i belongs to \mathcal{D} and has no star cutset. We apply on G^i the algorithm from [4] that finds a minimally-sided 2-join (the running time of this algorithm is $\mathcal{O}(n^3m)$). If no 2-join is found, then $i = p$, G^i is basic (by Theorem 2.1) and we stop the algorithm. If a 2-join (X_1^i, X_2^i) is found, then we build graphs G^{i+1} and G_B^{i+1} as in the definition of a 2-join decomposition tree. Note that, by Lemmas 2.4, 2.6 and 2.7 and Theorem 2.1, $G_B^{i+1} \in \mathcal{B}$ and G^{i+1} does not have a star cutset. Hence, (X_1^i, X_2^i) is an extreme 2-join of G^i (by Lemma 2.7).

Let T_G be the 2-join decomposition tree that is obtained using the algorithm we described. By Lemma 2.9 the depth of T_G is at most n and hence the running time of our algorithm is $\mathcal{O}(n^4m)$. \square

3 Induced cycles

In this section we consider an instance (G, \mathcal{V}) of the k -IN-A-CYCLE problem on \mathcal{C} , that is a graph $G \in \mathcal{C}$ and a set $\mathcal{V} = \{v_1, \dots, v_k\}$ of k distinct vertices of G . The problem is to decide whether there exists a chordless cycle that contains all vertices of \mathcal{V} .

2-in-a-Cycle

The 2-IN-A-CYCLE problem is to decide whether a graph G contains a chordless cycle through two specified vertices u and v of G . By Bienstock's construction, this problem is NP-complete for general graphs (see [2]). In this section we prove that for $G \in \mathcal{C}$ the only obstructions for the existence of such a cycle are clique cutsets that separate u and v . This leads to an algorithm of running time $\mathcal{O}(nm)$ for 2-IN-A-CYCLE for \mathcal{C} .

An additional property of P-graphs that we need in the proof of Theorem 3.2 below is the following: a P-graph is a graph $G \in \mathcal{B}$ with skeleton R and special clique K such that $K \neq \emptyset$ and each vertex of $L(R)$ that corresponds to an edge of R incident with a degree 1 vertex (in R), has a neighbor in K . We also need the following result from [20].

Lemma 3.1 (Lemma 3.4 in [20]) *Let R be a skeleton of a P-graph. If e_1 and e_2 are edges of R , then there exists a cycle of R that goes through e_1 and e_2 , or there exists a path in R whose endnodes are of degree 1 (in R) and that goes through e_1 and e_2 .*

Theorem 3.2 *Let G be a graph of \mathcal{C} , and u and v be non-adjacent vertices of G . Then there exists a hole of G that contains both u and v , or G admits a clique cutset that separates u and v .*

PROOF — Our proof is by induction on $|V(G)|$. By Theorem 2.1, it is enough to examine the following cases.

Case 1. G is a line graph of a triangle-free chordless graph.

Let R be the root graph of G , and let $e = x_1x_2$ and $f = y_1y_2$ be the edges of R that correspond to u and v (note that $\{x_1, x_2\} \cap \{y_1, y_2\} = \emptyset$). By Menger's theorem, in R there exist two vertex disjoint paths P and Q from $\{x_1, x_2\}$ to $\{y_1, y_2\}$, or there is a vertex z that separates $\{x_1, x_2\}$ from $\{y_1, y_2\}$. In the first case $P \cup Q$ is a hole of R (since R is triangle-free and chordless), and hence $L(P \cup Q)$ is a hole of G that contains u and v . In the second case the set $\{zz' : z' \in N_R(z)\} \setminus \{x_1x_2, y_1y_2\}$ correspond to a clique cutset of G that separates u and v .

Case 2. G is a P-graph.

Let R be the skeleton and K the special clique of G . First, let us consider the case when both u and v are in $L(R)$. We apply Lemma 3.1 to edges e_1 and e_2 that in R correspond to u and v . If a hole H is obtained, then $L(H)$ is a hole of G that contains u and v . If a

path P is obtained, then $L(P)$ together with neighbor(s) in K of its endnodes induces a hole in G that contains u and v .

So, we may assume that $u \in K$ and let $u' \in L(R)$ be a neighbor of u . Then $v \in L(R)$. Since u' is of degree 1 in $L(R)$, by Lemma 3.1, there exists a chordless path Q in $L(R)$ that contains both u' and v , and such that its other endnode w' ($w' \neq u'$) is of degree 1 in $L(R)$. Let w be the neighbor of w' in K . Then $V(Q) \cup \{u, w\}$ induces a desired hole in G .

Case 3. G admits a clique cutset.

Let (A, K, B) be a split of this cutset. If u and v are separated by this cutset, we are done, so we may assume that $u, v \in V(G_A)$. Then, by induction, there exists a hole H in G_A that contains both u and v , or a clique cutset K' of G_A that separates u and v . In the first case H is a hole of G that contains both u and v , and in the second K' is a clique cutset of G that separates u and v .

Case 4. G admits a 2-join.

Let $(X_1, X_2, A_1, A_2, B_1, B_2)$ be a 2-join of G . By Case 3, we may assume that G does not admit a clique cutset, and hence, by Lemma 2.5, that (X_1, X_2) is a consistent 2-join. So, by property (vi) of consistent 2-joins, for $i \in \{1, 2\}$ there is a chordless path Q^i in $G[X_i]$ that has exactly one vertex from both A_i and B_i . Also, by Lemma 2.6, blocks of decomposition G_1 and G_2 belong to \mathcal{D} .

First, let us assume that $u, v \in X_i$, for some $i \in \{1, 2\}$. Then, by induction, there is a hole H in G_i that contains u and v . If H is contained in $G[X_i]$, then we are done. Otherwise H contains the marker path P^{3-i} , and hence to obtain a desired hole it is enough to replace P^{3-i} with Q^{3-i} in H .

So, we may assume that $u \in X_1$ and $v \in X_2$. By induction there exists a hole H_1 (resp. H_2) in G_1 (resp. G_2) that contains u and c_2 (resp. v and c_1), where c_i is an internal vertex of P^i , for $i \in \{1, 2\}$. Let R^1 (resp. R^2) be the path obtained from H_1 (resp. H_2) by removing the vertices of the marker path P^2 (resp. P^1). Then $V(R^1) \cup V(R^2)$ induces a hole in G that contains both u and v . \square

Theorem 3.3 *There is an algorithm with the following specifications:*

Input: A graph $G \in \mathcal{C}$ and two vertices u and v of G .

Output: YES if there is a chordless cycle of G that contains both u and v , and NO otherwise.

Running time: $\mathcal{O}(nm)$.

PROOF — First, let us consider the case when uv is an edge of G . Let G' be the graph obtained from G by deleting uv . Now, if there exists a path from u to v in G' , then return YES, and otherwise return NO.

So, we may assume that u and v are not adjacent. We build a clique cutset decomposition tree T for G and remember (all) leaf nodes of this tree that contain some of the vertices u and v . By Theorem 2.2, this can be done in time $\mathcal{O}(nm)$. If there is a leaf node of T that contains both u and v , then there does not exist a clique cutset that separates u and v , and hence, by Theorem 3.2, we return YES. Otherwise, we return NO. The running time of this algorithm is $\mathcal{O}(nm)$. \square

k -in-a-Cycle

In this section we prove that the problem k -IN-A-CYCLE is fixed-parameter tractable, when parameterized by k , for graphs in \mathcal{C} .

We use the following result of Robertson and Seymour.

Theorem 3.4 ([22]) *The k -DISJOINT PATHS problem is fixed-parameter tractable, when parameterized by k . More precisely, there is a computable function h , that depends only on k , such that the k -DISJOINT PATHS problem can be solved in time $h(k)n^3$.*

Lemma 3.5 *For any fixed integer k , there is an algorithm with the following specifications:*

Input: *A graph $G \in \mathcal{B}$ and a set $\mathcal{V} = \{v_1, \dots, v_k\}$ of k distinct vertices from G .*

Output: *YES if the problem (G, \mathcal{V}) has a solution, and NO otherwise.*

Running time: $\mathcal{O}(n^5)$.

PROOF — Clearly, we may assume that G is connected. Recall that G is (wheel, diamond)-free. In time $\mathcal{O}(n^2m)$ we can find the set C of all centers of claws in G . If $|C| \geq 2$, then let K be the maximal (with respect to inclusion) clique that contains C (this can be done in $\mathcal{O}(n)$ -time since G is diamond-free). Otherwise, let $K = C$. So $G \setminus K$ is (wheel, diamond, claw)-free. By Lemma 2.4 in [7] it follows that $G \setminus K$ is a line graph of a triangle-free chordless graph. In $\mathcal{O}(n + m)$ -time we can compute graph R such that $G \setminus K = L(R)$ (see [17, 23]). It follows that K is a special clique and R a skeleton of G .

To solve the given problem it is enough to solve $k!$ problems (G, \mathcal{W}) , where $\mathcal{W} = \{(v_{\sigma(1)}, v_{\sigma(2)}), (v_{\sigma(2)}, v_{\sigma(3)}), \dots, (v_{\sigma(k-1)}, v_{\sigma(k)}), (v_{\sigma(k)}, v_{\sigma(1)})\}$ and σ is a permutation of $\{1, 2, \dots, k\}$ (note that these are not INDUCED DISJOINT PATH problems by our definition, but they are by the definition used in [13], where it is allowed that the paths in the solution share endnodes; see also Subsection 4.4). Hence, it is enough to show that each of them can be solved in $\mathcal{O}(n^5)$ time.

Case 1. $K = \emptyset$.

Then $G = L(R)$, so, as noted in [13, Lemma 3.7], to solve each of the given problems it is enough to solve 2^{2k} DISJOINT PATHS problems on R . These problems can be solved using the algorithm from Theorem 3.4. Since edges of R are vertices of G , graph R

has $\mathcal{O}(n)$ vertices (in the proof of [13, Lemma 3.7] a worse bound $\mathcal{O}(n^2)$ is used, which leads to a worse running time in that lemma), so the total running time in this case is $\mathcal{O}(n^3 + n^2m) = \mathcal{O}(n^4)$.

Case 2. $K \neq \emptyset$.

Let S be the set of all vertices of $L(R)$ that have a neighbor in K . For each pair (u, v) of vertices from S we build a graph $G_{u,v}$ in the following way: we start with $L(R)$, add to it the unique chordless (u, v) -path P_{uv} (of length 2 or 3) whose interior vertices are from K , and remove all vertices from $L(R) \setminus \{u, v\}$ that have a neighbor in the interior of P_{uv} . It is easy to see that (G, \mathcal{W}) has a solution if and only if $(L(R), \mathcal{W})$ or $(G_{u,v}, \mathcal{W})$, for some $u, v \in S$, has a solution. Each of the graphs $L(R)$ and $G_{u,v}$, for $u, v \in S$, is (wheel, diamond, claw)-free, and hence, by Lemma 2.4 in [7], it is the line graph of a triangle-free chordless graph. So, each of the problems $(L(R), \mathcal{W})$ and $(G_{u,v}, \mathcal{W})$, for $u, v \in S$, can be solved as in Case 1, and since $|S| = \mathcal{O}(n)$, this implies an $\mathcal{O}(n^2m + n^2 \cdot n^3) = \mathcal{O}(n^5)$ -time algorithm for solving the given problem. \square

Theorem 3.6 *For any fixed integer k , there is an algorithm with the following specifications:*

Input: A graph $G \in \mathcal{C}$ and a set $\mathcal{V} = \{v_1, \dots, v_k\}$ of k distinct vertices from G .

Output: YES if the problem (G, \mathcal{V}) has a solution, and NO otherwise.

Running time: $\mathcal{O}(n^6)$.

PROOF — In [20] an $\mathcal{O}(n^2m)$ -time algorithm is given for recognizing whether a graph belongs to \mathcal{B} . If $G \in \mathcal{B}$, then the problem (G, \mathcal{V}) can be solved in time $\mathcal{O}(n^5)$ using Lemma 3.5. So we may assume that $G \in \mathcal{C} \setminus \mathcal{B}$.

Next, we consider the case when $G \in \mathcal{D} \setminus \mathcal{B}$. Using Lemma 2.10 we build a 2-join decomposition tree T_G in time $\mathcal{O}(n^4m)$ (throughout the proof we use the notation from the definition of T_G). Let c_i (resp. c_i^B) be an internal vertex of the marker path of G^i (resp. G_B^i), for $1 \leq i \leq p$. By Theorem 2.1, graphs G_B^1, \dots, G_B^p, G^p are in \mathcal{B} .

Let $\mathcal{V}^0 = \mathcal{V}$. We now describe the problems (G^i, \mathcal{V}^i) and (G_B^i, \mathcal{V}_B^i) , $1 \leq i \leq p$, that we solve during our algorithm to obtain the solution of the problem (G, \mathcal{V}) .

We first introduce some notation. Let C^i be any chordless cycle of G^i , and Q_B^i (resp. Q^i) be the part of C^i contained in X_1^i (resp. X_2^i). Note that one of the following holds: (1) Q_B^i and Q^i are subpaths of C^i of length at least 1; or (2) Q^i is not a path of length at least 1; or (3) Q_B^i is not a path of length at least 1. Then we define C_B^{i+1} (resp. C^{i+1}) as the chordless cycle obtained from C^i in the following way: (1) by replacing Q^i (resp. Q_B^i) with the marker path of G_B^{i+1} (resp. G^{i+1}); (2) by replacing the vertices (if they exist) of $C^i \cap A_2^i$ or $C^i \cap B_2^i$ with endnodes a_2^i and b_2^i of the marker path of G_B^{i+1} (resp. C^{i+1} is empty); (3) C_B^{i+1} is empty (resp. C^{i+1} is equal to C^i). (Note that A_2^i and B_2^i are cliques, so in case (3) C^i is contained in X_2^i .)

Similarly, for a (non-empty) chordless cycle C_B^{i+1} (resp. C^{i+1}) of G_B^{i+1} (resp. G^{i+1}) that contains the marker path of G_B^{i+1} (resp. G^{i+1}) we define a chordless cycle C^i of G^i as follows. Let Q^i (resp. Q_B^i) be any chordless path from a vertex of A_1^i (resp. A_2^i) to a vertex of B_1^i (resp. B_2^i) in X_1^i (resp. X_2^i) (this path exists by Lemma 2.5). Then C^i is obtained from C_B^{i+1} (resp. C^{i+1}) by replacing the marker path of G_B^{i+1} (resp. G^{i+1}) by Q^i (resp. Q_B^i).

Now, we describe how our algorithm solves the problem (G^i, \mathcal{V}^i) . It is enough to consider the following 3 cases.

Case 1. $\mathcal{V}^i \subseteq X_1^i$.

We note that C^i is a solution of (G^i, \mathcal{V}^i) if and only if C_B^{i+1} is a solution of $(G_B^{i+1}, \mathcal{V}^i)$. So, in this case it is enough to decide whether $(G_B^{i+1}, \mathcal{V}^i)$ has a solution, which can be done using Lemma 3.5 in time $\mathcal{O}(n^5)$.

Case 2. $\mathcal{V}^i \subseteq X_2^i$.

We note that C^i is a solution of (G^i, \mathcal{V}^i) if and only if C^{i+1} is a solution of (G^{i+1}, \mathcal{V}^i) . So, we proceed recursively, by solving the problem (G^{i+1}, \mathcal{V}^i) .

Case 3. $\mathcal{V}^i \cap X_1^i \neq \emptyset$ and $\mathcal{V}^i \cap X_2^i \neq \emptyset$.

Let $\mathcal{V}_B^{i+1} = (\mathcal{V}^i \cap X_1^i) \cup \{c_B^{i+1}\}$ and $\mathcal{V}^{i+1} = (\mathcal{V}^i \cap X_2^i) \cup \{c^{i+1}\}$. We note that C^i is a solution of (G^i, \mathcal{V}^i) if and only if C^{i+1} is a solution of $(G^{i+1}, \mathcal{V}^{i+1})$ and C_B^{i+1} is a solution of $(G_B^{i+1}, \mathcal{V}_B^{i+1})$ (indeed, in this case the cycle C^{i+1} (resp. C_B^{i+1}) constructed from C^i contains the marker path of G^{i+1} (resp. G_B^{i+1})). So, we solve the problem $(G^{i+1}, \mathcal{V}^{i+1})$ using the algorithm from Lemma 3.5. If this algorithm returns NO, then we return NO and stop. Otherwise, we proceed recursively, by solving the problem $(G^{i+1}, \mathcal{V}^{i+1})$ (note that $|\mathcal{V}^{i+1}| \leq |\mathcal{V}^i|$).

The running time of the described algorithm is $\mathcal{O}(n^4m + n \cdot n^5) = \mathcal{O}(n^6)$, since there are at most $p + 1 = \mathcal{O}(n)$ calls to the algorithm from Lemma 3.5.

Finally, let us consider the general case, that is $G \in \mathcal{C}$. First, using Theorem 2.2 we build a clique cutset decomposition tree T of G in time $\mathcal{O}(nm)$ (in what follows we use the notation from the definition of T). Note that for each $1 \leq i \leq p$ a chordless cycle can not contain vertices from both A_i and B_i . So, for each $1 \leq i \leq p$ we check if both $A_i \cap \mathcal{V}$ and $B_i \cap \mathcal{V}$ are non-empty, and if this is the case we return NO. This can be done in time $\mathcal{O}(n)$. Hence, we may assume that \mathcal{V} is contained in G_j^B , for some $j \in \{1, 2, \dots, p + 1\}$. Then a desired cycle, if it exists, is also contained in G_j^B , so it is enough to solve the problem (G_j^B, \mathcal{V}) . Since $G_j^B \in \mathcal{D}$, this can be done in time $\mathcal{O}(n^6)$ using the previous part of the proof. Hence the running time of the algorithm is $\mathcal{O}(n + n^6) = \mathcal{O}(n^6)$. \square

Corollary 3.7 *For graphs in \mathcal{C} the k -IN-A-CYCLE problem is fixed-parameter tractable, when parameterized by k .*

PROOF — Let (G, \mathcal{V}) be an instance of the k -IN-A-CYCLE problem. By Theorem 3.4, the problem k -DISJOINT PATHS can be solved in time $h(k)n^3$, where h is a computable function

that depends only on k . Since Lemma 3.5 has at most $2^{2k}k!n^2$ calls to this algorithm, the problem (G, \mathcal{V}) can be solved in time $2^{2k}k!h(k)n^5$ for graphs in \mathcal{B} . Now, since for each $i \in \{0, 1, \dots, p-1\}$ the algorithm from Theorem 3.6 has at most one call to the algorithm from Lemma 3.5, and $p \leq n$ (by Lemma 2.10), we conclude that the problem (G, \mathcal{V}) can be solved in time $2^{2k}k!h(k)n^6$ for graphs in \mathcal{C} . \square

4 Induced disjoint paths

In this section, we consider an instance (G, \mathcal{W}) of the k -INDUCED DISJOINT PATHS problem, that is a graph G , a set $\mathcal{W} = \{(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)\}$ of pairs of vertices of G such that all $2k$ vertices are distinct and the only edges between these vertices are of the form $s_i t_i$, for some $1 \leq i \leq k$. We denote $W = \{s_1, \dots, s_k, t_1, \dots, t_k\}$. Vertices in W are the *terminals* of \mathcal{W} .

Recall that k is a fixed integer (that is not part of the input). We have to decide whether there exist k vertex-disjoint paths $P_i = s_i \dots t_i$, $1 \leq i \leq k$, such that there are no edges between vertices of paths P_i and P_j , for $i \neq j$.

Note that (G, \mathcal{W}) has a solution (say $P_i = s_i \dots t_i$, $1 \leq i \leq k$) if and only if (G, \mathcal{W}) has a solution so that all the paths in this solution are chordless (it is enough to take $P'_i = s_i \dots t_i$, $1 \leq i \leq k$, where P'_i is a chordless path of G contained in P_i). So, when solving (G, \mathcal{W}) we may add the condition that the paths $P_i = s_i \dots t_i$, $1 \leq i \leq k$, are chordless. Throughout the paper when we say that a set of paths \mathcal{P} is a solution of (G, \mathcal{W}) , we will assume that all paths in \mathcal{P} are chordless.

In Subsection 4.1, we study how clique cutset can be used. In Subsection 4.2, we study basic graphs. In Subsection 4.3, we study how 2-joins can be used, and we give the main algorithm. In Subsection 4.4, we study related problems.

4.1 Clique cutsets and induced paths

Let \mathcal{G} be a class of graphs that is closed under taking induced subgraphs, and let $\mathcal{G}_{\text{BASIC}} \subseteq \mathcal{G}$. Suppose that \mathcal{G} and $\mathcal{G}_{\text{BASIC}}$ satisfy the following:

If $G \in \mathcal{G}$, then $G \in \mathcal{G}_{\text{BASIC}}$ or G has a clique cutset.

Then we say that \mathcal{G} is $\mathcal{G}_{\text{BASIC}}$ -decomposable using clique cutsets.

Let \mathcal{G} be a class that is $\mathcal{G}_{\text{BASIC}}$ -decomposable using clique cutsets. In this section we show how an $\mathcal{O}(n^c)$ -time algorithm for k -INDUCED DISJOINT PATHS problem for graphs in $\mathcal{G}_{\text{BASIC}}$, where c is a constant (that does not depend on k), can be turned into an $\mathcal{O}(n^{2k+c})$ -time algorithm for k -INDUCED DISJOINT PATHS problem for graphs in \mathcal{G} .

Throughout the rest of the section, we consider an instance (G, \mathcal{W}) as described above for a graph $G \in \mathcal{G}$. Graphs from $\mathcal{G}_{\text{BASIC}}$ will be referred to as *basic graphs*.

Let T be a clique cutset decomposition tree of G of depth p (with all notations as in Section 2). The next lemma tells how a set obtained during the decomposition process behaves at the root level (that is in G).

Lemma 4.1 *Suppose that G is not basic (so $p \geq 1$). Let $i \in \{0, 1, \dots, p-1\}$ and $X \in \{A_i \cup K_i, K_i \cup B_i, K_p \cup B_p\}$. If C is a connected component of $G \setminus X$, then $N_G(C) = N_G(C) \cap X$ is a clique.*

PROOF — We prove the result by induction on p . If $p = 1$, then $i = 0$, (A_0, K_0, B_0) is a split of a clique cutset of G and either $X = A_0 \cup K_0 = K_1 \cup B_1$ or $X = K_0 \cup B_0$. In both cases, $N_G(C) \cap X \subseteq K_0$, so the conclusion holds.

Suppose $p > 1$. If $i = 0$, the conclusion holds as above. So, suppose $i \geq 1$. Observe that then $X \subseteq V(G_1) = A_0 \cup K_0$. Also, the tree obtained from T by deleting G_0 and G_1^B is a decomposition tree for G_1 with depth $p - 1$, and with exactly the same cutsets and splits as in T , so we may apply the induction hypothesis to it. Therefore, we know that for every connected component D of $G_1 \setminus X$, $N_{G_1}(D)$ is a clique.

Let C be a connected component of $G \setminus X$. Recall that (A_0, K_0, B_0) is a split of a clique cutset of G and $X \subseteq A_0 \cup K_0$. If $C \subseteq A_0$, then C is connected component of $G_1 \setminus X$, so $N_G(C) = N_{G_1}(C)$ is a clique by the induction hypothesis. Hence, we may assume that C contains at least one vertex in $K_0 \cup B_0$. In fact, since $G[K_0 \cup B_0]$ has no clique cutset (by the definition of a decomposition tree), it follows that B_0 is connected and every vertex of K_0 has a neighbor in B_0 , and hence $(B_0 \cup K_0) \setminus X \subseteq C$.

If $C = B_0$ (meaning in fact that $K_0 \subseteq X$), then $N_G(C) \subseteq K_0$. So suppose $B_0 \subsetneq C$. Hence, $C \cap K_0 \neq \emptyset$. Note that $C \setminus B_0$ is connected, because K_0 is a clique and every vertex in $C \setminus B_0$ can be linked by a path to some vertex in K_0 . Hence, $C \setminus B_0$ is a connected component of $G_1 \setminus X$, so $N_{G_1}(C \setminus B_0)$ is a clique K , and $K_0 \cap X \subseteq K$. Since $N_G(B_0) \subseteq K_0$, we have $N_G(C) \subseteq K$. \square

Lemma 4.2 *For every $i = 0, \dots, p$, $G[K_i \cup B_i]$ is basic.*

PROOF — By the definition of decomposition trees, $G[K_i \cup B_i]$ has no clique cutset. From the definition of $\mathcal{G}_{\text{BASIC}}$ -decomposable classes, $G[K_i \cup B_i]$ is therefore basic. \square

Let us assume that a *non-negative integer weight* function $w : V \rightarrow \mathbb{Z}_{\geq 0}$ is assigned to vertices of G (here $\mathbb{Z}_{\geq 0}$ is the set of non-negative integers). Then for $A \subseteq V$ we define $w(A) = \sum_{v \in A} w(v)$.

Lemma 4.3 *Suppose that G is not basic (so $p \geq 1$) and that a non-negative integer weight $w(v)$ is given to each vertex $v \in V(G)$. Then one of the following holds:*

- (i) *For some $i \in \{0, 1, \dots, p-1\}$, K_i is a clique cutset of G for which there exists a split (A, K_i, B) such that $w(A) \geq 2$ and $w(B) \geq 2$.*

(ii) For some $i \in \{0, 1, \dots, p\}$, every connected component C of $G \setminus (K_i \cup B_i)$, satisfies $w(C) \leq 1$.

PROOF — Suppose first that $w(B_0) \geq 2$. Then, we may assume that $w(A_0) \leq 1$ for otherwise, (i) holds with $K_i = K_0$ and the split (A_0, K_0, B_0) . Hence, (ii) holds for $i = 0$. So, we may assume that $w(B_0) \leq 1$.

It follows that every connected component C of $G \setminus (A_0 \cup K_0)$ satisfies $w(C) \leq 1$. Hence, it is well defined to consider the maximal index $\ell \in \{0, \dots, p-1\}$ such that every connected component C of $G \setminus (A_\ell \cup K_\ell)$ satisfies $w(C) \leq 1$. If $\ell = p-1$, then since $B_p = A_{p-1}$ and $K_p = K_{p-1}$, (ii) holds for $i = p$, so we may assume $\ell < p-1$. Therefore, $(A_{\ell+1}, K_{\ell+1}, B_{\ell+1})$ is a split of a clique cutset of $G_{\ell+1} = G[A_\ell \cup K_\ell]$. Since, by Lemma 4.1, for every connected component C of $G \setminus (A_\ell \cup K_\ell)$, $N_G(C)$ is a clique, there are three types of such components C :

- *type A*: connected components that have neighbors in $A_{\ell+1}$ but no neighbor in $B_{\ell+1}$ (possibly in $K_{\ell+1}$);
- *type B*: connected components that have neighbors in $B_{\ell+1}$ but no neighbor in $A_{\ell+1}$ (possibly in $K_{\ell+1}$);
- *type K*: connected component whose neighborhood (possibly empty) is included in $K_{\ell+1}$.

We denote by A (resp. B , K) the union of all connected components of type A (resp. B, K).

The connected components of $G \setminus (K_{\ell+1} \cup B_{\ell+1})$ are the connected components of type B or K (that all have weight at most 1) and the connected components of $G[A \cup A_{\ell+1}]$. Therefore, unless $K_{\ell+1} \cup B_{\ell+1}$ satisfies (ii), we may assume that $w(A \cup A_{\ell+1}) \geq 2$.

The connected components of $G \setminus (A_{\ell+1} \cup K_{\ell+1})$ are the connected components of type A or K (that all have weight at most 1) and the connected components of $G[B \cup B_{\ell+1}]$. Therefore, by the maximality of ℓ , we know that $w(B \cup B_{\ell+1}) \geq 2$.

Now, we observe that $(A_{\ell+1} \cup A \cup K, K_{\ell+1}, B_{\ell+1} \cup B)$ is the split of a clique cutset of G that satisfies (i). \square

Theorem 4.4 *Let \mathcal{G} be a class that is $\mathcal{G}_{\text{BASIC}}$ -decomposable using clique cutsets. Furthermore, let us assume that there is an $\mathcal{O}(n^c)$ -time algorithm for k -INDUCED DISJOINT PATHS problem for graphs in $\mathcal{G}_{\text{BASIC}}$, where $c \geq 1$ is a constant (that does not depend on k). Then there is an algorithm that solves in $\mathcal{O}(n^{2k+c})$ time the k -INDUCED DISJOINT PATHS problem for every instance (G, \mathcal{W}) such that $G \in \mathcal{G}$.*

PROOF — Our proof is by induction on k . If $k = 1$, then we can solve our problem in time $\mathcal{O}(n + m) = \mathcal{O}(n^2)$. So, we may assume that $k \geq 2$.

We build a clique cutset decomposition tree T for G . Let us assume that it is of depth p . By Theorem 2.2, this can be done in time $\mathcal{O}(nm) = \mathcal{O}(n^3)$. Clearly, we may assume that $p \geq 1$. We give weight 1 to every vertex in W , weight 0 to every vertex in $V(G) \setminus W$, and then we apply Lemma 4.3. This leads to two cases.

Case 1: For some $\ell \in \{0, 1, \dots, p-1\}$, K_ℓ is a clique cutset of G for which there exists a split (A, K_ℓ, B) such that $|A \cap W| \geq 2$ and $|B \cap W| \geq 2$.

Note that this situation can be detected in time $\mathcal{O}(n^3)$ by computing connected components of $G \setminus K_\ell$, for each $\ell \in \{0, 1, \dots, p-1\}$ (by Theorem 2.2, $p = \mathcal{O}(n)$).

We set $K = K_\ell$. Note that no two paths from a solution of (G, \mathcal{W}) can have a vertex in K , so if for some $i \neq j$ both $\{s_i, t_i\}$ and $\{s_j, t_j\}$ have a non-empty intersection with both $A \cup K$ and $B \cup K$, then we return NO and stop the algorithm. Hence, we may assume that at most one pair from \mathcal{W} has non-empty intersection with both $A \cup K$ and $B \cup K$.

Let $\mathcal{W}_A = \{(s_i, t_i) \mid s_i, t_i \in A\}$ and $\mathcal{W}_B = \{(s_i, t_i) \mid s_i, t_i \in B\}$. By the previous remark and the assumption in this case, both \mathcal{W}_A and \mathcal{W}_B are non-empty and $|\mathcal{W} \setminus (\mathcal{W}_A \cup \mathcal{W}_B)| \leq 1$.

We observe that for every solution \mathcal{P} of (G, \mathcal{W}) at most two vertices of K are contained in paths of \mathcal{P} , and if two vertices of K are members of these paths, then they are in the same path of \mathcal{P} .

We now build several pairs of instances of the INDUCED DISJOINT PATHS problem that we use to solve (G, \mathcal{W}) . The construction depends on $|\mathcal{W} \setminus (\mathcal{W}_A \cup \mathcal{W}_B)|$ and the position of vertices from the pair from $\mathcal{W} \setminus (\mathcal{W}_A \cup \mathcal{W}_B)$ (if $|\mathcal{W} \setminus (\mathcal{W}_A \cup \mathcal{W}_B)| = 1$). By symmetry, it is enough to examine the following cases.

Case 1.1: If $|\mathcal{W} \setminus (\mathcal{W}_A \cup \mathcal{W}_B)| = 0$, then these instances are:

- $(G[A], \mathcal{W}_A)$ and $(G[B], \mathcal{W}_B)$;
- for every $x \in K$, $(G[A \cup \{x\}], \mathcal{W}_A)$ and $(G[B \setminus N(x)], \mathcal{W}_B)$;
- for every $x \in K$, $(G[A \setminus N(x)], \mathcal{W}_A)$ and $(G[B \cup \{x\}], \mathcal{W}_B)$;
- for every distinct $x, y \in K$, $(G[A \cup \{x, y\}], \mathcal{W}_A)$ and $(G[B \setminus (N(x) \cup N(y))], \mathcal{W}_B)$;
- for every distinct $x, y \in K$, $(G[A \setminus (N(x) \cup N(y))], \mathcal{W}_A)$ and $(G[B \cup \{x, y\}], \mathcal{W}_B)$.

Case 1.2: If $|\mathcal{W} \setminus (\mathcal{W}_A \cup \mathcal{W}_B)| = 1$, then without loss of generality $\{(s_1, t_1)\} = \mathcal{W} \setminus (\mathcal{W}_A \cup \mathcal{W}_B)$.

In case $s_1 \in A$ and $t_1 \in B$, we build the following pairs of instances:

- for every $x \in K$, $(G[A \cup \{x\}], \mathcal{W}_A \cup \{(s_1, x)\})$ and $(G[B \cup \{x\}], \mathcal{W}_B \cup \{(x, t_1)\})$;
- for every distinct $x, y \in K$, $(G[(A \setminus N(y)) \cup \{x\}], \mathcal{W}_A \cup \{(s_1, x)\})$ and $(G[(B \setminus N(x)) \cup \{y\}], \mathcal{W}_B \cup \{(y, t_1)\})$.

In case $s_1 \in A$ and $t_1 \in K$, we build the following pairs of instances:

- $(G[A \cup \{t_1\}], \mathcal{W}_A \cup \{(s_1, t_1)\})$ and $(G[B \setminus N(t_1)], \mathcal{W}_B)$;
- for every $x \in K$, $(G[A \cup \{x\}], \mathcal{W}_A \cup \{(s_1, x)\})$ and $(G[B \setminus (N(x) \cup N(t_1))], \mathcal{W}_B)$.

In case $s_1, t_1 \in K$, we build the following pair of instances:

- $(G[A \setminus (N(s_1) \cup N(t_1))], \mathcal{W}_A)$ and $(G[B \setminus (N(s_1) \cup N(t_1))], \mathcal{W}_B)$.

In each case it is straightforward to check that (G, \mathcal{W}) has a solution if and only if for some pair of instances that we built both of them have a solution. So, we run recursively our algorithm for all these instances.

Case 2: For some $\ell \in \{0, 1, \dots, p\}$, every connected component C of $G \setminus (K_\ell \cup B_\ell)$ satisfies $|C \cap W| \leq 1$.

Note that this case can be detected in time $\mathcal{O}(n^3)$ by computing connected components of $G \setminus (K_\ell \cup B_\ell)$, for $l \in \{0, 1, \dots, p\}$ (by Theorem 2.2, $p = \mathcal{O}(n)$).

We first delete all connected components C of $G \setminus (K_\ell \cup B_\ell)$ that satisfies $|C \cap W| = 0$. This is correct because a path in a solution for (G, \mathcal{W}) cannot contain a vertex from such a connected component (by Lemma 4.1).

Now, for each connected component C of $G \setminus (K_\ell \cup B_\ell)$ (that is left after previous deletions) there exists a unique s_i or t_i in C and we set: $S_i = N(C)$ if $s_i \in C$, and $T_i = N(C)$ if $t_i \in C$. Let $\mathcal{I}_S \subseteq \{1, 2, \dots, k\}$ (resp. $\mathcal{I}_T \subseteq \{1, 2, \dots, k\}$) be the set of all i for which S_i (resp. T_i) is defined.

Choose $s'_i \in S_i$, for $i \in \mathcal{I}_S$, and $t'_i \in T_i$, for $i \in \mathcal{I}_T$, and let $s'_i = s_i$ for $i \in \{1, 2, \dots, k\} \setminus \mathcal{I}_S$, and $t'_i = t_i$ for $i \in \{1, 2, \dots, k\} \setminus \mathcal{I}_T$ (note that $i \in \{1, 2, \dots, k\} \setminus \mathcal{I}_S$ implies $s_i \in K_\ell \cup B_\ell$). Furthermore, let $\mathcal{W}' = \{(s'_i, t'_i) \mid i \in \{1, 2, \dots, k\}\}$. It is now straightforward to check that (G, \mathcal{W}) has a solution if and only if $(G[K_\ell \cup B_\ell], \mathcal{W}')$ has a solution for at least one \mathcal{W}' constructed in this way. Note that the graph $G[K_\ell \cup B_\ell]$ is basic.

Complexity analysis

Suppose that for some constant $q \geq 1$, the algorithm for the basic class runs in time at most qn^c . We claim that our algorithm then runs in time at most $T(n, k) \leq q3^k n^{2k+c} = \mathcal{O}(n^{2k+c})$ (because k is not part of the instance).

If we are in Case 2, this is direct, since we call at most n^{2k} times the algorithm for the basic class.

If we are in Case 1, we first observe that we run recursively the algorithm at most n^2 times on each side. Let $k_A = |\mathcal{W}_A|$ and $k_B = |\mathcal{W}_B|$. Then $1 \leq k_A, k_B \leq k - 1$.

Now, in Case 1.1 the running time $T(n, k)$ satisfies:

$$\begin{aligned}
T(n, k) &\leq n^2 T(n, k_A) + n^2 T(n, k_B) + n^3 \\
&\leq qn^2 3^{\max\{k_A, k_B\}} [n^{2k_A+c} + n^{2k_B+c} + n] \\
&\leq qn^2 3^{k-1} [n^{2k+c-2} + n^{2k+c-2} + n] \\
&\leq q3^k n^{2k+c}.
\end{aligned}$$

In Case 1.2 we have $k_A, k_B \leq k - 2$ and the running time $T(n, k)$ satisfies:

$$\begin{aligned}
T(n, k) &\leq n^2 T(n, k_A + 1) + n^2 T(n, k_B + 1) + n^3 \\
&\leq qn^2 3^{\max\{k_A+1, k_B+1\}} [n^{2k_A+2+c} + n^{2k_B+2+c} + n] \\
&\leq qn^2 3^{k-1} [n^{2k+c-2} + n^{2k+c-2} + n] \\
&\leq q3^k n^{2k+c},
\end{aligned}$$

which concludes our proof. \square

4.2 Basic graphs

In this section we provide a polynomial-time algorithm that solves the problem (G, \mathcal{W}) for graphs $G \in \mathcal{B}$.

Lemma 4.5 *For a fixed integer k , there is an algorithm with the following specifications:*

Input: *A graph $G \in \mathcal{B}$ and a set of pairs of vertices $\mathcal{W} = \{(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)\}$ from G , such that all $2k$ vertices are distinct and the only possible edges between these vertices are of the form $s_i t_i$, for some $1 \leq i \leq k$.*

Output: *YES if the problem (G, \mathcal{W}) has a solution, and NO otherwise.*

Running time: $\mathcal{O}(n^5)$.

PROOF — We use almost the same algorithm as in Lemma 3.5 – the only difference is that we do not construct $k!$ INDUCED DISJOINT PATH problems, but work directly with the problem (G, \mathcal{W}) . \square

With a more complicated algorithm the running time in the above lemma can be reduced to $\mathcal{O}(n^2 m + n^3)$, but since this will not help improve the overall complexity of the algorithm in this section we do not include it here.

4.3 2-joins and induced paths

In this section we give a polynomial-time algorithm that solves the problem (G, \mathcal{W}) for graphs $G \in \mathcal{D}$. In fact, we solve a similar problem on certain structures, called *o-graphs*, which allow us to use 2-joins in a more convenient way.

Definition 4.6 *An o-graph $G_{\mathcal{F}, \mathcal{O}}$ is a triple $(G, \mathcal{F}, \mathcal{O})$, where G is a graph, \mathcal{F} is a set of some flat paths of G of length at most 7 and some vertices of G (viewed as paths of length 0), and \mathcal{O} is a set such that for each $\mathcal{W} \in \mathcal{O}$, (G, \mathcal{W}) is an instance of the INDUCED DISJOINT PATHS problem where every terminal vertex is contained in a path $P \in \mathcal{F}$.*

We say that an o-graph $G_{\mathcal{F},\mathcal{O}}$ is *linkable* if for at least one $\mathcal{W} \in \mathcal{O}$, the problem (G, \mathcal{W}) has a solution.

Note that an instance (G, \mathcal{W}) of the INDUCED DISJOINT PATHS problem has a solution if and only if the o-graph $(G, \mathcal{F}, \mathcal{O})$ is linkable, where \mathcal{F} is the set of terminals of \mathcal{W} and $\mathcal{O} = \{\mathcal{W}\}$. So, to solve (G, \mathcal{W}) , it is enough to decide whether $G_{\mathcal{F},\mathcal{O}}$ is linkable.

We do this with 2-join decompositions that are performed to reduce the problem to basic graphs. If $G \in \mathcal{D}$ is not basic, then it has a 2-join and we consider a minimally sided 2-join (X_1, X_2) where X_1 is a minimal side. So, the block of decomposition G^1 (that contains X_1) is a basic graph, and our algorithm solves a number of instances of the INDUCED DISJOINT PATHS problem in it, to know how disjoint paths may exist through X_1 . We then define an o-graph $G_{\mathcal{F}^2, \mathcal{O}^2}^2$, where G^2 is the block of decomposition that contains X_2 . The sets $\mathcal{F}^2, \mathcal{O}^2$ are designed using the information gained from the computations made in G_1 , so that $G_{\mathcal{F},\mathcal{O}}$ is linkable if and only if $G_{\mathcal{F}^2, \mathcal{O}^2}^2$ is linkable (see Lemma 4.10). The marker path of G^2 (or an extension of it) is sometimes added to \mathcal{F}^2 to record different types of interactions of the solution paths with the 2-join. Throughout all these steps of the algorithm, we can prove that $|\mathcal{F}|$ is bounded by the initial number of terminals, and this leads to an FPT algorithm. In the following lemma we prove that $|\mathcal{O}|$ is bounded by a function of $|\mathcal{F}|$.

Lemma 4.7 *Let $G_{\mathcal{F},\mathcal{O}}$ be an o-graph, such that $|\mathcal{F}| \leq t$. Then $|\mathcal{O}| \leq 2^{8t}(8t)!$.*

PROOF — Each path in \mathcal{F} has at most 8 vertices. Hence $|\mathcal{F}| \leq t$ implies that each set in \mathcal{O} has at most $4t$ pairs of vertices. So, there is at most 2^{8t} ways to choose the vertices that are going to be in the pairs of an element of \mathcal{O} . Once we have chosen these vertices, say $2s$ of them ($2s \leq 8t$), to obtain an element of \mathcal{O} we only need to group them in s (disjoint) pairs. This can be done in $\frac{1}{s!} \binom{2s}{2} \binom{2s-2}{2} \cdots \binom{2}{2} = \frac{(2s)!}{2^s s!}$ ways, and hence

$$|\mathcal{O}| \leq 2^{8t} \frac{(2s)!}{2^s s!} \leq 2^{8t}(8t)!.$$

□

Corollary 4.8 *Let t be a fixed integer. If $G_{\mathcal{F},\mathcal{O}}$ is an o-graph such that $G \in \mathcal{B}$ and $|\mathcal{F}| \leq t$, then there is an $\mathcal{O}(n^5)$ -time algorithm that decides whether $G_{\mathcal{F},\mathcal{O}}$ is linkable.*

PROOF — Follows from Lemmas 4.5 and 4.7. □

Let $G \in \mathcal{D}$ be a graph that has a 2-join (X_1, X_2) and (G, \mathcal{W}) an instance of the k -INDUCED DISJOINT PATHS problem. For a split $(X_1, X_2, A_1, A_2, B_1, B_2)$ of (X_1, X_2) we define:

- $\mathcal{I}_1 = \{1 \leq i \leq k : s_i, t_i \in X_1\}$;
- $\mathcal{I}_2 = \{1 \leq i \leq k : s_i, t_i \in X_2\}$;




Figure 1: G and its blocks of decomposition G^1 and G^2

- $\mathcal{J} = \{1, 2, \dots, k\} \setminus (\mathcal{I}_1 \cup \mathcal{I}_2)$;
- $\mathcal{W}'_1 = \{(s_i, t_i) : i \in \mathcal{I}_1\}$;
- $\mathcal{W}'_2 = \{(s_i, t_i) : i \in \mathcal{I}_2\}$.

Furthermore, we may assume that for $j \in \mathcal{J}$, $s_j \in X_1$ and $t_j \in X_2$.

Recall that we build a block G^j , for $j \in \{1, 2\}$, by replacing X_{3-j} with a chordless path $P^{3-j} = a_{3-j}a'_{3-j}c_{3-j}d_{3-j}b'_{3-j}b_{3-j}$ (called the marker path) such that a_{3-j} (resp. b_{3-j}) is complete to A_j (resp. B_j). See Fig. 1.

Next, we list a number of pairs of problems $(\mathcal{S}_1, \mathcal{S}_2)$. Problem \mathcal{S}_1 is going to be solved on G^1 and problem \mathcal{S}_2 is going to be solved on G^2 . Each pair $(\mathcal{S}_1, \mathcal{S}_2)$ corresponds to a possible way a solution of (G, \mathcal{W}) interacts with the 2-join. We aim to prove Lemma 4.9 showing that (G, \mathcal{W}) has a solution if and only if for one of the pairs $(\mathcal{S}_1, \mathcal{S}_2)$, both \mathcal{S}_1 and \mathcal{S}_2 have a solution. We therefore call these pairs *potential solutions* (for example, in (2.3) a potential solution is described and we will refer to it as *potential solution (2.3)*).

Our definition of pairs $(\mathcal{S}_1, \mathcal{S}_2)$ depends on the *type* of interaction of \mathcal{W} with the 2-join (X_1, X_2) . We list these types and the potential solutions in what follows. (Note: it is best suited to examine the potential solutions while reading the proof of Lemma 4.9. Also, see Fig. 2, 3, 4 and 5.)

Type 1: $\mathcal{J} = \emptyset$.

$$(1.1) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(a_2, b_2), (c_2, d_2)\}), \quad \mathcal{S}_2 = (G^2, \mathcal{W}'_2);$$

$$(1.2) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(a_2, a'_2), (b_2, b'_2)\}), \quad \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(c_1, d_1)\});$$

$$(1.3) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(b_2, b'_2), (a'_2, c_2)\}), \quad \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a_1, a'_1)\});$$

$$(1.4) \quad A_2 = \{a\}, \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(b_2, b'_2)\}), \quad \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a, a_1)\});$$

$$(1.5) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(a_2, a'_2), (b'_2, d_2)\}), \quad \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(b_1, b'_1)\});$$




Figure 2: Type 1 potential solutions (1.1)–(1.6)




Figure 3: Type 1 potential solutions (1.7)–(1.11)

$$(1.6) \quad B_2 = \{b\}, \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(a_2, a'_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(b, b_1)\});$$

$$(1.7) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(a'_2, b'_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a_1, a'_1), (b_1, b'_1)\});$$

$$(1.8) \quad A_2 = \{a\}, \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(c_2, b'_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a, a_1), (b_1, b'_1)\});$$

$$(1.9) \quad B_2 = \{b\}, \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(a'_2, d_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(b, b_1), (a_1, a'_1)\});$$

$$(1.10) \quad A_2 = \{a\}, B_2 = \{b\}, \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(c_2, d_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a, a_1), (b, b_1)\});$$

$$(1.11) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a_1, b_1), (c_1, d_1)\}).$$

Type 2: $\mathcal{J} = \{j_1\}$.

$$(2.1) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, c_2), (b_2, b'_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a_1, t_{j_1}), (b'_1, d_1)\});$$

$$(2.2) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, d_2), (a_2, a'_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(b_1, t_{j_1}), (a'_1, c_1)\});$$

$$(2.3) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, a'_2), (b'_2, d_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a_1, t_{j_1}), (b_1, b'_1)\});$$

$$(2.4) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, b'_2), (a'_2, c_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(b_1, t_{j_1}), (a_1, a'_1)\});$$

$$(2.5) \quad B_2 = \{b\}, \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, a_2), (c_2, d_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a_1, t_{j_1}), (b, b_1)\});$$

$$(2.6) \quad A_2 = \{a\}, \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, b_2), (c_2, d_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(b_1, t_{j_1}), (a, a_1)\}).$$

Type 3: $\mathcal{J} = \{j_1, j_2\}$.

$$(3.1) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, a_2), (s_{j_2}, b_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(a_1, t_{j_1}), (b_1, t_{j_2})\});$$

$$(3.2) \quad \mathcal{S}_1 = (G^1, \mathcal{W}'_1 \cup \{(s_{j_1}, b_2), (s_{j_2}, a_2)\}), \mathcal{S}_2 = (G^2, \mathcal{W}'_2 \cup \{(b_1, t_{j_1}), (a_1, t_{j_2})\}).$$

In what follows, for a set of paths \mathcal{P} , we denote $V(\mathcal{P}) = \bigcup_{P \in \mathcal{P}} V(P)$.

Lemma 4.9 *Let $(X_1, X_2, A_1, A_2, B_1, B_2)$ be a split of a minimally-sided 2-join of $G \in \mathcal{D}$ such that X_1 is the minimal side. Then (G, \mathcal{W}) has a solution if and only if in one of the potential solutions listed above both \mathcal{S}_1 and \mathcal{S}_2 have a solution.*

PROOF — Let R^j , for $j \in \{1, 2\}$, be a chordless path in $G[X_j]$ whose one endnode is in A_j , the other in B_j and no interior node is in $A_j \cup B_j$ (these paths exist by Lemma 2.5). Furthermore, let $\mathcal{I}_1, \mathcal{I}_2, \mathcal{J}, \mathcal{W}'_1$ and \mathcal{W}'_2 be defined as above. If $|\mathcal{J}| \geq 3$, then (G, \mathcal{W}) has no solution. So, we may assume that $|\mathcal{J}| \leq 2$. Depending on how \mathcal{W} interacts with the 2-join (X_1, X_2) we define different options for \mathcal{S}_1 and \mathcal{S}_2 as above. For $i = 1, 2$, we denote by \mathcal{W}_i the set of terminal pairs of the problem \mathcal{S}_i .

By Lemma 2.4 and Lemma 2.7(i) we have $|A_1|, |B_1| \geq 2$, and consequently, by Lemma 2.5, A_2 and B_2 are both cliques, and if A_1 (resp. B_1) is not a clique then $|A_2| = 1$ (resp. $|B_2| = 1$).




Figure 4: Type 2 potential solutions




Figure 5: Type 3 potential solutions

(\Rightarrow): Let $\mathcal{P} = \{P_1, \dots, P_k\}$ be a solution of (G, \mathcal{W}) (where for $i = 1, \dots, k$, P_i is a path from s_i to t_i) and let $\mathcal{P}_j = \{P_i \in \mathcal{P} : i \in \mathcal{I}_j\}$, for $j \in \{1, 2\}$. Note that since A_2 and B_2 are cliques, if a path from \mathcal{P}_2 contains a vertex from X_1 , then it contains vertices from both A_1 and B_1 . We now consider the following cases (that correspond to different types of interaction of \mathcal{W} with 2-join (X_1, X_2) from the definition of \mathcal{S}_1 and \mathcal{S}_2).

Case 1: $\mathcal{J} = \emptyset$.

First, let us assume that a path P_i from \mathcal{P}_2 contains a vertex from X_1 . Then P_i contains vertices from both A_1 and B_1 . Let P'_i be the subpath of P_i contained in $G[X_1]$. Then $\mathcal{P}_1 \cup \{P'_i, c_2 d_2\}$ is a solution of the problem \mathcal{S}_1 from (1.1), where P'_i is the path induced by $V(P'_i) \cup \{a_2, b_2\}$, and $(\mathcal{P}_2 \cup \{P'_i\}) \setminus \{P_i\}$ is a solution of the problem \mathcal{S}_2 from (1.1), where P''_i is the path induced by $(V(P_i) \setminus V(P'_i)) \cup \{a_1, a'_1, b_1, b'_1, c_1, d_1\}$. So, in what follows, we may assume that $V(\mathcal{P}_2) \cap X_1 = \emptyset$.

Next assume that $V(\mathcal{P}_1) \cap X_2 = \emptyset$. If $V(\mathcal{P}_1) \cap (A_1 \cup B_1) = \emptyset$, then $\mathcal{P}_1 \cup \{a_2 a'_2, b_2 b'_2\}$ is a solution of \mathcal{S}_1 from (1.2) and $\mathcal{P}_2 \cup \{c_1 d_1\}$ is a solution of \mathcal{S}_2 from (1.2). If $V(\mathcal{P}_1) \cap A_1 \neq \emptyset$ and $V(\mathcal{P}_1) \cap B_1 = \emptyset$, then no path from \mathcal{P}_2 has a vertex in A_2 and hence $\mathcal{P}_1 \cup \{b_2 b'_2, a'_2 c_2\}$ is a solution of \mathcal{S}_1 from (1.3) and $\mathcal{P}_2 \cup \{a_1 a'_1\}$ is a solution of \mathcal{S}_2 from (1.3). Similarly, if $V(\mathcal{P}_1) \cap A_1 = \emptyset$ and $V(\mathcal{P}_1) \cap B_1 \neq \emptyset$, then both \mathcal{S}_1 and \mathcal{S}_2 from (1.5) have a solution. If $V(\mathcal{P}_1) \cap A_1 \neq \emptyset$ and $V(\mathcal{P}_1) \cap B_1 \neq \emptyset$, then no path from \mathcal{P}_2 has a vertex in $A_2 \cup B_2$ and hence $\mathcal{P}_1 \cup \{a'_2 c_2 d_2 b'_2\}$ is a solution of \mathcal{S}_1 from (1.7) and $\mathcal{P}_2 \cup \{a_1 a'_1, b_1 b'_1\}$ is a solution of \mathcal{S}_2 from (1.7).

We may now assume that $V(\mathcal{P}_1) \cap X_2 \neq \emptyset$. First, we examine the case when $|V(\mathcal{P}_1) \cap X_2| = 1$. Then $V(\mathcal{P}_1) \cap X_2 = \{x\}$, and $x \in A_2$ or $x \in B_2$. Suppose that $x \in A_2$. Then A_1 is not a clique, and hence $A_2 = \{x\}$. Also, no path from \mathcal{P}_2 contains x and no vertex of a path from \mathcal{P}_2 is adjacent to x . Let P_i be the path of \mathcal{P}_1 that contains x , and let P'_i be the

path of G^1 obtained from P_i by replacing x with a_2 . Also, let $\mathcal{P}'_1 = (\mathcal{P}_1 \setminus \{P_i\}) \cup \{P'_i\}$. If $V(\mathcal{P}_1) \cap B_1 = \emptyset$, then $\mathcal{P}'_1 \cup \{b_2b'_2\}$ is a solution of \mathcal{S}_1 from (1.4) and $\mathcal{P}_2 \cup \{a_1x\}$ is a solution of \mathcal{S}_2 from (1.4). If $V(\mathcal{P}_1) \cap B_1 \neq \emptyset$, then $V(\mathcal{P}_2) \cap B_2 = \emptyset$ and hence $\mathcal{P}'_1 \cup \{c_2d_2b'_2\}$ is a solution of \mathcal{S}_1 from (1.8) and $\mathcal{P}_2 \cup \{a_1x, b_1b'_1\}$ is a solution of \mathcal{S}_2 from (1.8). So we have shown that if $x \in A_2$ then either both \mathcal{S}_1 and \mathcal{S}_2 from (1.4) have a solution, or both \mathcal{S}_1 and \mathcal{S}_2 from (1.8) have a solution. By symmetric argument, if $x \in B_2$ then either both \mathcal{S}_1 and \mathcal{S}_2 from (1.6) have a solution, or both \mathcal{S}_1 and \mathcal{S}_2 from (1.9) have a solution.

Next suppose that $V(\mathcal{P}_1) \cap X_2 = \{a, b\}$, where ab is not an edge. Without loss of generality, let $a \in A_2$ and $b \in B_2$. It follows, as above, that A_1 and B_1 are not cliques, and hence $A_2 = \{a\}$ and $B_2 = \{b\}$. Also, no vertex of a path from \mathcal{P}_2 is adjacent to a or b . Let P_i (resp. P_j) be the path of \mathcal{P}_1 that contains a (resp. b), and let P'_i (resp. P'_j) be the path of G^1 obtained from P_i (resp. P_j) by replacing a (resp. b) with a_2 (resp. b_2). (Note that in this case possibly $P_i = P_j$, i.e. the path contains both a and b . In this case $P'_i = P'_j$ is the path obtained from $P_i = P_j$ by replacing a with a_2 and b with b_2 .) Then $(\mathcal{P}_1 \setminus \{P_i, P_j\}) \cup \{P'_i, P'_j, c_2d_2\}$ is a solution of \mathcal{S}_1 from (1.10), and $\mathcal{P}_2 \cup \{aa_1, bb_1\}$ is a solution of \mathcal{S}_2 from (1.10).

Finally, suppose that a path P_i from \mathcal{P}_1 contains an edge of X_2 . Let P'_i be the subpath of P_i contained in $G[X_2]$, P''_i the path induced by $(V(P_i) \setminus V(P'_i)) \cup \{a_2, a'_2, b_2, b'_2, c_2, d_2\}$, and P'''_i the path induced by $V(P'_i) \cup \{a_1, b_1\}$. Then $(\mathcal{P}_1 \setminus \{P_i\}) \cup \{P''_i\}$ is a solution of \mathcal{S}_1 from (1.11) and $\mathcal{P}_2 \cup \{P'''_i, c_1d_1\}$ is a solution of \mathcal{S}_2 from (1.11).

Case 2: $|\mathcal{J}| = 1$.

Let $\mathcal{J} = \{j\}$. So $P_j \in \mathcal{P}$ is a path from s_j to t_j . We consider the case when $V(P_j) \cap A_1$ and $V(P_j) \cap A_2$ are both non-empty, and show that in that case in one of the cases (2.1), (2.3) or (2.5), both \mathcal{S}_1 and \mathcal{S}_2 have a solution. The case when $V(P_j) \cap B_1$ and $V(P_j) \cap B_2$ are both non-empty is handled similarly and leads to \mathcal{S}_1 and \mathcal{S}_2 both having a solution in one of the cases (2.2), (2.4) or (2.6). So assume that both $V(P_j) \cap A_1$ and $V(P_j) \cap A_2$ are non-empty. Let P_j^1 (resp. P_j^2) be the s_ja' -subgraph (resp. $a''t_j$ -subpath) of P_j , where $V(P_j) \cap A_1 = \{a'\}$ (resp. $V(P_j) \cap A_2 = \{a''\}$). Furthermore, let P'_j be the path induced by $V(P_j^1) \cup \{a_2, a'_2, c_2\}$ and P''_j be the path induced by $V(P_j^2) \cup \{a_1\}$.

First, let us assume that no path from $\mathcal{P}_1 \cup \{P_j^1\}$ has a vertex in B_2 . If no path from $\mathcal{P}_1 \cup \{P_j^1\}$ has a vertex in B_1 , then $\mathcal{P}_1 \cup \{P'_j, b_2b'_2\}$ is a solution of \mathcal{S}_1 from (2.1) and $\mathcal{P}_2 \cup \{P''_j, b'_1d_1\}$ is a solution of \mathcal{S}_2 from (2.1). If some path from $\mathcal{P}_1 \cup \{P_j^1\}$ has a vertex in B_1 , then no path from \mathcal{P}_2 has a vertex in B_2 . Hence, $\mathcal{P}_1 \cup \{P'_j \setminus \{c_2\}, b'_2d_2\}$ is a solution of \mathcal{S}_1 from (2.3) and $\mathcal{P}_2 \cup \{P''_j, b_1b'_1\}$ is a solution of \mathcal{S}_2 from (2.3).

Let us now assume that a path from $\mathcal{P}_1 \cup \{P_j^1\}$ has a vertex in B_2 . Then this path has two vertices in B_1 , which implies that B_1 is not a clique and hence that B_2 has a single vertex b . So, no vertex from a path from \mathcal{P}_2 is adjacent to b . Let P_i be the path of $\mathcal{P}_1 \cup \{P_j^1\}$ that contains b , and let P'_i be the path of G^1 obtained from P_i by replacing b with b_2 . Now, if $P_i \neq P_j^1$, then $(\mathcal{P}_1 \cup \{P'_j \setminus \{a'_2, c_2\}, P'_i, c_2d_2\}) \setminus \{P_i\}$ is a solution of \mathcal{S}_1

from (2.5); if $P_i = P_j^1$, then $\mathcal{P}_1 \cup \{P'_i \setminus \{a'_2, c_2\}, c_2 d_2\}$ is a solution of \mathcal{S}_1 from (2.5). Clearly, $\mathcal{P}_2 \cup \{P''_j, bb_1\}$ is a solution of \mathcal{S}_2 from (2.5).

Case 3: $|\mathcal{J}| = 2$.

Let $\mathcal{J} = \{j_1, j_2\}$. So $P_{j_1}, P_{j_2} \in \mathcal{P}$ are paths from s_{j_1} to t_{j_1} and from s_{j_2} to t_{j_2} , respectively. Then no path from $\mathcal{P} \setminus \{P_{j_1}, P_{j_2}\}$ has a vertex in both X_1 and X_2 . For $s, r \in \{1, 2\}$, let $P_{j_r}^s$ be the subpath of P_{j_r} contained in $G[X_s]$. Let P'_{j_r} (resp. P''_{j_r}), for $r \in \{1, 2\}$, be the path induced by $V(P_{j_r}^1) \cup \{a_2\}$ (resp. $V(P_{j_r}^2) \cup \{a_1\}$) if an endnode of $P_{j_r}^1$ (resp. $P_{j_r}^2$) is in A_1 (resp. A_2), or the path induced by $V(P_{j_r}^1) \cup \{b_2\}$ (resp. $V(P_{j_r}^2) \cup \{b_1\}$) if an endnode of $P_{j_r}^1$ (resp. $P_{j_r}^2$) is in B_1 (resp. B_2). So, if an endnode of $P_{j_1}^1$ is in A_1 (resp. B_1), then $\mathcal{P}_1 \cup \{P'_{j_1}, P'_{j_2}\}$ is a solution of \mathcal{S}_1 from (3.1) (resp. (3.2)) and $\mathcal{P}_2 \cup \{P''_{j_1}, P''_{j_2}\}$ is a solution for \mathcal{S}_2 from (3.1) (resp. from (3.2)).

(\Leftarrow): Let \mathcal{P}_1 be a solution of \mathcal{S}_1 and \mathcal{P}_2 a solution of \mathcal{S}_2 of some potential solution. We consider the following cases.

Case 1: $\mathcal{J} = \emptyset$.

Suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (1.1). Let $P_i \in \mathcal{P}_1$ be the path from a_2 to b_2 . Since $c_2 d_2 \in \mathcal{P}_1$, P_i contains a vertex of A_1 and a vertex of B_1 . So no path from $\mathcal{P}_1 \setminus \{P_i, c_2 d_2\}$ contains a vertex from $A_1 \cup B_1$. Hence, if no path from \mathcal{P}_2 contains a vertex from $\{a_1, a'_1, b_1, b'_1, c_1, d_1\}$, then $(\mathcal{P}_1 \cup \mathcal{P}_2) \setminus \{P_i, c_2 d_2\}$ is a solution of (G, \mathcal{W}) . So we may assume that $P_j \in \mathcal{P}_2$ contains a vertex of $\{a_1, a'_1, b_1, b'_1, c_1, d_1\}$. Since A_2 and B_2 are cliques it follows that P_j contains all vertices of $\{a_1, a'_1, b_1, b'_1, c_1, d_1\}$. Let P'_j be the subgraph of P_j contained in $G[X_2]$. Furthermore, let P'_i be the path induced by $V(P_i) \setminus \{a_2, b_2\}$, and P''_j be the path induced by $V(P'_i) \cup V(P'_j)$. Then $(\mathcal{P}_1 \setminus \{P_i, c_2 d_2\}) \cup (\mathcal{P}_2 \setminus \{P_j\}) \cup \{P''_j\}$ is a solution for (G, \mathcal{W}) .

If \mathcal{S}_1 and \mathcal{S}_2 are from (1.2), then $V(\mathcal{P}_1) \setminus \{a_2, a'_2, b_2, b'_2\} \subseteq X_1 \setminus (A_1 \cup B_1)$ and $V(\mathcal{P}_2) \setminus \{c_1, d_1\} \subseteq X_2$ (as A_2 and B_2 are cliques), and hence $(\mathcal{P}_1 \cup \mathcal{P}_2) \setminus \{c_1 d_1, a_2 a'_2, b_2 b'_2\}$ is a solution of (G, \mathcal{W}) .

If \mathcal{S}_1 and \mathcal{S}_2 are from (1.3) (resp. (1.5)), then $V(\mathcal{P}_1) \setminus \{a'_2, b_2, b'_2, c_2\} \subseteq X_1 \setminus B_1$ (resp. $V(\mathcal{P}_1) \setminus \{a_2, a'_2, b'_2, d_2\} \subseteq X_1 \setminus A_1$) and no path from \mathcal{P}_2 has a vertex in A_2 (resp. B_2). Hence $(\mathcal{P}_1 \cup \mathcal{P}_2) \setminus \{a_1 a'_1, b_2 b'_2, a'_2 c_2\}$ (resp. $(\mathcal{P}_1 \cup \mathcal{P}_2) \setminus \{b_1 b'_1, a_2 a'_2, b'_2 d_2\}$) is a solution of (G, \mathcal{W}) .

Suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (1.4). Then no path of $\mathcal{P}_2 \setminus \{aa_1\}$ contains the vertex of A_2 nor a vertex that is adjacent to the vertex of A_2 , and no path of \mathcal{P}_1 contains a vertex of B_1 . So, if $V(\mathcal{P}_1) \cap \{a_2\} = \emptyset$, then clearly $(\mathcal{P}_1 \cup \mathcal{P}_2) \setminus \{aa_1, b_2 b'_2\}$ is a solution of (G, \mathcal{W}) . Now suppose that a path $P_i \in \mathcal{P}_1$ contains a_2 . Let P'_i be the path obtained from P_i by replacing a_2 by a . Then $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P'_i\}) \setminus \{P_i, aa_1, b_1 b'_1\}$ is a solution of (G, \mathcal{W}) . Similarly, if \mathcal{S}_1 and \mathcal{S}_2 are from (1.6), then (G, \mathcal{W}) has a solution.

If \mathcal{S}_1 and \mathcal{S}_2 are from (1.7), then no path from \mathcal{P}_2 has a vertex in $A_2 \cup B_2$ and hence $(\mathcal{P}_1 \cup \mathcal{P}_2) \setminus \{a_1 a'_1, b_1 b'_1, P_t\}$ is a solution of (G, \mathcal{W}) , where $P_t \in \mathcal{P}_1$ is the path from a'_2 to b'_2 .

Suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (1.8). Then no path of $\mathcal{P}_2 \setminus \{aa_1, b_1b'_1\}$ contains the vertex of A_2 nor a vertex that is adjacent to the vertex of A_2 ; also, no path of $\mathcal{P}_2 \setminus \{aa_1, b_1b'_1\}$ contains a vertex of B_2 . Let P_t be the path of \mathcal{S}_1 from c_2 to b'_2 . If $V(\mathcal{P}_1) \cap \{a_2\} = \emptyset$, then clearly $(\mathcal{P}_1 \cup \mathcal{P}_2) \setminus \{aa_1, b_1b'_1, P_t\}$ is a solution of (G, \mathcal{W}) . So, suppose that a path $P_i \in \mathcal{P}_1$ contains a_2 . Let P'_i be the path obtained from P_i by replacing a_2 by a . Then $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P'_i\}) \setminus \{P_i, aa_1, b_1b'_1, P_t\}$ is a solution of (G, \mathcal{W}) . Similarly, if \mathcal{S}_1 and \mathcal{S}_2 are from (1.9), then (G, \mathcal{W}) has a solution.

Suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (1.10). Then no path of \mathcal{P}_2 contains a vertex of $A_2 \cup B_2$ nor a vertex that is adjacent to a vertex of $A_2 \cup B_2$. If a_2 (resp. b_2) is contained in some path of \mathcal{P}_1 then replace it by a (resp. b). Let \mathcal{P}'_1 be the resulting set of paths. Then $(\mathcal{P}'_1 \cup \mathcal{P}_2) \setminus \{aa_1, bb_1, c_2d_2\}$ is a solution of (G, \mathcal{W}) .

Finally suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (1.11). Let P' be the path of \mathcal{P}_2 from a_1 to b_1 . If $V(\mathcal{P}_1) \cap \{a_2, a'_2, c_2, d_2, b'_2, b_2\} = \emptyset$, then $\mathcal{P}_1 \cup (\mathcal{P}_2 \setminus \{P', c_1d_1\})$ is a solution of (G, \mathcal{W}) . If some path of \mathcal{P}_1 contains a_2 (resp. b_2) and no other vertex of the marker path, then replace a_2 (resp. b_2) in that path by a vertex $a \in A_2$ (resp. $b \in B_2$), and let \mathcal{P}'_1 be the resulting family of paths. Note that by Lemma 2.5 2-join (X_1, X_2) is consistent and hence we may choose a and b to be non-adjacent. Then $\mathcal{P}'_1 \cup (\mathcal{P}_2 \setminus \{P', c_1d_1\})$ is a solution of (G, \mathcal{W}) . So we may assume that some path P_i of \mathcal{P}_1 contains all of the vertices $a_2, a'_2, c_2, d_2, b'_2, b_2$. Let P'_i be the path of G induced by $(V(P_i) \cap X_1) \cup (V(P') \setminus \{a_1, b_1\})$. Then $(\mathcal{P}_1 \setminus \{P_i\}) \cup (\mathcal{P}_2 \setminus \{P', c_1d_1\}) \cup \{P'_i\}$ is a solution for (G, \mathcal{W}) .

Case 2: $|\mathcal{J}| = 1$.

Let $\mathcal{J} = \{j\}$, and $P_j^1 \in \mathcal{P}_1$ be the path from s_j to the appropriate vertex of the marker path P^2 . First, suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (2.1) (resp. (2.2)), and let $P_j^2 \in \mathcal{P}_2$ be the path from t_j to a_1 (resp. b_1). Furthermore, let P_j be the path induced by $(V(P_j^1) \cup V(P_j^2)) \setminus \{a_2, a'_2, c_2, a_1\}$ (resp. $(V(P_j^1) \cup V(P_j^2)) \setminus \{b_2, b'_2, d_2, b_1\}$). Then no path from \mathcal{P}_1 has a vertex in B_1 (resp. A_1), and hence $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P_j\}) \setminus \{P_j^1, P_j^2, b'_1d_1, b_2b'_2\}$ (resp. $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P_j\}) \setminus \{P_j^1, P_j^2, a'_1c_1, a_2a'_2\}$) is a solution for (G, \mathcal{W}) .

Next, suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (2.3) (resp. (2.4)), and let $P_j^2 \in \mathcal{P}_2$ be the path from t_j to a_1 (resp. b_1). Furthermore, let P_j be the path induced by $(V(P_j^1) \cup V(P_j^2)) \setminus \{a_2, a'_2, a_1\}$ (resp. $(V(P_j^1) \cup V(P_j^2)) \setminus \{b_2, b'_2, b_1\}$). Then no path from \mathcal{P}_2 has a vertex in B_2 (resp. A_2), and hence $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P_j\}) \setminus \{P_j^1, P_j^2, b_1b'_1, b'_2d_2\}$ (resp. $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P_j\}) \setminus \{P_j^1, P_j^2, a_1a'_1, a'_2c_2\}$) is a solution for (G, \mathcal{W}) .

Suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (2.5), and let $P_j^2 \in \mathcal{P}_2$ be the path from t_j to a_1 . If P_j^1 contains b_2 then let P_j be the path induced by $(V(P_j^1) \cup V(P_j^2) \setminus \{a_2, b_2, a_1\}) \cup \{b\}$, and otherwise let P_j be the path induced by $(V(P_j^1) \cup V(P_j^2)) \setminus \{a_2, a_1\}$. If no path of $\mathcal{P}_1 \setminus P_j^1$ contains b_2 then $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P_j\}) \setminus \{P_j^1, P_j^2, bb_1, c_2d_2\}$ is a solution for (G, \mathcal{W}) . So suppose that $P_i \in \mathcal{P}_1 \setminus \{P_j^1\}$ contains b_2 . Let P'_i be the path obtained from P_i by replacing b_2 with b . Then $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P'_i, P_j\}) \setminus \{P_i, P_j^1, P_j^2, bb_1, c_2d_2\}$ is a solution for (G, \mathcal{W}) . Similarly, if \mathcal{S}_1 and \mathcal{S}_2 are from (2.6), then (G, \mathcal{W}) has a solution.

Case 3: $|\mathcal{J}| = 2$.

Let $\mathcal{J} = \{j_1, j_2\}$. Suppose that \mathcal{S}_1 and \mathcal{S}_2 are from (3.1). Let $P_{j_1}^1, P_{j_2}^1 \in \mathcal{P}_1$ be the paths from s_{j_1} to a_2 and from s_{j_2} to b_2 , respectively, and let $P_{j_1}^2, P_{j_2}^2 \in \mathcal{P}_2$ be the paths from t_{j_1} to a_1 and from t_{j_2} to b_1 , respectively. Furthermore, let P_{j_1} be the path induced by $(V(P_{j_1}^1) \cup V(P_{j_1}^2)) \setminus \{a_1, a_2\}$ and P_{j_2} be the path induced by $(V(P_{j_2}^1) \cup V(P_{j_2}^2)) \setminus \{b_1, b_2\}$. Then $(\mathcal{P}_1 \cup \mathcal{P}_2 \cup \{P_{j_1}, P_{j_2}\}) \setminus \{P_{j_1}^1, P_{j_1}^2, P_{j_2}^1, P_{j_2}^2\}$ is a solution for (G, \mathcal{W}) . Similarly, if \mathcal{S}_1 and \mathcal{S}_2 are from (3.2), then (G, \mathcal{W}) has a solution. \square

By the previous lemma, in order to solve (G, \mathcal{W}) it is enough to solve \mathcal{S}_1 and \mathcal{S}_2 in each of the potential solutions explained above. This will be recorded in the corresponding o-graph. Further, we note that in our main algorithm the graph G^1 is going to be basic (we use the decomposition tree T_G), so all the problems \mathcal{S}_1 can be solved using Lemma 4.5.

Let $G_{\mathcal{F}, \mathcal{O}}$ be an o-graph such that $G \in \mathcal{D}$ has a 2-join (X_1, X_2) . Further, we assume that (X_1, X_2) is a minimally-sided 2-join of G such that X_1 is the minimal side. Let G^1 and G^2 be the blocks of decompositions with respect to this 2-join and P^1 and P^2 the marker paths used to build these blocks (P^j is contained in G^{3-j} , for $j \in \{1, 2\}$). Let the *extended marker path* P_{ext}^1 of G^2 be the path obtained by adding to P^1 vertex a if it is the unique vertex of A_2 and b if it is the unique vertex of B_2 . Note that, by this definition, P_{ext}^1 is a flat path of length at most 7. In what follows we define the o-graph $G_{\mathcal{F}^2, \mathcal{O}^2}^2$.

For $j \in \{1, 2\}$, let \mathcal{F}'_j be the set of flat paths \mathcal{F} that are contained in X_j , together with the set of paths $P \in \mathcal{F}$ of length 0 contained in X_j . Let $\mathcal{F}^0 = \mathcal{F} \setminus (\mathcal{F}'_1 \cup \mathcal{F}'_2)$. So, \mathcal{F}^0 contains only flat paths. By Lemma 2.7 and Remark 2.8, for every $P \in \mathcal{F}^0$, the set $P \cap X_2$ is of size 1 and contained in $A_2 \cup B_2$. Furthermore, since P is a flat path of length greater than 1, if $P \cap A_2 \neq \emptyset$, then $|A_2| = 1$ and similarly if $P \cap B_2 \neq \emptyset$, then $|B_2| = 1$.

Let us now define \mathcal{F}^2 and \mathcal{O}^2 . If $\mathcal{F}'_1 \cup \mathcal{F}^0 = \emptyset$, then $\mathcal{F}^2 = \mathcal{F}'_2$ and otherwise $\mathcal{F}^2 = \mathcal{F}'_2 \cup \{P_{\text{ext}}^1\}$. Next, for every $\mathcal{W} \in \mathcal{O}$ and every potential solution associated with the problem (G, \mathcal{W}) we solve the corresponding problem \mathcal{S}_1 and if it has a solution we add the set of terminal pairs of the corresponding problem \mathcal{S}_2 to \mathcal{O}^2 , except when \mathcal{S}_1 from (1.1) has a solution. In this last case, we add the set of terminal pairs of the corresponding problem \mathcal{S}_2 to \mathcal{O}^2 and we disregard all other potential solutions for \mathcal{W} . By construction, all terminal vertices of \mathcal{S}_2 are in the paths from $\mathcal{F}'_2 \cup \{P_{\text{ext}}^1\}$. Further, if each terminal vertex of \mathcal{W} is in a path from \mathcal{F}'_2 (in particular when $\mathcal{F}'_1 \cup \mathcal{F}^0 = \emptyset$), then all terminal vertices of a set that is added to \mathcal{O}^2 are in the paths from \mathcal{F}'_2 . Indeed, then in potential solution (1.1) we have $\mathcal{S}_1 = (G^1, \{(a_2, b_2), (c_2, d_2)\})$, which has a solution since (X_1, X_2) is a consistent 2-join (by Lemma 2.5). Hence, only the set of terminals for \mathcal{S}_2 from the potential solution (1.1) is added to \mathcal{O}^2 , and this set contains only vertices from paths of \mathcal{F}'_2 . So, $(G^2, \mathcal{F}^2, \mathcal{O}^2)$ is a well-defined o-graph.

Lemma 4.10 *Let $G_{\mathcal{F}, \mathcal{O}}$ be an o-graph such that $G \in \mathcal{D}$, and let (X_1, X_2) be a minimally sided 2-join of G such that X_1 is the minimal side. Further, let G^1 and G^2 be the blocks*

of decompositions with respect to this 2-join and $G_{\mathcal{F}^2, \mathcal{O}^2}^2$ o-graph constructed above. Then $G_{\mathcal{F}, \mathcal{O}}$ is linkable if and only if $G_{\mathcal{F}^2, \mathcal{O}^2}^2$ is linkable.

PROOF — Suppose $G_{\mathcal{F}^2, \mathcal{O}^2}^2$ is linkable. Then $G_{\mathcal{F}, \mathcal{O}}$ is linkable as shown by Lemma 4.9 and the definition of $G_{\mathcal{F}^2, \mathcal{O}^2}^2$.

So, suppose that $G_{\mathcal{F}, \mathcal{O}}$ is linkable, that is some problem (G, \mathcal{W}) with $\mathcal{W} \in \mathcal{O}$ has a solution. Then the converse also follows from Lemma 4.9, except in the case when for (G, \mathcal{W}) the problem \mathcal{S}_1 from (1.1) has a solution. Recall that in this case we add the set of terminal pairs of problem \mathcal{S}_2 from (1.1) to \mathcal{O}^2 and disregard all other potential solutions for \mathcal{W} . Since \mathcal{W} is of type 1 and (G, \mathcal{W}) has a solution, by Lemma 4.9 problem \mathcal{S}'_2 from (1.i), for some $i = 1, \dots, 11$ has a solution. This implies that \mathcal{S}_2 from (1.1) has a solution since the set of pairs of terminals for \mathcal{S}_2 is a subset of the one for \mathcal{S}'_2 . Hence, $G_{\mathcal{F}^2, \mathcal{O}^2}^2$ is linkable. \square

Theorem 4.11 *Let $c \in \mathbb{N}$ be a constant. There is an algorithm with the following specifications:*

Input: An o-graph $G_{\mathcal{F}, \mathcal{O}}$, where $|\mathcal{F}| \leq c$ and $G \in \mathcal{D}$.

Output: YES if $G_{\mathcal{F}, \mathcal{O}}$ is linkable, and NO otherwise.

Running time: $\mathcal{O}(n^6)$.

PROOF — In [20] an $\mathcal{O}(n^2m)$ -time algorithm is given for recognizing whether a graph belongs to \mathcal{B} . If $G \in \mathcal{B}$, then the problem can be solved in time $\mathcal{O}(n^5)$ using Corollary 4.8. So we may assume that $G \in \mathcal{D} \setminus \mathcal{B}$. By Theorem 2.1, G has a 2-join, and by Lemma 2.4 G has no star cutset. Using Lemma 2.10 we build a 2-join decomposition tree T_G in time $\mathcal{O}(n^4m)$ and use the notation from the definition of T_G .

Description and correctness of the algorithm. We now process the decomposition tree T_G from its root $G^0 = G$ and the associated o-graph $G_{\mathcal{F}^0, \mathcal{O}^0}^0$ (where $\mathcal{F}^0 = \mathcal{F}$ and $\mathcal{O}^0 = \mathcal{O}$). For each $0 \leq i \leq p-1$ we build the o-graph $(G^{i+1}, \mathcal{F}^{i+1}, \mathcal{O}^{i+1})$. Then, by Lemma 4.10, $G_{\mathcal{F}, \mathcal{O}}$ is linkable if and only if $(G^p, \mathcal{F}^p, \mathcal{O}^p)$ is linkable. So, it is enough to check whether $(G^p, \mathcal{F}^p, \mathcal{O}^p)$ is linkable; since $G^p \in \mathcal{B}$, this can be done using Corollary 4.8 (in the next paragraph we will verify that all hypotheses to Corollary 4.8 hold).

Complexity of the algorithm. Let $|\mathcal{F}| = t$. We prove that for each $i \in \{0, 1, \dots, p\}$, $|\mathcal{F}^i| \leq t$. Our proof is by induction on i . So, suppose that this is true for $i \leq p-1$ and let us prove it for $i+1$. Note that \mathcal{F}^{i+1} is obtained from \mathcal{F}^i by adding at most one element (that corresponds to the extended marker path of G^{i+1}) and removing all elements from \mathcal{F}^i that are in the corresponding set $\mathcal{F}'_1 \cup \mathcal{F}^0$. So, $|\mathcal{F}^{i+1}| \leq |\mathcal{F}^i| \leq t$, unless in $(G^i, \mathcal{F}^i, \mathcal{O}^i)$ the set of corresponding paths $\mathcal{F}'_1 \cup \mathcal{F}^0$ is empty. But then the extended marker path of G^{i+1}

is not an element of \mathcal{F}^{i+1} , and hence $\mathcal{F}^{i+1} \subseteq \mathcal{F}^i$, which implies $|\mathcal{F}^{i+1}| \leq t$. In particular, since $|\mathcal{F}| \leq c$, we have $|\mathcal{F}^i| \leq c$, for $0 \leq i \leq p$.

Let us now examine the complexity of our algorithm. By Lemma 4.7 and previous paragraph, for each $i \in \{0, 1, \dots, p-1\}$, to build o-graphs $(G^{i+1}, \mathcal{F}^{i+1}, \mathcal{O}^{i+1})$ from the o-graph $(G^i, \mathcal{F}^i, \mathcal{O}^i)$ it is enough to solve at most $11 \cdot |\mathcal{O}^i| \leq 11 \cdot 2^{8c} (8c)!$ INDUCED DISJOINT PATHS problems on G_B^{i+1} (since for each problem (G^i, \mathcal{W}) , where $\mathcal{W} \in \mathcal{O}^i$ we need to solve at most 11 problems from the potential solutions) and each of these problems has at most $4|\mathcal{F}^i| \leq 4c$ terminal pairs. So, by Lemma 4.5, we can build o-graphs $(G^{k+1}, \mathcal{F}^{k+1}, \mathcal{O}^{k+1})$, for $i \in \{0, 1, \dots, p-1\}$, in time $\mathcal{O}(p \cdot n^5) = \mathcal{O}(n^6)$ (since $p = \mathcal{O}(n)$ by Lemma 2.10). Finally, using Corollary 4.8 we can check if $(G^p, \mathcal{F}^p, \mathcal{O}^p)$ is linkable in time $\mathcal{O}(n^5)$. \square

Theorem 4.12 *For a fixed integer k , there is an algorithm with the following specifications:*

Input: *A graph $G \in \mathcal{D}$ and a set of pairs $\mathcal{W} = \{(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)\}$ of vertices of G such that all $2k$ vertices are distinct and the only possible edges between these vertices are of the form $s_i t_i$, for some $1 \leq i \leq k$.*

Output: *YES if the problem (G, \mathcal{W}) has a solution, and NO otherwise.*

Running time: $\mathcal{O}(n^6)$.

PROOF — Let $G_{\mathcal{F}, \mathcal{O}}$ be the o-graph defined with $\mathcal{F} = \{s_i : 1 \leq i \leq k\} \cup \{t_i : 1 \leq i \leq k\}$ and $\mathcal{O} = \{\mathcal{W}\}$. Then $G_{\mathcal{F}, \mathcal{O}}$ is linkable if and only if (G, \mathcal{W}) has a solution. Hence, to solve the given problem it is enough to apply Theorem 4.11 for $G_{\mathcal{F}, \mathcal{O}}$. \square

Corollary 4.13 *For graphs in \mathcal{D} the k -INDUCED DISJOINT PATHS problem is fixed-parameter tractable, when parameterized by k .*

PROOF — Let (G, \mathcal{W}) be an instance of the k -INDUCED DISJOINT PATHS problem. As in the proof of Corollary 3.7, we conclude that the problem (G, \mathcal{W}) can be solved in time $2^{2k} h(k) n^5$ for graphs in \mathcal{B} . Now, for each $i \in \{0, 1, \dots, p\}$ the algorithm from Theorem 4.12 has at most $11 \cdot 2^{16k} (16k)!$ calls to the algorithm from Lemma 4.5 (here $c = 2k$), $p \leq n$ (by Lemma 2.10) and in each call of Lemma 4.5 we solve a problem with at most $8k$ terminal pairs. We conclude that the problem (G, \mathcal{W}) can be solved in time

$$(p+1) \cdot 11 \cdot 2^{16k} (16k)! 2^{16k} h(8k) n^5 \leq 22 \cdot 2^{32k} (16k)! h(8k) n^6$$

for graphs in \mathcal{D} . \square

4.4 Induced disjoint paths on \mathcal{C} and some related problems

Theorems 2.1, 4.4 and 4.12 directly imply the following theorem.

Theorem 4.14 *For a fixed integer k , there is an algorithm with the following specifications:*

Input: *A graph $G \in \mathcal{C}$ and a set of pairs $\mathcal{W} = \{(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)\}$ of vertices of G such that all $2k$ vertices are distinct and the only possible edges between these vertices are of the form $s_i t_i$, for some $1 \leq i \leq k$.*

Output: *YES if the problem (G, \mathcal{W}) has a solution, and NO otherwise.*

Running time: $\mathcal{O}(n^{2k+6})$.

In what follows we consider several problems (for graphs in \mathcal{C}) that are related to the k -INDUCED DISJOINT PATHS problem. The first is in fact its generalization.

Let k be a fixed integer (that is not part of the input), G a graph and $\mathcal{W} = \{(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)\}$ a multiset of pairs of vertices of G (so, pairs need not be disjoint, there may be edges between vertices from $W = \bigcup_{i=1}^k \{s_i, t_i\}$ that are not of the form $s_i t_i$ and we allow $(s_i, t_i) = (s_j, t_j)$, for $i \neq j$). Again, we say that vertices of W are *terminals* of \mathcal{W} .

We consider the following problem: decide whether there exist k paths $P_i = s_i \dots t_i$, $1 \leq i \leq k$, that are vertex-disjoint except that they can have a common endnode, and such that there are no edges between vertices of these paths except those on the paths and the ones from $G[W]$. For this problem we use the same notation as for the k -INDUCED DISJOINT PATHS problem, that is, we denote it with (G, \mathcal{W}) .

Theorem 4.15 *For a fixed integer k , there is an algorithm with the following specifications:*

Input: *A graph $G \in \mathcal{C}$ and a multiset $\mathcal{W} = \{(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)\}$ of pairs of vertices of G .*

Output: *YES if the problem (G, \mathcal{W}) has a solution, and NO otherwise.*

Running time: $\mathcal{O}(n^{4k+6})$.

PROOF — Let $W = \bigcup_{i=1}^k \{s_i, t_i\}$ be the set of terminals of \mathcal{W} in G . First, let \mathcal{Q} be the multiset of all pairs of type (v, v) from \mathcal{W} . Furthermore, let $Q = \bigcup_{(v,v) \in \mathcal{Q}} \{v\}$ and $N = \left(Q \cup \bigcup_{v \in Q} N(v)\right) \setminus W'$, where W' is the set of terminals of the multiset $\mathcal{W} \setminus \mathcal{Q}$. Then the problem (G, \mathcal{W}) is equivalent to $(G \setminus N, \mathcal{W} \setminus \mathcal{Q})$. So, we may assume that $s_i \neq t_i$, for $1 \leq i \leq k$.

Next, let \mathcal{F} be the multiset of all pairs of type (v, w) from \mathcal{W} such that vw is an edge of G . Furthermore, let $F = \bigcup_{(v,w) \in \mathcal{F}} \{v, w\}$ and $N = (\bigcup_{v \in F} N(v)) \setminus W'$, where W' is the set of terminals of the multiset $\mathcal{W} \setminus \mathcal{F}$. Then the problem (G, \mathcal{W}) is equivalent to $(G \setminus N, \mathcal{W} \setminus \mathcal{F})$. So, we may also assume that $s_i t_i$ is not an edge of G , for $1 \leq i \leq k$.

Let $W = \{w_1, w_2, \dots, w_s\}$ (here W is a set), $W_j = \{i \mid s_i = w_j \text{ or } t_i = w_j\}$ and $k_j = |W_j|$, for $1 \leq j \leq s$. Now, let us assume that (G, \mathcal{W}) has a solution \mathcal{P} . Then for each j , w_j is an endnode of k_j paths from \mathcal{P} and the set of neighbors N_j of w_j on these paths is contained in $N(w_j)$ (note that $|N_j| = k_j$). Let $N = \bigcup_{j=1}^s N_j$. If $v \in N_j \cap N_i$, for some $j \neq i$, then v is on path of \mathcal{P} of length 2, v has no other neighbors in N and is not contained in N_t , for $t \notin \{i, j\}$; if for $v', v'' \in N$, $v'v''$ is an edge of G , then v' and v'' are on the same path of a solution of (G, \mathcal{W}) (these two conditions are later called *conditions* $(*)$). Let S be the set of vertices that are in more than one of the sets N_i . Then a solution of (G, \mathcal{W}) gives a solution of (G', \mathcal{W}') , where G' is the graph obtained from G by removing all vertices from $W \cup S$ and all neighbors of vertices from S , and \mathcal{W}' is certain set of pairs of vertices of $N \setminus S$, that satisfies conditions of Theorem 4.14. So, now it is clear that to solve (G, \mathcal{W}) it is enough to do the following. For every $1 \leq j \leq k$ choose a k_j -element subset N_j of $N(w_j)$, check if these subsets satisfy the conditions $(*)$, build the corresponding problem (G', \mathcal{W}') and use Theorem 4.14 to solve it. If for at least one choice of the subsets N_j the corresponding problem (G', \mathcal{W}') has a solution we return YES, and otherwise return NO.

We have $|N(w_j)| = \mathcal{O}(n)$ and $|N_j| = k_j$, so the subset N_j can be chosen in $\mathcal{O}(n^{k_j})$ ways. Hence, all subsets N_j , for $1 \leq j \leq s$, can be chosen in $\mathcal{O}(n^{2k})$ ways, since $\sum_{j=1}^s k_j = 2k$. So, in our algorithm we solve $\mathcal{O}(n^{2k})$ problems of type (G', \mathcal{W}') (using Theorem 4.14), and hence its running time is $\mathcal{O}(n^{4k+6})$. \square

Now, we list some problems that can be reduced to the problem (G, \mathcal{W}) from the previous theorem. These reductions are simple and given in more detail in [9] and [13] (there the reductions are given for claw-free graphs, but they work equally well in general). Throughout k is fixed.

4.4.1 k -in-a-Path

The problem k -IN-A-PATH is to decide whether there is a chordless path in G that contains given vertices v_1, v_2, \dots, v_k of G . To solve this problem it is enough to solve $k!$ problems (G, \mathcal{W}) , where $\mathcal{W} = \{(v_{\sigma(1)}, v_{\sigma(2)}), (v_{\sigma(2)}, v_{\sigma(3)}), \dots, (v_{\sigma(k-1)}, v_{\sigma(k)})\}$ and σ is a permutation of $\{1, 2, \dots, k\}$. So, by Theorem 4.15 the problem k -IN-A-PATH can be solved in time $\mathcal{O}(n^{4k+6})$ for graphs in \mathcal{C} .

4.4.2 H -Anchored Induced Topological Minor

Let H be a fixed graph with vertex set $\{x_1, x_2, \dots, x_k\}$, l edges and s isolated vertices (then $l \leq \binom{k}{2}$ and $s \leq k$). The problem H -ANCHORED INDUCED TOPOLOGICAL MINOR is to decide whether for the given vertices v_1, v_2, \dots, v_k of G there ex-

ists an induced subgraph of G that is isomorphic to a subdivision of H such that the isomorphism maps v_i to x_i , for $1 \leq i \leq k$. We may assume that if x_i is an isolated vertex of H , then v_i is not adjacent to v_j for $j \neq i$, since otherwise the problem clearly does not have a solution. This problem is equivalent to (G, \mathcal{W}) , where $\mathcal{W} = \{(v_i, v_j) \mid x_i x_j \text{ is an edge of } H\} \cup \{(v_i, v_i) \mid x_i \text{ is an isolated vertex of } H\}$, so by Theorem 4.15 it can be solved in time $\mathcal{O}(n^{4(l+s)+6})$ for graphs in \mathcal{C} .

4.4.3 H -Induced Topological Minor

Let H be a fixed graph with vertex set $\{x_1, x_2, \dots, x_k\}$, l edges and s isolated vertices (then $l \leq \binom{k}{2}$ and $s \leq k$). The problem H -INDUCED TOPOLOGICAL MINOR is to decide whether G contains (as an induced subgraph) a subdivision of H . To solve this problem it is enough to solve H -ANCHORED INDUCED TOPOLOGICAL MINOR problem for every k distinct vertices of G . So this problem can be solved in time $\mathcal{O}(n^{k+4(l+s)+6})$ for graphs in \mathcal{C} .

4.4.4 k -in-a-Tree

The problem k -IN-A-TREE is to decide whether there is an induced tree of G that contains given vertices v_1, v_2, \dots, v_k of G . The problem is trivial for $k = 1$, so we may assume that $k \geq 2$. Let us suppose that this problem has a solution, and let T' be a minimal tree (with respect to inclusion) that contains vertices v_1, v_2, \dots, v_k . Then all leaves of T' are from $\{v_1, v_2, \dots, v_k\}$, and hence T' is an induced subdivision of a tree T that satisfies: all leaves and vertices of degree 2 of T are from $\{v_1, v_2, \dots, v_k\}$ and all vertices from $\{v_1, v_2, \dots, v_k\}$ are vertices of T . Note that T has at most $2k - 2$ vertices. Indeed, if T has e edges, v vertices, a leaves and b vertices of degree 2, then $2v - 2 = 2e = \sum_{u \in V(T')} \deg u \geq a + 2b + 3(v - a - b)$, and hence $v \leq 2a + b - 2 \leq 2k - 2$.

So, to solve k -IN-A-TREE for a graph $G \in \mathcal{C}$ (and its vertices $\{v_1, v_2, \dots, v_k\}$) it is enough to do the following. First, we find the set \mathcal{S} of all non-isomorphic trees T with at least k , but at most $2k - 2$ vertices, and such that the total number of leaves and degree 2 vertices of T is at most k . Then $|\mathcal{S}|$ is bounded by a constant (depending on k), and hence the set \mathcal{S} can be found in constant time. Now, for each $T \in \mathcal{S}$ we do the following. First, we choose $s - k$ vertices from $V(G) \setminus \{v_1, v_2, \dots, v_k\}$, where s is the number of vertices of T , and label them with $v_{k+1}, v_{k+2}, \dots, v_s$ arbitrarily. This can be done in time $\mathcal{O}(n^{s-k}) = \mathcal{O}(n^{k-2})$. Let a be the number of leaves and b the number of degree 2 vertices of T . Now, we assign labels $\{1, 2, \dots, s\}$ to vertices of T (we build several instances of such assignments) and solve the appropriate T -ANCHORED INDUCED TOPOLOGICAL MINOR problem. First, we choose a set A of a distinct vertices from $\{v_1, v_2, \dots, v_k\}$ and then label the leaves of T with a permutation of the appropriate a indices. Next, we choose a set B of b distinct vertices from $\{v_1, v_2, \dots, v_k\} \setminus A$ and then label the degree 2 vertices of T with a permutation of the appropriate b indices. The remaining vertices of $V(T)$ (which

are all of degree at least 3) are then label with a permutation of indices of elements from $\{v_1, v_2, \dots, v_s\} \setminus (A \cup B)$. So, the total number of T -ANCHORED INDUCED TOPOLOGICAL MINOR problem that is obtained is bounded by a constant (depending on k), and each of them can be solved in time $\mathcal{O}(n^{4(s-1)+6}) = \mathcal{O}(n^{8k-6})$.

Hence, the problem k -IN-A-TREE can be solved in time $\mathcal{O}(n^{k-2} \cdot n^{8k-6}) = \mathcal{O}(n^{9k-8})$ for graphs in \mathcal{C} .

5 When k is part of the input

In the previous section we proved that the problems k -INDUCED DISJOINT PATHS, k -IN-A-PATH and k -IN-A-CYCLE are polynomially solvable on \mathcal{C} for any fixed k . Using the reductions similar to the ones used in [9], we prove that these problems are NP-complete in the class of line graphs of triangle-free chordless graphs when k is part of the input, and hence in \mathcal{C} .

Theorem 5.1 *The INDUCED DISJOINT PATHS problem is NP-complete for the class of line graphs of triangle-free chordless graphs.*

PROOF — We can check in polynomial time if a given collection of paths is a solution of an instance of the INDUCED DISJOINT PATHS problem. Hence, this problem is in NP. To prove the NP-completeness of this problem in the class of line graphs of triangle-free chordless graphs, we reduce from the DISJOINT PATHS problem, which is NP-complete [15].

Let G be a graph and $(s_1, t_1), (s_2, t_2), \dots, (s_k, t_k)$ disjoint pairs of vertices from G that are terminals of an instance \mathcal{S} of the DISJOINT PATHS on G . Also, let G' be the graph obtained from G by subdividing each of its edges once. Then G' is chordless and triangle-free and \mathcal{S} is equivalent with the instance \mathcal{S}' of the DISJOINT PATHS problem on G' with the same terminals. Let G'' be the graph obtained from G' by adding new vertices s'_i and t'_i , and edges $s'_i s_i$ and $t_i t'_i$, for $1 \leq i \leq k$ (the graph G'' remains chordless and triangle-free). Then \mathcal{S}' has a solution if and only if the problem $(L(G''), \mathcal{W})$, where $\mathcal{W} = \{(s'_i s_i, t'_i t_i) \mid 1 \leq i \leq k\}$, has a solution. This completes our reduction, since $L(G'')$ is the line graph of a triangle-free chordless graph. \square

A graph is *cubic* if the degree of each of its vertices is 3. A path of a graph G is *Hamiltonian* if it contains all vertices of G . The HAMILTONIAN PATH problem is to decide whether the given graph has a Hamiltonian path.

Theorem 5.2 *The k -IN-A-PATH problem and the k -IN-A-CYCLE problem are NP-complete in the class of line graphs of triangle-free chordless graphs, when k is part of the input.*

PROOF — We can check in polynomial time if a given path (resp. cycle) is a solution of an instance of the k -IN-A-PATH (resp. k -IN-A-CYCLE) problem, and hence both problems are in NP. To prove the NP-completeness of these problems in the class of line graphs of

triangle-free chordless graphs, we reduce from the HAMILTONIAN PATH problem, which is NP-complete for cubic graphs (see [11, problem GT39]).

Let G be a cubic graph and \mathcal{S} the instance of the HAMILTONIAN PATH problem for G . We build the graph G' from G as follows. For each $u \in V(G)$ we build a triangle $u_1u_2u_3$ of G' , and $\{u_i \mid 1 \leq i \leq 3, u \in V(G)\}$ is the set of vertices of G' . Next, for each edge uv of G we build exactly one edge u_iv_j , for some $i, j \in \{1, 2, 3\}$, such that each vertex of G' is of degree 3 (so, for $u \in V(G)$, $i \in \{1, 2, 3\}$, u_i is adjacent to u_j , for $j \in \{1, 2, 3\} \setminus \{i\}$, and to a vertex v_s for some $v \in N(u)$ and $s \in \{1, 2, 3\}$). It is proved in [9] that \mathcal{S} has a solution if and only if G' has a path that passes through all edges from $\{u_1u_2 \mid u \in V(G)\}$. We call the latter problem \mathcal{S}' . Now, let G'' be the graph obtained from G' by subdividing each edge of G' , that is, each edge xy of G' is replaced with the path $xv_{xy}y$ (where v_{xy} is of degree 2 in G''). Note that G'' is triangle-free and chordless. Then it is clear that \mathcal{S}' has a solution if and only if G'' has a path that passes through all edges from $\{u_1v_{u_1u_2}, u_2v_{u_1u_2} \mid u \in V(G)\}$. We call the later problem \mathcal{S}'' . Since the paths of G'' are in one-to-one correspondence with induced paths of $L(G'')$, the problem \mathcal{S}'' has a solution if and only if there is an induced path of $L(G'')$ that passes through all vertices from $\{u_1v_{u_1u_2}, u_2v_{u_1u_2} \mid u \in V(G)\} \subseteq V(L(G''))$. This completes our reduction, since $L(G'')$ is the line graph of a triangle-free chordless graph.

For the k -IN-A-CYCLE problem analogous reduction is made from the HAMILTONIAN CYCLE problem, which is again NP-complete for cubic graphs (see [11]). \square

References

- [1] R. Belmonte, P.A. Golovach, P. Heggernes, P. van 't Hof, M. Kaminski, and D. Paulusma. Detecting fixed patterns in chordal graphs in polynomial time. *Algorithmica*, 69(3):501–521, 2014.
- [2] D. Bienstock. On the complexity of testing for odd holes and induced odd paths. *Discrete Mathematics*, 90:85–92, 1991. See also Corrigendum by B. Reed, *Discrete Mathematics*, 102:109, 1992.
- [3] H. Bruhn and A. Saito. Clique or hole in claw-free graphs. *Journal of Combinatorial Theory, Series B*, 102(1):1–13, 2012.
- [4] P. Charbit, M. Habib, N. Trotignon, K. Vušković. Detecting 2-joins faster. *Journal of Discrete Algorithms*, 17: 60-66, 2012.
- [5] M. Chudnovsky and P.D. Seymour. The three-in-a-tree problem. *Combinatorica*, 30(4):387–417, 2010.
- [6] N. Dehry, C. Picouleau, and N. Trotignon. The four-in-a-tree problem in triangle-free graphs. *Graphs and Combinatorics*, 25:489–502, 2009.

- [7] E. Diot, M. Radovanović, N. Trotignon, K. Vušković. The (theta,wheel)-free graphs Part I: only-prism and only pyramid graphs. To appear in *Journal of Combinatorial Theory, Series B*. arXiv:1504.01862
- [8] M.R. Fellows, J. Kratochvíl, M. Middendorf, F. Pfeiffer. The complexity of induced minors and related problems. *Algorithmica*, 13(3):266–282, 1995.
- [9] J. Fiala, M. Kamiński, B. Lidický, D. Paulusma. The k -in-a-Path Problem for Claw-free Graphs. *Algorithmica*, 62: 499-519, 2012.
- [10] F.V. Fomin and P. Kaski, editors. *Algorithm Theory - SWAT 2012 - 13th Scandinavian Symposium and Workshops, Helsinki, Finland, July 4-6, 2012. Proceedings*, volume 7357 of *Lecture Notes in Computer Science*. Springer, 2012.
- [11] M.R. Garey, D.S. Johnson. *Computers and Intractability*. Freeman, New York ,1979.
- [12] P.A. Golovach, D. Paulusma, and E.J. van Leeuwen. Induced disjoint paths in AT-free graphs. In Fomin and Kaski [10], pages 153–164.
- [13] P.A. Golovach, D. Paulusma, E.J. van Leeuwen. Induced disjoint paths in claw-free graphs. *SIAM Journal of Discrete Mathematics*, 29 (1): 348-375, 2015.
- [14] P.A. Golovach, D. Paulusma, and E.J. van Leeuwen. Induced disjoint paths in circular-arc graphs in linear time. *Theor. Comput. Sci.*, 640:70–83, 2016.
- [15] R.M. Karp. On the complexity of combinatorial problems. *Networks*, 5: 45-68, 1975.
- [16] K. Kawarabayashi and Y. Kobayashi. A linear time algorithm for the induced disjoint paths problem in planar graphs. *J. Comput. Syst. Sci.*, 78(2):670–680, 2012.
- [17] P.G.H. Lehot. An optimal algorithm to detect a line graph and output its root graph. *Journal of the Association for Computing Machinery*, 21(4):569-575, 1974.
- [18] B. Lévêque, D. Lin, F. Maffray, and N. Trotignon. Detecting induced subgraphs. *Discrete Applied Mathematics*, 157:3540–3551, 2009.
- [19] W. Liu and N. Trotignon. The k -in-a-tree problem for graphs of girth at least k . *Discrete Applied Mathematics*, 158:1644–1649, 2010.
- [20] M. Radovanović, N. Trotignon, K. Vušković. The (theta,wheel)-free graphs Part II: structure theorem. To appear in *Journal of Combinatorial Theory, Series B*. arXiv:1703.08675
- [21] M. Radovanović, N. Trotignon, K. Vušković. The (theta,wheel)-free graphs Part III: cliques, stable sets and coloring. To appear in *Journal of Combinatorial Theory, Series B*. arXiv:1707.04205

- [22] N. Robertson and P.D. Seymour. Graph minors. XIII. The disjoint paths problem. *Journal of Combinatorial Theory, Series B*, 63: 65-110, 1995.
- [23] N.D. Roussopoulos. A $\max\{m, n\}$ algorithm for determining the graph H from its line graph G . *Information Processing Letters*, 2(4):108-112, 1973.
- [24] R.E. Tarjan. Decomposition by clique separators. *Discrete Mathematics*, 55: 221-232, 1985.
- [25] N. Trotignon, K. Vušković. Combinatorial optimization with 2-joins. *Journal of Combinatorial Theory, Series B*, 102:153-185, 2012.