

Congresso Nacional das Escolas Superiores Agrárias

14 e 15 de novembro de 2019

Escola Superior Agrária de Viseu | IPV



Livro de Resumos



Congresso Nacional das Escolas Superiores Agrárias 14 e 15 de novembro de 2019

Escola Superior Agrária de Viseu | IPV

## Ficha Técnica

Título: Livro de resumos do III Congresso Nacional das Escolas Superiors Agrárias Editores: Comissão organizadora do III Congresso Nacional das Escolas Superiores Agrárias

Data: 14 e 15 de novembro de 2019

Local: Instituto Politécnico de Viseu





Escola Superior Agrária de Viseu | IPV

## LIFE CYCLE ENVIRONMENTAL IMPACTS OF VINEYARD IN THE SOUTH PORTUGAL

ELLEN SILVA<sup>1</sup>, PEDRO PRESUMIDO<sup>2</sup>, ARTUR GONÇALVES<sup>2</sup>, AULUS BINELI<sup>3</sup>, ARTUR SARAIVA<sup>4</sup>, PEDRO OLIVEIRA E SILVA<sup>5</sup>, SOFIA RAMÔA<sup>5</sup> MARGARIDA OLIVEIRA<sup>4</sup>, MANUEL FELICIANO<sup>2</sup>

<sup>1</sup> Instituto Politécnico de Bragança (IPB), Bragança, Portugal.

<sup>2</sup> Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Bragança, Portugal.

<sup>3</sup> Department of Environmental Engineering, Universidade Tecnológica Federal do Paraná, Londrina, Brasil.

<sup>4</sup>ESAS, UIIPS, Instituto Politécnico Santarém, Santarém, Portugal.

<sup>5</sup> ESAB, Instituto Politécnico Beja, Beja, Portugal.

Abstract: Agricultural activities have multiple negative impacts on the environment. In the wine sector, several studies have used the lifecycle approach to identify and measure these environmental impacts. The main aim of this study was to determine the environmental impact of two vineyards located in southern Portugal, considering 1 kg of grape yield, using the life cycle assessment (LCA) methodology (ISO 14040). The system boundary studied was "from cradle to gate" including grape production and harvesting. The methodology included the use of GaBi software for the assessment of different impact categories (CML 2001 Methods), including the global warming potential (GWP), acidification potential (AP), eutrophication potential (EP) and abiotic depletion potential (AD). The results show that the contribution in the AP and EP categories were associated with the use of agricultural machinery, while the AD was mostly due to external inputs, namely the production of pesticides. On the other hand, the GWP category, received a very equitable contribution among all production processes. This study allowed for the identification of which production processes may contribute the most to the different environmental impacts, thus prompting the identification of improvement opportunities for more sustainable production in the vineyards stages of the wine production.

Keywords: Vineyard; Life cycle assessment (LCA); Environmental impacts

Acknowledgements: This study received funding from COMPETE 2020 - Programa Operacional Competitividade e Internacionalização (POCI-0145-FEDER-023360), Wine WATERFootprint project. The authors would also like to thank the Foundation for Science and Technology (FCT, Portugal) and FEDER under the PT2020 Program for the financial support to CIMO [UID / AGR / 00690/2019]. Our thanks must also go to the farmers, companies and specialists in the wine sector for their kind collaboration in this study, through the provision of relevant data and information.

