



DR. VÍTOR RAMOS (Orcid ID : 0000-0002-5824-3633)

DR. PEDRO LEÃO (Orcid ID : 0000-0001-5064-9164)

Article type : Regular Article

The extremophile *Endolithella mcmurdoensis* gen. et sp. nov. (Trebouxiophyceae, Chlorellaceae), a new *Chlorella*-like endolithic alga from Antarctica¹

*Teresa P. Martins,[#] Vitor Ramos,^{#,§} Guilherme S. Hentschke,[#] Raquel Castelo-Branco,
Adriana Rego*

Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Terminal de
Cruzeiros do Porto de Leixões, 4450-208 Matosinhos, Portugal

Maria Monteiro

Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Terminal de
Cruzeiros do Porto de Leixões, 4450-208 Matosinhos, Portugal

School of Science, University of Waikato, Hamilton, New Zealand

Ângela Brito

Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Rua Alfredo Allen 208, 4200-
135 Porto, Portugal

Instituto de Biologia Molecular e Celular, Universidade do Porto, Rua Alfredo Allen 208, 4200-135
Porto, Portugal

Paula Tamagnini

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JPY.12940-19-090](https://doi.org/10.1111/JPY.12940-19-090)

This article is protected by copyright. All rights reserved

Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Rua Alfredo Allen 208, 4200-135 Porto, Portugal

Instituto de Biologia Molecular e Celular, Universidade do Porto, Rua Alfredo Allen 208, 4200-135 Porto, Portugal

Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal
S. Craig Cary

International Centre for Terrestrial Antarctic Research, University of Waikato, Hamilton, New Zealand

School of Science, University of Waikato, Hamilton, New Zealand
Vitor Vasconcelos

Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Terminal de Cruzeiros do Porto de Leixões, 4450-208 Matosinhos, Portugal

Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal
Lothar Krienitz

Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Department of Experimental Limnology, D-16775 Stechlin, Germany
Catarina Magalhães

Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Terminal de Cruzeiros do Porto de Leixões, 4450-208 Matosinhos, Portugal

Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal
Pedro N. Leão²

Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Terminal de Cruzeiros do Porto de Leixões, 4450-208 Matosinhos, Portugal

¹Received

#these authors contributed equally to this work.

§Present address: Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

²Author for correspondence: e-mail pleao@ciimar.up.pt, (+351) 22 340 18 00

Running title: Endolithella mcmurdoensis gen. et sp. nov.

Editorial Responsibility: H. Verbruggen (Associate Editor)

Accepted Article

Abstract

The McMurdo Dry Valleys constitute the largest ice-free region of Antarctica and one of the most extreme deserts on Earth. Despite the low temperatures, dry and poor soils and katabatic winds, some microbes are able to take advantage of endolithic microenvironments, inhabiting the pore spaces of soil and constituting photosynthesis-based communities. We isolated a green microalga, *Endolithella mcmurdoensis* gen. et sp. nov from an endolithic sandstone sample collected in the McMurdo Dry Valleys (Victoria Land, East Antarctica) during the K020 expedition, in January 2013. The single non-axenic isolate (*E. mcmurdoensis* LEGE Z-009) exhibits cup-shaped chloroplasts, electron-dense bodies and polyphosphate granules but our analysis did not reveal any diagnostic morphological characters. On the basis of phylogenetic analysis of the 18S rRNA (SSU) gene, the isolate was found to represent a new genus within the family Chlorellaceae.

Key index words: 18S rRNA, extremophiles, LEGEcc, genus, microalgae, sandstone

Abbreviations: BI, bayesian inference; DAPI, 4',6-diamidino-2-phenylindole; LEGEcc, Blue Biotechnology and Ecotoxicology culture collection; OTU, operational taxonomic unit; QIIME, Quantitative insights into microbial ecology

Introduction

The remarkable diversity of green algae reflects a great adaptive capacity, allowing for survival in the most extreme environments on Earth such as marine deep waters (Zechman et al. 2010, Hodač et al. 2016, Spalding et al. 2016), deserts (Friedmann and Ocampo 1976, Lewis and Lewis 2005), hypersaline habitats (Vinogradova and Darienko 2008) or habitats with temperatures below freezing point (Morgan-Kiss et al. 2008). Even in these extreme settings, green microalgae play an important ecological role (Lyon and Mock 2014), but their diversity is far from being understood, especially in unexplored polar environments (Hodač et al. 2016). The life-limiting environmental conditions that occur in Antarctica are triggers for speciation and development of endemic species, making this a particularly promising location for the exploration of microbial diversity (Vincent 2000).

In Antarctica, trebouxiophycean microalgae are known since the description of *Chlorella antarctica* (Wille 1928), and many studies have highlighted the considerable diversity of this group in cold environments (Friedmann and Ocampo 1976, Friedmann 1982, Carvalho et al. 2017, Nedbalová et al. 2017). The McMurdo Dry Valleys constitute the largest ice-free region of Antarctica (Horowitz et al. 1972) and one of the world's most extreme deserts. The average annual temperature is about -22°C, winter temperatures typically of -60°C, precipitation levels are lower than 100 mm water equivalent, the land is battered by katabatic winds and intense UV radiation and the soils are poor, with low levels of carbon, nitrogen and small amounts of clay (Wynnwilliams 1990, Ugolini and Bockheim 2008, Cary et al. 2010). Despite all these extreme conditions, life exists. Soil and rocks act as refuge, protecting life from intense solar radiation and desiccation providing milder microclimate conditions that ease the growth of microbial life (Pointing et al. 2009) with significant diversity (Wood et al. 2008). In fact, some microbes are able to inhabit porous rocks such as sandstones (Friedmann 1982, Hughes and Lawley 2003), constituting photosynthesis-based endolithic communities (Friedmann and Ocampo 1976) that rely heavily on cyanobacteria and microalgae. In Antarctica land environments, green algae, especially *Chlorella*-like forms, are particularly abundant and diverse (Hodač et al. 2016) and have an important role in nutrient cycling and as primary producers, providing organic matter in the oligotrophic soils (Tscherko et al. 2003).

Herein, we describe the new trebouxiophycean microalga *Endolithella mcmurdoensis* gen. et sp. nov, isolated from an endolithic habitat in the McMurdo Dry Valleys, Antarctica, that was found by pyrosequencing analysis to be dominated by cyanobacteria.

Methods

Sampling, isolation and culturing.

Endolithic samples (Fig. 1, A and B) were collected during the K020 Mission, that took place in January 2013, on Victoria Valley, located in McMurdo Dry Valleys region, in South Victoria Land, Antarctica (77°20.502 S; 161°39.306 E). Collected samples were preserved at -80°C in LifeGuard Solution (MoBio). Isolate LEGE Z-009 was obtained from an isolation process originally intended for isolating cyanobacteria (Rego et al. 2019). An aliquot of the endolithic sample was washed and centrifuged three times (4500g, 3 min) with nitrogen-free BG11₀ medium (Rippka 1988). It was then inoculated into a 100 mL Erlenmeyer flask with liquid Z8 medium (Kotai 1972) as an initial enrichment culture and placed at 19°C, under a 12:12 h light (25 μmol photons · m⁻² · s⁻¹) : dark cycle. When visible growth was detected, an aliquot was transferred and streaked onto a solid BG11₀ medium plate (1.5% agar supplemented with 25 μL · L⁻¹ of cycloheximide), and placed at 12°C. Single colonies were then microscopically inspected, selected and re-streaked aseptically in the same medium and/or in Z8, until isolation was achieved. The isolated strain was deposited in the LEGE Culture Collection in CIIMAR, and is maintained cryopreserved and at 12°C and 19°C, in liquid Z8 medium, under the identification LEGE Z-009.

Pyrosequencing (16S rRNA-based) analysis of the endolithic community.

Environmental DNA (eDNA) was extracted from the same Victoria Valley sample used for isolation of the microalga strain, by using a modification of the CTAB extraction protocol (Barrett et al. 2006). The 16S small subunit (SSU) rRNA gene was initially amplified by PCR using the universal primer pair 27F/1492R (Weisburg et al. 1991) and then sequenced by pyrosequencing technology. Briefly, the 16S rRNA gene was amplified for the V3/V4 hypervariable region with barcoded fusion primers containing the Roche-454 A and B Titanium sequencing adapters, an eight-base barcode sequence, the forward (5'– ACTCCTACGGGAGGCAG-3') and reverse (5'– TACNVRRGTHHTCTAATYC -3')

primers (Wang and Qian 2009). Amplicons were quantified by fluorimetry with PicoGreen (Invitrogen), pooled at equimolar concentrations and sequenced in the A direction with GS 454 FLX Titanium chemistry, according to the manufacturer's instructions (Roche, 454 Life Sciences) at Biocant (Cantanhede, Portugal). The 454-machine-generated FASTA (.fna) and quality score (.qual) files were processed using the QIIME (Quantitative insights into microbial ecology) pipeline (Caporaso et al. 2010). Initially, raw reads were demultiplexed and subjected to a quality filtering – sequences with a quality score below 25 were removed. The next step, Pick OTUs (Operational Taxonomic Units; Sneath and Sokal 1973) was performed using the pick_open_reference_otus.py workflow (open-reference method; i.e., unmatched sequences are clustered de novo and included in the downstream analysis). No action was undertaken to remove eventual plastid-derived sequences from the dataset. The OTU table obtained from the open-reference method was selected to the downstream analyses. Essentially, all sequences (average length 428 bp) were clustered into OTUs at 97% sequence similarity using UCLUST (Edgar 2010) and the reads aligned to the Greengenes v13_8 (GG; DeSantis et al. 2006) database using PyNAST. For the taxonomic assignment, the RDP Classifier 2.2 (Wang et al. 2007) was used with the UCLUST method. Sequencing reads from this study were deposited in the European Nucleotide Archive (ENA) under the accession number PRJEB34393.

Microscopy.

The isolated strain was observed and photographed under a Leica DMLB microscope. Measurements of 40 cells from a growing-phase culture were performed using a dedicated software (Leica LAS EZ, Leica Microsystems). For TEM studies, BG11- and N-depleted BG11₀-grown microalgal cultures were centrifuged and the cells were fixed in 2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M sodium cacodylate buffer (pH 7.2) for at least two h, followed by a buffer wash (5 min) and overnight incubation with osmium tetroxide (2% w/v). A new buffer wash was performed, and samples were incubated overnight with 1% uranyl acetate (w/v). Dehydration was performed using an ethanol series (25-100%; v/v), and twice propylene oxide (10 min each). Subsequently, samples were embedded in mixtures of propylene oxide and Epon resin, followed by Epon for at least 24 h, before being placed in embedding moulds with Epon, and allowed to polymerize at 65°C for 48 h. Ultrathin sections were

cut with an ultramicrotome Leica Reichert SuperNova, placed on electron microscopy grids, and visualized using an electron microscope (JEOL JEM-1400Plus).

Polyphosphate granules and DNA were visualized with DAPI staining according to previous standardized protocol (Mukherjee and Ray 2015). Preparations were visualized under Laser Scanning Confocal Microscope Leica SP2 AOBS SE (Leica Microsystems) using the following wavelength channels: 430-460 nm for DNA staining, 520-530 nm for polyphosphate granules and 640-680 nm for chlorophyll epifluorescence. Images were merged using Adobe Photoshop software.

DNA extraction, PCR amplification and cloning.

Genomic DNA extraction was performed with the DNeasy® Plant Mini Kit (QIAGEN) following the manufacturer's instructions for plant DNA extraction. Plastid-encoded 16S small subunit (SSU) rDNA was amplified by PCR using the primers CYA106F, CYA359F, CYA781R (Nubel et al. 1997) and 1492R (Weisburg et al. 1991). Nuclear-encoded 18S SSU rDNA amplification was performed using the internal primers 18S402F, 18S895F, 18S919R, 18S1339R (Katana et al. 2001) and the external 18SF and 18SR primers (Moon-van der Staay et al. 2000). The nuclear internal transcribed spacer region (ITS1-5.8S-ITS2) was amplified using primers 18S895F and ITS055R (Marin et al. 2003). All PCRs were performed according with the conditions originally reported for the used oligonucleotides. PCR products were separated in a 1.5% (w/v) agarose gel stained with GelRed™ (Biotium) and DNA fragments with the expected size were excised from the gel and purified using the NucleoSpin™ Gel and PCR Clean-up Kit (Macherey-Nagel). The purified 16S rDNA PCR product was sequenced (GATC Biotech) with CYA106F/1492R primers while purified 18S rDNA and ITS PCR products were TA cloned using a pGEM®-T Easy Vector System Kit (Promega), and transformed into *Escherichia coli* ONE SHOT® TOP10 chemically competent cells (Invitrogen). Colonies were selected by blue-white screening, and the presence of the appropriate insert was evaluated by colony PCR. Colonies with an insert of the expected size were grown overnight at 37°C in liquid LB medium supplemented with 100 µg · mL⁻¹ of ampicillin, with shaking. Plasmid DNA was isolated using the NZY Miniprep kit (NYZtech) and sequenced (GATC Biotech) using M13 primers. All nucleotide sequences were manually inspected for quality, trimmed and assembled using the Geneious 8.1.9. software package (Biomatters Limited). Sequences obtained in this study were

deposited in the NCBI GenBank database under accession numbers MN394916 (16S rRNA gene), MK682757 (18S rRNA gene) and MK682753 (ITS).

Phylogenetic analysis.

The 18S rRNA gene Maximum Likelihood (ML) and Bayesian inference (BI) trees presented here were constructed in two rounds. In the first round, to place our species within the Trebouxiophyceae tree, we performed a ML phylogeny with our sequence and with sequences of reference strains representing Chlorellales, Prasiolales and Trebouxiales. Then, we included sequences retrieved in GenBank by BLAST. In the second round, to obtain trees with higher phylogenetic support, we selected the phylogenetically most closely related taxa to *Endolithella mcmurdoensis* and performed ML and BI analysis. In the ITS2 phylogenies (ML and BI), we included the OTUs with available ITS2 sequences from our second round 18S rDNA phylogenies and sequences retrieved from GenBank by BLAST.

For all generated trees, sequences were aligned by ClustalW, and the GTR+G+I evolutionary model was selected by MEGA 7.0 (Kumar et al. 2016). The robustness of ML trees was estimated by bootstrap percentages, using 1000 replications in MEGA 7.0 (Kumar et al. 2016). Bayesian trees were constructed in two independent runs, with four chains each, for 5×10^5 generations using MrBayes (Ronquist and Huelsenbeck 2003) in Cipres Gateway (Miller et al. 2010). The outgroup used was *Oocystis marssonii* (AF228688).

Internal transcriber spacer 2 (ITS2) structures.

The ITS2 region of *Endolithella mcmurdoensis* MK682757 (280 bp) and the phylogenetically closest related sequences: *Auxenochlorella protothecoides* FN298931 (354 bp) and *A. protothecoides* LN610704 (354 bp) were annotated using ITS2 database (Koetschan et al. 2010) and folded with Mfold (Zuker 2003). Images were built using 4SALE (Seibel et al. 2006).

Results

Diversity in the endolithic community and isolation of Endolithella mcmurdoensis LEGE Z-009.

To obtain an overview of the microbial composition of the endolithic sample, 16S rRNA-based pyrosequencing analysis was performed and revealed that the endolithic community collected from the McMurdo Dry Valleys sample (Fig. 1, A and B) was dominated by Cyanobacteria (44%) and Actinobacteria (30%), followed by lower abundances of Chloroflexi (10%) and Proteobacteria (6%). The photosynthetic microbial community was clearly dominated by an *Acaryochloris*-like unicellular cyanobacterium representing 99% of the cyanobacterial/chloroplast OTUs present in the sample (Fig. S1 in the Supporting Information), which is in line with our own microscopic examination that revealed a clear dominance by coccoid cyanobacteria (Fig. S2 in the Supporting Information). These results are in accordance with other endolithic communities previously characterized (Walker et al. 2007), where oxygenic phototrophs, especially cyanobacteria, represented the major fraction of community, followed by Actinobacteria, Bacteroidetes, Chloroflexi, Firmicutes, Proteobacteria, and Crenarchaeota. In addition, the dominance of *Acaryochloris* suggested a high level of adaptation to this type of environment, which may result from its high chlorophyll d content (Partensky et al. 2018).

Although, from our pyrosequencing analysis, we found no evidence for chloroplast-derived 16S rRNA gene OTUs in the original endolithic sample, the strain *Endolithella mcmurdoensis* LEGE Z-009 was isolated from the sample, following repeated streaking in agar plates under laboratory conditions.

Morphology.

Endolithella mcmurdoensis was not detected by microscopic observation of the endolithic community sample; therefore, morphological analysis and taxon description were performed based only on the cultured isolate strain *E. mcmurdoensis* LEGE Z-009.

Description. *Endolithella* T.Martins, V.Ramos, Hentschke & Krienitz gen. nov.

Cells living in endolithic layers, solitary, ovoid, uninucleate. Cell wall two-layered. Asexual reproduction by autospores. The genus differs from other members of Chlorellaceae by its extreme habitat in the polar region, endolithic life style, 18S rRNA gene and ITS2 secondary structure. TYPE SPECIES: *Endolithella mcmurdoensis* T.Martins, V.Ramos, Hentschke & Krienitz.

ETYMOLOGY: The generic name reflects the endolithic habitat.

Endolithella mcmurdoensis T. Martins, V. Ramos, Hentschke & Krienitz sp. nov.

Figure 1, C-K

Individuals solitary, sometimes forming small chains with two to four cells. Mature vegetative cells spherical or sub-spherical, 3.2–5.2 μm in diameter (average: 4.42 μm), with double-layered cell wall covered by mucilage (Fig. 1C). Chloroplast cup-shaped, without pyrenoids (Fig. 1, D-F). From TEM analysis (Fig. 1, D-F), electron-dense bodies and polyphosphate granules present, the latter confirmed by fluorescence microscopy (Fig. 1, G-K). Reproduction by two (rarely) or four ovoid autospores with 2.0-3.8 μm in diameter (average: 3.1 μm). Sporangia spherical or sub-spherical, 4.4-6.6 μm in diameter (average: 5.3 μm).

Comments: There is no diagnostic morphological character in *Endolithella*. The taxonomic identification is only possible considering the typical extreme habitat (endolithic in Antarctica) and the molecular data presented below.

HOLOTYPE: authentic strain permanently cryopreserved, *Endolithella mcmurdoensis* LEGE Z-009, deposited in the Blue Biotechnology and Ecotoxicology Culture Collection (LEGEcc) at CIIMAR, Porto, Portugal ([http:// lege.ciimar.up.pt](http://lege.ciimar.up.pt)), collected 01/23/2013, leg. Catarina Magalhães. The strain has been deposited in the same collection as an active culture.

ISOTYPE: an air-dried herbarium specimen of the authentic strain LEGE Z-009 deposited at the Botanical Museum Berlin-Dahlem under the designation B 40 0043196.

TYPE LOCALITY: Antarctica, South Victoria Land, Victoria Valley, located in McMurdo Dry Valleys region.

HABITAT: Endolithic.

ETYMOLOGY: The specific epithet refers to the geographic origin of the type strain.

Phylogenetic position.

The nuclear-encoded 18S rRNA gene (1796 bp) and the ITS2 were used for the molecular characterization of *Endolithella mcmurdoensis* LEGE Z-009. Blast searches against the GenBank database indicated best identity scores for *Pseudochlorella pringsheimii* KY364701 (95.89%) and *Nannochloris* AS 2-10 AY195968 (95.84%). Both sequences were included in the first round trees, but proved to be not phylogenetically closely related to *E. mcmurdoensis*, reinforcing supporting the description of the new genus.

An initial 18S rRNA gene phylogenetic tree (Fig. S3 in the Supporting Information), with a total of 87 OTUs and 1487 nucleotide analyzed positions and containing genera of Trebouxiales, Chlorellales and Prasiolales, was constructed. In this tree, *Endolithella mcmurdoensis* LEGE Z-009 is positioned within the family of Chlorellaceae, weakly supported at the base of a large clade containing (sequentially) the *Nannochloris* (bootstrap value 62), *Parachlorella* (bootstrap value 99) and *Chlorella* (bootstrap value 49) clades. Basal to this group (*E. mcmurdoensis* included) are the *Auxenochlorella* clade, as well as genera of Prasiolales and Trebouxiales. *Endolithella mcmurdoensis* LEGE Z-009 is positioned outside any of the already known clades of Chlorellaceae.

In subsequently constructed trees (Fig. 2, Fig. S4 in the Supporting Information), only the most phylogenetically closely-related taxa to *Endolithella mcmurdoensis* were included in the analysis. Again, the new taxon proved not to be phylogenetically closely related to any currently known clades of Trebouxiophyceae. In the ML tree, the new genus is positioned basal to *Nannochloris* (bootstrap value 99), *Parachlorella* (bootstrap value 100) and *Chlorella* (bootstrap value 100) clades, as also observed for the first round tree, but now with stronger phylogenetic support (Fig. S4). In the BI tree (Fig. 2), *E. mcmurdoensis* LEGE Z-009 is basal to the *Auxenochlorella*-clade and again, not phylogenetically related to the other clades of Trebouxiophyceae. The average standard deviation of split frequencies for this analysis was 0.009191 and the average potential scale reduction factor (PSRF) for parameter values was 0.998. In all first and second round trees, all genera present strong phylogenetic support. The ITS2 ML and BI phylogenies corroborate with these results showing *Endolithella* as a weak supported sister clade of *Auxenochlorella* (Fig. S5 in the Supporting Information).

ITS2 secondary structure.

The ITS2 secondary structures (Figs. 3, S6 and S7 in the Supporting Information) are consistent with the usual structures for eukaryotes with four helices, a pyrimidine-pyrimidine mismatch in helix II, an A-rich region between helices II and III and a TGGT motif 5' in helix III, the longest one (Schultz et al. 2005). Comparisons between *Endolithella mcmurdoensis* LEGE Z-009 ITS2 secondary structure (280 bp, from MK682757) with the closest phylogenetic sequences showed that *Endolithella* presents a unique ITS2, differing in length and structure from *Auxenochlorella protothecoides* CCAP 211/7A (354 bp, from FN298931) and *A. protothecoides* CCAP 211/17 (354 bp, from LN610704). This analysis corroborates our 18S rDNA and ITS2 phylogenies in supporting *Endolithella* as a new genus.

Discussion

We described the new chlorellacean genus and species *Endolithella mcmurdoensis* based on 18S rRNA gene and ITS2 data from the strain LEGE Z-009. The isolate was obtained after repeated streaking in solid media, despite chloroplast sequences being absent from the pyrosequencing dataset, underpinning the importance of culture-dependent approaches to retrieve non-abundant biodiversity (Rego et al. 2019). From the morphological description presented here, *E. mcmurdoensis* LEGE Z-009 can be related to *Auxenochlorella*, *Pumilliosphaera* and *Meyerella* by the absence of a pyrenoid. This character distinguished *E. mcmurdoensis* from *Chlorella*-like pyrenoid-bearing genera (Gaysina et al. 2013, Neustupa et al. 2013). *Endolithella mcmurdoensis* LEGE Z-009 presents electron-dense bodies and polyphosphate granules that play important roles as environmental stress adaptation structures. Electron-dense bodies in green algae, previously documented in *Mychonastes homosphaera* from lake Kinneret (Hanagata et al. 1999), are related to metabolite deposition, working usually as poly-P accumulation structures (Ota and Kawano 2018). For instance, under early sulfur-depletion conditions, poly-P is accumulated in electron-dense bodies (Ota et al. 2016b), being hydrolyzed to Pi in a later stage, and released to the cytosol, acting as a protein-protective chaperone (Gray et al. 2014, Gray and Jakob 2015). The accumulation of polyphosphate and other nutrients such as lipids is known to be considerably high in members of Chlorellaceae (Ota et al. 2016a), working as survival strategy in harsh, adverse conditions, such as strong UV radiation and temperature

fluctuations (Achbergerova and Nahalka 2011), but also under N deprivation (Chu et al. 2013), which is precisely the condition under which this alga was isolated.

Despite the lack of a diagnostic morphological character, our phylogenies and ITS2 secondary structure data confirm that *Endolithella mcmurdoensis* LEGE Z-009 is a new taxon, with an uncertain position among the already known clades of Chlorellaceae. Comparing the topologies of the BI and ML trees (Fig. 2 and Fig. S4, respectively), *E. mcmurdoensis* LEGE Z-009 is basal to *Chlorella*, *Parachlorella* and *Nannochloris* clades in the ML trees, but in the BI tree the new taxon is more closely related to the extremophile *Auxenochlorella* clade. In both trees, the phylogenetic relationships between *E. mcmurdoensis* LEGE Z-009 and the other clades are only weakly supported but it was possible to infer that *Endolithella* is a distinct genus. We hypothesize that *E. mcmurdoensis* LEGE Z-009 could be a new lineage in the Chlorellaceae family, a proposition that has to be confirmed by further studies.

Herein we are presenting only the BI tree (which includes ML bootstrap values), because this tree shows the close relationship between *Endolithella mcmurdoensis* LEGE Z-009 and the extremophile clade of *Auxenochlorella* and *Prototheca*. The last two genera were described from sap of trees samples and were characterized as being dependent of exogenous organic compounds (Ueno et al. 2005, Darienko and Pröschold 2015). *Endolithella mcmurdoensis* LEGE Z-009 is also an extremophile taxon, living in Antarctica cold desert sandstones.

The potential of polar environments as collection sites for new microalgae taxa discoveries begins with the description of *Chlorella antarctica* in 1924 (Wille 1928). However, strains isolated from these environments were considered to be “non indigenous”, growing from spores that reached polar regions by wind dispersal. This view lasted until Friedmann and Ocampo (1976) reported endolithic microalgae growing in Antarctica. This was followed by the discovery of many microalgae species (including Cyanobacteria) in polar regions (Nedbalová et al. 2017, Davey et al. 2019). Our endolithic sample, presenting a conspicuous thick layer of different photosynthetic microorganisms inside a porous rock, is in line with an “indigenous” origin. Still, recent studies (Hodač et al. 2016) show that, within the same microalgae species, specimens from polar regions can show no Compensatory Base Changes in ITS2 secondary structures when compared to specimens from

temperate zones. In light of these findings, and despite the endolithic character of the LEGE Z-009 isolate, we cannot fully attest to the Antarctic endemic character of *Endolithella mcmurdoensis*.

The identification of this new genus represents an important taxonomic novelty and it is in line with recent studies that point towards a much more extensive and interesting biodiversity in Antarctica than previously thought (Chown et al. 2015).

Finally, we highlight the importance of studies on polar microalgae communities due to their potential in biotechnological applications like protection from UV radiation (Miao et al. 2002), energy production (Chen et al. 2012) or nanoparticle biosynthesis (Gallardo et al. 2014), considering their survival strategies and ability to grow and produce organic compounds in these extreme habitats with light, temperature and nutrient limitations.

Acknowledgments

We are sincerely grateful to Antarctica New Zealand for providing logistics support during K020 event. This campaign was conducted as part of the New Zealand Terrestrial Antarctic Biocomplexity Survey (nzTABS) through awards (UOWX0710, UOWX1401) from the New Zealand Foundation for Research and Technology (FRST), Antarctica New Zealand, and the New Zealand Ministry of Business, Innovation and Employment to SCC. The work was also supported by Portuguese Science and Technology Foundation (FCT) through grants UID/Multi/04423/2019 to CIIMAR, IF/01358/2014 and PTDC/MAR-BIO/2818/2012 to PNL, PTDC/CTA-AMB/30997/2017 to CM, and PhD scholarships to AR (SFRH/BD/140567/2018).

Conflicts of interest

The authors declare no conflicts of interest.

References

- Achbergerová, L. & Nahalka, J. 2011. Polyphosphate--an ancient energy source and active metabolic regulator. *Microb. Cell Fact.* 10:63.
- Ayesh, I., Sanders, M. D. & Friedmann, A. I. 1976. Retinitis pigmentosa and Coats's disease. *Br. J. Ophthalmol.* 60:775-7.
- Barrett, J. E., Virginia, R. A., Wall, D. H., Cary, S. C., Adams, B. J., Hacker, A. L. & Aislabie, J. M. 2006. Co-variation in soil biodiversity and biogeochemistry in northern and southern Victoria Land, Antarctica. *Antarct. Sci.* 18:535-48.
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Pena, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky, J. R., Tumbaugh, P. J., Walters, W. A., Widmann, J., Yatsunencko, T., Zaneveld, J. & Knight, R. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat. Meth.* 7:335-6.
- Carvalho, E. L., Maciel, L. F., Macedo, P. E., Dezordi, F. Z., Abreu, M. E. T., Victoria, F. C., Pereira, A. B., Boldo, J. T., Wallau, G. D. L. & Pinto, P. M. 2017. De novo assembly and annotation of the Antarctic alga *Prasiola crispa* transcriptome. *Front. Mol. Biosci.* 4:89.
- Cary, S. C., McDonald, I. R., Barrett, J. E. & Cowan, D. A. 2010. On the rocks: the microbiology of Antarctic Dry Valley soils. *Nat. Rev. Microbiol.* 8:129-38.
- Chen, Z., Gong, Y., Fang, X., Hu, H. 2012. *Scenedesmus* sp. NJ-1 isolated from Antarctica: a suitable renewable lipid source for biodiesel production. *World J. Microbiol. Biotechnol.* 28: 3219-25.
- Chu, F. F., Chu, P. N., Cai, P. J., Li, W. W., Lam, P. K. & Zeng, R. J. 2013. Phosphorus plays an important role in enhancing biodiesel productivity of *Chlorella vulgaris* under nitrogen deficiency. *Bioresour. Technol.* 134:341-6.
- Chown, S. L., Clarke, A., Fraser, C. I., Cary, S. C., Moon, K. L. & McGeoch, M. A. 2015. The changing form of Antarctic biodiversity. *Nature* 522:427-34.
- Darienko, T. & Pröschold, T. 2015. Genetic variability and taxonomic revision of the genus *Auxenochlorella* (Shihira et Krauss) Kalina et Punčochářová (Trebouxiophyceae, Chlorophyta). *J. Phycol.* 51:394-400.

- Davey, M. P., Norman, L., Sterk, P., Huete-Ortega, M., Bunbury, F., Loh, B. K. W., Stockton, S., Peck, L. S., Convey, P., Newsham, K. K. & Smith, A. G. 2019. Snow algae communities in Antarctica: metabolic and taxonomic composition. *New Phytol.* 222: 1242–55
- DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T., Dalevi, D., Hu, P. & Andersen, G. L. 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl. Environ. Microbiol.* 72:5069-72.
- Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26:2460-1.
- Friedmann, E. I. 1982. Endolithic microorganisms in the antarctic cold desert. *Science* 215:1045-53.
- Friedmann, E. I. & Ocampo, R. 1976. Endolithic blue-green algae in the dry valleys: primary producers in the antarctic desert ecosystem. *Science* 193:1247-9.
- Gallardo, C., Monras, J. P., Plaza, D. O., Collao, B., Saona, L. A., Duran-Toro, V., Venegas, F. A., Soto, C., Ulloa, G., Vasquez, C. C., Bravo, D., Perez-Donoso, J. M. 2014. Low-temperature biosynthesis of fluorescent semiconductor nanoparticles (CdS) by oxidative stress resistant Antarctic bacteria. *J. Biotechnol.* 187: 108-15.
- Gaysina, L., Němcová, Y., Škaloud, P., Sevciková, T. & Eliáš, M. 2013. *Chloropyrula uraliensis* gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a new green coccoid alga with a unique ultrastructure, isolated from soil in South Urals. *J. Syst. Evol.* 51:476-84.
- Gray, M. J. & Jakob, U. 2015. Oxidative stress protection by polyphosphate - new roles for an old player. *Curr. Opin. Microbiol.* 24:1-6.
- Gray, M. J., Wholey, W. Y., Wagner, N. O., Cremers, C. M., Mueller-Schickert, A., Hock, N. T., Krieger, A. G., Smith, E. M., Bender, R. A., Bardwell, J. C. A. & Jakob, U. 2014. Polyphosphate is a primordial chaperone. *Molecular Cell* 53:689-99.
- Hanagata, N., Malinsky-Rushansky, N. & Dubinsky, Z. 1999. Eukaryotic picoplankton, *Mychonastes homosphaera* (Chlorophyceae, Chlorophyta), in Lake Kinneret, Israel. *Phycol. Res.* 47:263-69.
- Hodač, L., Hallmann, C., Spitzer, K., Elster, J., Fasshauer, F., Brinkmann, N., Lepka, D., Diwan, V. & Friedl, T. 2016. Widespread green algae *Chlorella* and *Stichococcus* exhibit polar-temperate and tropical-temperate biogeography. *Fems Microbiol. Ecol.* 92.

- Horowitz, N. H., Cameron, R. E. & Hubbard, J. S. 1972. Microbiology of the dry valleys of antarctica. *Science* 176:242-5.
- Hughes, K. A. & Lawley, B. 2003. A novel Antarctic microbial endolithic community within gypsum crusts. *Environ. Microbiol.* 5:555-65.
- Katana, A., Kwiatowski, J., Spalik, K., Zakrys, B., Szalacha, E. & Szymanska, H. 2001. Phylogenetic position of *Koliella* (Chlorophyta) as inferred from nuclear and chloroplast small subunit rDNA. *J. Phycol.* 37:443-51.
- Koetschan, C., Forster, F., Keller, A., Schleicher, T., Ruderisch, B., Schwarz, R., Muller, T., Wolf, M. & Schultz, J. 2010. The ITS2 Database III--sequences and structures for phylogeny. *Nucleic Acids Res.* 38:D275-9.
- Kotai, J. 1972. Instructions for Preparation of Modified Nutrient Solution Z8 for Algae. *Norwegian Institute for Water Research, Oslo, Blindern, Oslo* 11/69: 5 pp
- Kumar, S., Stecher, G. & Tamura, K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* 33:1870-4.
- Lewis, L. A. & Lewis, P. O. 2005. Unearthing the molecular phylodiversity of desert soil green algae (Chlorophyta). *Systematic Biol.* 54:936-47.
- Lyon, B. R. & Mock, T. 2014. Polar microalgae: new approaches towards understanding adaptations to an extreme and changing environment. *Biology* 3:56-80.
- Marin, B., Palm, A., Klingberg, M. & Melkonian, M. 2003. Phylogeny and taxonomic revision of plastid-containing euglenophytes based on SSU rDNA sequence comparisons and synapomorphic signatures in the SSU rRNA secondary structure. *Protist* 154:99-145.
- Miao, J., Miao, J., Li, G., Hou, X., Zhang, Y., Jiang, Y., Wang, B., Zhang, B. 2002. Study on induced synthesis of anti-UV substances in the Antarctic algae. *High Technol. Lett.* 6: 179-83.
- Miller, M. A., Pfeiffer, W. & Schwartz, T. 2010. Creating the CIPRES science gateway for inference of large phylogenetic trees. *In Proceedings of the Gateway Computing Environments Workshop (GCE), 2010*, New Orleans, November 14, 2010.
- Moon-van der Staay, S. Y., van der Staay, G. W. M., Guillou, L., Vaultot, D., Claustre, H. & Medlin, L. K. 2000. Abundance and diversity of prymnesiophytes in the picoplankton community from

the equatorial Pacific Ocean inferred from 18S rDNA sequences. *Limnol. Oceanogr.* 45:98-109.

Morgan-Kiss, R. M., Ivanov, A. G., Modla, S., Czymmek, K., Huner, N. P., Priscu, J. C., Lisle, J. T. & Hanson, T. E. 2008. Identity and physiology of a new psychrophilic eukaryotic green alga, *Chlorella* sp., strain BI, isolated from a transitory pond near Bratina Island, Antarctica. *Extremophiles* 12:701-11.

Mukherjee, C. & Ray, K. 2015. An improved DAPI staining procedure for visualization of polyphosphate granules in cyanobacterial and microlagal cells. 7 July 2015, protocol (Version 1) available at Protocol Exchange (<https://doi.org/10.1038/protex.2015.066>)

Nedbalová, L., Mihál, M., Kvéřerová, J., Procházková, L., Řezanka, T. & Elster, J. 2017. Identity, ecology and ecophysiology of planktic green algae dominating in ice-covered lakes on James Ross Island (northeastern Antarctic Peninsula). *Extremophiles* 21:187-200.

Neustupa, J., Němcová, Y., Vesela, J., Steinová, J. & Škaloud, P. 2013. *Leptochlorella corticola* gen. et sp. Nov. and *Kalinella apyrenoidosa* sp. nov.: two novel *Chlorella*-like green microalgae (Trebouxiophyceae, Chlorophyta) from subaerial habitats. *Int. J. Syst. Evol. Micr.* 63:377-87.

Nubel U, M. G., Garcia-pichel F, Muyzer G 1997. PCR primers to amplify 16S rRNA genes from cyanobacteria PCR primers to amplify 16S rRNA genes from cyanobacteria. *Microbiology* 63: 3327-32.

Ota, S. & Kawano, S. 2018. Three-dimensional ultrastructure and hyperspectral imaging of metabolite accumulation and dynamics in *Haematococcus* and *Chlorella*. *Microscopy* 68:57-68.

Ota, S., Oshima, K., Yamazaki, T., Kim, S., Yu, Z., Yoshihara, M., Takeda, K., Takeshita, T., Hirata, A., Bisova, K., Zachleder, V., Hattori, M. & Kawano, S. 2016a. Highly efficient lipid production in the green alga *Parachlorella kessleri*: draft genome and transcriptome endorsed by whole-cell 3D ultrastructure. *Biotechnol. Biofuels* 9:13-22.

Ota, S. H., Yoshihara, M., Yamazaki, T., Takeshita, T., Hirata, A., Konomi, M., Oshima, K., Hattori, M., Bisova, K., Zachleder, V. & Kawano, S. 2016b. Deciphering the relationship among phosphate dynamics, electron-dense body and lipid accumulation in the green alga *Parachlorella kessleri*. *Sci. Rep.* 6:25731.

- Partensky, F., Six, C., Ratin, M., Garczarek, L., Vaultot, D., Probert, I., Calteau, A., Gourvil, P., Marie, D., Grebert, T., Bouchier, C., Le Panse, S., Gachenot, M., Rodriguez, F., Garrido, J. L. 2018. A novel species of the marine cyanobacterium *Acaryochloris* with a unique pigment content and lifestyle. *Sci Rep* 8: 1–13.
- Pointing, S. B., Chan, Y. K., Lacap, D. C., Lau, M. C. Y., Jurgens, J. A. & Farrell, R. L. 2009. Highly specialized microbial diversity in hyper-arid polar desert. *Proc. Natl. Acad. Sci. USA* 106:19964-69.
- Rego, A., Raio, F., Martins, T., Ribeiro, H., Sousa, A., Silva, J., Baptista, M., Lee, C., Cary, C., Ramos, V., Carvalhos, F., Leão, P. & Magalhães, C. 2019. Actinobacteria and cyanobacteria diversity in terrestrial Antarctic microenvironments using culture-dependent and independent methods. *Front. Microbiol.* 10:1018.
- Rippka, R. 1988. Isolation and purification of cyanobacteria. *Methods Enzymol.* 167:3-27.
- Ronquist, F. & Huelsenbeck, J. P. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572-4.
- Schultz, J., Maisel, S., Gerlach, D., Muller, T. & Wolf, M. 2005. A common core of secondary structure of the internal transcribed spacer 2 (ITS2) throughout the Eukaryota. *RNA* 11:361-4.
- Seibel, P. N., Muller, T., Dandekar, T., Schultz, J. & Wolf, M. 2006. 4SALE--a tool for synchronous RNA sequence and secondary structure alignment and editing. *BMC Bioinformatics* 7:498.
- Sneath, P. H. A. & Sokal, R. R. 1973. *Numerical Taxonomy the principles and practice of numerical classification*. 1st edition. W H Freeman and Co., San Francisco, CA, 573 pp.
- Spalding, H. L., Conklin, K. Y., Smith, C. M., O'Kelly, C. J. & Sherwood, A. R. 2016. New Ulvaceae (Ulvophyceae, Chlorophyta) from mesophotic ecosystems across the Hawaiian Archipelago. *J. Phycol.* 52:40-53.
- Tscherko, D., Bolter, M., Beyer, L., Chen, J., Elster, J., Kandeler, E., Kuhn, D. & Blume, H. P. 2003. Biomass and enzyme activity of two soil transects at King George Island, maritime Antarctica. *Arct. Antarct. Alp. Res.* 35:34-47.
- Ueno, R., Hanagata, N., Urano, N. & Suzuki, M. 2005. Molecular phylogeny and phenotypic variation in the heterotrophic green algal genus *Prototheca* (Trebouxiophyceae, Chlorophyta). *J. Phycol.* 41:1268-80.

- Ugolini, F. C. & Bockheim, J. G. 2008. Antarctic soils and soil formation in a changing environment: A review. *Geoderma* 144:1-8.
- Vincent, W. F. 2000. Evolutionary origins of Antarctic microbiota: invasion, selection and endemism. *Antarct. Sci.* 12:374-85.
- Vinogradova, O. M. & Darienko, T. M. 2008. Terrestrial algae of hypersaline environments of the Central Syvash islands (Kherson Region, Ukraine). *Biologia* 63:813-23.
- Walker, J. J. & N. R. Pace 2007. Phylogenetic composition of Rocky Mountain endolithic microbial ecosystems. *Appl. Environ. Microbiol.* 73: 3497-3504
- Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* 73:5261-7.
- Wang, Y. & Qian, P. Y. 2009. Conservative fragments in bacterial 16S rRNA genes and primer design for 16S ribosomal DNA amplicons in metagenomic studies. *PLoS ONE* 4:e7401.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. 1991. 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.* 173:697-703.
- Wille, N. 1928. Süßwasseralgen von der deutschen Südpolar-Expedition auf dem Schiff "Gauss". In: Drygalski, E.V.E. [Ed.] *Deutsche Südpolar-Expedition 1901-1903*. G. Reimer, Berlin, pp. 373-444.
- Wood, S. A., Rueckert, A., Cowan, D. A. & Cary, S. C. 2008. Sources of edaphic cyanobacterial diversity in the Dry Valleys of Eastern Antarctica. *ISME J.* 2:308-20.
- Wynnwilliams, D. D. 1990. Ecological aspects of antarctic microbiology. *Adv. Microb. Ecol.* 11:71-146.
- Zechman, F. W., Verbruggen, H., Leliaert, F., Ashworth, M., Buchheim, M. A., Fawley, M. W., Spalding, H., Pueschel, C. M., Buchheim, J. A., Verghese, B. & Hanisak, M. D. 2010. An unrecognized ancient lineage of green plants persists in deep marine waters. *J. Phycol.* 46:1288-95.
- Zuker, M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.* 31: 3406–15.

Figure legends

Fig. 1 – A,B -Colonization of the cryptoendolith-bearing sandstone in McMurdo Dry Valleys. C - Light photomicrographs of the chlorophyte microalga LEGE Z-009 (1000x magnification). D-F – TEM photomicrographs of the chlorophyte microalga LEGE Z-009 culture: th, thylakoids; db, electron-dense body; cw, cell wall; en, cell envelope; n, nucleus, vg – vegetative cell, sp - sporangium. G-K – Fluorescence photomicrographs of *Endolithella mcmurdoensis* LEGE Z-009 with DAPI staining; chloroplast epifluorescence (G), DAPI blue-fluorescence emission upon DNA binding (H), DAPI green-fluorescence emission (1000× magnification) upon polyphosphate granules binding (I), merged G, H and I (J) and merged H and I (K).

Fig. 2 –18S rRNA gene Bayesian phylogenetic analysis. Posterior probabilities and Bootstrap values (BI/ML, respectively) are indicated at nodes. “–“ denotes no support.

Fig. 3 - ITS2 secondary structure of *Endolithella mcmurdoensis* LEGE Z-009 (280 bp, from MK682757) visualized with 4SALE. ITS2 helices are numbered I–IV and typical motifs are highlighted (arrows): the pyrimidine–pyrimidine mismatch in helix II; the A-rich region between helices II and III, as well as the UGGU motif 5' to the apex of helix III.

Figure S1 – (A) Taxonomy summary bar chart of relative frequency of 10 most abundant phyla in the initial sample, based on 16S rRNA gene sequences (B) Summary bar chart of the most abundant taxonomic frequency distributions at Genus level for the oxygenic photosynthetic community (both taxa are Cyanobacteria).

Figure S2 – Environmental sample photomicrographs. Bright field image (A) of the same optical field as in the fluorescence image (B).


Figure S3 –18S rRNA gene Maximum Likelihood phylogenetic analysis of trebouxyophycan strains. Bootstrap values are indicated at nodes. The analysis involved 87 nucleotide sequences and there were a total of 1487 positions in the final dataset.

Figure S4 – 18S rRNA gene Maximum Likelihood phylogenetic analysis of a selection of chlorellacean strains. Bootstrap values are indicated at nodes. The analysis involved 26 nucleotide sequences and there were a total of 1664 positions in the final dataset.

Figure S5 - ITS2 secondary structure of *Auxenochlorella protothecoides* FN298931, visualized with 4SALE.

Figure S6 - ITS2 secondary structure of *Auxenochlorella protothecoides* LN610704, visualized with 4SALE.

Figure S7 - ITS2 secondary structure of *Auxenochlorella protothecoides* LN610704, visualized with 4SALE.



jpy_12940-19-090_f1.tif

