



Fernandes, G. S., Hammerton, G. L., Abeysekera, K. W. M., Mahedy, L. L., A, E., Lewis, G., Hickman, M., & Heron, J. E. (2020). Alcohol consumption and internalising disorders in young adults of ALSPAC: a population-based study. *Journal of Epidemiology and Community Health*. <https://doi.org/10.1136/jech-2020-213922>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1136/jech-2020-213922](https://doi.org/10.1136/jech-2020-213922)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via BMJ at <https://jech.bmj.com/content/early/2020/07/05/jech-2020-213922>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research


General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



OPEN ACCESS

Alcohol consumption and internalising disorders in young adults of ALSPAC: a population-based study

Gwen Sascha Fernandes ¹, Gemma Lewis,² Gemma Hammerton,¹ Kushala Abeysekera,¹ Liam Mahedy,¹ Alexis Edwards,³ Glyn Lewis,² Matthew Hickman,⁴ Jonathan Heron⁴

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jech-2020-213922>).

¹Population Health Sciences, University of Bristol School of Social and Community Medicine, Bristol, UK

²UCL Division of Psychiatry, London, UK

³Virginia Institute for Psychiatric and Behavior Genetics, Richmond, Virginia, USA

⁴School of Social and Community Medicine, University of Bristol, Bristol, UK

Correspondence to

Gwen Sascha Fernandes, Population Health Sciences, Barley House, Oakfield Grove, Bristol BS8 2BN, UK; gwen.fernandes@bristol.ac.uk

Received 13 February 2020

Revised 8 April 2020

Accepted 2 June 2020

ABSTRACT

Introduction Depression and harmful alcohol consumption contribute significantly to the global health burden, but in young adults, this relationship is under-researched and conflicted. The aim of this study was to determine the sex-based prevalence and the association between internalising disorders such as depression and alcohol use disorders.

Method Using the Avon Longitudinal Study of Parents and Children, we assessed the sex-specific prevalence of International Classification of Diseases, Tenth Revision diagnosed generalised anxiety disorder (GAD), depression and fear-based anxieties (FBA) at 24 years (n=3572). We examined the association between internalising disorders and alcohol consumption using the Alcohol Use Disorder Identification Test for Consumption 5+ threshold and Diagnostic and Statistical Manual on Mental Disorders defined criteria for alcohol dependence.

Results Women reported more GAD (11.6% vs 6.5%), depression (13.4% vs 6.9%) and FBA (1.3% vs 0.5%) than men (p<0.001). Harmful drinking, after adjustment for sex and socioeconomic status, was associated with a higher prevalence of depression (OR 1.8, 95% CI 1.3 to 2.4, p<0.001), anxiety (OR 1.4, 95% CI 1.0 to 2.0, p<0.001) and FBA (OR 2.4, 95% CI 1.04 to 5.56, p=0.009) compared with lower-risk drinkers. In contrast, hazardous drinking was associated with a lower prevalence of GAD (OR 0.69, 95% CI 0.54 to 0.88) and depression (OR 0.68, 95% CI 0.54 to 0.86) compared with lower-risk drinkers.

Conclusions Young adults in the UK who drink harmfully are more likely to have depression and other internalising disorders. Further research should test whether there is a J-shaped relationship between alcohol consumption and mental health in young people and whether this varies across the life course.

disorders.^{5–7} This is due in part to alcohol consumption leading to depression, and individuals with depressive symptoms being more likely to consume alcohol frequently and in higher quantities—the ‘self-medication’ hypothesis.⁸ Some older adult, non-UK-based studies indicate that alcohol misuse and anxiety/depressive symptoms are extensively associated with each other, although the nature of the association is often inconsistent by gender or age groups.^{3 9 10} In the UK, results from the Whitehall II prospective cohort study found that mental health is the primary indicator of change in the relationship between mental health and alcohol consumption in middle-aged adults. Poor mental health predicted increasing alcohol consumption among lower-risk drinkers and was a maintaining factor for heavy alcohol consumption later along the life course.¹¹ The conflicting evidence has been attributed to small sample sizes; heterogenous sample characteristics (university vs general populations); variations in study designs; or varying assessments of alcohol use and severity of use (validated tools and thresholds).^{2 3 10}

There is a need for evidence on the relationship between internalising disorders and alcohol consumption from a large contemporary general population data set, specifically among young adults. Given this under-researched and controversial topic,^{12 13} our aim was to investigate the burden of drinking and internalising disorders in a young adult, community-based sample in the UK. The study will use data from the Avon Longitudinal Study of Parents and Children (ALSPAC) and first, aims to presents sex-based prevalences of internalising disorders such as anxiety and depression and alcohol use in 24-year-olds. Second, the study will test the association between these internalising disorders and alcohol consumption with adjustment for potential covariates such as sex and socioeconomic factors.

INTRODUCTION

Early adulthood (aged 18 years and over) is a developmental period marked by seminal life changes (eg, leaving home, finding a job, etc) and is considered a high-risk period for the onset of internalising disorders and/or alcohol-related problems.^{1–3} In the UK, excessive alcohol use remains the leading cause of premature mortality and can have negative consequences on mental health particularly an increased risk of depression and anxiety.⁴ Previous studies have shown consistent associations between alcohol use disorders (AUD) and major depressive

METHODS

Sample

Data were drawn from ALSPAC, the most phenotyped birth cohort in the world, and detailed information about our ALSPAC sample is presented as online supplementary material 1.

Of the 10 018 ALSPAC study participants who were invited for a clinical assessment, 40.1% (n=4021/10 018) attended for clinical assessment (denoted TF5 visit) (mean age of 24 years (22–26 years; SD 0.8; 1507 males, 2510 females¹⁴).



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

To cite: Fernandes GS, Lewis G, Hammerton G, et al. *J Epidemiol Community Health* Epub ahead of print: [please include Day Month Year]. doi: 10.1136/jech-2020-213922

Informed consent for use of data collected via questionnaires and clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law Committee at the time.

A flow chart of participants in this study is presented in online supplementary appendix 1.

MEASURES

Internalising disorders

Generalised anxiety disorder (GAD)

GAD was assessed using a self-administered online version of the Clinical Interview Schedule-Revised (CIS-R) based on the extensively used International Classification of Diseases, Tenth Revision (ICD-10) criteria^{15 16} and created a binary variable indicating presence or absence of GAD over the past 1-week period.

Depression

We assessed depression using the CIS-R and ICD-10 criteria based on the previous 1-month period.¹⁵ For this study, we used a binary variable indicating the presence of depression (merged mild, moderate and severe depression) or not over the past 1-week period.

Fear-Based Anxieties (FBA)

Given that social anxieties associate with harmful and hazardous alcohol consumption,^{17 18} we created a binary FBA variable by grouping three ICD-10 anxiety disorders from CIS-R: agoraphobia, social phobia and panic disorder over the past 1 week due to interrelated and overlapping cluster of phobic symptoms between them.

The use of this standardised version of the CIS-R within a community setting such as ALSPAC has been previously validated and demonstrated that lay assessors such as research nurses were as reliable as psychiatrists and did not exhibit bias in their use of CIS-R.¹⁹ The CIS-R assigns a total weighted score according to the frequency and severity of symptoms during the week immediately preceding the interview.

Alcohol measures

Within the ALSPAC sample, alcohol problems were assessed from responses to a computer-assisted standardised clinical interview. We used Alcohol Use Disorder Identification Test for Consumption (AUDIT-C) to assess hazardous drinking in the past 1 year.^{20 21} With the AUDIT-C data, we used a validated (≥ 5) cut-point to derive a binary measure as an indication of hazardous alcohol use.²² ALSPAC also collected data on alcohol use and alcohol dependence on administering using the Diagnostic and Statistical Manual on Mental Disorders—Fourth Edition (DSM-IV) tool (consisting of four items of alcohol abuse and seven items of alcohol dependence) and refers to drinking behaviour in the past 1 year. This measure of AUD comprises all individuals who indicate alcohol abuse and/or alcohol dependence, a maladaptive pattern of alcohol use leading to clinically significant impairment or distress and corresponds with harmful drinking behaviour that causes serious consequences physical and mental health.^{23 24}

The Office for National Statistics (2018) reports that almost half the population between the ages of 16–24 years consumed alcohol in the past week and of these, almost 30% of all drinkers exceeded the recommended alcohol units on the heaviest

drinking days. Given these findings, heavy drinking to a hazardous level in a young adult population²⁵ is not uncommon and needs to be separated from more harmful drinking patterns—indicating alcohol abuse or alcohol dependence. Therefore, we created a three-category alcohol phenotype (lower-risk drinkers, hazardous drinkers and harmful drinkers) by combining the AUD with (AUDIT-C). The following three drinking categories were generated:

1. AUDIT-C < 5 and no AUD=lower-risk drinkers,
2. AUDIT-C ≥ 5 + no AUD=hazardous drinkers and
3. AUDIT-C ≥ 5 + AUD-positive=harmful drinkers.

A sensitivity analysis was undertaken in which an alternative classification of alcohol use was created where all individuals with an 'extremely hazardous' AUDIT-C drinking score (10–12) were included in the harmful drinker category. The following three categories were used:

1. AUDIT-C < 5 and no AUD=lower-risk drinkers,
2. AUDIT-C ≥ 5 –9 + no AUD=hazardous drinkers and,
3. AUDIT-C ≥ 10 OR AUD-positive=harmful drinkers.

Statistical analysis

Prevalence of internalising disorders and alcohol use was presented by gender, and associations were tested using Pearson's χ^2 test (with associated p value). ORs and their corresponding 95% CIs were derived from logistic regressions with the lower-risk drinking group as the referent. We presented a crude model and models adjusted for sex (model 2) and for sex and socioeconomic status, parental income and maternal education (model 3) using imputed data. Further multiple imputation details are in online supplementary material 2. We used STATA 15.1 for all our analyses.

RESULTS

A total of 3572 participants attended for clinical assessment and provided complete data on alcohol use and internalising disorders. Table 1 shows that 9.7% of the cohort presented with GAD, 10.9% presented with depression and 1% with FBA. There were differences in GAD (11.6 vs 6.5%, $p < 0.001$), depression (13.4% vs 6.9%, $p < 0.001$) and FBA (1.3% vs 0.5%, $p = 0.02$) by sex, with women indicating more internalising disorders than men. Overall, for alcohol use, 379 participants (10.6%) met criteria for harmful drinking with women more likely to be lower-risk drinkers compared with men (44.8% vs 32.2%) while men were more

Table 1 The prevalence of generalised anxiety disorder (GAD), depression and fear-based anxieties and alcohol phenotypes by sex

24-year clinic (TF5) n=3572	Anxiety GAD (n, %)	Depression (n, %)	Fear-based anxieties* (n, %)	Alcohol phenotype** (n, %)		
				0	1	2
Overall	347 (9.7%)	391 (10.9%)	37 (1.0%)	1431 (40.1%)	1762 (49.3%)	379 (10.6%)
Female (n=2236)	260 (11.6%)	299 (13.4%)	30 (1.3%)	1001 (44.8%)	1044 (46.7%)	191 (8.5%)
Males (n=1336)	87 (6.5%)	92 (6.9%)	7 (0.5%)	430 (32.2%)	718 (53.7%)	188 (14.1%)
χ^2 , p value	25, $p < 0.001$	36.1, $p < 0.001$	5.5, $p = 0.02$	65.6, $p < 0.001$		

*Fear-based anxieties is an amalgamation of panic disorder, agoraphobia and social phobia.

**Alcohol phenotype groups: 0=lower-risk drinkers, 1=hazardous drinkers, 2=harmful drinkers.

Table 2 Logistic regression model outputs based on imputed data set when comparing hazardous drinkers and harmful drinkers to lower-risk drinkers

24-year clinic	Prevalence (n, %)	Crude (model 1) OR (95% CI, p value)	Model 2* OR (95% CI, p value)	Model 3* OR (95% CI, p value)
Anxiety, GAD				
Lower-risk drinkers	163 (11.39)	–	–	–
Hazardous drinkers	133 (7.55)	0.64 (0.50–0.80)	0.67 (0.53–0.86)	0.69 (0.54–0.88)
Harmful drinkers	51 (13.46)	1.20 (0.86–1.69) p<0.001	1.36 (0.97–1.91) p<0.001	1.39 (0.99–1.96) p<0.001
Depression				
Low-risk drinkers	180 (12.58)	–	–	–
Hazardous drinkers	144 (8.17)	0.62 (0.49–0.79)	0.66 (0.52–0.84)	0.68 (0.54–0.86)
Harmful drinkers	67 (17.68)	1.49 (1.09–2.02) p<0.001	1.72 (1.26–2.36) p<0.001	1.77 (1.29–2.43) p<0.001
Fear-based anxieties				
Low-risk drinkers	17 (1.19)	–	–	–
Hazardous drinkers	11 (0.62)	0.52 (0.24–1.12)	0.57 (0.27–1.22)	0.59 (0.27–1.27)
Harmful drinkers	9 (2.37)	2.02 (0.89–4.58) p=0.01	2.40 (1.05–5.47) p=0.006	2.41 (1.04–5.56) p=0.009

*Model 1 (crude). Model 2 adjusted for sex. Model 3 adjusted for sex, social status, income and maternal education.

**Estimates are accompanied by omnibus p values with two df to assess differences in odds across all alcohol groups.

likely to be hazardous drinkers (53.7% vs 46.75) and harmful (14% vs 8.5%) drinkers (p<0.001).

Table 2 shows an association between alcohol consumption and internalising problems with harmful drinkers indicating more GAD (13.46% vs 11.39%), depression (17.68% vs 12.58%) and FBA (2.37% vs 1.19%) compared with lower-risk drinkers. Harmful drinking, after adjustment for sex and socioeconomic status, was associated with a higher prevalence of depression (OR 1.8, 95% CI 1.3 to 2.4, p<0.001), anxiety (OR 1.4, 95% CI 1.0 to 2.0, p<0.001) and FBA (OR 2.4, 95% CI 1.04 to 5.56, p=0.009) compared with lower-risk drinkers. In contrast, hazardous drinkers presented with a lower prevalence of GAD (7.55%), depression (8.17%) and FBA (0.62%) compared with harmful and lower-risk drinkers. After adjustment, hazardous drinking was associated with a lower prevalence of GAD (OR 0.69, 95% CI 0.54 to 0.88) and depression (OR 0.68, 95% CI 0.54 to 0.86) but not FBA (OR 0.59, 95% CI 0.27 to 1.27) compared with lower-risk drinkers. We found similar results when analysing alcohol consumption and individual, non-co-morbid internalising disorders, and these results are presented in online supplementary appendix 3.

Lastly, to explore the robustness of our findings with an alternative classification of alcohol misuse, a sensitivity analysis was conducted in which we found little difference to the distribution of alcohol use and misuse by sex and by internalising symptoms (online supplementary appendix 4).

DISCUSSION

Main findings

Our main findings show an association between all internalising disorders (GAD, depression and FBA) and harmful alcohol consumption with harmful drinkers more likely to present with GAD (14% vs 11%), depression (18% vs 13%) and FBA (2% vs 1%) compared with lower-risk drinkers. After adjustment for sex and socioeconomic status, harmful drinkers were more likely to present with anxiety (by 40%), depression (by 80%) and FBA (by 141%) compared with lower-risk drinkers. We found a J-shaped distribution with lower rates of internalising disorders particularly GAD (OR 0.69, 95 CI 0.54 to 0.88) and depression (OR 0.68, 95% CI 0.54 to 0.86) in hazardous drinkers compared with lower-risk drinkers.

Comparison with previous studies

Depression was more marked in women (13.4%) than in men (6.9%) consistent with previous work on sex differences in depression in early adulthood.^{2 26 27} Similarly, anxiety was more marked in women (11.6%) than in men (6.5%). Although anxiety and depression are comorbid conditions which share genetic markers, neural mechanisms and treatment effects,²⁸ each of these internalising disorders have individual characteristics that correspond different to the use and misuse of alcohol as our results depict.

Previous studies have reported higher levels of internalising disorders both in non-drinkers and harmful drinkers using AUD from the DSM-IV criteria.^{29–31} Some studies suggest a dose–response relationship between internalising symptoms and alcohol consumption⁶ while others indicate a consistent, elevated relationship across all levels of alcohol use.³² In contrast, we found a potential J-shaped relationship between alcohol use and internalising symptoms in young adults with hazardous drinking associated with lower levels of internalising compared with lower-risk drinkers and harmful drinkers.

A Brisbane-based maternal cohort found similar J-shaped relationships between alcohol consumption and internalising disorders at age 30 years and linear relationships at ages 25 and 40 years,³³ consistent with previous work particularly pertaining to Australian men.³⁴ Results from a New Zealand-based 35-year birth cohort showed little evidence of J-shaped relationship over the life course.⁷ Inverse relationships of higher mental ill-health in lower-risk drinkers or abstainers compared to harmful drinkers defined using DSM-IV criteria have also been reported,^{33 35} adding greater complexity to understanding the nature of this relationship across varying levels of alcohol consumption. It has been suggested that the increased risk of depression and anxiety found among both abstainers and those who consume large amounts of alcohol (harmful drinkers) may be an indicator of poor overall health,³³ or ‘sick-quitters’,³⁶ that is, individuals who have experienced psychiatric disorders as a result of their alcohol consumption and quit drinking—which may explain a higher prevalence of depression and anxiety in the lower-risk drinking group.³⁷ Other studies with longitudinal data found that abstinence may be a true predictor of poor mental health and is an early indicator of the relationship between abstinence and mortality.³⁸ Other potential explanations for the association between hazardous drinking and lower rates of internalising disorders include an increasing social or peer interaction as part of the UK’s wider ‘drinking culture’,³ which may mitigate or preclude the effects of mental ill-health in a young adult population. Personality traits and temperament may also explain behaviour with alcohol as novelty seeking, harm avoidance and reward dependence on consuming alcohol may

Short report

shape the way in which AUD develops.^{39 40} This may be unique to adolescents and young adults as these periods are often associated with higher prevalence of alcohol use and associated problems, but this transitional period is also characterised by normative personality changes leading to psychological maturity and increasing emotional stability.⁴¹ It could also be that lower-risk drinkers may have differing social characteristics (although we adjust for socioeconomic status and the association persists),⁷ or simply unmeasured confounding such as genetic predisposition to alcohol tolerance.⁴²

Limitations

The cross-sectional nature of the study design is a limitation as we cannot determine the nature or direction of associations between alcohol use and internalising disorders. Researchers^{6 7} also suggest that J-shaped relationships between these are largely reported from cross-sectional study designs while longitudinal analyses may better capture the variation of alcohol consumption and associated internalising disorders across the life course. Second, attrition of participants and missing data from several waves of data collection is an issue for any longitudinal cohort such as ALSPAC. When we compared clinic attenders with non-attenders (online supplementary appendix 2), we found differences in sex, social status, income levels and maternal education. More women attended clinic (62.6%) than did not in the non-attender group (44.83%) and in those who attended, 45.4% of achieved educational qualifications beyond school while this was only 27.9% of the non-attenders. In order to mitigate some of these differences, we used multiple imputation techniques using a large number of auxiliary variables in our analyses. While this does not completely address the high attrition rate in ALSPAC, it does help maximise the model power. Third, the CIS-R which was used to determine internalising symptoms focuses on the past 1 week only and may not include on all internalising disorder 'cases'. However, CIS-R has demonstrated good validity and reliability in diagnosing common mental disorders and depressive episodes according to ICD-10 and compares well with other self-administered tools such as the General Health Questionnaire and Centre for Epidemiologic Studies Depression Scale.⁴³ Lastly, the analysis excludes additional covariates factors such as personal relationships or exposures to early childhood adversities that may independently explain the presence of internalising disorders⁴⁴ and alcohol misuse,⁴⁵ but may also mediate the associations between these disorders.⁴⁶

Implications

The results suggest that young adults who drink harmfully do present with increased internalising symptoms, and future work should focus on unpicking the complex and reverse relationships found between hazardous drinking and harmful drinking and, internalising disorders. The J-shaped results suggest a varying relationship between different levels of alcohol consumption and internalising disorders being subject to unmeasured covariates (eg, personality temperaments or genetic predisposition or exposure to early childhood adversities) and relationship itself further varying across the life course. Future work should consist of longitudinal analyses where the nature and direction of the relationship is explored with specific consideration for personal- and societal-level factors.

What is already known on this subject

- Depression and anxiety often associate with harmful alcohol consumption. Early adulthood is a developmental period marked by significant life changes and is considered a high-risk period for the onset of both, internalising disorders and alcohol-related problems. Internalising disorders and alcohol problems have a bi-directional relationship and have both been associated with longer-term mental ill-health and alcohol misuse.

What this study adds

- The study reports that young adults, from a UK-based birth cohort (ALSPAC), who drink harmfully, are more likely to have internalising disorders such as depression, anxiety and fear-based anxieties. Women report more internalising disorders than men. After adjusting for sex and socioeconomic status, harmful drinking was associated with a higher prevalence of internalising disorders compared with lower-risk drinkers. However, future research should focus on unpicking the complex and reverse relationships found between hazardous drinking and harmful drinking and, internalising disorders.

Twitter KushAbey.

Acknowledgements We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. This publication is the work of GSF and JH who will serve as guarantors for the contents of this paper.

Contributors GH, JH and MH were responsible for the initial study design. GSF performed the statistical analysis and GSF wrote the initial draft of the manuscript. GH and KA corroborated the analysis. JH, LM, AE, Gemma Lewis, GL and MH contributed to and approved the final manuscript.

Funding This research was specifically funded by the UK Medical Research Council and Alcohol Research UK Grant (MR/L022206/1). The UK Medical Research Council and Wellcome Trust (Grant Ref: 092731) and the University of Bristol provide core support for ALSPAC. We also acknowledge support from The Centre for the Development and Evaluation of Complex Interventions for Public Health Improvement (DECIPHer), a UKCRC Public Health Research Centre of Excellence (joint funding (MR/K023233/1) from the British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, the Welsh Government and the Wellcome Trust, under the auspices of the UK Clinical Research Collaboration), the NIHR School of Public Health Research, NIHR Health Protection Research Unit in Evaluation and NIHR Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol. LM is supported by the Elizabeth Blackwell Institute for Health Research, University of Bristol and the Wellcome Trust Institutional Strategic Support Fund (204813/Z/16/Z). GH is supported by a Sir Henry Wellcome Postdoctoral Fellowship (209138/Z/17/Z).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Gwen Sascha Fernandes <http://orcid.org/0000-0003-0203-7053>

REFERENCES

- 1 Crum RM, Green KM, Storr CL, *et al*. Depressed mood in childhood and subsequent alcohol use through adolescence and young adulthood. *Arch Gen Psychiatry* 2008;65:702–12.

- 2 Edwards AC, Joinson C, Dick DM, *et al.* The association between depressive symptoms from early to late adolescence and later use and harmful use of alcohol. *Eur Child Adolesc Psychiatry* 2014;23:1219–30.
- 3 Pedrelli P, Shapero B, Archibald A, *et al.* Alcohol use and depression during adolescence and young adulthood: a summary and interpretation of mixed findings. *Curr Addict Rep* 2016;3:91–7.
- 4 World Health Organisation. Global status report on alcohol and health. 2018. Available <https://apps.who.int/iris/bitstream/handle/10665/274603/9789241565639-eng.pdf?ua=1> (accessed 2nd Jan 2020)
- 5 Rehm J, Gmel GE, Sr, Gmel G, *et al.* The relationship between different dimensions of alcohol use and the burden of disease—an update. *Addiction (Abingdon, England)* 2017;112:968–1001.
- 6 Boden JM, Fergusson DM. Alcohol and depression. *Addiction* 2011;106:906–14.
- 7 Boden JM, Foulds JA, Horwood LJ. Examination of a possible J-shaped relationship between alcohol consumption and internalizing disorders in a longitudinal birth cohort. *Drug Alcohol Depend* 2016;162:88–91.
- 8 Bolton JM, Robinson J, Sareen J. Self-medication of mood disorders with alcohol and drugs in the national epidemiologic survey on alcohol and related conditions. *J Affect Disord* 2009;115:367–75.
- 9 Frojd S, Ranta K, Kaltiala-Heino R, *et al.* Associations of social phobia and general anxiety with alcohol and drug use in a community sample of adolescents. *Alcohol* 2011;46:192–9.
- 10 Chung T, Creswell KG, Bachrach R, *et al.* Adolescent binge drinking. *Alcohol Res* 2018;39:5–15.
- 11 Bell S, Britton A. An exploration of the dynamic longitudinal relationship between mental health and alcohol consumption: a prospective cohort study. *BMC Med* 2014;12:91.
- 12 McCambridge J, McAlaney J, Rowe R. Adult consequences of late adolescent alcohol consumption: a systematic review of cohort studies. *PLoS Med* 2011;8:e1000413.
- 13 Public Health England. The public health burden of alcohol and the effectiveness and cost-effectiveness of alcohol control policies. An evidence review. 2016. Available https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf (accessed 1st Jan 2020)
- 14 Northstone K, Lewcock M, Groom A, *et al.* The Avon Longitudinal Study of Parents and Children (ALSPAC): an update on the enrolled sample of index children in 2019. *Wellcome Open Res* 2019;4:51.
- 15 World Health Organization, 2016. International Classification Of Diseases, Tenth Revision. Available <http://apps.who.int> ((last accessed)12 July 2016).
- 16 Crocq MA. The history of generalized anxiety disorder as a diagnostic category. *Dialogues Clin Neurosci* 2017;19:107–16.
- 17 Dahne J, Banducci AN, Kurdziel G, *et al.* Early adolescent symptoms of social phobia prospectively predict alcohol use. *J Stud Alcohol Drugs* 2014;75:929–36.
- 18 Torvik FA, Rosenstrom TH, Gustavson K, *et al.* Explaining the association between anxiety disorders and alcohol use disorder: a twin study. *Depress Anxiety* 2019;36:522–32.
- 19 Lewis G, Pelosi AJ, Araya R, *et al.* Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. *Psychol Med* 1992;22:465–86.
- 20 Gordon AJ, Maisto SA, McNeil M, *et al.* Three questions can detect hazardous drinkers. *J Fam Pract* 2001;50:313–20.
- 21 Fujii H, Nishimoto N, Yamaguchi S, *et al.* The Alcohol Use Disorders Identification Test for Consumption (AUDIT-C) is more useful than pre-existing laboratory tests for predicting hazardous drinking: a cross-sectional study. *BMC Public Health* 2016;16:379.
- 22 Khadjesari Z, White IR, McCambridge J, *et al.* Validation of the AUDIT-C in adults seeking help with their drinking online. *Addict Sci Clin Pract* 2017;12:2.
- 23 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th edn. Washington, DC: American Psychiatric Association, 1994.
- 24 Takahashi T, Lapham G, Chavez LJ, *et al.* Comparison of DSM-IV and DSM-5 criteria for alcohol use disorders in VA primary care patients with frequent heavy drinking enrolled in a trial. *Addict Sci Clin Pract* 2017;12:17.
- 25 Office for national statistics. Adult drinking habits in Great Britain. 2018. Available <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/druguse/alcoholandsmoking/bulletins/opinionsandlifestylesurveyadultdrinkinghabitsingreatbritain/2017> (accessed 5th Mar 2020)
- 26 Marmorstein NR, Iacono WG, Malone SM. Longitudinal associations between depression and substance dependence from adolescence through early adulthood. *Drug Alcohol Depend* 2009;107:154–60.
- 27 Albert PR. Why is depression more prevalent in women? *J Psychiatry Neurosci* 2015;40:219–21.
- 28 Goodwin GM. The overlap between anxiety, depression, and obsessive-compulsive disorder. *Dialogues Clin Neurosci* 2015;17:249–60.
- 29 Dawson D, Grant B, Stinson F, *et al.* Psychopathology associated with drinking and alcohol use disorders in the college and general adult populations. *Drug Alcohol Depend* 2005;77:139–50.
- 30 Brière F, Rohde P, Seeley J, *et al.* Comorbidity between major depression and alcohol use disorder from adolescence to adulthood. *Compr Psychiatry* 2014;55:526–33.
- 31 Buchanan JL. Prevention of depression in the college student population: a review of the literature. *Arch Psychiatr Nurs* 2012;26:21–42.
- 32 Keyes KM, Hamilton A, Patrick ME, *et al.* Diverging trends in the relationship between binge drinking and depressive symptoms among adolescents in the U.S. from 1991 through 2018. *J Adolesc Health* 2019.
- 33 Alati R, Najman JM, Kinner SA, *et al.* Early predictors of adult drinking: a birth cohort study. *Am J Epidemiol* 2005;162:1098–107.
- 34 Caldwell TM, Rodgers B, Jorm AF, *et al.* Patterns of association between alcohol consumption and symptoms of depression and anxiety in young adults. *Addiction* 2002;97:583–94.
- 35 Lucas N, Windsor TD, Caldwell TM, *et al.* Psychological distress in non-drinkers: associations with previous heavy drinking and current social relationships. *Alcohol Alcohol* 2009;45:95–102.
- 36 Degenhardt L, Hall W, Lynskey M. Testing hypotheses about the relationship between cannabis use and psychosis. *Drug Alcohol Depend* 2003;71:37–48.
- 37 Stockwell T, Zhao J, Panwar S, *et al.* Do “moderate” drinkers have reduced mortality risk? A systematic review and meta-analysis of alcohol consumption and all-cause mortality. *J Stud Alcohol Drugs* 2016;77:185–98.
- 38 Power C, Rodgers B, Hope S. U-shaped relation for alcohol consumption and health in early adulthood and implications for mortality. *Lancet* 1998;352:877.
- 39 Cloninger CR, Sigvardsson S, Bohman M. Type I and type II alcoholism: an update. *Alcohol Res Health* 1996;20:18.
- 40 Helle AC, Watts AL, Trull TJ, *et al.* Alcohol use disorder and antisocial and borderline personality disorders. *Alcohol Res: Curr Rev* 2019;40:arcr.v40.1.05.
- 41 Roberts BW, Walton KE, Viechtbauer W. Patterns of mean-level change in personality traits across the life course: a meta-analysis of longitudinal studies. *Psychol Bull* 2006;132:1–25.
- 42 Holmes MV, Dale CE, Zuccolo L, *et al.* Association between alcohol and cardiovascular disease: Mendelian randomisation analysis based on individual participant data. *BMJ* 2014;349:g4164.
- 43 Head J, Stansfeld SA, Ebmeier KP, *et al.* Use of self-administered instruments to assess psychiatric disorders in older people: validity of the General Health Questionnaire, the Center For Epidemiologic Studies Depression Scale and the Self-completion Version of the Revised Clinical Interview Schedule. *Psychol Med* 2013;43:2649–56.
- 44 Poole JC, Dobson KS, Pusch D. Childhood adversity and adult depression: the protective role of psychological resilience. *Child Abuse Negl* 2017;64:89–100.
- 45 Dube SR, Miller JW, Brown DW, *et al.* Adverse childhood experiences and the association with ever using alcohol and initiating alcohol use during adolescence. *J Adolesc Health* 2006;38:444.e441–410.
- 46 Kim Y, Kim K, Chartier KG, *et al.* Adverse childhood experience patterns, major depressive disorder, and substance use disorder in older adults. *Aging Ment Health* 2019;1–8.