



Hardy, Christopher, Sayyed, Imam, Leslie, Andrew and Dittrich, Alex DK. (2020) Effectiveness of insecticides, physical barriers and size of planting stock against damage by the pine weevil (*Hylobius abietis*). *Crop Protection* . p. 105307.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/5606/>

***Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.***

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

**provided that**

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
  - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

**You may not**

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing [insight@cumbria.ac.uk](mailto:insight@cumbria.ac.uk).

1 Effectiveness of insecticides, physical barriers and size of planting stock  
2 against damage by the pine weevil (*Hylobius abietis*).  
3

4 Christopher Hardy<sup>a</sup>, Imam Sayyed<sup>a)</sup>, Andrew Dunbar Leslie<sup>b\*</sup> and Alex DK Dittrich<sup>c</sup>

5 <sup>a</sup> Maelor Forest Nurseries Ltd, Bronington, Wales, SY13 3HZ, UK

6 <sup>b</sup> National School of Forestry, University of Cumbria, Ambleside, LA22 9BB, UK

7 <sup>c</sup> Department of Science, Natural Resources and Outdoor Studies, University of Cumbria, Carlisle, UK

8 \* Corresponding author  
9

## 10 Abstract

11 A series of five trials was established in Great Britain to test the effectiveness of fourteen treatments  
12 and a control on reducing mortality and damage by pine weevil (*Hylobius abietis*) on recently  
13 replanted Sitka spruce (*Picea sitchensis*). Overall percentage mortality and damage was significantly  
14 different between trials, varying from a median of 24% to 100%. The most effective treatments in  
15 reducing mortality were insecticides and physical barriers, with insecticides being most cost  
16 effective. Low volume applications of insecticides were found to be as effective as higher doses.  
17 Using larger trees did not reduce mortality compared to the control, nor did application of a  
18 controlled release fertiliser. At Auchencairn, the trial where mortality was highest no treatment  
19 provided protection and so there is a need to develop an effective and integrated approach to  
20 reducing damage by pine weevil.

21 **Keywords:**

22

23 *Hylobius*, control measures, conifer plantation, seedling damage, *Picea sitchensis*

## 24 **1. Introduction**

25 In the UK, pine weevils (*Hylobius abietis*) are one of the most damaging pests on recently planted  
26 conifer restock sites (Willoughby *et al.*, 2017) and the only insect pest where routine preventative  
27 measures are taken (Leather *et al.* 1999). Maturation feeding by adult weevils damages the young  
28 trees, through chewing of the bark on the lower stem, causing girdling and death (Willoughby *et al.*,  
29 2017). Without some form of intervention, in the UK mortality of young trees averages 50%  
30 (Willoughby *et al.* 2017) ranging between 30% and 100% (Leather *et al.*, 1999). The pine weevil also  
31 feeds on branches of mature trees and can be a vector of damaging fungal pathogens (Leather *et al.*,  
32 1999).

33 Despite pine weevil being recognised as a serious pest since the beginning of the last century  
34 (Leather *et al.* 1999) it remains a highly damaging agent to commercial forestry in the UK (Willoughby  
35 *et al.*, 2017). There are no recent assessments of the cost of this damage but it was estimated in the  
36 late 1990s that pine weevil cost the forestry sector £4 million per year, for direct control measures  
37 alone, not including the replacement of killed trees (Leather *et al.*, 1999). Furthermore, future  
38 damage from pine weevil is likely to increase due to the heightened temperatures predicted as a  
39 result of climate change (Inward *et al.*, 2012).

40

41 The pine weevil is an important pest elsewhere in Europe (Leather *et al.*, 1999), and a wide range of  
42 approaches have been adopted or are being investigated to control damage. These can be grouped  
43 into nine broad categories: (1) chemical control (Luoranen & Viiri, 2005), (2) biological control

44 (Williams *et al.*, 2013), (3) physical barriers (Petersson *et al.*, 2004), (4) site preparation (Wallertz *et*  
45 *al.* 2018), (5) use of larger (Thorsen *et al.* 2001, Nordlander *et al.*, 2011) or smaller planting material  
46 (Petersson *et al.*, 2008), (6) delayed planting (Moore 2004), (7) use of antifeedants (Unelius *et al.*,  
47 2018), (8) enhanced plant defences (Lundborg *et al.*, 2016, Zas *et al.* 2014) and (9) genetic  
48 improvement (Zas *et al.*, 2017). There have also been attempts to develop systems that predict the  
49 risk of damage to the young trees through characterisation of sites (Heritage & Moore 2000,  
50 Nordlander *et al.*, 2017, Lopez-Villamor *et al.*, 2019) and population monitoring systems have also  
51 been developed (Wainhouse *et al.*, 2007, Forest Research 2019). There has been interest in  
52 combinations of these approaches to create an integrated pest management system for pine weevil  
53 in the UK (Evans *et al.*, 2003). The trials that this paper reports were focused on chemical  
54 treatments, physical barriers, the use of larger plantings stock and fertiliser. **These treatments in**  
55 **combination have not been formally compared before in the UK and other than one treatment, the**  
56 **polymer, the methods used could all be adopted currently to protect trees on planting sites.**

57 Use of insecticide treated trees has been shown to be a cost effective and practical approach to  
58 reducing damage. **In 2016 it was used to protect 40% of trees planted in the EU** (Norsk Wax, 2016).  
59 Leather *et al.*, (1999) describe the ideal insecticide for pine weevil as one that is systemic and kills  
60 the adults but also masks or alters attractive host volatiles. Some insecticides have these two  
61 attributes; a study by Rose *et al.*, (2005) showed that application of **either a pyrethroid or a**  
62 **neonicotinoid insecticide to young Scots pine (*Pinus sylvestris*) trees inhibited feeding on them by**  
63 **pine weevil.**

64 **While insecticides are effective and cheap, there is pressure to minimise the use of pesticides in**  
65 **forestry (Willoughby *et al.*, 2004), while under the UK Woodland Assurance Scheme the use of**  
66 **pesticides, biological control measures and fertilisers are to be reduced (UKWAS, 2018). There are**  
67 **also international influences encouraging reduction of pesticide use in forests, such as forest**

68 certification (FSC, 2019) and regional policies such as the 2009 European Commission Directive on  
69 the Sustainable Use of Pesticides (European Commission, 2019).

70 Three insecticides are currently used in the UK to treat stock in the nursery, prior to planting to  
71 protect it from pine weevil damage. These are alpha cypermethrin and two neonicotinoid  
72 pesticides: imidacloprid, marketed as Merit Forest and acetamiprid, marketed as Gazelle (The  
73 Advisory Service for Weed & Pest Control in Forestry & Amenity, 2018). For top up spraying in the  
74 forest two insecticides are used: cypermethrin, marketed as Forester and Gazelle (The Advisory  
75 Service for Weed & Pest Control in Forestry & Amenity, 2018). In a trial in Sweden, Merit Forest was  
76 shown by Petersson *et al.*, (2006) to be less effective than cypermethrin as a pre-planting treatment  
77 with significantly higher losses in the second year. Evidence shows that Gazelle is potentially much  
78 less harmful than alpha cypermethrin (Willoughby *et al.*, 2017).

79 An alternative to using chemicals are physical barriers to pine weevil. Various guards have been  
80 tested and design is known to affect their efficacy. Petersson *et al.*, (2004) tested two broad types of  
81 guards, ones with a collar (a structure at the top of the guard preventing weevils from climbing up  
82 and over the guard) and ones without a collar. They found those with a collar to be effective at  
83 controlling damage by pine weevil. A later piece of research by Petersson *et al.*, (2006) in Sweden  
84 showed Clipstop, a collared guard to be as effective as permethrin over three years.

85 Clipstops were tested in informal trials in the UK in 2005 but did not reduce damage (Leslie &  
86 Liddon, 2017) and other barriers have been tried including paper guards and those fabricated from  
87 ladies' stockings but none have been successful (Leslie & Liddon, 2017). Despite this unpromising  
88 experience, there are other guards available in the UK, including plastic Biosleeves, Multipro Sleeves  
89 and Weenets (Willoughby *et al.*, 2017).

90 A different approach involves the application of protective coatings on the lower part of the stem of  
91 young trees. These include Flexcoat, a polysaccharide coating (Harlin and Eriksson, 2010), and  
92 Conniflex, a sand and glue based coating (Nordlander *et al.*, 2009). Conniflex has proved to be

93 effective in protecting Norway spruce (*Picea abies*) and Scots pine (*Pinus sylvestris*) against pine  
94 weevil damage in a trial in Sweden (Nordlander *et al.*, 2009) and to date has protected 320 million  
95 container grown stock trees (Svenska-Skogsplantor ,2019). Another treatment, Norsk Wax has been  
96 developed in Scandinavia (Norsk Wax 2016). Field trials in Sweden of this product were encouraging  
97 with 1.3% of treated trees being killed compared with 29% of those unprotected (Ohrn &  
98 Nordlander, 2015). The use of protective coatings has largely replaced the use of insecticides in  
99 Sweden with an intention that 2020 will be the last year where insecticides will be applied (Sveriges  
100 Lantbruksuniversitet, 2018).

101 Larger trees have been shown to be less attractive to pine weevil and to exhibit lower mortality  
102 (Nordlander *et al.*, 2011). Thorsen *et al.* (2001) identified a threshold basal diameter of greater than  
103 8mm and scarification around the planting position were needed for high seedling survival of  
104 Norway spruce (*Picea abies*) in Sweden. However, a study (Petersson *et al.*, 2008) showed mini-  
105 trees, 10 week old containerised seedlings of Norway spruce to be less damaged (3.5% vs 55%) than  
106 conventional planting stock. When the mini-trees were damaged by pine weevil they released larger  
107 concentrations of limonene, a chemical known to be repellent to pine weevil, while conventional  
108 planting stock released the attractant alpha-pinene.

109 In April 2017 Maelor Forest Nurseries Ltd. in collaboration with Scottish Woodlands, Flintshire  
110 Woodlands Ltd and Tilhill Forestry established a series of five trials across Great Britain with the aim  
111 of testing the effectiveness of fifteen treatments (including a control) in reducing the mortality and  
112 damage by pine weevil on newly planted Sitka spruce (*Picea sitchensis*). The treatments included  
113 chemical control measures, physical barriers and planting materials that had greater girth and  
114 greater height than standard planting material. Specifically, the aims of the trials were to:

- 115 1. Identify the most effective control measures in reducing mortality and damage
- 116 2. **Relate** the effectiveness of the control measures to their cost

- 117 3. Compare effectiveness of treatments across sites.
- 118 4. Relate the characteristics of the sites to damage from pine weevil

119 An analysis of results at two years is presented in this paper and recommendations made for future  
120 work. The cumulative damage over two growing seasons, rather than one was used as it gives a  
121 better indication of the effects of treatments on damage and mortality over the establishment  
122 period.

123

## 124 2. Material and Methods

125

### 126 2.1 Site descriptions

127

128 Five trials were established in April 2017 in the uplands of the UK, with four located in south west  
129 Scotland and the fifth, Cwm Henog, in southern Wales. The locations are described in Table 1 with  
130 details on the climate and suitability for Sitka spruce as generated by the Ecological Site  
131 Classification (Forest Research 2020). Sitka spruce was selected as it is the most common species in  
132 commercial softwood forestry plantations in the UK (Forestry Commission, 2019). Heritage and  
133 Moore (2000) identified site characteristics that strongly influenced damage by pine weevil and the  
134 sites are compared against these factors. All comprised stands of monoculture Sitka spruce or  
135 stands with a preponderance of Sitka spruce which had been clear felled recently. Furthermore,  
136 the sites were not close to areas of serious windthrow, the crop had not been thinned within six  
137 years previously, however all had older trees within 1-2km distance, and all had standing trees  
138 nearby. There was exposed mineral soil around the trees on all sites, reducing risk but conversely

139 the brash and other harvesting residues had not been burned and stumps had not been removed.

140 Characteristics that differed between the trials are described in Table 2.

## 141 2.2 Treatments

142

143 We tested 14 treatments and a control. For Auchencairn, Cwm Henog, Newton Stewart and  
144 Ramshaw Rig sites, Sitka spruce came from A12 Seed Orchard material. For Lamloch, vegetatively  
145 propagated Sitka spruce was used from PF80, a full sib family with material vegetatively propagated.

146 All trees were bare root and were graded for size and quality before being treated. There were four  
147 broad categories of treatment: physical barriers, chemical insecticides, size of planting stock and  
148 application of controlled release fertiliser.

149 The physical barriers comprised four treatments. Biosleeves (Greenerpol Ltd, UK) and Multipro  
150 sleeves (Svenska Skogplantor, Sweden) were commercially available protection, fitted around the  
151 stems of the young trees at the nursery. Multipro sleeves were made of waxed cardboard and are  
152 known to be biodegradable. Biosleeves, which are no longer available were made from an  
153 experimental biodegradable plastic, with an estimated lifetime in the forest of 2-5 years. The other  
154 two barrier treatments used an experimental polymer, applied to the tree stem in the nursery,  
155 covering either 85% or 50% of the stem. Application was carried out by dipping plants into a  
156 container of liquid polymer, taking care to avoid contact with roots or branches of the plant. The  
157 plants were then left to air dry for 1 hour before packing. Roots were regularly misted with water to  
158 prevent them drying out.

159 The chemical treatments used three insecticides: Gazelle (Nisso Chemical Europe GmbH: 0.037g per  
160 tree acetamiprid), Alpha C6 ED (Techneat Ltd: 0.006g per tree alpha-cypermethrin) and Coragen  
161 (Dupont: 0.013g per tree chlorantraniliprole). Coragen is not currently used on an operational scale  
162 to control pine weevil damage in Great Britain. The Alpha C 6 ED treatment was applied using an  
163 Electrodyn machine (Techneat Engineering Ltd.), which applies electrically charged insecticide



164 accurately to individual root collars of trees. The trees are earthed in the machine by passing over an  
165 earthing wire, ensuring that the insecticide is attracted to exactly the right location on the plant. The  
166 treatments of Coragen and Gazelle were applied using ultra low volume (ULV) application nozzles in  
167 bundles of ten trees but is now applied to individual trees using a purpose built machine. Due to the  
168 low volumes of water used in this process, the plants could be packed immediately after treatment.  
169 Conventional application was by knapsack sprayer to trees laid out on a clean, protected ground  
170 surface. Trees were turned over during the process to ensure good coverage. The trees were left to  
171 air dry for one hour before being packed. Roots were regularly misted with water to prevent them  
172 drying out.

173 For all trials except Lamloch, seed orchard PSI A12 material was planted. Four treatments using  
174 larger trees than conventional planting stock (1 +1, 30-50 cm tall, 5mm root collar diameter) were  
175 tested in the trials. Planting stock of 2+1 (30-50cm tall, 6mm root collar diameter), with a larger  
176 girth and 2+1 50-70cm with a larger girth (8mm root collar diameter) and height were included. At  
177 Lamloch, vegetatively propagated (PF80) planting stock was planted. The 1+1 treatment was  
178 replaced by S+1 (25-50cm tall, 5.5mm root collar diameter), while the 2+1 treatment was replaced  
179 by a S+2 (25-50cm tall, 6.5mm root collar diameter) and the 2+1 50-70cm with a larger girth  
180 treatment was replaced with S+2 (50-70cm tall, 8.5mm root collar diameter).

181

182 For each of two larger tree treatments and for the standard sized trees there were trees with and  
183 without application of Treeboost, a controlled release fertiliser (CRF) developed by Maelor Forest  
184 Nurseries Ltd. This contains the following elements at the following concentrations: 13%N, 11% P,  
185 8% K + 2% CaO, 0.10% B, 0.015% Cu, 0.20% Fe, 0.08% Mn and 0.025% Zn. A summary of the  
186 treatments is described in Table 3.

187 For each plot 25 trees were packed into an individual labelled plastic bag. Four bags of the same  
188 treatment type were then packed into a larger bag which was then sealed. These bags were then  
189 kept in a cold store at approximately -2° C until point of planting.

## 190 2.3 Experimental Design and Measurements

191

192 The trials were established using a randomised complete block design with 15 treatments and four  
193 replicates. Each plot contained 25 trees in a 5 tree by 5 tree layout. Stocking density was the  
194 commercial stocking rate used in the UK of 2,700 stems ha<sup>-1</sup>. Assessments were made in July 2017,  
195 November 2017 and July 2018. For each plot each individual tree was scored using the following five  
196 categories:

A = Undamaged

B = Damaged by weevil, but tree likely to survive

C = Tree dead/dying due to weevil damage

D = Tree dead for reasons other than weevil damage

X = Tree missing

197 Damage was attributed to pine weevil if the characteristic damage to the bark of the stem of the  
198 young tree was present. It is the 2018 assessment that is reported on in this paper.

199

200 Statistical analysis

201

202 Percentage mortality and percentage damaged by pine weevil per plot were combined and  
203 calculated by:

204

205 
$$\text{Percentage mortality and damaged} = \frac{100 \cdot (B + C)}{(A + B + C)}$$

206 Originally damage and death were to be analysed separately. However, these two variables were  
207 combined for two reasons. The first was that it has been shown that damage by pine weevil reduced  
208 subsequent survival and growth (Leather *et al.*, 1999) and second separating damage and mortality  
209 for analysis presented confusing results. For example, a trial with high mortality may have low  
210 damage, because there are few trees alive to be damaged. This was particularly notable at  
211 Auchencairn, the trial with the highest mortality.

212 The unstandardised residuals for percentage mortality and damaged were tested for normality. For  
213 the treatments a Shapiro Wilkes test was used due to the small number of observations per  
214 treatment, whereas for the comparison between trials, the larger number of observations for each  
215 treatment meant a Kolmogorov- Smirnov test was used. For the data for all trials the residuals from  
216 mortality and damage data before and after an Arcsine transformation did not conform to normality  
217 so non-parametric Kruskal Wallis tests have been used to test significance of differences. For the  
218 mortality and damage data for individual trials all but Auchencairn were normally distributed but the  
219 variances were unequal for all trials except Cwym Henog. Non-parametric Kruskal Wallis tests were  
220 therefore used to test significance of differences.

221 To identify which treatments were significantly different a post-hoc Dunn's test was employed with  
222 pairwise comparison applying the Bonferroni adjustment for multiple comparisons. This approach  
223 was applied to the comparison of death and damage between trials. However, this is overly  
224 conservative when applied to large numbers of comparisons (McDonald, 2015) such as when testing  
225 significance between the 15 treatments. In this case, the Benjamini-Hochberg procedure was  
226 applied, accepting a potential false-positive rate of one in ten (McDonald, 2015).

## 227 3. Results

228

### 229 3.1 Overview of Results

230

231 The results of damage by the five damage categories assessed are presented in Figure 1 for the trials  
232 and Figure 2 for the treatments. The percentage of trees killed by agents other than pine weevil was  
233 as much as 25%, but the percentage of trees missing was low at 10% or less. Detailed results of  
234 damage and mortality are described in the following sections.

### 235 3.2 Mortality and damage by trial

236

237 There were significant differences ( $p < 0.001$ ) in the median percentage mortality and damage  
238 between trials (Figure 3). Level of damage and mortality at Lamloch (median = 35%), Ramshaw Rig  
239 (median = 36%) and Newton Stewart (median = 24%) were lower than the other trials and not  
240 significantly different. There was a high level of damage and mortality at Cwym Henog  
241 (median = 72%) and complete mortality and damage at Auchencairn (median = 100%).

### 242 3.3 Mortality and damage by treatment

243

244 Figure 4 (a-e) describe the death and damage caused by pine weevil for each of the treatments.  
245 Results from Auchencairn are not shown as death and damage for all treatments was 100%.  
246 Combining the data from all trials including Auchncairn (Figure 4e) showed significant differences  
247 between treatments ( $p < 0.00001$ ) and that chemical treatments and barriers were as effective as  
248 Gazelle, the exception being the 50% polymer treatment. Larger planting stock and larger planting  
249 stock with controlled release fertiliser were no more effective than the control. Highly significant

250 differences between treatments were found at all trials, except Auchencairn (Lamloch  $p=0.003$ ,  
251 Ramshaw Rig  $p=0.008$ , Newton Stewart  $p=0.0001$  and Cwym Henog  $p=0.007$ ). Despite the variation  
252 in overall levels of death and damage between trials the general findings were that chemicals and  
253 barriers were most effective and that larger plants, with or without fertiliser were no more effective  
254 than the control. The efficacy of even the effective treatments **was** much reduced in trials where  
255 overall damage was high, such as Cwym Henog (Figure 4a) and Auchencairn, where all treatments  
256 exhibited 100% dead or damaged.

## 257 4. Discussion

258 There were significant differences in the level of **damage and** mortality between trials. The highest  
259 level of mortality was at Auchencairn and the next highest was at Cwym Henog. The percentage  
260 mortality and damage was significantly different between these trials and between them and the  
261 other three trials (**Figure 1, Figure 3**). At Auchencairn the previous Sitka spruce stand was felled over  
262 a protracted period, which may have allowed damaging populations of pine weevil to build up on  
263 the site. **Also, there was heavy grass growth on that site, which is known to increase pine weevil**  
264 **damage (Orlander and Nordlander, 2003). Furthermore,** the new trees were 'hot planted'  
265 immediately after harvesting and the extended harvesting of the previous crop may have acted to  
266 attract pine weevil onto the site through a continuous supply of fresh stumps and brash. Lamloch  
267 was the only site where vegetatively propagated material was planted and this has been shown to  
268 be **less** attractive to pine weevil than seedling stock (Kennedy et al., 2006). **Lamloch was the trial**  
269 **site that sustained least mortality and damage from pine weevil and using vegetatively propagated**  
270 **stock could be a factor. The treatments that were effective at reducing mortality and damage at**  
271 **other trials were also effective at Lamloch.**

272 Using the data pooled from all five trials, significant differences in percentage mortality were  
273 detected between treatments. Broadly, the treatments can be divided into two groups, those that  
274 **effectively protected the young trees and those that were ineffective (no better than the control).**

275 Chemical treatments and the physical barriers provided by the Multipro sleeves, Biosleeves and the  
276 85% polymer treatment proved effective in reducing damage and death. The Multipro sleeves and  
277 Biosleeves provided excellent protection but their effectiveness was compromised by poor  
278 installation in some cases, allowing access to trees' stems by the pine weevils. The fitting of Multipro  
279 sleeves and Biosleeves onto trees was labour intensive, and relatively slow compared to other  
280 treatments which adds to their cost (Table 3, Figure 5). However, the additional cost of the guards  
281 may be offset by the need in some cases for top up spraying on insecticide treated trees. Application  
282 of guards was hindered by bigger branches on larger seedlings which reduced clearance for the  
283 sleeve. Proper burial of the lower part of the sleeve base was impeded by stony ground on some  
284 sites, allowing weevils to access the tree from the base. Furthermore, on exposed sites, it has been  
285 observed that movement of the tree shifted the position of the sleeve on the stem, allowing access  
286 by pine weevils. Weevils that were found inside sleeves were observed to cause significant damage  
287 to the tree.

288 The novel polymer barrier treatment gave mixed results, with death and damage for the 85% stem  
289 coverage treatment being significantly different from the control using pooled data from all trials. In  
290 contrast, death and damage using the 50% stem coverage was not significantly different from the  
291 control. It was only at Cwym Henog, a trial with high levels of damage that the difference in death  
292 and damage is significantly lower for the 85% stem coverage treatment than the 50% treatment.  
293 The mixed performance of the polymer may be due to poor adhesion to the stem in some cases,  
294 reducing its protection to the young tree. The success of stem coatings in Scandinavia in reducing  
295 pine weevil damage (Norsk Wax, 2016) and their widespread use (Sveriges Lantbruksuniversitet,  
296 2018) suggests this is an approach worth further investigation. However, problems were  
297 experienced in trials in the UK with the protective wax cracking and exposing the stem to damage  
298 (Leslie and Liddon ,2017). The cost of the experimental polymer was not known but is assumed to be  
299 similar to that of Norsk Wax, which costs €0.05 (£0.04) to €0.09 (£0.08) per tree (Norsk Wax, 2016)  
300 (Table3, Figure 5).

301 Using greater stem coverage, such as in the 85% treatment may have drawbacks over the 50%  
302 treatment as research in Sweden showed that greater coverage of the stem and needles reduced  
303 growth (Norsk Wax, 2016) and the importance of protection higher up the stem is not clear as  
304 weevils prefer to feed on the lower part of the stem where there is more cover (Nordlander *et al.*,  
305 2005). In these trials, mortality from causes other than pine weevil was greater in the 85% treatment  
306 than others (Figure 2) which suggests extensive covering of the stem reduces survival in addition to  
307 growth.

308 Using the pooled data from all trials and those from individual trials, chemical treatments and  
309 physical barriers were both effective in terms of their reduction in damage and mortality, with  
310 Gazelle, Gazelle ULV, Electrodyne, Coragen and Coragen ULV ranked highly, except at Cwym Henog.  
311 Chemical treatments were also more cost effective (Table 3, Figure 5) than the physical barriers.  
312 However, there is a global aim to reduce the use of chemicals used in forestry, driven by concerns  
313 about the toxic effect of insecticides on animals and by encouragement to reduce pesticide use by  
314 certification bodies (FSC, 2019). Coragen is less toxic to non-target organisms than Gazelle (Roubos  
315 *et al.*, 2014) but is currently not used on an operational scale in Great Britain to control pine weevil.  
316 The results were promising. Two treatments, the Gazelle ULV and the Coragen ULV applied  
317 insecticide at much lower volumes. The lower volume applications were as effective as those applied  
318 at conventional volumes.

319 Across all trials, using larger planting stock did not reduce damage and mortality. Thorsen *et al.*  
320 (2001) found that larger stock with a root collar diameter of 8mm on scarified sites and 10mm on  
321 other sites showed negligible damage in Scandinavia. The root collar diameter of the larger planting  
322 stock employed in this trial was at 8 to 8.5mm. Populations of pine weevil are higher in the UK than  
323 in Scandinavia (Willoughby *et al.*, 2017) and it may be that using larger trees would be an effective  
324 strategy at lower population densities, if this could reliably be predicted. However, using larger trees  
325 was not a successful control measure at the two trials with low overall damage and mortality and

326 use of the seedling planting stock; Newton Stewart and Ramshaw Rig and so based on these data it  
327 is not an effective strategy to reduce mortality and damage.

328 Applying controlled release fertiliser did not influence death and damage from pine weevil,  
329 compared to the same sized planting stock without fertiliser applied. This contrasts with the findings  
330 of Zas et al. (2006) who found fertiliser application increased damage by pine weevil.

331 The lack of complete protection from any of the treatments when under high levels of damage, such  
332 as at Auchencairn is an encouragement for the development of an integrated pest management  
333 system for pine weevil. There are methods for predicting the population size of pine weevil, such as  
334 that devised by Nordlander *et al.*, (2017) in Sweden and the Hylobius Management Support System  
335 (Willoughby *et al.*, 2017) from the UK but these will need to be combined with an effective means of  
336 protection at high population densities.

337

## 338 5. Conclusions

339 The five trials tested the success of fourteen treatments against a control in reducing damage and  
340 mortality from pine weevil. While there were significant differences in the damage and mortality  
341 caused by pine weevil across the sites, overall the following conclusions can be made:

342 At these trials, insecticides and physical barriers were equally effective. However, physical barriers  
343 have been known to be less effective when used in certain site conditions.

344 Ultra low volume insecticide treatments were as effective as using higher volume applications.

345 The new insecticide Coragen was as effective as Gazelle and alpha cypermethrin.

346 Planting larger trees did not reduce mortality and damage.

347 Application of fertiliser in the field did not reduce mortality and damage.



348 At Auchencairn, where there were extreme levels of damage and mortality, none of the treatments  
349 were effective and this supports the need to develop an integrated solution to pine weevil.

350

## 351 Acknowledgements

352

353 The authors would like to thank the many people and organisations who have collaborated in the  
354 establishment and assessment of the pine weevil trials. Trial sites were kindly provided by Tim  
355 Liddon at Tillhill Forestry (Auchencairn, Lamloch and Ramshaw Rig) and Stuart Wilkie of Scottish  
356 Woodlands Ltd (Cwym Henog and Newton Stewart). Planting and assessment was undertaken by  
357 the following people; Kerstin Leslie and Ben Wallbank of Tillhill Forestry Ltd (Auchencairn, Lamloch,  
358 Ramshaw Rig), Andrew Maclachlan and Euan Wilkie of Tillhill Forestry Ltd (Newton Stewart) and  
359 Charles Gittins of Flintshire woodlands Ltd (Cwym Henog) and Flintshire Woodlands Ltd. Finally , we  
360 would also like to acknowledge the input of colleagues at Maelor Forest Nurseries Ltd. These include  
361 Mike Harvey, the Managing Director of Maelor Forest Nurseries Ltd for general support, Steve  
362 Smout and Damien McGrade for chemical treatment of the trees and Lorenza Pozzi for data  
363 collation, organisation and management. Finally, the authors would like to thank the anonymous  
364 referees for constructive feedback on the draft manuscript.

## 365 References

366

367 European Commission (2019) Report from the Commission to the European Parliament and the  
368 Council on Member State National Action Plans and on progress in the implementation of Directive  
369 2009/128/EC on the sustainable use of pesticides. European Commission, Brussels.

370 Evans H, Moore R, Heritage S, Wainhouse D (2003). Developments in the integrated management of  
371 pine weevil, a pest of restocking in conifer plantations. Forest Research annual report and  
372 accounts, 2004: 78-87.

373 Forestry Commission (2019) Forestry Statistics 2019, A compendium of statistics about woodland,  
374 forestry and primary wood processing in the United Kingdom [online 24 February 2020] URL:  
375 [https://www.forestresearch.gov.uk/documents/7435/Complete\\_FS2019\\_zluGlog.pdf](https://www.forestresearch.gov.uk/documents/7435/Complete_FS2019_zluGlog.pdf)

376 Forest Research (2019). Hylobius Management Support System (MSS) [online 14 June 2019] URL:  
377 [https://www.forestresearch.gov.uk/tools-and-resources/tree-health-and-protection-](https://www.forestresearch.gov.uk/tools-and-resources/tree-health-and-protection-services/hylobius-management-support-system/)  
378 [services/hylobius-management-support-system/](https://www.forestresearch.gov.uk/tools-and-resources/tree-health-and-protection-services/hylobius-management-support-system/)

379 Forest Research (2020) Ecological Site Classification 4 [online 4 June 2020] URL:  
380 <http://www.forestdss.org.uk/geoforestdss/>

381 FSC (2019) FSC Pesticides Policy, FSC-POL-30-001 V3-0 EN. Forest Stewardship Council, Bonn.

382 Harlin C and Eriksson S (2010) Mekaniska plantskydd mot snytbaggeskador, anlagt 2007. Report 1 -  
383 Swedish University of Agricultural Sciences.

384 Heritage S, Moore R (2000). The assessment of site characteristics as part of a management strategy  
385 to reduce damage by *Hylobius*. Forestry Commission Information Note 38, Forestry Commission,  
386 Edinburgh, UK, pp.6.

387 Inward DJ, Wainhouse D, Peace A (2012). The effect of temperature on the development and life  
388 cycle regulation of the pine weevil *Hylobius abietis* and the potential impacts of climate  
389 change. Agricultural and Forest Entomology, 14(4): 348-357.

390 Kennedy S, Cameron A, Thoss V, Wilson M (2006). Role of monoterpenes in *Hylobius abietis* damage  
391 levels between cuttings and seedlings of *Picea sitchensis*. Scandinavian Journal of Forest  
392 Research 21(4): 340-344.

393 Leather SR, Day KR, Salisbury AN (1999). The biology and ecology of the large pine weevil, *Hylobius*  
394 *abietis* (Coleoptera: Curculionidae): a problem of dispersal? Bulletin of Entomological Research 89  
395 (1): 3-16.

396 Leslie K, Liddon T (2017) An integrated pest management strategy for *Hylobius* - the holy grail of  
397 forestry? Forest and Timber News, April 2017: 44-45.

398 López-Villamor A, Carreño S, López-Goldar X, Suárez-Vidal E, Sampedro L, Nordlander G, Björklund N,  
399 Zas R (2019). Risk of damage by the pine weevil *Hylobius abietis* in southern Europe: Effects of  
400 silvicultural and landscape factors. Forest Ecology and Management 444: 290-298.

401 Lundborg L, Fedderwitz F, Björklund N, Nordlander G, Borg-Karlson AK (2016). Induced defences  
402 change the chemical composition of pine seedlings and influence meal properties of the pine weevil  
403 *Hylobius abietis*. Phytochemistry 130: 99-105.

404 Luoranen J, Viiri H (2005). Insecticides sprayed on seedlings of *Picea abies* during active growth:  
405 Damage to plants and effect on pine weevils in bioassay. Scandinavian Journal of Forest  
406 Research, 20(1): 47-53.

407 McDonald JH (2015). Multiple Comparisons, Handbook of Biological Statistics.[online 14 June 2019]  
408 URL: <http://www.biostathandbook.com/multiplecomparisons.html>

409 Moore R (2004). Managing the threat to restocking posed by the large pine weevil, *Hylobius abietis*:  
410 The importance of time of felling of spruce stands. Forestry Commission Information Note 61.  
411 Forestry Commission, Edinburgh, UK, pp 4.

412 Nordlander G, Bylund H, Björklund N (2005). Soil type and microtopography influencing feeding  
413 above and below ground by the pine weevil *Hylobius abietis*. Agricultural and Forest  
414 Entomology, 7(2):107-113.

415 Nordlander G, Nordenhem H, Hellqvist C (2009). A flexible sand coating (Conniflex) for the  
416 protection of conifer seedlings against damage by the pine weevil *Hylobius abietis*. Agricultural and  
417 Forest Entomology, 11(1): 91-100.

418 Nordlander G, Hellqvist C, Johansson K, Nordenhem H (2011). Regeneration of European boreal  
419 forests: effectiveness of measures against seedling mortality caused by the pine weevil *Hylobius*  
420 *abietis*. Forest Ecology and Management, 262 (12): 2354-2363.

421 Nordlander G, Mason EG, Hjelm K, Nordenhem H, Hellqvist C (2017). Influence of climate and forest  
422 management on damage risk by the pine weevil *Hylobius abietis* in northern Sweden. Silva Fennica  
423 51 (5): 1-20.

424 Norsk Wax (2016) Substitution of insecticides with wax. [online 14 June 2019] URL:  
425 <http://www.metla.fi/tapahtumat/2016/taimitarhapaivat/Pettersen.pdf>

426 Ohrn P, Nordlander G (2015) WeevilSTOP -Field activities in Sweden 2015 [online 20 June 2019] URL:  
427 [https://irp-](https://irp-cdn.multiscreensite.com/9a5e28a2/files/uploaded/WeevilStop.field%20activities.2015.pptx)  
428 [cdn.multiscreensite.com/9a5e28a2/files/uploaded/WeevilStop.field%20activities.2015.pptx](https://irp-cdn.multiscreensite.com/9a5e28a2/files/uploaded/WeevilStop.field%20activities.2015.pptx)

429 Örländer G and Nordlander G (2003). Effects of field vegetation control on pine weevil (*Hylobius*  
430 *abietis*) damage to newly planted Norway spruce seedlings. Annals of Forest Science, 60 (7): 667-  
431 671.

432 Petersson M, Örländer G, Nilsson U (2004). Feeding barriers to reduce damage by pine weevil  
433 (*Hylobius abietis*). Scandinavian Journal of Forest Research, 19 (1): 48-59.

434 Petersson M, Eriksson S, Zettergvist F (2006) Mekaniska plantskydd mot snytbaggskador, anlagt  
435 2003 -slutrapport. Sveriges Lantbruksuniversitet [online 25 July 2019] URL:  
436 <https://www.yumpu.com/sv/document/read/30822748/slutrapport-pdf>

437 Pettersson M, Kännaste A, Lindström A, Hellqvist C, Stattin E, Långström B, Borg-Karlson AK (2008).  
438 Mini-seedlings of *Picea abies* are less attacked by *Hylobius abietis* than conventional ones: is plant  
439 chemistry the explanation? *Scandinavian Journal of Forest Research*, 23(4): 299-306.

440 Rose D, Leather SR, Matthews GA (2005). Recognition and avoidance of insecticide-treated Scots  
441 Pine (*Pinus sylvestris*) by *Hylobius abietis* (Coleoptera: Curculionidae): implications for pest  
442 management strategies. *Agricultural and Forest Entomology*, 7(3): 187-191.

443 Svenska-Skogsplantor (2019) Svenska Skogsplantor to Stop Using Chemical Pine Weevil Treatments  
444 [online 17 January 2019] [https://news.cision.com/svenska-skogsplantor/r/svenska-skogsplantor-to-](https://news.cision.com/svenska-skogsplantor/r/svenska-skogsplantor-to-stop-using-chemical-pine-weevil-treatments,c2789643)  
445 [stop-using-chemical-pine-weevil-treatments,c2789643](https://news.cision.com/svenska-skogsplantor/r/svenska-skogsplantor-to-stop-using-chemical-pine-weevil-treatments,c2789643)

446 Sveriges Lantbruksuniversitet (2018) Stem coatings on conifer plants replace insecticides [online 16  
447 January 2020] <http://snyttagge.slu.se/stem-coatings-on-conifer-plants-replace-insecticides.php>

448 The Advisory Service for Weed & Pest Control in Forestry & Amenity (2018) HIFIC Presentation -  
449 Inverness 15th March 2018 [online 29 May 2020]  
450 <http://www.forestryscotland.com/media/377766/weeds,%20weevils%20and%20worries.pdf>

451 Thorsen ÅA, Mattsson S, Weslien J (2001) Influence of stem diameter on the survival and growth of  
452 containerized Norway spruce seedlings attacked by pine weevils (*Hylobius* spp.). *Scandinavian*  
453 *Journal of Forest Research*, 16(1):54-66.

454 UKWAS (2018) The UK Woodland Assurance Standard [online 13 March 2020] <http://ukwas.org.uk/>

455 Unelius C., Bohman B and Nordlander G (2018). Comparison of Phenylacetates with Benzoates and  
456 Phenylpropanoates as Antifeedants for the Pine Weevil, *Hylobius abietis*. *Journal of Agricultural and*  
457 *Food Chemistry*, 66(44):11797-11805.

458 Wainhouse D, Brough S, Greenacre B (2007). Managing the pine weevil on lowland pine. Forestry  
459 Commission. Forestry Commission Practice Note 14. Forestry Commission, Edinburgh, UK. pp.12.

460 Wallertz K, Björklund N, Hjelm K, Petersson M, Sundblad L-G (2018). Comparison of different site  
461 preparation techniques: quality of planting spots, seedling growth and pine weevil damage. *New*  
462 *Forests* 49: 705-722.

463 Williams CD, Dillon AB, Harvey CD, Hennessy R, Mc Namara L, Griffin CT (2013). Control of a major  
464 pest of forestry, *Hylobius abietis*, with entomopathogenic nematodes and fungi using eradicator and  
465 prophylactic strategies. *Forest Ecology and Management*, 305: 212-222.

466 Willoughby I, Evans H, Gibbs J, Pepper H, Gregory S, Dewar J, Nisbet T, Pratt J, McKay H, Siddons R,  
467 Mayle, B (2004). Reducing pesticide use in forestry: practice guide. Reducing pesticide use in  
468 forestry: practice guide. Forestry commission, Edinburgh, UK. pp138.

469 Willoughby I, Moore R, Nisbet T (2017). Interim guidance on the integrated management of *Hylobius*  
470 *abietis* in UK forestry. Forest Research, Farnham, UK. pp 28.

471 Zas R, Björklund N, Nordlander G, Cendán C, Hellqvist C, Sampedro L (2014) Exploiting jasmonate-  
472 induced responses for field protection of conifer seedlings against a major forest pest, *Hylobius*  
473 *abietis*. *Forest Ecology and Management* 313: 212-223.

474 Zas R, Björklund N, Sampedro L, Hellqvist C, Karlsson B, Jansson S, Nordlander G (2017). Genetic  
475 variation in resistance of Norway spruce seedlings to damage by the pine weevil *Hylobius*  
476 *abietis*. *Tree Genetics & Genomes*, 13(5): 111-123.

477 Zas R, Sampedro, L, Prada E, Lombardero MJ and Fernández-López J (2006). Fertilization increases  
478 *Hylobius abietis* L. damage in *Pinus pinaster* Ait. seedlings. *Forest Ecology and Management*, 222(1-  
479 3): 137-144.

480

481 **Table 1** Location of trials. Climatic data and suitability for Sitka spruce generated by the Ecological  
 482 Site Classification v4 (Forest Research 2020). AT = Accumulated temperature above 5°C, MD =  
 483 moisture deficit and DAMS = exposure index where 3 is sheltered and 22+ is too exposed for  
 484 productive forestry.

| Site           | Latitude         | Longitude         | AT (day degrees above 5°C) | MD (mm) | DAMS | Species suitability |
|----------------|------------------|-------------------|----------------------------|---------|------|---------------------|
| Auchencairn    | 55° 13' 50.82" N | 003° 39' 14.65" W | 1103                       | 75      | 17   | Suitable            |
| Cwm Henog      | 52° 07' 12.99" N | 003° 43' 55.94" W | 1057                       | 54      | 17   | Suitable            |
| Lamloch        | 55° 14' 41.17" N | 004° 20' 14.62" W | 1179                       | 79      | 16   | Suitable            |
| Newton Stewart | 55° 01' 16.79" N | 004° 40' 06.86" W | 1387                       | 107     | 17   | Marginal            |
| Ramshaw Rig    | 55° 15' 55.83" N | 003° 18' 39.04" W | 1022                       | 66      | 15   | Marginal            |

485

486 **Table 2** Description of trials addressing the main factors influencing pine weevil damage as described  
 487 in Heritage and Moore (2000).

| Site name                                      | Auchencairn              | Cwym Henog                    | Lamloch                 | Newton Stewart     | Ramshaw Rig              |
|--|--------------------------|-------------------------------|-------------------------|--------------------|--------------------------|
| <b>Site management</b>                         |                          |                               |                         |                    |                          |
| Month site was clearfelled                     | February 2016-March 2017 | April - July 2015             | May 2015- November 2015 | March- August 2016 | July 2016- December 2016 |
| Period between clearfell and planting (months) | 0-11                     | 18-21                         | 17-23                   | 8-13               | 4-8                      |
| Exposed mineral soil around plants? (Y/N)      | Yes                      | Yes                           | Yes                     | Yes                | Yes                      |
| Site burned after clearfell? (Y/N)             | No                       | No                            | No                      | No                 | No                       |
| Stumps removed or destroyed? (Y/N)             | No                       | No                            | No                      | No                 | No                       |
| Vegetation, woody? (light/moderate/heavy)      | light                    | Light regen of willow / rowan | moderate                | Light              | light                    |
| Vegetation grasses? (light/moderate/heavy)     | heavy                    | Light – moderate              | light                   | Moderate           | light                    |
| <b>Plant Details</b>                           |                          |                               |                         |                    |                          |
| Nursery code                                   | PSI A12                  | PSI A12                       | PF80                    | PSI A12            | PSI A12                  |
| Type   | seedling                 | seedling                      | vegetatively propagated | seedling           | Seedling                 |
| Size   | 30-50                    | 30-50                         | 25-50                   | 30-50              | 30-50                    |

488

489

490 **Table 3** Details of treatments (\* based on cost of Norsk Wax, a similar coating (Norsk wax 2016)).


| Treatment name          | Details   | Additional Cost per tree |
|-------------------------|---|--------------------------|
| Control                 | No treatment applied to the tree.   | None                     |
| Electrodyne             | alpha-cypermethrin – applied by electrodyne   | £0.11                    |
| Gazelle spray           | Nisso Chemical Europe GmbH Gazelle - Applied by knapsack sprayer.   | £0.11                    |
| Gazelle ULV             | New ULV nozzles, improved machine.  | £0.11                    |
| Coragen spray           | Dupont coragen - Applied by knapsack sprayer.   | £0.11                    |
| Coragen ULV             | New ULV nozzles, improved machine.  | £0.11                    |
| Multipro Sleeves        | Multipro cardboard sleeve fixed around the stem of the tree before planting.  | £0.30                    |
| Biosleeves              | Biosleeve (biodegradable plastic) fitted around the stem of the tree before planting.   | £0.35                    |
| Polymer barrier 50 %    | A new polymer applied to 50% of the lower length of the stem.   | £0.06*                   |
| Polymer barrier 85 %    | A new polymer applied to 85% of the lower length of the stem.   | £0.06*                   |
| 1+1 stock + CRF         | 1+1 stock, 10g Treeboost controlled release fertiliser applied to roots before heeling in.                                      | £0.05                    |
| 2+1 stock               | 2+1 stock (greater girth)   | £0.06                    |
| 2+1 stock + CRF         | 2+1 stock, 10g Treeboost controlled release fertiliser applied to roots before heeling in.                                      | £0.11                    |
| 50-70cm 2+1 stock       | 50-70cm tall, 2+1 stock (greater height + girth).   | £0.06                    |
| 50-70cm 2+1 stock + CRF | 50-70cm tall, 2+1 stock (greater height + girth)10g Treeboost controlled release fertiliser applied to roots before heeling in. | £0.11                    |

491


492

493






496 **Figure 1** Percentage of trees in each of the five damage categories in each of the trials.




498 **Figure 2** Percentage of trees in each of the five damage categories for each treatment.



499


500 **Figure 3** Median percentage mortality and damage by trial. The same letter above the bars denote  
 501 **groupings of sites** with no significant difference. Boxplots show the median values as the dark  
 502 horizontal lines; 25th and 75th percentiles as the top and bottom of the boxes. The vertical lines  
 503 show the maximum and minimum values, excluding outliers shown as a circle.

(a) Cwym Henog




504

(b) Lamloch




505

(c) Newton Stewart




506

(d) Ramshaw Rig




507



508

509 **Figure 4** Median percentage mortality and damage by treatment. The same letter above the bars  
 510 denote groupings of treatments with no significant difference. Boxplots show the median values as  
 511 the dark horizontal lines; 25th and 75th percentiles as the top and bottom of the boxes. The vertical  
 512 lines show the maximum and minimum values, excluding outliers shown as a circle.

513



514

515 **Figure 5** Comparison of treatments by cost (£) and median percentage damage. 1=control,  
 516 2=Electrodyne, 3=Gazelle spray, 4=Gazelle ULV, 5=Coragen spray, 6=Coragen ULV, 7=MultiPro  
 517 sleeves, 8=Polymer barrier 50%, 9=Polymer barrier 85%, 10=Biosleeves, 11=1+1 stock + CRF, 12=2+1  
 518 stock, 13=2+1 stock+CRF, 14=50-70 2+1 stock and 15=50-70 2+1 stock+CRF

519