



CASE REPORT

Acta Orthop Traumatol Turc 2014;48(4):459-462
doi: 10.3944/AOTT.2014.3107

Endovascular treatment of an iliac artery rupture caused by invasive *Salmonella* spondylodiscitis

Ebru KURŞUN¹, Tuba TURUNÇ¹, Uğur ÖZKAN², Yusuf Ziya DEMİROĞLU¹¹Department of Infectious Diseases, Adana Practice and Research Center, Faculty of Medicine, Başkent University, Adana, Turkey;²Department of Radiology, Adana Practice and Research Center, Faculty of Medicine, Başkent University, Adana, Turkey

The rate of non-typhoid *Salmonella* infections has increased remarkably in recent years. Endovascular system infection is one of the most serious forms of extraintestinal *Salmonella* infection. The abdominal aorta is frequently involved, while bone and joint involvement are rarely seen. We present a rare case of successful endovascular treatment of a left iliac artery rupture and pseudoaneurysm both occurring due to the direct invasion of lumbar spondylodiscitis caused by *Salmonella typhimurium*.

Key words: Endovascular treatment; pseudoaneurysm; *Salmonella typhimurium*; spondylodiscitis; stent graft.

The rate of non-typhoid *Salmonella* infections has gradually increased worldwide. *Salmonella* bacteria usually originate from food and cause gastroenteritis, bacteremia and focal infections.^[1] Advanced age, changes in endogenous intestinal flora, diabetes, cancer, autoimmune disorders, HIV infection, and therapeutic immunosuppression are risk factors for non-typhoid salmonellosis and bacteremia.^[2] Endovascular system infection is one of the most serious forms of extraintestinal infection and endovascular *Salmonella* infections usually result in mycotic abdominal aortic aneurysms.^[3] While successful endovascular treatment of *Salmonella* mycotic aneurysms has been reported,^[4-9] endovascular treatment of iliac artery ruptures and pseudoaneurysms caused by *Salmonella* invasion have not.

We present a rare case in which an iliac artery rupture caused by *S. typhimurium* spondylodiscitis received successful endovascular treatment.

Case report

A 57-year-old hypertensive and diabetic male patient presented to our emergency service with walking difficulty. The patient had experienced back pain and numbness in his left leg for 15 days. His anamnesis revealed that he had used various antibiotics due to high fever. During follow-up treatment for a urinary infection in another center, an aortic aneurysm was detected on his abdominal ultrasonography and he was transferred to our hospital.

On physical examination, the patient's body temperature was 38°C, blood pressure was 130/70 mmHg and pulse was 112 beats/min. His overall condition was stable and he was conscious, cooperative and oriented. Paresthesia was observed in the left lower extremity and no pulse was present. Laboratory results were as follows; white blood cell count: 10,700/mm³; hemoglobin: 9.0 g/dL; thrombocyte count: 359,000/mm³; C-reactive

Correspondence: Ebru Kursun, MD. Başkent Üniversitesi Hastanesi Dadaloğlu Mah. Serinevler 2591 Sk. 4/A, 01250, Yüreğir, Adana, Turkey.

Tel: +90 533 – 268 10 68 e-mail: ebru.kursun@hotmail.com

Submitted: November 11, 2012 **Accepted:** June 10, 2013

©2014 Turkish Association of Orthopaedics and Traumatology

Available online at
www.aott.org.tr

doi: 10.3944/AOTT.2014.3107

QR (Quick Response) Code






Fig. 1. (a) Abdominal contrast-enhanced CT image shows left iliac artery rupture and pseudoaneurysm (arrow). (b) Lumbar contrast-enhanced MR image shows contrast enhancement of corpus of L4 and L5 vertebrae due to spondylodiscitis (arrows).

protein: 205 mg/L; erythrocyte sedimentation rate: 72 mm/hour; and creatinine: 0.7 mg/dL. Abdominal contrast-enhanced computed tomography images revealed a ruptured left main iliac artery and a 6x10 cm left retroperitoneal pseudoaneurysm (Fig. 1a). Lumbar magnetic resonance imaging (MRI) was performed on the patient to confirm spondylodiscitis. The lumbar MRI revealed pathological signal change and contrast uptake on the anterior aspect of the L4 and L5 vertebral corpus and an approximately 10x6 cm hematoma in the left psoas muscle (Fig. 1b).

Implantation of a stent graft was performed in the angiography unit by interventional radiologists. Single-wall punctures of the bilateral common femoral arteries (CFA) were performed with an 18-G needle under ultrasound guidance. After the first puncture, 6-F vascular sheaths were inserted. The patient received a bolus of 5000 IU of heparin, and diagnostic angiograms were obtained. After placing the sutures of the Prostar XL closure devices (Abbott Vascular, Redwood, CA, USA), large vascular sheaths, compatible with the stent graft system, were introduced into the common femoral arter-




Fig. 2. (a) Preoperative abdominal aortography reveals left iliac artery rupture and pseudoaneurysm. (b) After implantation of aortoiliac stent graft, abdominal aortography shows successful treatment of left iliac artery rupture and pseudoaneurysm.

ies. The left iliac artery rupture and related pseudoaneurysm were successfully closed by placing a bifurcated aortoiliac stent graft (The Gore® Excluder®, W.L. Gore & Associates, Inc., Flagstaff, AZ, USA). After deployment of the stent graft system, the suture knot was prepared. Preoperative and postoperative angiographies are shown in Figure 2. A drainage catheter was placed into the pseudoaneurysm in the same session.

Empirical treatment with intravenous teicoplanin was commenced at a dose of 800 mg/day after the hematoma and blood cultures were obtained and maintained at 400 mg/day in combination with a 3 g/day dose of intravenous meropenem. The hematoma and blood cultures revealed *S. typhimurium* growth. Antibiotic treatment was halted in response to culture antibiogram results and treatment with intravenous ciprofloxacin was applied at a dose of 2x400 mg. A consultation was requested from the physical therapy and rehabilitation department and the patient was enrolled in an exercise program to build strength in the left leg and improve his ability to walk. By the seventh day of his hospital stay, the patient's condition improved and body temperature decreased and he was discharged with 750 mg ciprofloxacin tablets to be taken twice daily. The patient was kept on antibiotics for six months. After five months, a lumbar MRI was performed which confirmed a considerable decrease in the dimensions of the hematoma, now approximately 83x47 mm in the left psoas muscle. A CT scan showed that the aortoiliac stent graft was patent.

Discussion

Non-typhoid *Salmonella* bacteria originate from contaminated food and cause gastroenteritis, bacteremia and focal infections.^[1] The prevalence of these bacteria has been gradually increasing worldwide. Local bone infections account for less than 1% of *Salmonella* infections. According to the literature, spondylodiscitis is a rare local manifestation of *Salmonella* infection.^[10-15] However, *S. typhimurium* should still be considered as a possible causative agent in bone and joint infections. Local *Salmonella* infections are rarely seen in those with normal immune systems. Advanced age, changes in endogenous intestinal flora, sickle cell anemia, diabetes mellitus, cancer, autoimmune disorders, HIV infection, and therapeutic immunosuppression are the main risk factors for non-typhoid salmonellosis and bacteremia.^[2,10] In the current case, the patient's sole risk factor was diabetes mellitus.

Endovascular system infection is one of the most serious forms of extraintestinal *Salmonella* infection and frequently involves the abdominal aorta.^[3] *Salmonella* is a common causative bacteria for inducing infectious aortic

aneurysm and spondylodiscitis at the contiguous spine.^[16] When the spine or aorta is the first to be infected, the septic embolus settles in the vasa vasorum and the infection travels along the outside of the neighboring blood vessels to the neighboring organs. Infectious aortic aneurysms and spondylodiscitis occur simultaneously in many cases.^[13] When *Salmonella* cases with vascular involvement were analyzed, it was observed that antibiotherapy was generally commenced together with surgical intervention. However, endovascular interventions have also been reported in case reports.^[4-10] Two cases with mycotic aneurysm have been reported in the literature. Mahlfeld et al.^[12] reported three spondylodiscitis cases caused by *S. typhimurium* and detected an abdominal aortic aneurysm in one. Since the other case Mahlfeld et al. reported had undergone an operation due to an aortic aneurysm three years prior, either the aneurysm or the spine could have been the source of infection. Learch et al.^[17] reported a case of spondylodiscitis associated with an abdominal aortic aneurysm and paravertebral abscess caused by *S. enteritidis*. The patient in their case underwent antibiotherapy and exploratory laparotomy with mycotic aneurysm resection, closure of the proximal aorta and common iliac arteries and insertion of an axillary bifemoral bypass.

The present case is the first reported case of a *S. typhimurium* infection causing spondylodiscitis and iliac artery rupture. Our patient had diabetes and hypertension, which may have led to vascular pathologies. Unlike the surgical interventions performed on a few number of cases with iliac artery involvement in the literature, our patient underwent successful endovascular intervention. Weber and Pirovino reported successful surgical intervention and antibiotic treatment of a right iliac artery aneurysm caused by *S. typhimurium*.^[18] Ben et al.^[19] reported an aneurysm of the right common iliac artery infected by *Salmonella* group C1. Antibiotic treatment and surgical intervention with an aneurysmectomy and suprapubic crossover femorofemoral artery graft resulted in a good recovery and outcome. Despite limited experience, antibiotherapy together with endovascular intervention might provide a lower-risk alternative to conventional surgery while similarly retaining normal vascular anatomy. We are of the opinion that the present report contributes to the literature a unique and intriguing case.

In conclusion, *S. typhimurium* may result in spondylodiscitis and vascular involvement and should be kept in mind as a causative agent given such clinical conditions. Medical treatment combined with endovascular stent grafts may serve as an alternative option to surgical treatment.

Conflicts of Interest: No conflicts declared.

References

1. Hohmann EL. Nontyphoidal salmonellosis. *Clin Infect Dis* 2001;32:263-9.
2. Hsu RB, Tsay YG, Chen RJ, Chu SH. Risk factors for primary bacteremia and endovascular infection in patients without acquired immunodeficiency syndrome who have nontyphoid salmonellosis. *Clin Infect Dis* 2003;36:829-34.
3. Drinković D, Taylor SL, Lang S. Five cases of non-typhoidal *Salmonella* endovascular infection. *Intern Med J* 2004;34:641-5.
4. Berchtold C, Eibl C, Seelig MH, Jakob P, Schönleben K. Endovascular treatment and complete regression of an infected abdominal aortic aneurysm. *J Endovasc Ther* 2002;9:543-8.
5. Koeppel TA, Gahlen J, Diehl S, Prosst RL, Dueber C. Mycotic aneurysm of the abdominal aorta with retroperitoneal abscess: successful endovascular repair. *J Vasc Surg* 2004;40:164-6.
6. Ting AC, Cheng SW, Ho P, Poon JT. Endovascular repair for multiple *Salmonella* mycotic aneurysms of the thoracic aorta presenting with Cardiovascular syndrome. *Eur J Cardiothorac Surg* 2004;26:221-4.
7. Forbes TL, Harding GE. Endovascular repair of *Salmonella*-infected abdominal aortic aneurysms: a word of caution. *J Vasc Surg* 2006;44:198-200.
8. Kam MH, Toh LK, Tan SG, Wong D, Chia KH. A case report of endovascular stenting in *Salmonella* mycotic aneurysm: a successful procedure in an immunocompromised patient. *Ann Acad Med Singapore* 2007;36:1028-31.
9. Hartman V, Jiang H, Thomas B. Successful [Successful] endovascular repair of an abdominal mycotic aneurysm. A case report. *Acta Chir Belg* 2009;109:788-90.
10. Hook EW. *Salmonella* species (including typhoid fever). In: Mandel GL, Douglas RG, Bennett JE, editors. Principles and practice of infectious diseases. Edinburgh: Churchill Livingstone; 1990. p. 1700-10.
11. Tautenhahn J, Fahlke J, Halloul Z, Grasshoff H, Bürger T. Single intervention for treatment of *Salmonella* typhimurium-induced symptomatic abdominal aortic aneurysm with spondylitis. [Article in German] *Vasa* 2001;30:293-6. [Abstract]
12. Mahlfeld K, Franke J, Grasshoff H. Spondylitis due to *Salmonella* typhimurium. [Article in German] *Unfallchirurg* 2003;106:334-8. [Abstract]
13. Chang IC. *Salmonella* spondylodiscitis in patients without sickle cell disease. *Clin Orthop Relat Res* 2005; 430:243-7.
14. Winkelmann W, Schulitz KP. Spondylitis caused by *Salmonella* typhimurium (author's transl). [Article in German] *Z Orthop Ihre Grenzgeb* 1977;115:6-8. [Abstract]
15. Choukroun G, Quint L, Amoura Z, Boudes P, Guillemin L. *Salmonella* typhi murium spondylodiscitis in systemic lupus. [Article in French] *Ann Med Interne (Paris)* 1988;139:446-7. [Abstract]
16. Kim JM, Park YH, Kim D, Park MJ, Ahn HJ, Ryoo BY, Yang SH. *Salmonella* vertebral osteomyelitis with sepsis in healthy adult. *Korean J Med* 2005;69:1003-8.
17. Learch TJ, Sakamoto B, Ling AC, Donovan SM. *Salmonella* spondylodiscitis associated with a mycotic abdominal aortic aneurysm and paravertebral abscess. *Emerg Radiol* 2009;16:147-50.
18. Weber E, Pirovino M. Mycotic aneurysm of the iliac artery caused by *Salmonella* infection. A case report. [Article in German] *Helv Chir Acta* 1992;58:601-7. [Abstract]
19. Ben RJ, Peng MY, Hsu GJ, Chen SG, Lee FY, Chou MY, et al. Infected common iliac aneurysm due to *Salmonella*: a case report. [Article in Chinese] *Zhonghua Yi Xue Za Zhi (Taipei)* 1993;52:62-5. [Abstract]