

## Aberystwyth University

### *Subglacial water drainage, storage, and piracy beneath the Greenland ice sheet*

Lindbäck, K.; Pettersson, R.; Hubbard, Alun; Doyle, Sam; Van As, D.; Mikkelsen, A. B.; Fitzpatrick, Andrew  
Alexander William

*Published in:*  
Geophysical Research Letters

*DOI:*  
[10.1002/2015GL065393](https://doi.org/10.1002/2015GL065393)

*Publication date:*  
2015

*Citation for published version (APA):*

Lindbäck, K., Pettersson, R., Hubbard, A., Doyle, S., Van As, D., Mikkelsen, A. B., & Fitzpatrick, A. A. W. (2015). Subglacial water drainage, storage, and piracy beneath the Greenland ice sheet. *Geophysical Research Letters*, 42(18), 7606-7614. <https://doi.org/10.1002/2015GL065393>

**Document License**  
CC BY-NC

**General rights**

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)



## RESEARCH LETTER

10.1002/2015GL065393

## Key Points:

- We present a high-resolution subglacial hydrological analysis of the Greenland ice sheet
- Small subglacial lakes remain undetectable by methods using surface elevation or radar techniques
- We identify the first evidence for subglacial water piracy beneath the Greenland ice sheet

## Supporting Information:

- Supporting Information S1

## Correspondence to:

K. Lindbäck,  
katrin.lindback@geo.uu.se;  
katrin.lindback@gmail.com

## Citation:

Lindbäck, K., R. Pettersson, A. L. Hubbard, S. H. Doyle, D. van As, A. B. Mikkelsen, and A. A. Fitzpatrick (2015), Subglacial water drainage, storage, and piracy beneath the Greenland ice sheet, *Geophys. Res. Lett.*, 42, doi:10.1002/2015GL065393.

Received 15 JUL 2015

Accepted 24 AUG 2015

Accepted article online 26 AUG 2015

## Subglacial water drainage, storage, and piracy beneath the Greenland ice sheet

K. Lindbäck<sup>1</sup>, R. Pettersson<sup>1</sup>, A. L. Hubbard<sup>2,3</sup>, S. H. Doyle<sup>3</sup>, D. van As<sup>4</sup>, A. B. Mikkelsen<sup>5</sup>, and A. A. Fitzpatrick<sup>3</sup>

<sup>1</sup>Department of Earth Sciences, Air, Water, and Landscape Science, Uppsala University, Uppsala, Sweden, <sup>2</sup>Centre for Arctic Gas Hydrate, Environment and Climate, Department of Geology, Arctic University of Norway, Tromsø, Norway, <sup>3</sup>Centre for Glaciology, Department of Geography and Earth Sciences, Aberystwyth University, Aberystwyth, UK, <sup>4</sup>Geological Survey of Denmark and Greenland, Copenhagen, Denmark, <sup>5</sup>Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark

**Abstract** Meltwater drainage across the surface of the Greenland ice sheet (GrIS) is well constrained by measurements and modeling, yet despite its critical role, knowledge of its transit through the subglacial environment remains limited. Here we present a subglacial hydrological analysis of a land-terminating sector of the GrIS at unprecedented resolution that predicts the routing of surface-derived meltwater once it has entered the basal drainage system. Our analysis indicates the probable existence of small subglacial lakes that remain undetectable by methods using surface elevation change or radar techniques. Furthermore, the analysis suggests transient behavior with rapid switching of subglacial drainage between competing catchments driven by seasonal changes in the basal water pressure. Our findings provide a cautionary note that should be considered in studies that attempt to relate and infer future response from surface temperature, melt, and runoff from point measurements and/or modeling with measurements of proglacial discharge and ice dynamics.

### 1. Introduction

Mass loss from the Greenland ice sheet (GrIS) accelerated over the two last decades, and the GrIS is expected to be the largest cryospheric contributor to global sea level rise in the 21st century [Hanna *et al.*, 2013]. To predict the future contribution of the GrIS to sea level rise, we must understand the physical processes that govern ice sheet mass balance and dynamics. Despite this, the important role that subglacial water routing and storage plays in ice sheet dynamics remains poorly constrained. Several recent studies [e.g., Bartholomew *et al.*, 2010] have argued that hydrological processes on valley glacier systems could be scaled up to serve as analogues for ice sheets. While this analogue appears to hold for the ice sheet margin [e.g., Sundal *et al.*, 2011], the considerable differences in geometry between valley glaciers and ice sheets become pronounced further inland, where the development of efficient subglacial hydrology is hindered by low surface melt rates and gentle bed slopes [Meierbachtol *et al.*, 2013; Rennermalm *et al.*, 2013b; Dow *et al.*, 2014; Doyle *et al.*, 2014]. Direct observations of the basal hydrological system from boreholes or tracing experiments are limited to the lower ablation zone of the GrIS [Smeets *et al.*, 2012; Chandler *et al.*, 2013; Meierbachtol *et al.*, 2013; Andrews *et al.*, 2014; Doyle *et al.*, 2015; van de Wal *et al.*, 2015] while geophysical data covering the whole ice sheet (used for subglacial water routing) [e.g., Livingstone *et al.*, 2013] have only been available at a low spatial resolution (1 km grids of bed elevation data) [Bamber *et al.*, 2013a]. Nevertheless, several studies, which were based on a combination of high-resolution ice surface and low-resolution bed digital elevation models (DEMs), have concluded that there is substantial subglacial or englacial meltwater storage [Rennermalm *et al.*, 2013a; Smith *et al.*, 2015]. The lack of high-resolution bed topography data has also led previous comparative studies of ice surface runoff and proglacial discharge to extrapolate subglacial drainage catchments from ice surface DEMs [e.g., van de Wal and Russell, 1994; Mernild and Hasholt, 2009; Mernild *et al.*, 2010; Bartholomew *et al.*, 2011; Palmer *et al.*, 2011; van As *et al.*, 2012; Chandler *et al.*, 2013; Cowton *et al.*, 2013; Fitzpatrick *et al.*, 2014]. Yet when water leaves the supra-glacial system, its pathway and rates of flow to the ice sheet margin is largely unknown and the water does not necessarily follow the ice surface topography.

Hydraulic potential analysis enables subglacial drainage to be estimated from ice thickness and basal topography on spatial scales where the main control of the drainage is the geometry of the ice sheet [Shreve, 1972]. This method has been applied in Greenland [e.g., Lewis and Smith, 2009; Bamber *et al.*, 2013b; Banwell *et al.*, 2013;


*Livingstone et al., 2013; Karlsson and Dahl-Jensen, 2015*] and Antarctica [e.g., *Wingham et al., 2006; Wright and Siegert, 2012; Wolovick et al., 2013*] to study subglacial water flow and to identify the locations of subglacial lakes. Antarctic subglacial hydrological flow paths have shown to be highly sensitive to changes in ice surface elevation and may also exhibit highly unstable conditions [*Fricke et al., 2007; Wright et al., 2008; Allison et al., 2009*] depending on the subglacial water pressure regime. In some cases, ice flow has been observed to switch on and off owing to water competition between adjacent ice flow units, a behavior termed water piracy [*Anandakrishnan and Alley, 1997; Vaughan et al., 2008; Carter et al., 2013*]. Water piracy has not yet been considered in Greenland; this is surprising given its important role in marginal ice flow dynamics. Variations in subglacial water pressures are often large in Greenland and are caused by the seasonal evolution of the subglacial drainage system [*Meierbachtol et al., 2013; Andrews et al., 2014; van de Wal et al., 2015*] and short-term variations in surface melt runoff [*van de Wal et al., 2008; Bartholomew et al., 2012*]. Hence, we expect that water piracy may also have an effect on subglacial water routing beneath the GrIS. Here we apply hydraulic potential analysis to the Kangerlussuaq sector of the GrIS. We use high-resolution DEMs to (1) compare modeled ice surface runoff with measured proglacial discharge and (2) test the hypothesis that subglacial water piracy exerts an important control on GrIS dynamics and proglacial discharge.

## 2. Data and Methods

We applied hydraulic potential calculations to a  $\sim 8500 \text{ km}^2$  land-terminating area of the Kangerlussuaq sector of the GrIS to investigate subglacial water routing and storage. Hydraulic potential is a steady state proxy for routing of subglacial water [*Shreve, 1972*] and is the sum of the pressure potential from the overburden ice and the elevation potential. We used a 250 m resolution ice thickness DEM from the compiled radar data sets of *Lindbäck et al.* [2014], supplemented with the mass conservation model of *Morlighem et al.* [2014] at 150 m resolution. We calculated the bed elevation from the latter by subtracting the ice thickness from the GrIS Mapping Project (GIMP) surface elevation model at 30 m resolution [*Howat et al., 2014*]. To capture sufficient topographic detail and accuracy for our analysis, we used a DEM with a resolution of 150 m, which represents a compromise between the differing horizontal resolutions of the data sets. We used a single-direction flow algorithm and surface analysis (see Text S1 in the supporting information) to derive water flow networks, sinks, and drainage catchments on the hydraulic potential surface. Our calculated subglacial drainage pathways and sinks were compared with previous water drainage catchment delineations [*Mernild and Hasholt, 2009; Bartholomew et al., 2011; Fitzpatrick et al., 2014; Smith et al., 2015*], supraglacial lake locations [*Fitzpatrick et al., 2014*], supraglacial river locations [*Smith et al., 2015*], and ice flow velocity [*Joughin et al., 2010*]. Using our subglacial drainage catchment delineations, we have calculated ice sheet surface runoff (melt minus refreezing in snow and firn) using an updated version of the *van As et al.* [2012] surface energy budget model. The runoff was compared to discharge gauged using different techniques for both the Isortoq River and the Watson River. Discharge data from the Isortoq River were measured by *Smith et al.* [2015] using field-calibrated time-lapse photography of braid plain inundation area and a power law correlation between water surface area and discharge [*Gleason et al., 2015*]. The discharge of the Watson River was measured using observations of river stage and rating curves based on the method of *Hasholt et al.* [2013]. We investigated changes in subglacial catchments by varying the subglacial water pressure between 50 and 110% of ice overburden pressure. Further details of the methods employed are given in the supporting information Text S1.

## 3. Results

Using hydraulic potential we have characterized subglacial catchments, flow networks, and sinks up to 1750 m above sea level (asl),  $\sim 120 \text{ km}$  from the ice margin (Figure 1). Our study area contains two main subglacial catchments that extend up to the ice divide (Table 1, Figure 1, and supporting information Figure S1), which we refer to as the (1) Isunnguata Sermia and (2) Kangerlussuaq catchments. The Isunnguata Sermia catchment drains into the Isortoq River. The Kangerlussuaq catchment consists of three subcatchments: (2a) the Sandflugtsdalen subcatchment, with the Russell and Leverett outlet glaciers; (2b) the Ørkendalen subcatchment, with the Ørkendalen and Isorlersuup outlet glaciers; and (2c) the Point 660 subcatchment. The rivers emerging from these three subcatchments converge into the Watson River. The calculated subglacial water networks show a dendritic pattern controlled largely by basal topography and which flows from the interior of the ice sheet toward the margin (Figure 1 and supporting information Figures S1–S3). We located 523 subglacial hydrological sinks in the study area, with median and maximum areas of  $0.30 \text{ km}^2$  and  $12.7 \text{ km}^2$ , respectively (supporting information Figure S4). The total modeled runoff from 29 May to



**Figure 1.** Subglacial hydrology and ice flow. The Isunnguata Sermia and Kangerlussuaq catchments (with three subcatchments: Ørkendalen, Sandflugtsdalen, and Point 660) are delineated with white lines for subglacial water pressures equal to ice overburden. Automatic weather stations [van As et al., 2012] used in the runoff calculations and previous borehole measurement sites [Meierbachtol et al., 2013; van de Wal et al., 2015] are shown. See supporting information Figure S1 for an extended map of the drainage catchments up to the ice divide. (a) Supraglacial lakes (maximum extent 2002 to 2012) [Fitzpatrick et al., 2014] and streams (2012) [Smith et al., 2015] recur in subglacial depressions. See supporting information Figure S2 for a higher-resolution image. (b) Previous delineations of supraglacial [Mernild and Hasholt, 2009; Bartholomew et al., 2011; Fitzpatrick et al., 2014] and subglacial (from hydraulic potential) [Smith et al., 2015] catchments correlate spatially with the subglacial drainage catchments close to the ice margin but differ considerably farther inland. Annual ice surface velocities (2005/2006) are from Joughin et al. [2010], and surface elevations are from Howat et al. [2014]. Elevation is given as height above the WGS-1984 ellipsoid.


10 September 2012 equals  $5.4 \pm 0.5 \text{ km}^3$  for the Isunnguata Sermia catchment and  $6.5 \pm 0.7 \text{ km}^3$  for the Kangerlussuaq catchment (Figure 2 and Table S1). Observed total discharge from the Isortoq River was between 2.7 and  $5.4 \text{ km}^3$  (with linear interpolation used to account for 32 missing observations) [Smith et al., 2015]. Observed total discharge from the Watson River was  $6.4 \pm 1.0 \text{ km}^3$  from 29 May to 10 September 2012, which

**Table 1.** Comparison of the Area of Our Subglacial Catchments With Those Defined by Previous Studies Assuming Subglacial Water Pressures Equal to Ice Overburden<sup>a</sup>


Catchment	Area up to ~ 1750 m asl (km <sup>2</sup> )	Area up to Ice Divide ~ 2600 m asl (km <sup>2</sup> )
1 Isunnguata Sermia catchment	3,200	15,900
2 Kangerlussuaq catchment	2,800	12,000
<i>van As et al.</i> [2012]	ND	12,547
<i>Fitzpatrick et al.</i> [2014]	ND	9,743
2a Sandflugtsdalen subcatchment	NA	900
<i>Bartholomew et al.</i> [2011]	NA	1,145
2b Ørkendalen subcatchment	1,800	11,100
2c Point 660 subcatchment	NA	40
<i>Rennermalm et al.</i> [2013a]	NA	31–60

<sup>a</sup>ND, no data; NA, not applicable.

compares well to the modeled runoff for the Kangerlussuaq catchment of  $6.5 \pm 0.7 \text{ km}^3$ . Finally, the modeled runoff from the Sandflugtsdalen subcatchment from 29 May to 29 August 2012 totaled  $2.3 \pm 0.2 \text{ km}^3$ , which compares favorably to the measured discharge of  $2.2 \text{ km}^3$  from the outlet river of Leverett Glacier [Tedstone et al., 2013]. Our results (Figure 3 and supporting information Figures S5 and S6) also reveal a significant transience in the delineation of drainage catchments by varying the subglacial water pressure (expressed by factor  $k$  in equation (1); see supporting information Text S1), especially within  $\pm 10\%$  of the ice overburden pressures ( $k=0.9$  to  $1.1$ ). This range in subglacial water pressure is consistent with borehole measurements in the area; Meierbachtol et al. [2013] measured subglacial water pressure factors within the Isunnguata Sermia catchment ranging from  $k=0.88$  to above 1 in the S3 and S4 boreholes, and van de Wal et al. [2015] measured water pressures in the Kangerlussuaq catchment ranging from  $k \approx 0.95$  to above 1 in the SHR borehole (Figure 1a).



**Figure 2.** Daily totals of modeled supraglacial runoff and measured proglacial discharge from 29 May to 10 September 2012. (a) The Isunnguata Sermia catchment and the Isortoq River, with maximum and minimum discharge values [Smith et al., 2015]. (b) The Kangerlussuaq catchment and the Watson River, with  $\pm 15\%$  uncertainty in the discharge calculations in error bars. (c) Stacked discharge for the Isortoq (linearly interpolated to fill the 32 missing observations) and Watson proglacial rivers. (d) Stacked runoff for the Kangerlussuaq and Isunnguata Sermia catchment for subglacial water pressures equal to ice overburden.



**Figure 3.** (a) Spatial changes in subglacial drainage delineations at different subglacial water pressures (see equation (1) in the supporting information Text S1) for  $k = 0.9$  (dotted line),  $1.0$  (solid line), and  $1.1$  (dashed line): the Isunnguata Sermia catchment (red), the Kangerlussuaq catchment (blue), and overlapping areas (purple). See also supporting information Figure S5. (b) Changes in subglacial catchment area with changing subglacial water pressures for  $k = 0.5$  to  $1.1$ . The Kangerlussuaq catchment shifts northward for  $k = 1.1$  (without changing significantly in size) and takes a large part ( $\sim 30\%$ ) of the area from the Isunnguata Sermia catchment. The area changes are calculated up to  $1750$  m asl. A cubic polynomial regression was applied to the catchment areas and the coefficient of determination ( $R^2$ ) was  $0.95$  for Isunnguata Sermia and  $0.38$  for Kangerlussuaq (with  $0.79$  for the Ørkendalen subcatchment and  $0.91$  for the Sandflugtsdalen subcatchment).

#### 4. Discussion

The hydraulic and ice thickness gradients along the main flow paths in the ablation zone are on the order of  $-10^1$  to  $-10^2$  m km $^{-1}$  (supporting information Figure S3 and Table S2), which is high in comparison with flow paths in Antarctica, where typical values of  $-10^{-4}$  to  $-10^{-3}$  m km $^{-1}$  have been measured for the hydraulic head and  $-10^{-3}$  to  $-10^{-2}$  m km $^{-1}$  for the ice thickness [Wolovick *et al.*, 2013]. This can be explained by the relatively rough bed topography [Lindbäck and Pettersson, 2015] and steep ice surface slopes in the marginal areas of our study area. Consistent with previous work [Joughin *et al.*, 2013] our results suggest that the spatial pattern of enhanced summer ice flow can be attributed to surface and bed slopes acting in unison to determine subglacial hydrological pathways. Our identified sinks in the hydraulic potential surface (Figure 1a and supporting information Figures S2 and S4) are possible locations for subglacial lakes, and expectedly the density of sinks is much higher than found by previous studies from the GrIS [e.g., Livingstone *et al.*, 2013] that used lower resolution DEMs. Although many of the sinks are small (median area  $0.30$  km $^2$ ), they have a combined area of  $290$  km $^2$  and an estimated combined storage capacity of  $4.1$  km $^3$ . The preponderance of sinks indicates that small subglacial lakes ( $<1$  km $^2$ ) are likely to exist in our study area but remain undetected by remote sensing data owing to a negligible surface expression created by subglacial lakes of this size [e.g., Langley *et al.*, 2011]. In many locations the sinks are mirrored at the surface by supraglacial lakes ( $\sim 40\%$  overlap; Figure 1a), which provides support for the modeling work of Sergienko [2013], where supraglacial and subglacial lakes are colocated originating from the same basal features.

Previous studies that delineated hydrological catchments using ice surface topography [e.g., Mernild and Hasholt, 2009; Mernild *et al.*, 2010; Bartholomew *et al.*, 2011; van As *et al.*, 2012; Chandler *et al.*, 2013; Cowton *et al.*, 2013; Fitzpatrick *et al.*, 2014] agree well with our subglacial catchments close to the margin ( $<50$  km distance and  $<1200$  m asl), but large differences occur farther inland (Figure 1b). In accordance with previous studies [e.g., Budd and Carter, 1971; Gudmundsson, 2003], we suggest that the large differences between previous catchment delineations above  $1200$  m asl may, in part, be explained by thicker ice having a larger influence on the hydrostatic pressure as well as muting the surface expression of the bed, making accurate delineation of the subglacial drainage from surface topography more difficult. Large discrepancies also occur between our subglacial catchment delineation and the subglacial catchments calculated from the

1 km gridded *Bamber et al.*'s [2013a] DEM [e.g., *Smith et al.*, 2015]. We attribute the differences to our higher data density, which provides greater detail in our ice thickness DEM that is especially important in regions where subglacial water routing is sensitive to small variations in elevation. Moreover, the geostatistical interpolation model that we used to generate the DEM was adapted to the regional setting whereas the *Bamber et al.*'s [2013a] DEM was optimized for the whole GrIS.

The high level of agreement between modeled runoff ( $6.5 \pm 0.7 \text{ km}^3$ ) for the Kangerlussuaq catchment and measured bulk proglacial discharge ( $6.4 \pm 1.0 \text{ km}^3$ ) in the Watson River suggests that a limited volume of water ( $\sim 0.1 \text{ km}^3$ ) is retained after the melt season in this catchment. The coherence between our observed proglacial discharge and modeled runoff compared to previous studies [e.g., *van As et al.*, 2012; *Rennermalm et al.*, 2013a; *Fitzpatrick et al.*, 2014] can be explained by the reduced uncertainty in catchment delineation, especially in the upper part of the ablation zone ( $>1200 \text{ m asl}$ ) from where a significant part of the surface runoff originates (65% in 2012) and the majority of supraglacial lakes are located [*Fitzpatrick et al.*, 2014; *Leeson et al.*, 2014]. The discharge estimates of the Isortoq River have larger uncertainties (discharge with lower and upper bounds ranging from 2.7 to  $5.4 \text{ km}^3$ ) [*Smith et al.*, 2015], due to a different gauging technique (see section 2). Our modeled runoff ( $5.4 \pm 0.5 \text{ km}^3$ ) is close to the upper limit of the measured discharge but exceeds the lower limit by 50%. Previous findings of *Smith et al.* [2015] measured Isortoq discharge to total 37 to 75% of modeled runoff in 2012, suggesting substantial subglacial water storage in the Isunnguata Sermia catchment, as opposed to the evidence of low storage in the Kangerlussuaq catchment. *Rennermalm et al.* [2013a] also found evidence for meltwater retention (up to 54% of the modeled runoff between 2008 and 2012) in the Point 660 subcatchment, which is small (100 times smaller than the Kangerlussuaq catchment) and therefore has larger relative uncertainty in delineation. The speculated missing water may indeed be retained, as hypothesized by these studies, but alternative explanations also exist, such as groundwater losses and subglacial water piracy, the latter of which is discussed below. Water can be stored interannually when water recharges subglacial sinks that have been flushed out during the melt season. Such storage transience is revealed in our hydraulic potential analysis where subglacial sinks decrease in size with higher subglacial water pressures, decreasing with  $\sim 25\%$  in total size (equating to an area of  $\sim 70 \text{ km}^2$ ) from  $k = 1$  to  $k = 1.1$  (supporting information Figure S5). Comparing daily differences between measured discharge and modeled runoff shows that  $\sim 1.4 \text{ km}^3$  of water has a delayed release within the melt season for both catchments. We attribute this to the temporary retention of water in snow, supraglacial lakes, and the subglacial drainage system.

The bed topography in the Kangerlussuaq sector of the GrIS shows highly variable subglacial trough systems, channelizing the ice flow and enhancing ice deformation. The close agreement between ice flow patterns [*Joughin et al.*, 2010; *Palmer et al.*, 2011] and subglacial water networks in our study area indicates that basal water flow has a primary impact on ice dynamics within the troughs. The extent and magnitude of enhanced summer ice flow and measured discharge varies considerably between the adjacent outlets despite similar geometry and climatic control in the area [e.g., *Fitzpatrick et al.*, 2013]. Outlet glaciers within the Kangerlussuaq catchment accelerate with high magnitude at the beginning of the melt season. By contrast, Isunnguata Sermia has relatively low rates of ice flow and proglacial discharge at the melt season onset compared to later in the season (Figure 2). The low discharge values and ice velocities measured for the Isunnguata Sermia catchment cannot be linked to differences in modeled surface melt in the adjacent Kangerlussuaq catchment, and we hypothesize that this is instead caused by water piracy. The Watson River receives a larger fraction ( $\sim 75\%$ ) of the discharge than the Isortoq River before peak runoff on 13 July 2012, whereas after peak runoff the two proglacial rivers each discharge approximately equal amounts (53% and 47%, respectively). Hence, the relatively larger volumes of water draining through the Kangerlussuaq catchment during melt season onset may explain the pronounced early-season ice flow response of the outlet glaciers of this catchment compared to the Isunnguata Sermia catchment. Instead, Isunnguata Sermia accelerates later in the melt season possibly owing to an expanding catchment area (water piracy) driving increased subglacial water routing and discharge. We test the hypothesis of water piracy by varying subglacial water pressures in our subglacial catchment delineation and modeled runoff (expressed by factor  $k$ ; Figure 3 and supporting information Figures S5 and S6). Fluctuations in water pressure are common in the study area during the melt season both on temporal and spatial scales [e.g., *Meierbachtol et al.*, 2013; *Andrews et al.*, 2014], and the assumption of uniform subglacial water pressure is not accurate over the whole region or over time. At subglacial water pressures below 80% of the ice overburden pressures

( $k = 0.5$  to  $0.8$ ), which is expected close ( $< 50$  km) to the ice margin [e.g., Meierbachtol *et al.*, 2013], the catchments do not change significantly in size (Figure 3b). The largest changes are for  $k$  values from  $0.9$  to  $1.1$  and occur where the subglacial valleys are shallow, or where subglacial valleys converge (supporting information Figure S5). Remarkably, the Kangerlussuaq catchment migrates northward for  $k = 1.1$  but remains similar in area and captures  $\sim 30\%$  of the area from the Isunnguata Sermia catchment (Figure 3 and supporting information Figure S5). As the net area of the Kangerlussuaq catchment does not significantly change, the total discharge into Watson River neither increased drastically. On the other hand, Isunnguata Sermia catchment does not migrate northward suffers from reduced area and hence reduced discharge. Catchment migration (or water piracy) caused by changes in subglacial water pressure ( $k$  values) over, for example, a summer melt season could therefore provide alternative hypothesis to that of meltwater retention. At this time regional basal water pressures would be high and observations of surface uplift indicate that transient water pressures in excess of overburden do occur [Shepherd *et al.*, 2009]. Individual drainage catchments may therefore fluctuate significantly in size in response to relatively small changes in basal water pressure. Hence, the subglacial routing of water may be influenced not only by the basal topography but also by varying subglacial water pressure; both play a key role in governing subglacial water routing and therefore the ice flow regime of individual outlet glaciers. Our findings suggest that a more sophisticated subglacial hydrology modeling to predict variable subglacial water pressure [e.g., Werder *et al.*, 2013] is warranted especially in sensitive areas in the subglacial topography (i.e., areas that are potential switching points) to understand the nonlinear and transient relationships between subglacial water flow (i.e., its routing and discharge) and ice dynamics over diurnal, seasonal, and annual time scales.

#### Acknowledgments

The ice thickness and bed topography DEMs used in this study are available at doi:10.1594/pangaea.830314. The mass conservation DEM was provided by Mathieu Morlighem and is available at <http://sites.uci.edu/morlighem/dataproducts/mass-conservation-dataset/>. The GIMP data set was provided by the BPRC Glacier Dynamics Research Group, Ohio State University, and is available at <http://research.bprc.osu.edu/GDG/gimpdem.php>. Data from the three KAN automatic weather stations used for the runoff calculations are available through the Programme for Monitoring the Greenland Ice Sheet (PROMICE) at <http://www.promice.dk>. The ice velocity data set was provided by the NASA MEaSUREs program and is available at [http://nsidc.org/data/docs/measures/nsidc0478\\_joughin/](http://nsidc.org/data/docs/measures/nsidc0478_joughin/). This project, including the gathering of radar data and weather station data, was funded by the Greenland Analogue Project (GAP), the UK Natural Environment Research Council (NERC) grants NE/G005796/1, NE/G010595/1, and NE/H024204/1, the Royal Geographical Society (Gilchrist Educational Trust), and the Aberystwyth University Research Fund. K.L. was supported by additional field and travel support from the Nordic Centre of Excellence SVALI, the Swedish Society for Anthropology and Geography (SSAG), and the Geographical Society of Uppsala. A.L.H. was supported by funding from the Research Council of Norway grant 223259 and S.H.D. and A.A.F. were supported by Aberystwyth University doctoral scholarships. We thank Colin J. Gleason and Laurence C. Smith for providing the Isortoq discharge data and Kang Yang for providing the supraglacial river map. SPOT images were provided by the SPIRIT Program © CNES 2008–2009 and Spot Image 2008.

The Editor thanks two anonymous reviewers and Laurence C. Smith for their assistance in evaluating this paper.

## 5. Conclusions

The presence of water beneath ice sheets has a fundamental impact on ice flow due to its role as a lubricant either between the ice and its base or between grains of subglacial sediment, and hence, water delivery to the bed plays a key role on the rate of dynamic mass loss to global sea level. We present a high-resolution subglacial hydrological analysis of the land-terminating Kangerlussuaq sector of the GrIS, characterizing subglacial catchments, flow networks, and hydrological sinks. The agreement between observed proglacial discharge and modeled melt runoff exceeds that of previous studies using similar runoff calculations, due to reduced uncertainty in our catchment delineation. Nevertheless, our analysis indicates that many small subglacial lakes may occur in this region and yet would remain undetectable by standard methods using coarse-resolution DEMs, ice surface elevation change, or radar techniques. Furthermore, our findings indicate that drainage catchments may vary not only on glacial time scales of  $10^1$  to  $10^3$  years, as determined for ice streams in Antarctica, but also on subseasonal time scales in response to relatively small perturbations in the subglacial water pressure. These findings should be considered in studies that attempt to relate estimates of surface runoff from energy balance models with measurements of proglacial discharge and ice dynamics. Our study highlights the need not only for accurate high-resolution DEMs but also for regionally optimized interpolation when conducting detailed hydrological studies of the GrIS. Continued efforts targeting the hydrological system of the ice sheet should over time result in finer spatial coverage, allowing for a broader understanding of the ice sheet's hydrological and dynamic response to climate change and ultimately its contribution to global sea level rise.

## References

- Allison, I., R. B. Alley, H. A. Fricker, R. H. Thomas, and R. C. Warner (2009), Ice sheet mass balance and sea level, *Antarct. Sci.*, 21(05), 413, doi:10.1017/S0954102009990137.
- Anandakrishnan, S., and R. B. Alley (1997), Stagnation of ice stream C, West Antarctica by water piracy, *Geophys. Res. Lett.*, 24(3), 265–268, doi:10.1029/96GL04016.
- Andrews, L., G. Catania, M. J. Hoffman, J. D. Gullely, M. P. Lüthi, C. Ryser, R. L. Hawley, and T. A. Neumann (2014), Direct observations of evolving subglacial drainage beneath the Greenland ice sheet, *Nature*, 514, doi:10.1038/nature13796.
- Bamber, J. L., et al. (2013a), A new bed elevation dataset for Greenland, *Cryosphere*, 7(2), 499–510, doi:10.5194/tc-7-499-2013.
- Bamber, J. L., M. J. Siegert, J. A. Griggs, S. J. Marshall, and G. Spada (2013b), Paleofluvial mega-canyon beneath the central Greenland ice sheet, *Science*, 341(6149), 997–999, doi:10.1126/science.1239794.
- Banwell, A. F., I. C. Willis, and N. S. Arnold (2013), Modeling subglacial water routing at Paakitsoq, W Greenland, *J. Geophys. Res. Earth Surf.*, 118, 1282–1295, doi:10.1002/jgrf.20093.
- Bartholomew, I., P. Nienow, D. Mair, A. Hubbard, M. A. King, and A. Sole (2010), Seasonal evolution of subglacial drainage and acceleration in a Greenland outlet glacier, *Nat. Geosci.*, 3(6), 408–411, doi:10.1038/ngeo863.



- Bartholomew, I., P. Nienow, A. Sole, D. Mair, T. Cowton, S. Palmer, and J. Wadham (2011), Supraglacial forcing of subglacial drainage in the ablation zone of the Greenland ice sheet, *Geophys. Res. Lett.*, *38*, L08502, doi:10.1029/2011GL047063.
- Bartholomew, I., P. Nienow, A. Sole, D. Mair, T. Cowton, and M. A. King (2012), Short-term variability in Greenland ice sheet motion forced by time-varying meltwater drainage: Implications for the relationship between subglacial drainage system behavior and ice velocity, *Geophys. Res. Lett.*, *117*, F03002, doi:10.1029/2011JF002220.
- Budd, W. F., and D. B. Carter (1971), An analysis of the relation between the surface and bedrock profiles of ice caps, *J. Glaciol.*, *10*(59), 197–209.
- Carter, S. P., H. A. Fricker, and M. R. Siegfried (2013), Evidence of rapid subglacial water piracy under Whillans Ice Stream, West Antarctica, *J. Glaciol.*, *59*(218), 1147–1162, doi:10.3189/2013JoG13J085.
- Chandler, D. M., et al. (2013), Evolution of the subglacial drainage system beneath the Greenland ice sheet revealed by tracers, *Nat. Geosci.*, *6*(3), 195–198, doi:10.1038/ngeo1737.
- Cowton, T., P. Nienow, A. Sole, J. Wadham, G. Lis, I. Bartholomew, D. Mair, and D. Chandler (2013), Evolution of drainage system morphology at a land-terminating Greenlandic outlet glacier, *J. Geophys. Res. Earth Surf.*, *118*, 29–41, doi:10.1029/2012JF002540.
- Dow, C. F., B. Kulesa, I. C. Rutt, S. H. Doyle, and A. Hubbard (2014), Upper bounds on subglacial channel development for interior regions of the Greenland ice sheet, *J. Glaciol.*, *60*(224), 1044–1052, doi:10.3189/2014JoG14J093.
- Doyle, S. H., A. Hubbard, A. A. W. Fitzpatrick, D. van As, A. B. Mikkelsen, R. Pettersson, and B. Hubbard (2014), Persistent flow acceleration within the interior of the Greenland ice sheet, *Geophys. Res. Lett.*, *41*, 899–905, doi:10.1002/2013GL058933.
- Doyle, S. H., et al. (2015), Amplified melt and flow of the Greenland ice sheet driven by late-summer cyclonic rainfall, *Nat. Geosci.*, doi:10.1038/ngeo2482.
- Fitzpatrick, A. A. W., A. Hubbard, I. Joughin, D. J. Quincey, D. van As, A. P. B. Mikkelsen, S. H. Doyle, B. Hasholt, and G. A. Jones (2013), Ice flow dynamics and surface meltwater flux at a land-terminating sector of the Greenland ice sheet, *J. Glaciol.*, *59*(216), 687–696, doi:10.3189/2013JoG12J143.
- Fitzpatrick, A. A. W., A. L. Hubbard, J. E. Box, D. J. Quincey, D. van As, A. P. B. Mikkelsen, S. H. Doyle, C. F. Dow, B. Hasholt, and G. A. Jones (2014), A decade (2002–2012) of supraglacial lake volume estimates across Russell Glacier, West Greenland, *Cryosphere*, *8*(1), 107–121, doi:10.5194/tc-8-107-2014.
- Fricker, H. A., T. Scambos, R. Bindenschadler, and L. Padman (2007), An active subglacial water system in West Antarctica mapped from space, *Science*, *315*(5818), 1544–1548, doi:10.1126/science.1136897.
- Gleason, C. J., L. C. Smith, D. C. Finnegan, A. L. LeWinter, L. H. Pitcher, and V. W. Chu (2015), Technical Note: Semi-automated effective width extraction from time-lapse RGB imagery of a remote, braided Greenlandic river, *Hydrol. Earth Syst. Sci.*, *19*, 2963–2969, doi:10.5194/hess-19-2963-2015.
- Gudmundsson, G. H. (2003), Transmission of basal variability to a glacier surface, *J. Geophys. Res.*, *108*(B5, 2253), doi:10.1029/2002JB002107.
- Hanna, E., et al. (2013), Ice-sheet mass balance and climate change, *Nature*, *498*(7452), 51–9, doi:10.1038/nature12238.
- Hasholt, B., A. Bech Mikkelsen, M. Holtegaard Nielsen, and M. Andreas Dahl Larsen (2013), Observations of runoff and sediment and dissolved loads from the Greenland ice sheet at Kangerlussuaq, West Greenland, 2007 to 2010, *Zeitschrift für Geomorphol. Suppl. Issues*, *57*(2), 3–27, doi:10.1127/0372-8854/2012/S-00121.
- Howat, I. M., A. Negrete, and B. E. Smith (2014), The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *Cryosphere*, *8*(4), 1509–1518, doi:10.5194/tc-8-1509-2014.
- Joughin, I., B. E. Smith, I. M. Howat, T. Scambos, and T. Moon (2010), Greenland flow variability from ice-sheet-wide velocity mapping, *J. Glaciol.*, *56*(197), 415–430, doi:10.3189/002214310792447734.
- Joughin, I., S. B. Das, G. E. Flowers, M. D. Behn, R. B. Alley, M. A. King, B. E. Smith, and J. L. Bamber (2013), Influence of ice-sheet geometry and supraglacial lakes on seasonal ice-flow variability, *Cryosphere*, *7*, 1185–1192, doi:10.5194/tc-7-1185-2013.
- Karlsson, N. B., and D. Dahl-Jensen (2015), Response of the large-scale subglacial drainage system of Northeast Greenland to surface elevation changes, *Cryosphere*, *9*, 1465–1479, doi:10.5194/tc-9-1465-2015.
- Langley, K., J. Kohler, K. Matsuoka, A. Sinisalo, T. Scambos, T. Neumann, A. Muto, J. G. Winther, and M. Albert (2011), Recovery Lakes, East Antarctica: Radar assessment of subglacial water extent, *J. Geophys. Res.*, *38*, L05501, doi:10.1029/2010GL046094.
- Leeson, A. A., A. Shepherd, K. Briggs, I. Howat, X. Fettweis, M. Morlighem, and E. Rignot (2014), Supraglacial lakes on the Greenland ice sheet advance inland under warming climate, *Nat. Clim. Change*, 1–5, doi:10.1038/NCLIMATE2463.
- Lewis, S., and L. C. Smith (2009), Hydrologic drainage of the Greenland ice sheet, *Hydrol. Process.*, *23*, 2004–2011, doi:10.1002/hyp.
- Lindbäck, K., and R. Pettersson (2015), Spectral roughness and glacial erosion of a land-terminating section of the Greenland ice sheet, *Geomorphology*, *238*, 149–159, doi:10.1016/j.geomorph.2015.02.027.
- Lindbäck, K., R. Pettersson, S. H. Doyle, C. Helanow, P. Jansson, S. S. Kristensen, L. Stenseng, R. Forsberg, and A. L. Hubbard (2014), High-resolution ice thickness and bed topography of a land-terminating section of the Greenland ice sheet, *Earth Syst. Sci. Data*, *6*, 331–338, doi:10.5194/essd-6-331-2014.
- Livingstone, S. J., C. D. Clark, J. Woodward, and J. Kingslake (2013), Potential subglacial lake locations and meltwater drainage pathways beneath the Antarctic and Greenland ice sheets, *Cryosphere*, *7*(6), 1721–1740, doi:10.5194/tc-7-1721-2013.
- Meierbachtol, T., J. Harper, and N. Humphrey (2013), Basal drainage system response to increasing surface melt on the Greenland ice sheet, *Science*, *341*(6147), 777–9, doi:10.1126/science.1235905.
- Mernild, S. H., and B. Hasholt (2009), Observed runoff, jökulhlaups and suspended sediment load from the Greenland ice sheet at Kangerlussuaq, West Greenland, 2007 and 2008, *J. Glaciol.*, *55*(193), 855–858.
- Mernild, S. H., G. E. Liston, K. Steffen, M. van den Broeke, and B. Hasholt (2010), Runoff and mass-balance simulations from the Greenland ice sheet at Kangerlussuaq (Søndre Strømfjord) in a 30-year perspective, 1979–2008, *Cryosphere*, *4*(2), 231–242, doi:10.5194/tc-4-231-2010.
- Morlighem, M., E. Rignot, J. Mougino, H. Seroussi, and E. Larour (2014), Deeply incised submarine glacial valleys beneath the Greenland ice sheet, *Nat. Geosci.*, *7*, 18–22, doi:10.1038/NGL02167.
- Palmer, S., A. Shepherd, P. Nienow, and I. Joughin (2011), Seasonal speedup of the Greenland ice sheet linked to routing of surface water, *Earth Planet. Sci. Lett.*, *302*(3–4), 423–428, doi:10.1016/j.epsl.2010.12.037.
- Rennermalm, A. K., L. C. Smith, V. W. Chu, J. E. Box, R. R. Forster, M. R. Van den Broeke, D. Van As, and S. E. Moustafa (2013a), Evidence of meltwater retention within the Greenland ice sheet, *Cryosphere*, *7*(5), 1433–1445, doi:10.5194/tc-7-1433-2013.
- Rennermalm, A. K., et al. (2013b), Understanding Greenland ice sheet hydrology using an integrated multi-scale approach, *Environ. Res. Lett.*, *8*, 015017, doi:10.1088/1748-9326/8/1/015017.
- Sergienko, O. V. (2013), Glaciological twins: Basally controlled subglacial and supraglacial lakes, *J. Glaciol.*, *59*(213), 3–8, doi:10.3189/2013JoG12J040.
- Shepherd, A., A. Hubbard, P. Nienow, M. King, M. McMillan, and I. Joughin (2009), Greenland ice sheet motion coupled with daily melting in late summer, *Geophys. Res. Lett.*, *36*, L01501, doi:10.1029/2008GL035758.

- Shreve, R. L. (1972), Movement of water in glaciers, *J. Glaciol.*, *11*(62), 205–214.
- Smeets, C. J. P. P., W. Boot, A. Hubbard, R. Pettersson, F. Wilhelms, M. R. van den Broeke, and R. van de Wal (2012), Instruments and methods—A wireless subglacial probe for deep ice applications, *J. Glaciol.*, *58*(211), 841–848, doi:10.3189/2012JoG11J130.
- Smith, L. C., V. W. Chu, K. Yang, C. J. Gleason, L. H. Pitcher, and A. K. Rennermalm (2015), Efficient meltwater drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, *Proc. Natl. Acad. Sci. U.S.A.*, *112*(4), 1001–1006, doi:10.1073/pnas.1413024112.
- Sundal, A. V., A. Shepherd, P. Nienow, E. Hanna, S. Palmer, and P. Huybrechts (2011), Melt-induced speed-up of Greenland ice sheet offset by efficient subglacial drainage, *Nature*, *469*(7331), 521–524, doi:10.1038/nature09740.
- Tedstone, A. J., P. W. Nienow, A. J. Sole, D. W. F. Mair, T. R. Cowton, I. D. Bartholomew, and M. A. King (2013), Greenland ice sheet motion insensitive to exceptional meltwater forcing, *Proc. Natl. Acad. Sci. U.S.A.*, *110*, 19,719–19,724, doi:10.1073/pnas.1315843110.
- van As, D., A. L. Hubbard, B. Hasholt, A. B. Mikkelsen, M. R. van den Broeke, and R. S. Fausto (2012), Large surface meltwater discharge from the Kangerlussuaq sector of the Greenland ice sheet during the record-warm year 2010 explained by detailed energy balance observations, *Cryosphere*, *6*(1), 199–209, doi:10.5194/tc-6-199-2012.
- van de Wal, R. S. W., and A. J. Russell (1994), A comparison of energy balance calculations, measured ablation and meltwater runoff near Sondre Stromfjord, West Greenland, *Global Planet. Change*, *9*, 29–38, doi:10.1016/0921-8181(94)90005-1.
- van de Wal, R. S. W., W. Boot, M. R. van den Broeke, C. J. P. P. Smeets, C. H. Reijmer, J. J. A. Donker, and J. Oerlemans (2008), Large and rapid melt-induced velocity changes in the ablation zone of the Greenland ice sheet, *Science*, *321*(5885), 111–3, doi:10.1126/science.1158540.
- van de Wal, R. S. W., et al. (2015), Self-regulation of ice flow varies across the ablation area in south-west Greenland, *Cryosphere*, *9*(2), 603–611, doi:10.5194/tc-9-603-2015.
- Vaughan, D. G., H. F. J. Corr, A. M. Smith, H. D. Pritchard, and A. Shepherd (2008), Flow-switching and water piracy between Rutford ice stream and Carlson inlet, West Antarctica, *J. Glaciol.*, *54*(184), 41–48, doi:10.3189/002214308784409125.
- Werder, M. A., I. J. Hewitt, C. G. Schoof, and G. E. Flowers (2013), Modeling channelized and distributed subglacial drainage in two dimensions, *J. Geophys. Res. Earth Surf.*, *118*, 2140–2158, doi:10.1002/jgrf.20146.
- Wingham, D. J., M. J. Siegert, A. Shepherd, and A. S. Muir (2006), Rapid discharge connects Antarctic subglacial lakes, *Nature*, *440*(7087), 1033–6, doi:10.1038/nature04660.
- Wolovick, M. J., R. E. Bell, T. T. Creyts, and N. Frearson (2013), Identification and control of subglacial water networks under Dome A, Antarctica, *J. Geophys. Res. Earth Surf.*, *118*, 1–15, doi:10.1002/2012JF002555.
- Wright, A., and M. Siegert (2012), A fourth inventory of Antarctic subglacial lakes, *Antarct. Sci.*, *24*(6), 1–6, doi:10.1017/S095410201200048X.
- Wright, A. P., M. J. Siegert, A. M. Le Brocq, and D. B. Gore (2008), High sensitivity of subglacial hydrological pathways in Antarctica to small ice-sheet changes, *Geophys. Res. Lett.*, *35*, L17504, doi:10.1029/2008GL034937.