



Aberystwyth University

Firn air-content of Larsen C Ice Shelf, Antarctic Peninsula, from seismic velocities, borehole surveys and firn modelling

Kulesa, Bernd; Brisbourne, Alex; Booth, Adam ; Kuipers Munneke, Peter; Bevan, Suzanne; Luckman, Adrian; Hubbard, Bryn; Gourmelen, Noel; Palmer, Steve; Holland, Paul; Ashmore, David; Shepherd, Andrew

Publication date:

2016

Citation for published version (APA):

Kulesa, B., Brisbourne, A., Booth, A., Kuipers Munneke, P., Bevan, S., Luckman, A., Hubbard, B., Gourmelen, N., Palmer, S., Holland, P., Ashmore, D., & Shepherd, A. (2016). *Firn air-content of Larsen C Ice Shelf, Antarctic Peninsula, from seismic velocities, borehole surveys and firn modelling*. Poster session presented at European Geosciences Union General Assembly 2016, Vienna, Austria.

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk



Firn air-content of Larsen C Ice Shelf, Antarctic Peninsula, from seismic velocities, borehole surveys and firn modelling

Bernd Kulesa (1), Alex Brisbourne (2), Adam Booth (3), Peter Kuipers Munneke (1,4), Suzanne Bevan (1), Adrian Luckman (1), Bryn Hubbard (5), Noel Gourmelen (6), Steve Palmer (7), Paul Holland (2), David Ashmore (5), and Andrew Shepherd (3)

(1) College of Science, Swansea University, Swansea, UK, (2) British Antarctic Survey, Cambridge, UK, (3) School of Earth and Environment, University of Leeds, Leeds, UK, (4) Institute for Marine and Atmospheric Research Utrecht (IMAU), Utrecht University, Utrecht, The Netherlands, (5) Geography and Earth Sciences, Aberystwyth University, Aberystwyth, UK, (6) School of GeoSciences, University of Edinburgh, Edinburgh, UK, (7) Department of Geography, University of Exeter, Exeter, UK

The rising surface temperature of Antarctic Peninsula ice shelves is strongly implicated in ice shelf disintegration, by exacerbating the compaction of firn layers. Firn compaction is expected to warm the ice column and, given sufficiently wet and compacted layers, to allow meltwater to penetrate into surface crevasses and thus enhance hydrofracture potential. Integrating seismic refraction surveys with borehole neutron and firn core density logging, we reveal vertical and horizontal changes in firn properties across Larsen C Ice Shelf. Patterns of firn air-content derived from seismic surveys are broadly similar to those estimated previously from airborne radar and satellite data. Specifically, these estimates show greater firn compaction in the north and landward inlets compared to the south, although spatial gradients in seismic-derived air-contents are less pronounced than those previously inferred. Firn thickness is less than 10 m in the extreme northwest of Larsen C, in Cabinet Inlet, yet exceeds 40 m in the southeast, suggesting that the inlet is a focus of firn compaction; indeed, buried layers of massive refrozen ice were observed in 200 MHz GPR data in Cabinet and Whirlwind Inlets during a field campaign in the 2014-15 austral summer. Depth profiles of firn density provide a reasonable fit with those derived from closely-located firn cores and neutron probe data.

Our model of firn structure is driven by RACMO and includes a 'bucket'-type hydrological implementation, and simulates the depth-density profiles in the inlets well. Discrepancies between measured and modelled depth-density profiles become progressively greater towards the ice-shelf front. RACMO incorrectly simulates the particular leeward (sea-ice-influenced) microclimate of the shallow boundary layer, leading to excess melt and/or lack of snowfall.

The spatial sampling density of our seismic observations will be augmented following a further field campaign in the 2016-17 austral summer, allowing quantitative comparison of our firn air-content estimates (and their hydrostatic implications) versus those derived from airborne radar data. We suggest that our seismic data may provide more realistic profiles of firn density and air-content in the southeastern sector of Larsen C, where the ice shelf is heavily fractured.