



## Aberystwyth University

### *The Subglacial Access and Fast Ice Research Experiment (SAFIRE)*

Hubbard, Bryn; Doyle, Samuel; Christoffersen, Poul; Young, Tun Jan; Hofstede, Coen; Hubbard, Alun; Box, Jason; Todd, Joe; Bougamont, Marion

*Publication date:*  
2016

*Citation for published version (APA):*

Hubbard, B., Doyle, S., Christoffersen, P., Young, T. J., Hofstede, C., Hubbard, A., Box, J., Todd, J., & Bougamont, M. (2016). *The Subglacial Access and Fast Ice Research Experiment (SAFIRE): 1. Borehole-based englacial and subglacial measurements from a rapidly-moving tidewater glacier: Store Glacier, Greenland*. Poster session presented at European Geosciences Union General Assembly 2016, Vienna, Austria.

#### **General rights**

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

## **The Subglacial Access and Fast Ice Research Experiment (SAFIRE): 1. Borehole-based englacial and subglacial measurements from a rapidly-moving tidewater glacier: Store Glacier, Greenland**

Bryn Hubbard (1), Samuel Doyle (1), Poul Christoffersen (2), Tun Jan Young (2), Coen Hofstede (3), Alun Hubbard (1,5), Jason Box (4), Joe Todd (2), and Marion Bougamont (2)

(1) Geography and Earth Sciences, Aberystwyth University, Aberystwyth, UK (byh@aber.ac.uk), (2) Scott Polar Research Institute, University of Cambridge, Cambridge, UK, (3) Alfred Wegener Institute, Bremerhaven, Germany, (4) Geological Survey of Denmark and Greenland, Copenhagen, Denmark, (5) Department of Geology, University of Tromsø, Tromsø, Norway

As part of the Subglacial Access and Fast Ice Research Experiment (SAFIRE) pressurised hot water was used to drill four 603-616 m-long boreholes to the bed of the Greenland Ice Sheet at a site located 30 km from the calving front of fast-flowing, marine-terminating Store Glacier (70 degrees N, ~1000 m elevation). Four wired sensor strings were successfully installed in three of the boreholes. These included a thermistor string to obtain the englacial temperature profile installed in the same borehole as a string of tilt sensors to measure borehole deformation, and two sets of combined water pressure, electrical conductivity and turbidity sensors installed just above the bed in separate, adjacent boreholes. The boreholes made a strong hydrological connection to the bed during drilling, draining rapidly to ~80 m below the ice surface. The connection of subsequent boreholes was observed as a perturbation in water pressure and temperature recorded in neighbouring boreholes, indicating an effective hydrological connection between them. The sensors, which were wired to data-loggers at the surface, operated for between ~30 and >80 days from late summer into autumn before the cables stretched and snapped, with the lowermost sensors failing first. The records obtained from these sensors reveal (i) subglacial water pressures that were close to overburden but which generally increased through the period of measurement and varied diurnally by ~0.3 m, (ii) a minimum englacial temperature of -21 degrees C underlain by a zone of temperate ice, some tens of m thick, located immediately above the bed, and (iii) high rates of internal deformation and strain that increased towards the bed. These borehole observations are complemented by GPS measurements of ice motion, meteorological data, and seismic and radar surveys.