

## Aberystwyth University

### *Indications of a pan-hemispheric bi-partition of the Younger Dras Stadial from Lake Suigetsu, Japan*

Schlolaut, Gordon; Brauer, Achim; Nakagawa, Takeshi; Lamb, Henry; Marshall, Michael Henry; Kato-Saito, Megumi; Staff, Richard ; Ramsey, Christopher Bronk; Bryant, Charlotte

*Publication date:*  
2016

*Citation for published version (APA):*

Schlolaut, G., Brauer, A., Nakagawa, T., Lamb, H., Marshall, M. H., Kato-Saito, M., Staff, R., Ramsey, C. B., & Bryant, C. (2016). *Indications of a pan-hemispheric bi-partition of the Younger Dras Stadial from Lake Suigetsu, Japan*. Poster session presented at European Geosciences Union General Assembly 2016, Vienna, Austria.

#### **General rights**

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)



## **Indications of a pan-hemispheric bi-partition of the Younger Dryas Stadial from Lake Suigetsu, Japan**

Gordon Schlolaut (1), Achim Brauer (2), Takeshi Nakagawa (3), Henry Lamb (4), Michael Marshall (4), Megumi Kato-Saito (5), Richard Staff (6), Christopher Bronk Ramsey (6), and Charlotte Bryant (7)

(1) Center for Ocean Drilling Science (ODS), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 3173-25 Showa-machi, Kanazawa-ku, Yokohama 236-0001, Japan, (2) Section 5.2: Climate Dynamics and Landscape Evolution, GFZ German Research Center for Geosciences, Telegrafenberg, 14473 Potsdam, Germany, (3) Research Centre for Palaeoclimatology, Ritsumeikan University, 1-1-1 Noji-Higashi Kusatsu, Shiga 525-8577, Japan, (4) Institute of Geography and Earth Sciences, Aberystwyth University, SY23 3DB, UK, (5) Department of Geology and Paleontology, National Museum of Nature and Science, Tsukuba-shi, Ibaraki 305-0005, Japan, (6) Research Laboratory for Archaeology & the History of Art, University of Oxford, Oxford, UK, (7) NERC Radiocarbon Laboratory, Scottish Enterprise Technology Park, Rankine Avenue, East Kilbride, UK

The Younger Dryas Stadial marks the final succession of climatic fluctuations of the last Glacial. Whilst well studied in records from Europe and Greenland, few high resolution records are available from East Asia.

Here we present a high resolution, multi-proxy study of the Lake Suigetsu (Japan) sediments using the 'SG06' composite profile. Utilising microfacies,  $\mu$ XRF, pollen and diatom analysis we characterise changes occurring in the timeframe corresponding to the Younger Dryas Stadial.

Firstly, our results show that the climatic equivalent of the Younger Dryas at Lake Suigetsu shows no major lead or lag in comparison to records from the North Atlantic region, which was postulated by an earlier project on the Suigetsu sediments ('SG93'). Reason for this disagreement between the SG06 and SG93 core is that the SG93 core/chronology was compromised by gaps between individual cores and varve count uncertainties.

Furthermore, some of the analysed proxies from the SG06 core show a sub-division of the Younger Dryas Stadial. The timing of this sub-division is similar to the bi-partition of the Younger Dryas Stadial observed in a number of European records (e.g. Lane et al., 2013). This bi-partition was related to a northward shift of the westerly wind jet in the North Atlantic region. Our findings imply that the underlying climatic mechanism operated on a hemispheric rather than just on a regional scale.

### References:

Lane et al. 2013, Volcanic ash reveals time-transgressive abrupt climate change during the Younger Dryas, *Geology* 41, 1251-1254