



Aberystwyth University

Co-production of 11-hydroxyprogesterone and ethanol using recombinant yeast expressing fungal steroid hydroxylases

Hull, Claire M. ; Warrilow, Andrew G. S. ; Rolley, Nicola J. ; Price, Claire L. ; Donnison, Iain; Kelly, Diane E. ; Kelly, Steven L.

Published in:

Biotechnology for Biofuels

DOI:

[10.1186/s13068-017-0904-z](https://doi.org/10.1186/s13068-017-0904-z)

Publication date:

2017

Citation for published version (APA):

Hull, C. M., Warrilow, A. G. S., Rolley, N. J., Price, C. L., Donnison, I., Kelly, D. E., & Kelly, S. L. (2017). Co-production of 11-hydroxyprogesterone and ethanol using recombinant yeast expressing fungal steroid hydroxylases. *Biotechnology for Biofuels*, 10, [226]. <https://doi.org/10.1186/s13068-017-0904-z>

Document License

CC BY

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400

email: is@aber.ac.uk

RESEARCH

Open Access



Co-production of 11 α -hydroxyprogesterone and ethanol using recombinant yeast expressing fungal steroid hydroxylases

Claire M. Hull¹, Andrew G. S. Warrilow¹, Nicola J. Rolley¹, Claire L. Price¹, Iain S. Donnison², Diane E. Kelly^{1*} and Steven L. Kelly^{1*} 

Abstract

Background: Bioethanol production from sustainable sources of biomass that limit effect on food production are needed and in a biorefinery approach co-products are desirable, obtained from both the plant material and from the microbial biomass. Fungal biotransformation of steroids was among the first industrial biotransformations allowing corticosteroid production. In this work, the potential of yeast to produce intermediates needed in corticosteroid production is demonstrated at laboratory scale following bioethanol production from perennial ryegrass juice.

Results: Genes encoding the 11 α -steroid hydroxylase enzymes from *Aspergillus ochraceus* (11 α -SH^{Aoch}) and *Rhizopus oryzae* (CYP509C12) transformed into *Saccharomyces cerevisiae* for heterologous constitutive expression in p425TEF. Both recombinant yeasts (AH22:p11 α -SH^{Aoch} and AH22:p509C12) exhibited efficient progesterone bioconversion (on glucose minimal media containing 300 μ M progesterone) producing either 11 α -hydroxyprogesterone as the sole metabolite (AH22:p11 α -SH^{Aoch}) or a 7:1 mixture of 11 α -hydroxyprogesterone and 6 β -hydroxyprogesterone (AH22:p509C12). Ethanol yields for AH22:p11 α -SH^{Aoch} and AH22:p509C12 were comparable resulting in \geq 75% conversion of glucose to alcohol. Co-production of bioethanol together with efficient production of the 11-OH intermediate for corticosteroid manufacture was then demonstrated using perennial ryegrass juice. Integration of the 11 α -SH^{Aoch} gene into the yeast genome (AH22:11 α -SHAoch+K) resulted in a 36% reduction in yield of 11 α -hydroxyprogesterone to 174 μ mol/L using 300 μ M progesterone. However, increasing progesterone concentration to 955 μ M and optimizing growth conditions increased 11 α -hydroxyprogesterone production to 592 μ mol/L product formed.

Conclusions: The progesterone 11 α -steroid hydroxylases from *A. ochraceus* and *R. oryzae*, both monooxygenase enzymes of the cytochrome P450 superfamily, have been functionally expressed in *S. cerevisiae*. It appears that these activities in fungi are not associated with a conserved family of cytochromes P450. The activity of the *A. ochraceus* enzyme was important as the specificity of the biotransformation yielded just the 11-OH product needed for corticosteroid production. The data presented demonstrate how recombinant yeast could find application in rural biorefinery processes where co-production of value-added products (11 α -hydroxyprogesterone and ethanol) from novel feedstocks is an emergent and attractive possibility.

Keywords: Bioconversion, Bioethanol, Biorefinery, 11 α -Hydroxyprogesterone, Ryegrass, Yeast

*Correspondence: d.kelly@swansea.ac.uk; S.L.Kelly@swansea.ac.uk

¹ Institute of Life Science, Swansea University Medical School, Swansea SA2 8PP, Wales, UK

Full list of author information is available at the end of the article

Background

Biofuels such as bioethanol are a key focus for sustainability away from fossil fuels and in a biorefinery approach maximum value and product extraction is envisaged. For instance, plant protein or other products may be extracted prior to fermentation of carbohydrates. The microbial biomass also offers routes to co-products. Some of the first industrial biotransformations involved filamentous fungi for production of key intermediates in corticosteroid production that were hard to achieve by chemical synthesis. Ever since the first study reporting microbial 11α -hydroxylation of progesterone using *Rhizopus arrhizus* over 60 years ago [1], there has been sustained scientific and commercial interest in the application of bacteria [2–6] and filamentous fungi [7–9] to produce hydroxylated pharmaceuticals and intermediates. Structural hydroxylations are often necessary to functionalize steroid-based drugs that exhibit higher efficacy than their non-hydroxy analogs [10], and the ability to perform regio- and stereo-specific modifications using whole-cell microbial biocatalysis is invaluable for large-scale steroid manufacturing [11] not least because certain hydroxylations can prove difficult or even impossible to achieve via synthetic chemistry.

Unlike unicellular yeasts, filamentous fungi (e.g., *Rhizopus*, *Aspergillus*, and *Mucor* spp.) can hydroxylate the sterane skeleton (Fig. 1) at numerous positions (1β , 2β , 4, 5α , 6β , 7α , 7β , 9α , 10β , 11α , 11β , 12α , 12β , 14, 15α , 15β , 16α , 17α , 19, 22, 26) [12]. Alpha and beta hydroxylations at carbon 11 have for more than 50 years been performed commercially using filamentous fungi [11] and are particularly desirable for the synthesis of the glucocorticoid steroids that are commonly prescribed as anti-inflammatory, immunosuppressive, anti-allergic, and contraceptive drugs [13]. However, despite their industrial and biomedical value, species-specific information about 11α -steroid hydroxylases from filamentous fungi and data regarding their prevalence across the fungal kingdom remains limited; instead they are classified very broadly as monooxygenase enzymes of the cytochrome P450 (CYP) superfamily. In fact, sequencing data for just two fungal 11α -steroid hydroxylases from *Aspergillus ochraceus* [14] and *Rhizopus oryzae* [15] are currently available. Following historical neglect, it is hoped that the recent establishment of fungal genome databases and increasing interest in transcriptional data for filamentous species will now permit new biotechnological approaches that will harness the full biosynthetic potential of fungal cytochrome




Fig. 1 Steroid chemical structure. **a** Gonane, the steroid nucleus comprising three cyclohexane rings (rings A, B, and C) and one cyclopentane (ring D) labeled using the IUPAC-recommended ring-lettering and atom-numbering. **b** Hydroxylation of progesterone to 11 α -hydroxyprogesterone

P450 enzymes [16]. Filamentous fungal genomes can contain 100–200 cytochrome P450 genes, representing a large potential reservoir of biocatalysts.

It has long been known that the efficiency of using certain filamentous fungi (e.g., *Aspergillus fumigatus*) for steroid biotransformations can be compromised by the production of unwanted side-product steroids [9, 17], and there is much interest in the scope to optimize and apply recombinant yeast as ‘microbial cell factories’ in order to achieve specific hydroxylations [10, 13, 16]. Recombinant strains of the fission yeast *Schizosaccharomyces pombe* have been employed for steroid bioconversions [18] including the 11 α -hydroxylation of progesterone [15]. Designer strains of *Saccharomyces cerevisiae* that express mammalian steroid hydroxylases involved in xenobiotic metabolism (e.g., CYP3A4) have also been reported [19] that are capable of synthesizing hydrocortisone from glucose [20]. The scope of microbial biotechnology continues to expand and it is increasingly well recognized that the production of bio-based chemicals and biofuels from non-food carbohydrate feedstock using metabolically engineered yeast could offer a sustainable alternative to their petroleum-based production [21]. The application of recombinant yeast to assist the process development that is needed to overcome challenges in manufacturing and pharmaceutical industries has also been highlighted [22].

In the present study, we sought to develop a recombinant strain of *S. cerevisiae* that could hydroxylate progesterone at carbon 11 α (Fig. 1) and that could be applied to the rural biorefinery concept [23] to achieve co-production of 11 α -hydroxyprogesterone and bioethanol from grass juice feedstock [24]. We have previously shown that *S. pombe*, like *S. cerevisiae* cannot perform steroid hydroxylation in its native state [25] and it remains an attractive yeast host for heterologous protein expression. However, we chose to extend our work with *S. cerevisiae* [24, 26] because it is already employed in large-scale ethanol fermentations and because commercial and research ventures continue to yield more robust strains that are able to tolerate industrial conditions [21]. Grass juice sourced from *Lolium perenne* (ryegrass) represents one of several fractions of non-food plant biomass that are currently under investigation as feedstock for rural biorefinery and bio-based production processes in the United Kingdom [27–29]. It is rich in water-soluble carbohydrates (fructans, fructose, glucose, and sucrose) and its potential to support yeast growth and ethanol production has already been demonstrated [24, 26]. Therefore, producing a strain of *S. cerevisiae* that can produce both 11 α -hydroxyprogesterone and ethanol using grass juice as carbon source can increase the financial yield from the bioconversion. As the global market for pharmaceutical

steroid productions stands at around 10 billion USD [30] and the total cost of bioethanol is continually falling [31], using an integrative approach to coproduce a high-value platform chemical can increase the profitability of bioethanol production and agriculture.

Results from this study are the first to demonstrate functional expression of the 11 α -steroid hydroxylase enzyme, documented for *A. ochraceus* [14], in *S. cerevisiae* grown on grass juice feedstock. This underscores the potential to diversify the scope of the rural biorefinery concept through application of yeast biotechnology, which represents an emergent and valuable asset for industries of the low-carbon and green economy.

Results

Recombinant yeast: progesterone bioconversion

Metabolite profiles for yeast transformants grown on YM+His with 300 μ M progesterone indicate that the optimized gene sequences encode functional proteins that were constitutively expressed and conferred progesterone hydroxylating capabilities on AH22:p11 α -SH^{Aoch} and AH22:pCYP509C12 yeast (Table 1; Fig. 2). During initial growth experiments with YM+His with 300 μ M progesterone, almost total bioconversion of progesterone was achieved with AH22:p11 α -SH^{Aoch} which produced $\geq 90\%$ 11 α -hydroxyprogesterone as the sole metabolite after 48 h of growth (Table 1). AH22:pCYP509C12 also produced 11 α -hydroxyprogesterone, but with lower efficiency ($47.61 \pm 1.7\%$; t_{48h}) and lower purity as detectable quantities of 6 β -hydroxyprogesterone ($6.85 \pm 1.7\%$; t_{48h}), identified from fragmentation spectra containing a prominent fragment ion at m/z 346 (Fig. 3e), were present in ethyl acetate extracts recovered from cell-free AH22:pCYP509C12 culture supernatant. Control (AH22:p425TEF^{Ctrl}) yeast harboring empty p425TEF vector did not hydroxylate exogenous progesterone (Fig. 2), and only progesterone (Fig. 3a) or its trimethylsilyl (TMS) derivative (Fig. 3b) was detected in ethyl acetate extracts from AH22:p425TEF^{Ctrl} culture supernatant.

Recombinant yeast: growth and ethanol fermentation

In studies undertaken using the Bioscreen C plate reader with starting (t_0h) inoculum of 5×10^5 cells/mL and YM+His media (without progesterone supplementation), the growth parameters for AH22:p11 α -SH^{Aoch}, AH22:pCYP509C12, and AH22:p425TEF^{Ctrl} yeast were comparable (Table 2; Fig. 4). All constructs achieved similar final population densities (1.61 and 2.1×10^8 cells/mL) and exhibited uniform doubling times (typically 4–5 h) at room temperature (20 ± 2 °C). There were no significant differences between the ethanol yields recorded for AH22:p11 α -SH^{Aoch}, AH22:pCYP509C12, and AH22:p425TEF^{Ctrl} grown on YM+His (ANOVA,

Table 1 GC–MS parameters and diagnostic ions for progesterone (Pg) substrate and hydroxylated (hydroxyPg) products

Peak	Gas chromatography–mass spectrometry: steroid identification				Steroid: % total (t_{48h})		
	Steroid	RT (min)	[M] ⁺	Fragment ions m/z (relative intensity)	AH22:p425TEF ^{Ctrl}	AH22:p11 α -SH ^{Aoch}	AH22:pCYP509C12
a	Progesterone (Pg)	21.04	314	43 (999) 124 (887) 314 (701) 272 (430) 229 (362) 91 (329) 41 (314) 79 (313) 22 (238) 67 (234)	100	9.90 (\pm 1.2)	45.54 (\pm 3.1)
b	Pg + TMS (\times 1)	20.36	386	386 (999) 73 (820) 43 (365) 387 (311) 371 (145) 75 (141) 209 (132) 208 (118) 91 (101) 196 (97)			
c	11 α -hydroxyPg + TMS (\times 2)	21.61	474	73 (999) 474 (559) 43 (230) 192 (227) 475 (219) 75 (198) 193 (126) 208 (85) 74 (84) 476 (80)	0	90.10 (\pm 1.6)	47.61 (\pm 1.7)
d	11 α -hydroxyPg + TMS (\times 1)	23.29	402	73 (999) 43 (825) 75 (395) 402 (347) 91 (310) 312 (277) 197 (260) 143 (253) 169 (233) 105 (223)			
e	6 β -hydroxyPg + TMS (\times 1)	21.40	402	73 (999) 43 (906) 346 (712) 75 (688) 91 (386) 387 (347) 79 (295) 93 (262) 402 (240) 119 (226)	0	0	6.85 (\pm 1.7)
Maximum % bioconversion (Σ hydroxyPg)					0.00	91.70	57.86

Steroid (%) abundances are mean values from three replicates with associated standard deviations

TMS trimethylsilyl derivatized, RT retention time, [M]⁺ mass ion

$P < 0.05$, $n = 3$). All produced at least 16 mg/mL ethanol after 72 h fermentation; this equates to 78% of the theoretical maximum (20.4 mg/mL) for 4% glucose. In contrast, supplementation of YM+His with 300 μ M progesterone had an inhibitory effect on the growth of AH22:p425TEF^{Ctrl} cultures which exhibited slower population doubling times (9–10 h), reached lower population densities (1.6×10^8 cells/mL), and produced lower, more variable amounts of ethanol (12.5 ± 3.2 mg/mL) than those grown in the absence of progesterone (Table 2). Progesterone supplementation had no effect on the growth profiles or ethanol productivity of AH22:p11 α -SH^{Aoch} or AH22:pCYP509C12 yeast (Table 2). In view of their bioconversion efficiency and 11 α -hydroxyprogesterone specificity, only AH22:p11 α -SH^{Aoch} yeast was selected for subsequent process optimization studies.


Two-step (bioconversion–fermentation) batch culture studies

For AH22:p11 α -SH^{Aoch} yeast (Fig. 5), complete progesterone bioconversion to 11 α -hydroxyprogesterone was achieved after just 8 h when cultures were agitated continuously at 200 rpm at 20 °C. Following progesterone bioconversion (Step 1), decreasing the shaking regime from 200 to 30 rpm (at t_{10h}) was found to support optimal ethanol production of 18 ± 2 mg/mL (Fig. 5b) which

was 88% of the theoretical maximum ethanol yield attainable with 4% glucose and 10% higher than that achieved in Bioscreen studies. In summary, the use of higher t_0h inoculums (1×10^8 cells/mL) and vigorous culture agitation (200 rpm) facilitated progesterone bioconversion (Step 1). The higher cell densities (typically $\geq 3 \times 10^8$ cells/mL) present by t_{10h} and lower agitation speeds (30 rpm) employed during fermentation (Step 2) resulted in higher ethanol yields.

A biorefinery application for recombinant AH22:p11 α -SH^{Aoch}

Complete bioconversion of progesterone to 11 α -hydroxyprogesterone (Step 1) was achieved within 8 h when either grass juice with progesterone or 7% glucose YM+His with progesterone was used as feedstock for AH22:p11 α -SH^{Aoch} yeast maintained at 20 ± 2 °C with vigorous shaking (200 rpm). As expected, ethyl acetate extraction of 11 α -hydroxyprogesterone from the grass juice supernatant at the end (t_{10h}) of the bioconversion step (Fig. 6) resulted in the recovery of numerous phyto-chemicals that were identified using GC–MS (data not shown). The most abundant of these were 1-hexacosanol, phytol, α -tocopherol, 1-tetracosanol, 1-octacosanol, campesterol, stigmaterol, and β -sitosterol. Following resuspension of the AH22:p11 α -SH^{Aoch} cell pellet in fresh feedstock (without progesterone supplementation) and



after decreasing the culture agitation speed to 30 rpm for fermentation (Step 2), the maximum ethanol yields produced by AH22:p11 α -SH^{Aoch} yeast on both grass juice and 7% glucose YM+His were 30–32 mg/mL. These ethanol yields exceed 85% of the theoretical maximum (35.7 mg/mL) for 7% glucose (Fig. 6). At the end of the studies undertaken using grass juice feedstock, screening of yeast cells plated onto selective YM+His agar at the end of both Step 1 (bioconversion) and Step 2 (fermentation) recovered a high percentage (typically $\geq 95\%$) of colony forming AH22:p11 α -SH^{Aoch} yeast that had retained the plasmid.

Progesterone bioconversion obtained with 11 α -SH^{Aoch} integrated into *S. cerevisiae* clone

To establish whether expression of 11 α -SH^{Aoch} using the p425 plasmid was comparable to integrating the gene, an integrated expression cassette, 11 α -SH^{Aoch}+K, was produced. Initial growth experiments in YM+His with 300 μ M progesterone showed that AH22:11 α -SH^{Aoch}+K was able to achieve 58% conversion of progesterone to 11 α -hydroxyprogesterone (174 μ mol/L) after 48 h. Control AH22 cells did not hydroxylate progesterone as no steroid metabolites were detected upon extraction with ethyl acetate.

To investigate further whether 11 α -SH^{Aoch} was able to metabolize higher concentrations of progesterone, the starting concentration was increased from 300 to 954 μ M (0.3 g/L, in accordance with Žnidaršič et al. [32]) and the ratio of grass juice to YM, the inoculum size, and the incubation temperature were varied. Lowering the temperature from 30 to 24 $^{\circ}$ C increased the bioconversion of progesterone to 11 α -hydroxyprogesterone from 42–50 to 54–62% after 48 h (Table 3). Diluting the grass juice with YM and using a lower inoculum only marginally affected the bioconversion rates. The maximum conversion rate obtained using AH22:11 α -SH^{Aoch}+K of 62% (591 μ mol/L) was achieved after 48 h at 24 $^{\circ}$ C with a 1:1 mixture of grass juice and yeast minimal media (Table 3). This was a 3.4-fold increase in yield of 11 α -hydroxyprogesterone compared to that obtained using 300 μ M progesterone as substrate in YM+His and a 2.2-fold increase in yield compared to using clone AH22:p11 α -SH^{Aoch} in YM+His after 48 h at 30 $^{\circ}$ C with 300 μ M progesterone (Table 1).

Discussion

In situ hydroxylation of progesterone (Table 1) indicated both *A. ochraceus* 11 α -SH and *R. oryzae* CYP509C12 were native steroid hydroxylating enzymes. AH22:p11 α -SH^{Aoch} (Figs. 2, 3) produced 11 α -hydroxyprogesterone as the sole product and AH22:pCYP509C12 produced a 7:1 mixture of 11 α -hydroxyprogesterone and 6 β -hydroxyprogesterone (Fig. 5e); *A. ochraceus* 11 α -SH and *R. oryzae* CYP509C12 shared only 15% sequence identity (82 conserved amino acid residues) along the whole length of the two enzymes. However, NCBI BLAST2 analysis (http://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch&PROG_DEF=blastn&BLAST_PROG_DEF=megaBlast&BLAST_SPEC=blast2seq) identified 26% sequence identity (45 amino acid residues) in a 174 amino acid long region from the C-termini. The relatively low sequence identity between these two CYP enzymes suggests they belong to different CYP families. Native complementary DNA (cDNA) encoding the *A. ochraceus* 11 α -SH has previously been co-expressed with cDNA for human oxidoreductase in





Fig. 3 Fragmentation patterns (electron ionization) for progesterone substrate and hydroxylated metabolites. **a** Progesterone; **b** TMS-derivatized progesterone; **c** diTMS-derivatized 11 α -hydroxyprogesterone; **d** TMS-derivatized 11 α -hydroxyprogesterone; **e** TMS-derivatized 6 β -hydroxyprogesterone. (TMS = trimethylsilyl derivatized)

Table 2 Growth parameters and ethanol yields for recombinant yeast grown on 4% glcM^{his} ± Pg

Construct	ΔOD_{600} [t_{48h}]		Max. population doubling time (h)		Density: cells/mL [t_{72h}]		Ethanol: mg/mL [t_{72h}]	
	-Pg	+Pg	-Pg	+Pg	-Pg	+Pg	-Pg	+Pg
AH22;p425TEF ^{Ctrl}	1.61	0.98	4.6 (± 0.5)	9.0 (± 1.0)	2.1×10^8	1.6×10^8	15.9 (± 1.4)	12.5 (± 3.2)
AH22;p11 α -SH ^{Aoch}	1.65	1.63	4.3 (± 0.5)	4.5 (± 0.5)	2.2×10^8	2.1×10^8	16.4 (± 1.8)	16.2 (± 2.0)
AH22;pCYP509C12	1.61	1.62	4.4 (± 0.5)	4.6 (± 0.5)	2.0×10^8	2.1×10^8	16.5 (± 1.5)	16.0 (± 1.1)


Mean values from three replicates are shown along with the associated standard deviations. Underlined values are significantly lower than those from comparator experiments (ANOVA, $P < 0.05$; $n = 3$)




Spodoptera frugiperda (Sf-9) insect cells [14]. Similarly, cDNA encoding the *R. oryzae* CYP509C12 has also been co-expressed with its native redox partner (*R. oryzae* NAD[P]H-dependent reductase) in recombinant fission yeast, *S. pombe*, which were also able to perform 11 α - and 6 β -steroid hydroxylation [15]. It is noteworthy that homology modeling and site-directed-mutagenesis studies have recently proved successful in redirecting the regio-selectivity of CYP106A2 (a mammalian P450 with 15 β -hydroxylase and minor 11 α -hydroxylase activity)

from C15 to C11 [33]. The potential to bioengineer the specificity of fungal steroid hydroxylases (e.g., to remove the 6 β -activity of CYP509C12) also exists and advances in this area now demand much better understanding of the catalytic and structural features of fungal steroid hydroxylases.

The recombinant AH22 (*S. cerevisiae*) whole-cell system reported in the present study extends the scope and application of the technological achievements of previous authors [14, 15] and demonstrates the constitutive



expression of functional 11 α -steroid hydroxylating enzymes in Brewer's yeast. We chose to focus on the application of AH22:p11 α -SH^{Aoch} yeast after preliminary metabolite studies as this biotransformation was highly specific and active. The maximum bioconversion initially achieved using AH22:p11 α -SH^{Aoch} yeast and 300 μ M progesterone was very high (90.9% 11 α -hydroxyprogesterone or 273 μ mol/L) and did not require heterologous electron donors to cytochrome P450 such as NADPH cytochrome P450 reductase (Table 1). The inhibitory effect of exogenous progesterone on the growth and ethanol yields of AH22:p425TEF^{Ctrl} (Table 2; Fig. 4) was consistent with the effect of steroid toxicity on *S. cerevisiae* [34]. The growth rates and ethanol yields for AH22:p11 α -SH^{Aoch}



and AH22:pCYP509C12 grown on YM+His with or without progesterone were comparable suggesting that constitutive expression of *A. ochraceus* 11 α -SH and *R. oryzae* CYP509C12 served a protective role in recombinant AH22:p11 α -SH^{Aoch} and AH22:pCYP509C12 yeast. The ability of *Aspergillus* and *Rhizopus* spp. to hydroxylate and excrete steroids represents one mechanism through which filamentous species are believed to detoxify and so persist when exposed to steroids or similar molecules [35].

By increasing the starting inocula from 5×10^5 to 1×10^8 cells/mL in optimization studies undertaken with AH22:p11 α -SH^{Aoch} yeast (Fig. 5) and through initial application of vigorous culture agitation (Step 1: bioconversion, 200 rpm), we achieved 100% conversion of progesterone to 11 α -hydroxyprogesterone in just 8 h.

Table 3 Bioconversion of progesterone to 11 α -hydroxyprogesterone with AH22:11 α -SH^{Aoch}+K

Grass juice: yeast minimal	Inoculum (cells/mL)	Temperature (°C)	% product formed (μ mol/L)
1:0	1 \times 10 ⁷	30	47 (448)
1:0	1 \times 10 ⁷	24	54 (515)
1:0	1 \times 10 ⁵	30	44 (420)
1:0	1 \times 10 ⁵	24	58 (553)
1:1	1 \times 10 ⁷	30	44 (420)
1:1	1 \times 10 ⁷	24	58 (553)
1:1	1 \times 10 ⁵	30	44 (420)
1:1	1 \times 10 ⁵	24	62 (591)
1:3	1 \times 10 ⁷	30	42 (401)
1:3	1 \times 10 ⁷	24	59 (563)
1:3	1 \times 10 ⁵	30	50 (477)
1:3	1 \times 10 ⁵	24	60 (572)

Progesterone bioconversion rates obtained using the integrated clone AH22:11 α -SH^{Aoch}+K with 0.3 mg/mL (954 μ M) progesterone in growth media after 48 h

Here, it is likely that increased oxygenation of the culture medium favored the growth of AH22:p11 α -SH^{Aoch} yeast and facilitated the catalytic activity of constitutively expressed *A. ochraceus* 11 α -SH monooxygenase. During fermentation (Step 2), the presence of higher cell densities ($\geq 3 \times 10^8$ cells/mL) and the reduction of culture agitation speeds from 200 to 30 rpm (Fig. 5) improved the ethanol yields from 78% (Bioscreen studies) to 88%. These results reflect a reduction in the amount of glucose used for growth under semi-anaerobic conditions and an increase in the fermentation of glucose to ethanol; this is further evidenced by the lower ethanol yields recorded during fermentation (Step 2) regimes (100, 150 and 200 rpm) that increased the oxygenation of AH22:p11 α -SH^{Aoch} cultures (Fig. 5).

Agriculturally sourced grass juice feedstock was amenable to the two-step bioconversion–fermentation process developed using AH22:p11 α -SH^{Aoch} yeast (Fig. 6) presenting bioengineering opportunities within biorefinery settings in temperate regions [27–29]. Firstly, it is significant that high product (11 α -hydroxyprogesterone and ethanol) yields were achievable using grass juice feedstock at room temperature (20 \pm 2 °C). On an industrial scale, this could help reduce the need for energy-intensive heating and cooling regimes. Secondly, the solvent extraction of 11 α -hydroxyprogesterone product from the grass juice/progesterone feedstock resulted in the recovery of numerous phyto-chemicals including 1-hexacosanol, 1-tetracosanol, 1-octacosanol (all commercially relevant fatty alcohols), phytol and α -tocopherol (both vitamin E precursors with applications in cosmetics, detergents, and as antioxidants), and campesterol,

stigmasterol, and β -sitosterol (plant sterols with dietary health benefits). Commercial interest in high-value phyto-chemicals [23] strengthens the premise for using grass juice and other plant-based substrates as feedstock for progesterone bioconversion and the development of technologies that can optimize the recovery (extraction, separation, and purification) of multiple desirable products is an emergent and necessary area of research [36].

Further efforts towards yield optimization were focused on the production of the high-value platform chemical 11 α -hydroxyprogesterone by the 11 α -steroid hydroxylase enzyme from *A. ochraceus*. The enzyme was integrated into the *S. cerevisiae* genome to facilitate clone stability (no risk of losing the enzyme through plasmid loss over time). Increasing the progesterone concentration from 300 to 954 μ M and reducing the incubation temperature from 30 to 24 °C resulted in 2.2-fold greater bioconversion to 11 α -hydroxyprogesterone than obtained using the yeast strain containing the steroid hydroxylase on a plasmid (AH22:p11 α -SH^{Aoch}). The increased yield of product at lower temperature reduces the external energy requirements and therefore increases the cost effectiveness of the bioconversion of progesterone to 11 α -hydroxyprogesterone. *A. ochraceus* steroid 11 α -hydroxylase was also regiospecific, only hydroxylating progesterone at the 11 α - position, unlike *R. oryzae* steroid hydroxylase which hydroxylated progesterone at two positions (11 α - and 6 β -). Therefore, the use of *A. ochraceus* steroid 11 α -hydroxylase would be the more economical as further expense to separate hydroxyprogesterone isomers would not be required. However, further optimization is still required before process scale-up can take place [22]. High ethanol yields (14.9% v/v) have recently been achieved using pure inulin and soybean feedstock and an engineered yeast strain (*Saccharomyces* sp. W0) that expresses the inulinase gene from *Pichia guilliermondii* [37]. The use of grass juice feedstock extracted from high sugar varieties of ryegrass [29] and/or alternative (e.g., flocculating) host strains of *S. cerevisiae* that exhibit enhanced ethanol production [38] remains to be investigated. The influence of other factors including nutrient supplements, temperature, pH stress, and culture volume on ethanol yields from grass juice warrants investigation and the use of chemostat and bioreactor platforms represents the next step for research development [39].

In light of the successful bioconversion of progesterone to 11 α -hydroxyprogesterone by AH22:11 α -SH^{Aoch}+K, stable, genome-integrated designer yeasts that are able to exploit specific feedstocks for the co-production of high-value products provide an important avenue for commercial biotransformations of chemicals. For example, yeasts could be utilized for the production of dietary

mycoprotein, other myco-products (e.g., sterols, fatty acids, minerals, and B-complex vitamins), and specific platform chemicals (e.g., vitamin precursors, NADPH, and other cofactors required for growth) and opportunities for the development of industrial scale biotechnology exist here.

Conclusions

This study explored the potential for biotransformation for co-products to add value to a biofuel production process using a novel feedstock. Production of a key intermediate of corticosteroid production using fungal enzymes was demonstrated, a biotransformation of historical interest among biotransformations. Further biotransformations can be envisaged using the yeast biomass involved in bioethanol production as well as recovery of useful cellular components of yeast from the fermentation.

Methods

Sequence optimization

Genes encoding *A. ochraceus* 11 α -steroid hydroxylase (11 α -SH^{Aoch}) (Genbank accession number ABH71415.1) [14] and *R. oryzae* CYP509C12 (Genbank accession number RO3G_05077.1) [15] were designed with codon bias optimized for *S. cerevisiae* and the following modifications: (i) an AAA triplet (Kozak sequence) upstream of ATG initiation codons, (ii) external *NotI* single restriction sites (for the excision of whole fragments), and (iii) flanking *BamHI* and *XhoI* single restriction sites for the double-digest and direct ligation of genes into p425TEF (Dualsystems Biotech), a shuttle vector [ampicillin selection (bacteria) and *LEU2* auxotrophic marker (yeast)], that enables constitutive expression driven by the transcription elongation factor (TEF1) promoter [40]. Both genes were synthesized commercially (Eurofins MWG Operon) and supplied in pEX-K vector (kanamycin selection) for maintenance in *Escherichia coli*.

Plasmid construction and recombinant yeast

Optimized genes for *A. ochraceus* and *R. oryzae* were excised from pEX-K and ligated into pre-cut (*BamHI*, *XhoI*) p425TEF vector. The resulting plasmids (p11 α -SH^{Aoch} and pCYP509C12, respectively) were transformed into *E. coli* strain DH5 α (Novagen) that were selected and maintained at 37 °C using Luria–Bertani medium supplemented with sodium ampicillin (100 mg/L). Purified p11 α -SH^{Aoch} and pCYP509C12 plasmid DNA were isolated from *E. coli* cultures (Wizard Plus SV Minipreps, Promega) and transformed into *S. cerevisiae* strain AH22 (*MATa*, *leu2-3*, *112 his4-519 can1*) via electroporation to yield the experimental constructs AH22:p11 α -SH^{Aoch} and AH22:pCYP509C12. A control construct

(AH22:p425TEF^{Ctrl}) was created by transforming AH22 with p425TEF vector. All AH22 yeast transformants were selected and maintained at 30 °C using glucose-based yeast minimal (YM+His) plates containing (w/v) 2% agarose, 1.34% yeast nitrogen base without amino acids, 4% glucose, and L-histidine (100 mg/L).

Progesterone conversion

AH22:p11 α -SH^{Aoch}, AH22:pCYP509C12, and AH22:p425TEF^{Ctrl} transformants were screened for their ability to convert progesterone into hydroxylated products by inoculating 60 mL volumes of YM+His + 300 μ M progesterone with single colonies ($n = 3$) taken directly from transformation plates. Cells were grown in 100-mL flasks (30 \pm 2 °C, 200 rpm) and harvested by centrifugation (4000 rpm, 5 min) after 48 h. Residual progesterone substrate and hydroxyprogesterone products were extracted from 5 mL volumes of cell-free culture supernatant with 3 \times 2 mL volumes of ethyl acetate which were pooled and evaporated to dryness using a centrifugal evaporator (Heto Maxi dry plus). Extracts were stored at –20 °C prior to further analysis. Fresh 100 mM progesterone stocks (31.4 mg progesterone dissolved in 1 mL DMSO) were used to prepare YM+His + 300 μ M progesterone for all experimental work.

Gas chromatography–mass spectrometry (GC–MS) metabolite analysis

Prior to GC–MS analysis, steroid extracts were heated (70 °C, 10 min) to ensure complete dryness and derivatized using 100 μ L *N,O*-bis(trimethylsilyl)trifluoroacetamide-trimethylchlorosilane (BSTFA-TMCS [99:1]) and 100 μ L anhydrous pyridine heated at 70 °C for 2 h [41]. Trimethylsilyl (TMS)-derivatized sterols were analyzed using a 7890A GC–MS system (Agilent Technologies) with a DB-5MS fused silica column (30 m \times 0.25 mm \times 0.25 μ m film thickness; J&W Scientific). The oven temperature was initially held at 70 °C for 4 min, and then increased at 25 °C/min to a final temperature of 280 °C, which was held for a further 25 min. Samples were analyzed in splitless mode (1 μ L injection volume) using helium carrier gas, electron ionization (ion source temperature of 150 °C), and mass scanning from m/z 40 to 850 [41]. GC–MS data files were analyzed using MSD Enhanced ChemStation software (Agilent Technologies Inc.) to determine metabolite profiles for AH22:p11 α -SH^{Aoch}, AH22:pCYP509C12, and AH22:p425TEF^{Ctrl} and for derivation of integrated peak areas. Progesterone and hydroxyprogesterone metabolites were identified with reference to retention times and mass fragmentation patterns for known standards. Phyto-chemicals that were co-extracted from grass juice feedstock in 'Biorefinery pilot studies' were identified

using the NIST Mass Spectral Search Program (NIST/EPA/NIH Mass Spectral Library, Version 2.0 g, May 2011) and with reference to diagnostic fragmentation spectra.

Growth and ethanol production

Growth rate and ethanol productivity comparisons for AH22:p11 α -SH^{Aoch}, AH22:pCYP509C12, and AH22:p425TEF^{Ctrl} grown on YM+His with 300 μ M progesterone were undertaken in 100-well honeycomb microplates using a Bioscreen C (Oy Growth Curves Ab Ltd., Finland) as previously described [26]. Briefly, uniform starting (t_0h) culture densities were achieved by resuspending single colonies in YM+His (with or without 300 μ M progesterone) and diluting to achieve 5×10^5 cells/mL. Starting cultures were vortexed and dispensed into Bioscreen wells (3 \times 300 μ L replicates per colony suspension). All experiments were incubated at 20 ± 2 °C in the Bioscreen (no shaking regime) for 72 h, with optical density (at 600 nm) readings taken every 45 min. Growth data were exported from the Bioscreen in ASCII format prior to analysis using Excel (Microsoft Office 2003).

Growth parameters

In addition to manual (Neubauer haemocytometer) cell counts, ΔOD_{600} values (derived by subtracting the starting optical density from the final reading) were recorded. Maximum doubling times were estimated by dividing the natural logarithm of 2 by the fastest population growth rates (μ) derived using linear trend lines ($y = mx + c$, where y = vertical axis coordinate; m = gradient; x = horizontal axis coordinate and c = y axis intercept) fitted to log-transformed OD data. For ethanol analyses, Bioscreen measurements were suspended at specific time points (t_0h , t_{24h} , t_{48h} , and t_{72h}) and a 10 μ L volume of culture supernatant removed from representative experimental wells.

Ethanol content

Ethanol content was determined using a spectrophotometric ethanol assay kit (K-ETOH 11/06; Megazyme Ltd.) according to manufacturer's instructions. All samples were diluted 1000-fold with distilled water prior to analysis. Ethanol yields were calculated as a percentage of the maximal theoretical yield for glucose (0.51 g/g).

Two-step batch culture studies: optimization of shaking regime

Two-step (bioconversion–fermentation) batch culture studies were performed at 20 ± 2 °C with AH22:p11 α -SH^{Aoch} yeast grown in 100-mL conical flasks containing 60 mL volumes of YM+His with 300 μ M progesterone.

Step 1 (progesterone bioconversion) experiments were initiated with t_0h inoculums of 1×10^8 cells/mL that were agitated at speeds of 30, 100, 150, or 200 rpm for 48 h. One milliliter volumes were aseptically removed from experimental flasks at t_2h , t_4h , t_8h , t_{24h} , and t_{48h} in order to determine % conversion of progesterone to 11 α -hydroxyprogesterone. Step 2 (ethanol fermentation) optimization was pursued using AH22:p11 α -SH^{Aoch} that had been maintained (throughout Step 1) at 200 rpm on YM+His with 300 μ M progesterone. At t_{10h} , the shaking speed of these cultures was decreased to 150, 100, and 30 rpm or held at 200 rpm for a further 75 h. One milliliter volumes were removed from fermentation cultures for the determination of ethanol content at t_{24h} , t_{48h} , t_{60h} , and t_{72h} . The theoretical maximum ethanol yield for 4% (40 mg/mL) glucose (assuming 0.51 g/g) was calculated to be 20.4 mg/mL.

Biorefinery studies using grass juice feedstock

Biorefinery studies were performed using AH22:p11 α -SH^{Aoch} yeast and enzyme (*Lactobacillus paracasei* β -fructosidase) pretreated grass juice comprising 70 mg/mL monosaccharide [24] or synthetic 7% glucose YM+His (containing 70 mg/mL glucose equivalent) as feedstock. Grass juice was extracted from ryegrass *L. perenne* supplied by the Institute of Biological, Environmental Research and Rural Sciences (IBERS, Aberystwyth, UK), screened to remove large particulates, autoclaved (121 °C, 30 min), and pretreated with purified *L. paracasei* β -fructosidase [24] prior to use as a growth and fermentation substrate. Step 1 (progesterone bioconversion) was initiated with 60 mL volumes of grass juice + 300 μ M progesterone or 7% glucose YM+His + 300 μ M progesterone and t_0h inoculums of 1×10^8 cells/mL that were maintained in 100-mL conical flasks at 20 ± 2 °C and agitated at 200 rpm. After 10 h, AH22:p11 α -SH^{Aoch} cultures were harvested by centrifugation (4000 rpm, 5 min) and metabolites in the cell-free culture supernatant were extracted using ethyl acetate for GC–MS analysis. Cell pellets were then resuspended in 60 mL volumes of fresh grass juice or 7% glucose YM+His (without progesterone) and maintained for a further 120 h at 20 ± 2 °C and at a shaking speed of 30 rpm. One milliliter volumes were removed at t_{24h} , t_{48h} , t_{72h} , t_{96h} , and t_{120h} for the determination of ethanol content. The theoretical maximum ethanol yield for 7% (70 mg/mL) glucose (assuming 0.51 g/g) was calculated to be 35.7 mg/mL. For studies using grass juice feedstock, colony screening of AH22:p11 α -SH^{Aoch} cells (100 μ L of a 1×10^3 cells/mL dilution) plated onto selective YM+His agar was performed to calculate the recovery (%) of yeast that had retained the p11 α -SH^{Aoch} plasmid at the end of Step 1 (bioconversion; t_{10h}) and Step 2 (fermentation; t_{125h}).

Integration of 11 α -SH^{Aoch} into the genome of *S. cerevisiae*

The 11 α -SH^{Aoch} gene was synthesized (Eurofins MWG Operon) with optimization for *S. cerevisiae* and supplied in the vector pBSII SK(+). Forward and reverse PCR primers were prepared to produce a cassette for the transformation of *S. cerevisiae* AH22 cells. Primers were designed to include PCR priming sites for ends of the gene (how many bases?), 50-bp complementary sequences upstream of the *Ura3* gene and downstream of the *Ura3* gene, *Sall*/*SpeI* restriction sites for insertion of the KanMX marker, and *NotI* sites for excision from the plasmid. The PCR cassette was inserted into pBSII SK(+) at the *NotI* site (chosen as it had been altered to remove the *Sall* and *SpeI* sites) producing pBSII SK(+):11 α -SH^{Aoch}+K. Construct integrity was checked by DNA sequencing. A KanMX cassette was cut out of pAH3 vector using *Sall*–*SpeI* digest and inserted into the *Sall*/*SpeI* site of pBSII SK(+):11 α -SH^{Aoch}+K to give the final transformation cassette. pBSII SK(+):11 α -SH^{Aoch}+K was digested with *NotI* prior to transformation into chemically competent AH22 cells. Positive transformants were selected using geneticin in YPD and were checked by sequencing PCR products produced when using primers located upstream and downstream of the *Ura3* site, internal 11 α -SH^{Aoch} gene primers, and KanMx primers.

The ability of AH22:11 α -SH^{Aoch}+K to hydroxylate progesterone was assessed using the methodology described above, except bioconversion was also measure when 954 μ M (0.3 g/L) of progesterone was added to YM.

Abbreviations

BSTFA: *N,O*-bis(trimethylsilyl)trifluoroacetamide; DMSO: dimethylsulfoxide; EtOH: ethyl alcohol; GC–MS: gas chromatography–mass spectrometry; TMCS: trimethylchlorosilane; TMS: trimethylsilyl.

Authors' contributions

CMH undertook the experimental studies and drafted the manuscript initially. AGSW expressed and purified *Lactobacillus paracasei* β -fructosidase protein for the enzymatic pretreatment of grass juice used in biorefinery pilot studies and contributed to the preparation of the manuscript. NJR undertook the additional integration experimental studies. CLP performed the additional sample preparation and GCMS analysis. ISD and DEK participated in the design of the study and written work. DEK and SLK conceived the study, and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

Author details

¹ Institute of Life Science, Swansea University Medical School, Swansea SA2 8PP, Wales, UK. ² Institute of Biological, Environmental & Rural Sciences, Aberystwyth University, Gogerddan, Aberystwyth, Wales SY23 3EE, UK.

Acknowledgements

We thank Marcus Hull for technical support. Analytical facilities were provided by the EPSRC National Mass Spectrometry Service Centre (Swansea University, UK).

Competing interests

The authors declare that they have no competing interests.

Availability of supporting data

Not applicable.

Consent for publication

The authors hereby give consent for the publication of this manuscript in 'Biotechnology for Biofuels'.

Ethical approval and consent to participate

Not applicable.

Funding

Funding for this study was provided by the BEACON Convergence project supported by Welsh Government and the European Regional Development Fund (ERDF) of the European Union.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 13 July 2016 Accepted: 7 September 2017

Published online: 29 September 2017

References

- Murray HC, Peterson DH. Oxygenation of steroids by Mucorales fungi. U.S. Patent 2602769. 1952.
- Goodman JJ, Smith LL. 16 α -Hydroxy steroids. XI. 2 β -and 16 α -hydroxylation of 9 α -fluorohydrocortisone by strains of *Streptomyces roseochromogenes*. *Appl Microbiol*. 1961;9:372–5.
- Donova MV. Transformation of steroids by actinobacteria: a review. *Appl Biochem Microbiol*. 2007;43:1–14.
- Restaino OF, Marseglia M, De Castro C, Diana P, Forni P, Parrilli M, et al. Biotechnological transformation of hydrocortisone to 16 α -hydroxy hydrocortisone by *Streptomyces roseochromogenes*. *Appl Microbiol Biotechnol*. 2014;98:1291–9.
- Restaino OF, Marseglia M, Diana P, Borzacchiello MG, Finamore R, Vitiello M, et al. Advances in 16 α -hydroxy transformation of hydrocortisone by *Streptomyces roseochromogenes*. *Process Biochem*. 2016;51:1–8.
- Zhang W, Cui L, Wu M, Zhang R, Xie L, Wang H. Transformation of prednisolone to a 20 β -hydroxy prednisolone compound by *Streptomyces roseochromogenes* TS79. *Appl Microbiol Biotechnol*. 2011;92:727–35.
- Breskvar K, Hudnik-Plevnik T. Inducibility of Cytochrome P-450 and of NADPH-cytochrome C reductase in progesterone treated filamentous fungi *Rhizopus nigricans* and *Rhizopus arrhizus*. *J Steroid Biochem*. 1981;14:395–9.
- Samanta TB, Ghosh DK. Characterization of progesterone 11 α -hydroxylase of *Aspergillus ochraceus* TS: a cytochrome P-450 linked monooxygenase. *J Steroid Biochem*. 1987;28:327–32.
- Smith KE, Ahmed F, Williams RAD, Kelly SL. Microbial transformations of steroids—VIII. Transformation of progesterone by whole cells and microsomes of *Aspergillus fumigatus*. *J Steroid Biochem Mol Biol*. 1994;49:93–100.
- Donova MV, Egorova OV. Microbial steroid transformations: current state and prospects. *Appl Microbiol Biotechnol*. 2012;94:1423–47.
- Sambamurthy K, Kar A. Biotransformation process and its improvements with special reference to steroids. In: Sambamurthy K, Kar A, editors. *Pharm Biotechnology*. New Delhi: New Age International Ltd.; 2006. p. 295–304.
- Kristan K, Rižner TL. Steroid-transforming enzymes in fungi. *J Steroid Biochem Mol Biol*. 2012;129:79–91.
- Fernandes P, Cruz A, Angelova B, Pinheiro H, Cabral JM. Microbial conversion of steroid compounds: recent developments. *Enzym Microb Technol*. 2003;32:688–705.
- Bolten SL, Clayton RA, Easton AM, Engel LC, Messing DM, Ng JS, et al. *Aspergillus ochraceus* 11 α hydroxylase and oxidoreductase. U.S. Patent 7033807 B2. 2007.

15. Petrič Š, Hakki T, Bernhardt R, Žigon D, Črešnar B. Discovery of a steroid 11 α -hydroxylase from *Rhizopus oryzae* and its biotechnological application. *J Biotechnol*. 2010;150:428–37.
16. Kelly SL, Kelly DE. Microbial cytochromes P450: biodiversity and biotechnology. Where do cytochromes P450 come from, what do they do and what can they do for us? *Philos Trans R Soc Lond B Biol Sci*. 2013;368:20120476.
17. Mukherjee A, Banerjee S, Mahato SB. Metabolism of progesterone by *Aspergillus fumigatus*. *J Steroid Biochem*. 1982;17:443–6.
18. Dragan C, Zearo S, Hannemann F, Bernhardt R, Bureik M. Efficient conversion of 11-deoxycortisol to cortisol (hydrocortisone) by recombinant fission yeast *Schizosaccharomyces pombe*. *FEMS Yeast Res*. 2005;5:621–5.
19. Mehmood Z, Kelly DE, Kelly SL. Metabolism of the herbicide chlortoluron by human cytochrome P450 3A4. *Chemosphere*. 1995;31:4515–29.
20. Kelly D, Kelly S. Rewiring yeast for drug synthesis. *Nat Biotechnol*. 2003;21:133–4.
21. Nielsen J, Larsson C, van Maris A, Pronk J. Metabolic engineering of yeast for production of fuels and chemicals. *Curr Opin Biotechnol*. 2013;24:398–404.
22. Martinez CA, Rupasinghe SG. Cytochrome P450 bioreactors in the pharmaceutical industry: challenges and opportunities. *Curr Top Med Chem*. 2013;13:1470–90.
23. Clark JH, Luque R, Matharu AS. Green chemistry, biofuels, and biorefinery. *Annu Rev Chem Biomol Eng*. 2012;3:183–207.
24. Martel CM, Warrilow AGS, Jackson CJ, Mullins JGL, Togawa RC, Parker JE, et al. Expression, purification and use of the soluble domain of *Lactobacillus paracasei* β -fructosidase to optimise production of bioethanol from grass fructans. *Bioresour Technol*. 2010;101:4395–402.
25. Pajić T, Vitas M, Žigon D, Pavko A, Kelly SL, Komel R. Biotransformation of steroids by the fission yeast *Schizosaccharomyces pombe*. *Yeast*. 1999;15:639–45.
26. Martel CM, Parker JE, Jackson CJ, Warrilow AGS, Rolley N, Greig C, et al. Expression of bacterial levanase in yeast enables simultaneous saccharification and fermentation of grass juice to bioethanol. *Bioresour Technol*. 2011;102:1503–8.
27. Morris SM, Jackson CJ, Gallagher JA, Kelly RO, Donnison IS. High-sugar perennial ryegrass as a bioethanol feedstock. An alternative use for forage grasses. In: Proceedings 16th Europe Biomass Conference Exhibition ETA-Florence Renewable Energies. ETA-Florence Renewable Energies; 2008. p. 60–6. <http://cadair.aber.ac.uk/dspace/handle/2160/13953>.
28. Charlton A, Elias R, Fish S, Fowler P, Gallagher J. The biorefining opportunities in Wales: understanding the scope for building a sustainable, biorenewable economy using plant biomass. *Chem Eng Res Des*. 2009;87:1147–61.
29. Farrar K, Bryant DN, Turner L, Gallagher JA, Thomas A, Farrell M, et al. Breeding for bio-ethanol production in *Lolium perenne* L.: association of allelic variation with high water-soluble carbohydrate content. *Bioenergy Res*. 2012;5:149–57.
30. Bureik M, Bernhardt R. Steroid hydroxylation: microbial steroid biotransformations using cytochrome P450 enzymes. In: Schmid RD, Urlacher VB, editors. *Modern biooxidation enzymes reactions applications*. Weinheim: Wiley-VCH Verlag GmbH & Co. KGaA; 2007. p. 155–76.
31. Saini JK, Saini R, Tewari L. Lignocellulosic agriculture wastes as biomass feedstocks for second-generation bioethanol production: concepts and recent developments. *3 Biotech*. 2015;5:337–53.
32. Žnidaršič P, Komel R, Pavko A. Studies of a pelleted growth form of *Rhizopus nigricans* as a biocatalyst for progesterone 11 α -hydroxylation. *J Biotechnol*. 1998;60:207–16.
33. Nguyen KT, Virus C, Günnewich N, Hannemann F, Bernhardt R. Changing the regioselectivity of a P450 from C15 to C11 hydroxylation of progesterone. *ChemBioChem*. 2012;13:1161–6.
34. Cvelbar D, Žist V, Kobal K, Žigon D, Žakelj-Mavrič M. Steroid toxicity and detoxification in Ascomycetous fungi. *Chem Biol Interact*. 2013;202:243–58.
35. Črešnar B, Žakelj-Mavrič M. Aspects of the steroid response in fungi. *Chem Biol Interact*. 2009;178:303–9.
36. Znidarsic-Plazl P, Plazl I. Steroid extraction in a microchannel system-mathematical modelling and experiments. *Lab Chip*. 2007;7:883–9.
37. Zhang T, Chi Z, Chi Z, Parrou J-L, Gong F. Expression of the inulinase gene from the marine-derived *Pichia guilliermondii* in *Saccharomyces* sp. W0 and ethanol production from inulin. *Microb Biotechnol*. 2010;3:576–82.
38. Choi G-W, Kang H-W, Moon S-K. Repeated-batch fermentation using flocculent hybrid, *Saccharomyces cerevisiae* CHFY0321 for efficient production of bioethanol. *Appl Microbiol Biotechnol*. 2009;84:261–9.
39. Thiry M, Cingolani D. Optimizing scale-up fermentation processes. *Trends Biotechnol*. 2002;20:103–5.
40. Mumberg D, Müller R, Funk M. Yeast vectors for the controlled expression of heterologous proteins in different genetic backgrounds. *Gene*. 1995;156:119–22.
41. Martel CM, Parker JE, Bader O, Weig M, Gross U, Warrilow AGS, et al. A clinical isolate of *Candida albicans* with mutations in ERG11 (encoding sterol 14-demethylase) and ERG5 (encoding C22 desaturase) is cross resistant to azoles and amphotericin B. *Antimicrob Agents Chemother*. 2010;54:3578–83.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

