

## Aberystwyth University

### *Hydrology of debris-covered glaciers in High Mountain Asia*

Miles, Katie E.; Hubbard, Bryn; Irvine-fynn, Tristram D.I.; Miles, Evan S.; Quincey, Duncan J.; Rowan, Ann V.

*Published in:*  
Earth-Science Reviews

*DOI:*  
[10.1016/j.earscirev.2020.103212](https://doi.org/10.1016/j.earscirev.2020.103212)

*Publication date:*  
2020

*Citation for published version (APA):*

Miles, K. E., Hubbard, B., Irvine-fynn, T. D. L., Miles, E. S., Quincey, D. J., & Rowan, A. V. (2020). Hydrology of debris-covered glaciers in High Mountain Asia. *Earth-Science Reviews*, 207, [103212].  
<https://doi.org/10.1016/j.earscirev.2020.103212>

**Document License**  
CC BY-NC-ND

**General rights**

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

# Hydrology of debris-covered glaciers in High Mountain Asia

Author's post-print/accepted version  
(CC-BY-NC-ND)

DOI of published version (Earth-Science Reviews):

<https://doi.org/10.1016/j.earscirev.2020.103212>

# Hydrology of debris-covered glaciers in High Mountain Asia

Katie E. Miles<sup>1\*</sup>, Bryn Hubbard<sup>1</sup>, Tristram D. L. Irvine-Fynn<sup>1</sup>, Evan S. Miles<sup>2</sup>, Duncan J. Quincey<sup>3</sup>  
and Ann V. Rowan<sup>4</sup>

*1. Centre for Glaciology, Department of Geography and Earth Sciences, Aberystwyth University, Aberystwyth, UK*

*2. Swiss Federal Research Institute WSL, 8903 Birmensdorf, Switzerland*

*3. School of Geography, University of Leeds, Leeds, UK*

*4. Department of Geography, University of Sheffield, Sheffield, UK*

\* Correspondence to: [kam64@aber.ac.uk](mailto:kam64@aber.ac.uk)

## Key words


Glaciers; debris-covered glaciers; glacier hydrology; High Mountain Asia

## Abstract

The hydrological characteristics of debris-covered glaciers are known to be fundamentally different from those of clean-ice glaciers, even within the same climatological, geological, and geomorphological setting. Understanding how these characteristics influence the timing and magnitude of meltwater discharge is particularly important for regions where downstream communities rely on this resource for sanitation, irrigation, and hydropower, as in High Mountain Asia. The hydrology of debris-covered glaciers is complex: rugged surface topographies typically route meltwater through compound supraglacial-englacial systems involving both channels and ponds, as well as pathways that remain unknown. Low-gradient tongues that extend several kilometres retard water conveyance and promote englacial storage. Englacial conduits are frequently abandoned and reactivated as water supply changes, new lines of permeability are exploited, and drainage is captured due to high rates of surface and subsurface change. Seasonal influences, such as the monsoon, are superimposed on these distinctive characteristics, reorganising surface and subsurface drainage rapidly from one season to the next. Recent advances in understanding have mostly come from studies aimed at quantifying and describing supraglacial processes; little is known about the subsurface hydrology, particularly the nature (or even existence) of subglacial drainage. In this review, we consider in turn the supraglacial, englacial, subglacial, and proglacial hydrological domains of debris-covered glaciers in High Mountain Asia. We summarise different lines of evidence to establish the current state of knowledge and, in doing so, identify major knowledge gaps. Finally, we use this information to suggest six themes for future hydrological research at High Mountain Asian debris-covered glaciers in order to make timely long-term predictions of changes in the water they supply.

36 **1. Introduction**

37 Debris-covered glaciers have gained increased research attention over recent years, partly in  
38 recognition of their role as water sources for large parts of the world's population (Immerzeel et  
39 al., 2020; Scherler et al., 2011) and partly because they host a range of distinctive features, driven  
40 by processes that are largely absent from their clean-ice counterparts. Definitions of what  
41 constitutes a 'debris-covered glacier' vary widely (e.g. Anderson, 2000; Kirkbride, 2011), but here  
42 we define them to be glaciers with a largely continuous layer of supraglacial debris over most of  
43 the ablation area, typically increasing in thickness towards the terminus (Figure 1). Debris can be  
44 supplied to such glaciers by snow avalanches, rockfalls, and landslides from local mountainsides  
45 onto the glacier surface (Figure 2, 3A), melt-out of englacial debris, thrusting transporting debris  
46 from the glacier bed, dust blown from exposed moraines, or solifluction from (ice-cored) moraines  
47 (Dunning et al., 2015; Gibson et al., 2017b; Hambrey et al., 2008; Kirkbride and Deline, 2013;  
48 Rowan et al., 2015; van Woerkom et al., 2019). The surface debris layer can range in thickness  
49 from scattered particles to several metres, including large rocks and substantial boulders (Figure  
50 3C and D) (Inoue and Yoshida, 1980; McCarthy et al., 2017; Nicholson et al., 2018).



51

Figure 1 – Debris-covered glaciers in the Sagarmatha National Park, Nepal Himalaya, annotated with some of the features distinctive to High Mountain Asian debris-covered glaciers. **A)** Chola Glacier (image width is ~1.5 km across the glacier terminus and lake). **B)** Imja-Lhotse Shar Glacier, showing the terminus and calving front (~0.75 km width) into Imja Tsho, looking towards the accumulation area of the tributary Amphulapcha Glacier. **C)** Khumbu Glacier, showing the upper ablation area (clean-ice flowing from the Khumbu Icefall) to the left and the ~8 km long lower

*ablation area (debris-covered tongue) to the right; dashed yellow lines are 100 m contours. Image credit for A and B: Katie Miles; and C: Tristram Irvine-Fynn.*


52 Debris-covered glaciers are present in nearly all of Earth's glacierised regions, with a  
53 particularly large concentration in High Mountain Asia (Bolch et al., 2012; Kraaijenbrink et al.,  
54 2017; Scherler et al., 2018); sub-regional variability in the debris cover of which is presented in  
55 Brun et al. (2019) (their Figure 1). Debris-covered glaciers therefore contribute an important  
56 proportion of streamflow used for drinking water, irrigation, and hydroelectric power; this  
57 streamflow is particularly effective in reducing seasonal water shortages (Bolch et al., 2019;  
58 Immerzeel et al., 2020, 2010; Pritchard, 2019; Scott et al., 2019). Glacier mass loss in response to  
59 climate warming is currently increasing river discharge and contributions to sea level (Hock et al.,  
60 2019; Lutz et al., 2014; Radić et al., 2014; Shea and Immerzeel, 2016), but studies simulating future  
61 scenarios universally project long-term reductions in flow, perhaps as soon as 2050 in central Asia  
62 (Barnett et al., 2005; Bolch et al., 2012; Huss and Hock, 2018; Lutz et al., 2014; Ragettli et al., 2016b;  
63 Rounce et al., 2020; Sorg et al., 2012). Passing of 'peak water' threatens future water security in  
64 many regions, particularly across High Mountain Asia (Bolch et al., 2019; Eriksson et al., 2009;  
65 Hannah et al., 2005; Huss and Hock, 2018; Immerzeel et al., 2010; Winiger et al., 2005). A decrease  
66 in discharge from the Indus and Brahmaputra rivers alone is estimated to affect 260 million people  
67 (Immerzeel et al., 2010).

68 The long-term response of debris-covered glaciers to changing climatic conditions is non-  
69 linear and results from complexities relating to spatial variability in debris concentration and  
70 climatic controls integrated over at least several decades (Benn et al., 2012; Vaughan et al., 2013).  
71 A multidecadal trend of surface lowering, stagnation, and glacier mass loss has already been  
72 observed on many debris-covered glaciers across High Mountain Asia (Bolch et al., 2012, 2011;  
73 Brun et al., 2017; Dehecq et al., 2019; Hock et al., 2019; Käab et al., 2012; Pellicciotti et al., 2015;  
74 Scherler et al., 2011) as a result of warmer air temperatures and weaker monsoons (Pieczonka et  
75 al., 2013; Thakuri et al., 2014). However, predictions of mass loss from individual glacierised  
76 regions vary considerably. For example, in the Everest region of the Himalaya, estimates of ice  
77 mass loss by 2100 vary from ~10% (Rowan et al., 2015), through 50% (Soncini et al., 2016), to 99%  
78 in extreme scenarios (warming of ~3°C) (Shea et al., 2015). Model outputs also vary spatially at a  
79 regional scale (e.g. Chaturvedi et al., 2014; Kraaijenbrink et al., 2017; Zhao et al., 2014). Such  
80 projections depend on the future climate scenario used, but a number of key knowledge gaps also  
81 exist concerning the character of debris-covered glaciers and the processes influencing their varied  
82 geometrical response to climate change (Benn et al., 2012; Bolch et al., 2012; Huss, 2011; Scherler  
83 et al., 2011).

84 Understanding how meltwater is produced, transported, and stored within High Mountain  
85 Asian debris-covered glaciers is therefore imperative. There is growing recognition that the  
86 configuration and efficiency (i.e. bulk system transit velocity) of water routing across and through  
87 debris-covered ice is distinctively different from that of clean-ice glaciers, even within the same  
88 glacial system. This was first shown by a recent study on Miage Glacier, a debris-covered glacier in  
89 the Italian Alps (Fyffe et al., 2019b). Debris-covered glacier surfaces are complex, particularly those  
90 in High Mountain Asia, the ablation areas of which are often characterised by hummocky, rugged

91 topography atop a shallow (or even reversed) longitudinal surface gradient (Figures 1 and 2). This  
92 commonly results from an inverted mass-balance regime, where the greatest ablation rates are  
93 experienced in the middle, rather than lower, ablation area (King et al., 2017). Debris-covered  
94 ablation areas also exhibit bare ice cliffs and supraglacial ponds – depressions capable of storing  
95 meltwater for both short and long periods within nested catchments of varying spatial scales  
96 (Section 2) – and these glaciers frequently terminate in proglacial lakes (Section 5). Other unique  
97 characteristics of High Mountain Asian debris-covered glaciers include the accumulation areas  
98 often being at extremely high elevations, with a steep surface gradient (often an icefall)  
99 transporting ice into the ablation area (Figure 1). These features provide a setting that strongly  
100 influences the nature of hydrological systems in this region (Benn et al., 2017; Miles et al., 2019),  
101 but has restricted hydrological research due to the remoteness and inaccessibility of such glaciers.

102 In this review, we consider the current state of knowledge of debris-covered glacier  
103 hydrological systems in High Mountain Asia. Four hydrological domains are considered in turn:  
104 supraglacial (Section 2), englacial (Section 3), subglacial (Section 4), and proglacial (Section 5).  
105 Within each section, we summarise existing research and understanding of debris-covered glacier  
106 hydrological systems and then address key remaining knowledge gaps. Figure 2 provides a  
107 reference conceptual diagram of a (High Mountain Asian) debris-covered glacier, with each  
108 hydrological feature encompassing both known and unknown elements of each domain. Finally,  
109 in light of the review, we propose six future research themes concerning the hydrology of debris-  
110 covered glaciers (Section 6). This review is intended to complement existing reviews of clean-ice  
111 valley glacier hydrology (e.g. Fountain and Walder, 1998; Hubbard and Nienow, 1997; Irvine-Fynn  
112 et al., 2011; Jansson et al., 2003). We note that there are a number of differing climatic regimes  
113 across High Mountain Asia, with precipitation in particular varying closely with topography  
114 (Bookhagen and Burbank, 2006); these climatic regimes will influence the thermal regime,  
115 geometry, mass balance, and thus hydrology of the glaciers in each of these sub-regions. While  
116 our review draws on research carried out across High Mountain Asia, much of that research has  
117 been carried out in the monsoon-influenced Himalaya, particularly Nepal, from where the review  
118 and our illustrations of many of the key elements draw strongly.



119

*Figure 2 – A conceptual illustration of the main landscape and hydrological features of a typical debris-covered glacier. Features specific to debris-covered glaciers in High Mountain Asia are labelled in Figure 1.*


## 120 2. Supraglacial hydrology

### 121 2.1 Supraglacial zone

#### 122 2.1.1 Meltwater generation

123 Supraglacial meltwater is produced on debris-covered glaciers through ablation of surface ice and  
 124 snow, with the spatial pattern of melt complicated by the surface debris extent, thickness, and  
 125 lithological characteristics (Figures 1 and 3). A debris layer shallower than the critical thickness,  
 126 typically  $\sim 0.05$  m, decreases albedo and thus increases the ablation rate compared to debris-free  
 127 ice (Figure 4). The ablation rate peaks at a debris thickness of  $\sim 0.02\text{--}0.05$  m, known as the effective  
 128 thickness (Adhikary et al., 2000; Evatt et al., 2015; Inoue and Yoshida, 1980; Juen et al., 2014;  
 129 Kayastha et al., 2000; Lejeune et al., 2013; Nicholson and Benn, 2013, 2006; Østrem, 1959; Singh  
 130 et al., 2000; Takeuchi et al., 2000). The exact values of the critical and effective thickness strongly  
 131 depend on the debris thermal conductivity, which can vary widely both across a glacier surface  
 132 and in time according to whether the debris is wet or dry (Casey et al., 2012; Collier et al., 2015,  
 133 2014; Gibson et al., 2017b; Nicholson and Benn, 2013; Pelto, 2000). In contrast, a debris layer  
 134 thicker than the critical thickness of  $\sim 0.05$  m insulates the ice from incoming solar radiation,  
 135 inhibiting the receipt of surface energy at the ice-debris interface and thus reducing the melt rate  
 136 (Figure 4). Beneath a debris thickness of  $0.25\text{--}0.30$  m, ice becomes almost fully insulated from

137 daily surface energy fluxes, with only longer-term changes in surface energy balance reaching the  
 138 underlying debris-ice interface (Bocchiola et al., 2015; Brock et al., 2010; Conway and Rasmussen,  
 139 2000; Nicholson and Benn, 2013; Østrem, 1959; Reid and Brock, 2010). In addition, turbulent  
 140 energy fluxes have been shown to reduce net radiative fluxes at the debris surface (of a 0.75 m  
 141 thick debris layer) by 17% over a full melt season, further diminishing the energy available for melt  
 142 at the ice-debris interface (Steiner et al., 2018b). Variations in ablation according to these factors  
 143 represent an important first-order control on glacier surface morphology and are partially  
 144 responsible for the characteristic hummocky topography superimposed on a shallow or concave-  
 145 upward (reversed gradient) debris-covered glacier surface profile (Figure 1).




146

*Figure 3 – Images illustrating debris transport processes, englacial debris inclusions, and variations in supraglacial debris thickness on Khumbu Glacier, Nepal Himalaya: **A**) a landslide scar (yellow circle, ~500 m wide) and unstable rock faces (purple circle) providing debris to the glacier surface; image is taken looking east across the surface of Khumbu Glacier, and the debris layer above ice cliffs can also be seen. **B**) an ice cliff with entrained debris (green circle), debris-rich ice layers (orange circle), and a moderately thick (~1–2 m) surface debris layer; **C**) a thin (~0.20 m; red arrow) surface debris layer above ice adjacent to a supraglacial pond; and **D**) a thick (> 5 m; red arrow) surface debris layer above an ice cliff. Image credit for A: Duncan Quincey; and B–D: Katie Miles.*

147 Counteracting the influence of a thick surface debris layer, the ablation rate of debris-  
 148 covered glaciers is enhanced by the presence of supraglacial ponds (Section 2.1.2) and ice cliffs



149 (Figure 3B and D). The latter form by slumping of debris from steep slopes, calving at supraglacial  
 150 pond margins (Section 2.1.2), or the collapse of englacial voids (Section 3.1), all of which expose  
 151 steep, bare ice (Figure 3B) or thinly debris-covered faces (Figure 3D) at the glacier surface (Benn  
 152 et al., 2012, 2001; Sakai et al., 2002; Thompson et al., 2016). The melting of ice cliffs can be  
 153 responsible for a substantial proportion of debris-covered glacier ablation (Brun et al., 2016; Buri  
 154 et al., 2016b; Han et al., 2010; Juen et al., 2014; Reid and Brock, 2014; Sakai et al., 2002, 2000;  
 155 Thompson et al., 2016), accounting for 23–69% of the total ablation of debris-covered areas whilst  
 156 covering a small proportion of the total glacier area. The ice cliffs exhibit melt rates that are 3–14  
 157 times higher than beneath debris-covered ice (Brun et al., 2018; Immerzeel et al., 2014; Sakai et  
 158 al., 1998). Where ice cliffs are associated with supraglacial ponds, there is further potential for  
 159 increased melting through undercutting and calving processes (Brun et al., 2016; Buri et al., 2016a;  
 160 Miles et al., 2016; Röhl, 2008; Thompson et al., 2016). Taken together, ice cliff and pond systems  
 161 may contribute considerably to the surface lowering of debris-covered glaciers in the central  
 162 ablation area (King et al., 2017; Nuimura et al., 2012; Pellicciotti et al., 2015; Ragettli et al., 2016a;  
 163 Thompson et al., 2016; Watson et al., 2017), contributing to the inverted mass-balance regime  
 164 typical of High Mountain Asian debris-covered glaciers.



165

*Figure 4 – Østrem curve examples showing variations in the relationship between debris thickness and ice ablation on different glaciers. (a) notes the effective thickness, namely the debris thickness at which maximum melt occurs. (b) marks the critical thickness, the debris thickness at which melt becomes inhibited compared to that of clean ice on different glaciers (indicated on both for Isfjällsglaciaren). From Nicholson & Benn (2006).*

166 **2.1.2 Meltwater storage**

167 Supraglacial ponds (Figure 5), a term used here to include larger water bodies elsewhere  
 168 sometimes referred to as lakes, are common and important features on debris-covered glaciers.  
 169 Ponds are generally absent from clean-ice valley glaciers but are prevalent on low-gradient areas  
 170 of ice sheet margins (Chu, 2014; Sundal et al., 2009). Similarly for debris-covered glaciers, the most  
 171 important control on the location of supraglacial pond formation is a low glacier surface slope  
 172 (Miles et al., 2017b; Quincey et al., 2007; Reynolds, 2000; Sakai, 2012; Sakai et al., 2000; Sakai and

173 Fujita, 2010; Salerno et al., 2012). A surface gradient of  $\leq 2^\circ$  is considered to promote the  
174 development of larger ponds, while smaller isolated and transient ponds are considered more  
175 likely on steeper slopes (Miles et al., 2017b; Quincey et al., 2007; Reynolds, 2000). The upglacier  
176 slope has also been shown to have an influence, being inversely correlated to the total area of  
177 lakes downglacier (Salerno et al., 2012).

178           Glacier velocity and motion type also exert controls over supraglacial pond location. An  
179 increase in lake concentration is common towards the termini of debris-covered glaciers, areas  
180 that are typically characterised by low surface velocities (Kraaijenbrink et al., 2016b; Miles et al.,  
181 2017b; Quincey et al., 2007; Sakai, 2012; Salerno et al., 2015, 2012). A decrease in velocity towards  
182 both the glacier terminus and ice inflow at the confluences of flow units (Kraaijenbrink et al.,  
183 2016b) causes compressive flow, which tends to close crevasses and drive water back to the  
184 surface, as well as limiting effective drainage from the glacier surface (Kraaijenbrink et al., 2016b;  
185 Miles et al., 2017b). The thinning and stagnation of debris-covered glacier termini may also  
186 enhance meltwater production, further promoting the formation of ponds (Salerno et al., 2015;  
187 Thakuri et al., 2016).




Figure 5 – Examples of supraglacial pond size and temporal changes on Khumbu Glacier, Nepal Himalaya. Ponds range in diameter from: **A)** several metres; **B)** tens of metres (person circled in red for scale); **C)** and **D)** hundreds of metres; **E)** and **F)** several kilometres. A) and B) are located in the upper ablation area. C) and D) show the same pond-cliff-cave system in the mid-ablation area two years apart, with notable expansion of the cave via undercutting and calving. The pond, which has reduced in area (likely partly drained), was filled with a large amount of small, calved ice blocks in May 2019 and large cracks in the cliff system suggest further imminent large-scale calving. E) and F) show the expanding linked supraglacial pond chain at the terminus, also two years apart (green star indicates the same location as images were taken from slightly different positions). Pond growth and coalescence has progressively eroded the hummocks that formerly separated these

*ponds. Higher melt rates are indicated by the covering of ice cliffs in fine debris ('dirty ice'). Image credit for A–D and F: Katie Miles; and E: Evan Miles.*

188 Initial supraglacial pond growth occurs primarily through subaqueous melting at the base  
189 of any slight depression (Chikita et al., 1998; Mertes et al., 2016; Miles et al., 2016; Stokes et al.,  
190 2007; Thompson et al., 2012). Water accumulates and is heated by incoming solar radiation,  
191 causing the pond to warm. For example, Chikita et al. (1998) measured a maximum temperature  
192 of  $\sim 5^{\circ}\text{C}$  at a supraglacial pond surface on Trakarding Glacier, Nepal Himalaya. Excess energy is thus  
193 available for lateral and vertical ablation wherever pond water is in contact with ice, increasing the  
194 pond size, steepening marginal slopes and mobilising debris to expose bare ice (Figure 5E and F)  
195 (Stokes et al., 2007). Subaqueous pond melt rates are greatest when bare ice is exposed or covered  
196 in a thin layer of debris; layers of thick sediment at the base of ponds effectively terminate bottom  
197 deepening by preventing transfer of energy from the warmer pond water to the ice surface  
198 (Horodyskyj, 2015). Furthermore, mixing of pond stratification by inflowing meltwater on Koxkar  
199 Glacier, Tien Shan, has been shown to increase the temperature (by  $\sim 4^{\circ}\text{C}$ ) and density of the pond  
200 (Wang et al., 2012). Here, the warmed surface water sinks to the pond base and increases the  
201 potential for subaqueous melting; a process that can also be induced by wind-driven currents  
202 (Chikita et al., 1998).

203 Supraglacial ponds surrounded by ice cliffs tend to be larger and deeper than those without  
204 cliffs (Watson et al., 2018), as the ice cliffs facilitate pond growth by subaerial melting and  
205 backwasting, particularly during the monsoon melt season (Röhl, 2008; Steiner et al., 2019). Where  
206 warm surface pond water meets glacier ice, it can undercut the cliff beneath the waterline;  
207 progressive undercutting and thermo-erosional notch development may then lead to calving of  
208 the ice cliff and pond expansion (Figure 5C and D) (Chikita et al., 1998; Kirkbride and Warren, 1997;  
209 Mihalcea et al., 2006; Miles et al., 2016; Röhl, 2008, 2006; Sakai et al., 2009). Conversely, where  
210 the subaqueous and ice cliff melt rates are similar, the ice cliff will persist and backwaste stably  
211 (Brun et al., 2016; Buri et al., 2016a; Miles et al., 2016). Calving is most effective at larger ponds  
212 (Röhl, 2008), in particular where the fetch is greater than 20 m and the water temperature is 2–  
213  $4^{\circ}\text{C}$  (Sakai et al., 2009). Calving events cause further mixing of pond layers, driving warmer surface  
214 water towards the base and again enhancing basal melting; the greatest supraglacial pond  
215 deepening rates have been shown to occur adjacent to the tallest calving ice cliffs (Thompson et  
216 al., 2012). Although sedimentation from ice cliffs and inflowing water can reduce pond depth, this  
217 effect is commonly outstripped by ablation (Thompson et al., 2012), resulting in general long-term  
218 pond growth.

219 A pattern of supraglacial pond evolution into ice-marginal moraine-dammed lakes has been  
220 observed for some ponds on debris-covered glaciers in High Mountain Asia. Supraglacial ponds  
221 form initially as 'perched ponds', isolated above the englacial drainage network (Benn et al., 2012).  
222 As these ponds increase in area and depth, they evolve from perched to base-level features, where  
223 the base-level is determined by the height at which water leaves the glacial system (usually the  
224 elevation of a spillway through the terminal moraine or the glacier bed, if water is transported  
225 there) (Mertes et al., 2016; Thompson et al., 2012; Watanabe et al., 2009). However, differing sub-  
226 catchments may have differing base-levels defined by other hydrological features such as moulins,

227 resulting in a stepped hydrological cascade based on several local base-levels. Alternatively, the  
228 presence of a groundwater system can result in a regional base-level. Over an extended period of  
229 glacier recession, an increasing number of supraglacial ponds form and grow over time, creating a  
230 chain of terminus-base-level ponds that eventually coalesce (Figure 5E and F) (Sakai, 2012; Salerno  
231 et al., 2012). The growth of base-level ponds is not limited by periodic drainage, potentially  
232 allowing dramatic increases in area, particularly through calving (Benn et al., 2001; Sakai, 2012;  
233 Thompson et al., 2012). If meltwater cannot escape from the system, pond expansion and  
234 coalescence may eventually lead to the formation of a single base-level moraine-dammed  
235 proglacial lake at the glacier terminus (Section 5.1.1) (Mertes et al., 2016; Watanabe et al., 2009)  
236 that will continue to expand both upglacier and downwards by ice melt.

237 Various stages of this supraglacial pond evolution are simultaneously present on many High  
238 Mountain Asian debris-covered glaciers. An increase in supraglacial pond area and proglacial lake  
239 formation, assumed to be in response to a warmer climate and glacier surface lowering, has been  
240 observed in recent decades in, for example, the Tien Shan (Wang et al., 2013), Bhutan Himalaya  
241 (Ageta et al., 2000; Komori, 2008), and Nepal Himalaya (Benn et al., 2000; Watson et al., 2016).  
242 Within the Hindu-Kush Himalaya, a clear divide has appeared between the East, where there are  
243 a greater number of larger ponds that have grown between 1990–2009 and become increasingly  
244 proglacial, and the West, where already generally smaller supraglacial ponds have been decreasing  
245 further in area (Gardelle et al., 2011). However, local variations do occur and the pattern is not  
246 universal (e.g. Steiner et al., 2019).

247 As isolated perched ponds grow, they can deepen such that they become connected to the  
248 englacial system by intersecting englacial flow pathways, and drain (Benn et al., 2001; Liu et al.,  
249 2015; Röhl, 2008; Watson et al., 2018, 2016; Wessels et al., 2002), temporarily halting further pond  
250 expansion (Mertes et al., 2016). Pond drainage is promoted in zones of higher local surface velocity  
251 and strain rates, connecting the supraglacial and englacial drainage networks and resulting in  
252 smaller-sized ponds (Miles et al., 2017b). However, as noted above, ponds are generally more  
253 likely to form in areas with lower surface velocities. Ponds may also drain by preferentially  
254 exploiting inherited structural weaknesses such as (sediment-filled) crevasse traces, open  
255 crevasses, and englacial conduits that have been forced closed by longitudinal compression,  
256 allowing drainage by hydrofracture (the penetration of a water-filled crevasse through an ice mass  
257 assisted by the additional pressure of the water at the crevasse tip) (Benn et al., 2017, 2012, 2009;  
258 Gulley and Benn, 2007; Miles et al., 2017b). Alternatively, perched ponds may drain by overspilling,  
259 when a channel is melted into the downstream end of a pond. If, during drainage, such a channel  
260 incises faster than the pond lowers then unstable and potentially catastrophic drainage can result  
261 (Liu et al., 2015; Raymond and Nolan, 2000). However, analyses on Lirung Glacier, Nepal Himalaya,  
262 provided strong evidence for continuous inefficient drainage of supraglacial ponds, likely into  
263 debris-choked englacial conduits (Miles et al., 2017a).

264 A periodic cycle of supraglacial pond expansion and drainage may occur until the pond  
265 becomes large enough to become permanently connected to the englacial system, and thus  
266 stabilise due to inputs of meltwater from streams and other ponds located farther upglacier (Benn  
267 et al., 2001; Miles et al., 2017a; Wessels et al., 2002). An abundant supply of meltwater from the

268 ice surface or the wider drainage system is evidenced by ponds with a high suspended-sediment  
269 concentration (Takeuchi et al., 2012). A seasonal pattern of supraglacial pond filling and drainage  
270 has been observed at seven glaciers in the Tien Shan, with 94% of observed ponds draining during  
271 the monsoon every year between 2013–2015 (Narama et al., 2017). Similar cycles were reported  
272 for five glaciers in Langtang Valley, Nepal Himalaya, where the maximum ponded area between  
273 1999–2013 occurred early in the melt season, subsequently decreasing as ponds drained or froze  
274 (Miles et al., 2017b). Conversely, larger ponds have been observed to drain incompletely and  
275 separate into multiple smaller ponds, subsequently refilling to re-form one large pond (Benn et al.,  
276 2001; Miles et al., 2017b; Wessels et al., 2002). Warmer spring temperatures have been noted to  
277 correlate with a greater number of drainage events later the same year, likely due to larger  
278 meltwater inputs earlier in the year triggering redevelopment of the subsurface drainage system  
279 (Liu et al., 2015).


280           Supraglacial ponds are responsible for a large proportion of debris-covered glacier ablation,  
281 absorbing heat up to 14 times more quickly than even the debris-covered area. In the Langtang  
282 Valley, Nepal, this accounted for 12.5% of catchment ice loss (E. S. Miles et al., 2018b). However,  
283 linked supraglacial pond chains have been suggested to provide only a small proportion of total  
284 glacier proglacial discharge (Irvine-Fynn et al., 2017; Miles et al., 2019), primarily storing meltwater  
285 and thus increasing the potential for enhanced ablation. Ponds release  $\geq 50\%$  of their absorbed  
286 energy with the melt output from the pond, contributing to internal melting along supraglacial and  
287 englacial conduits (Miles et al., 2016; Sakai et al., 2000). This in turn may lead to englacial roof  
288 collapse and the formation of new ponds (Benn et al., 2012; Miles et al., 2017a; Sakai et al., 2000),  
289 resulting in a net glacier-wide increase in ablation. The growing presence of ponds has been  
290 described as the clearest indicator of the influence of climate change on debris-covered glaciers  
291 (Salerno et al., 2012).

### 292 2.1.3 Meltwater transport

293 Supraglacial streams (Figure 6) on High Mountain Asian debris-covered glaciers vary widely in  
294 prevalence, size, and length. To exist and persist, a supply catchment is required (Benn et al., 2017;  
295 Gulley et al., 2009a) and the rate of stream incision, driven by thermal erosion, must outpace the  
296 rate of surface lowering (Marston, 1983). Such conditions may be promoted beneath thicker debris  
297 that suppresses surface ablation in the lower ablation area (Benn et al., 2017), yet observations of  
298 streams in this region are rare, likely due to the hummocky topography both limiting the size of  
299 supraglacial catchments (Fyffe et al., 2019b) and preventing any streams that do form from  
300 persisting for long distances (Benn et al., 2017). Farther upglacier, often under conditions of strong  
301 longitudinal extension associated with ice falls, open crevasses are common and also suppress  
302 supraglacial stream development (Benn et al., 2017). Most supraglacial streams have therefore  
303 been observed in the upper to mid-ablation area (Figure 6A-D) (Gulley et al., 2009a; Miles et al.,  
304 2019), downglacier of crevasse fields but still upglacier of the pronounced hummocky topography  
305 and thick debris layer (Section 2.1.1).

306           A perennial supraglacial stream (Figure 6A-D) has been present in the upper ablation area  
307 of Khumbu Glacier, Nepal Himalaya, for over 14 years (Gulley et al., 2009a; Miles et al., 2019). This  
308 stream and its smaller tributaries originate just downglacier of the Khumbu icefall, where the mean

309 longitudinal surface gradient decreases dramatically (Figure 1), from  $\sim 23^\circ$  down the icefall to  $\sim 3^\circ$   
 310 just below the clean-ice region (estimated from Shean (2017) over one km segments of the  
 311 glacier's central flowline). The low surface gradient of the ablation area results in this channel  
 312 having a high sinuosity (Miles et al., 2019). As streams transfer meltwater downglacier, they can  
 313 incise effectively into the glacier surface (Figure 6B and C); one channel had melted 5–10 m deep  
 314 by the time it reached the lower ablation area (Gulley et al., 2009a; Iwata et al., 1980). Such incision  
 315 is evident where the channel sides and surrounding glacier surface have ablated more slowly than  
 316 the channel itself, leaving walls of horizontally notched ice showing former, less incised channel  
 317 elevations (Figure 6C). Supraglacial streams may drain into debris-covered glaciers through  
 318 crevasses or moulins (Gulley et al., 2009a; Iwata et al., 1980), or through channel 'cut-and-closure'  
 319 (see Section 3.1) (Gulley et al., 2009a; Jarosch and Gudmundsson, 2012). Relict channels  
 320 abandoned by continued incision can often be exposed on the surface as a result of spatially  
 321 variable surface lowering (Figure 6D).



322

*Figure 6 – Examples of supraglacial streams on Khumbu Glacier, Nepal Himalaya, in: **A-C**) the upper ablation area, incised into the ice beneath the debris layer. Blue arrows indicate water flow direction; yellow arrows indicate abandoned/relict channels. The supraglacial stream in A) is extensive, sinuous, and very well developed, transporting large volumes of meltwater efficiently. B) and C) are upstream of A) (white and black star, respectively): B) shows a relict, debris-filled meander bend which has been superseded by a more direct channel; C) shows multiple levels of stream incision (grooves indicated by red dashed line,  $\sim 1$  m high); **D**) the mid-ablation area, where the same incised channel becomes englacial through cut-and-closure after several hundred metres of progressive downcutting, visible from the multiple relict levels (channel drop in the image is  $\sim 10$  m); **E**) and **F**) the lower ablation area. The channel in E) is a short stretch between a supraglacial pond and a shallow moulin, flowing over the debris layer. The stream in F) flows into a breach in the lateral moraine to form the proglacial stream; here, it has eroded into the sand-like sediment across a basin that seasonally floods. Images in A, E, and F were taken during fluorescein dye*

*tracing experiments (Miles et al., 2019). Image credit for A: Duncan Quincey; B, C, and E: Katie Miles; D: Evan Miles; and F: Bryn Hubbard.*

323           Supraglacial streams can undergo rapid pathway changes. Figure 6B shows a debris-filled  
324 section of channel, abandoned as meltwater progressively took a more direct route, leaving a  
325 central island of protruding ice. This process may have been similar to the formation of an ox-bow  
326 lake in an abandoned terrestrial river meander. However, the abandoned channel section may be  
327 reactivated during times of high flow, evidenced by the presence of thick, evenly spread debris  
328 deposits in Figure 6B. Farther downglacier, where supraglacial stream observations are rarer,  
329 pathway changes have also been witnessed on short timescales (Miles et al., 2019). In Figure 6E,  
330 the stream flows into a shallow moulin, yet within 10 days this moulin had collapsed and been  
331 abandoned, with the stream routing into a new moulin just upstream. Moulin collapse has been  
332 attributed to the highly spatially variable surface lowering and ablation rates on debris-covered  
333 glaciers (Miles et al., 2019), while the short timescale indicates that the new moulin exploited an  
334 existing weakness in the ice.

## 335 2.2 Supraglacial knowledge gaps

336 Predictions of future mass balance regimes on High Mountain Asian debris-covered glaciers are  
337 uncertain. Surface lowering is leading to both an overall increase in debris thickness (Gibson et al.,  
338 2017a) and an upglacier emergence of a thin supraglacial debris layer. These processes will likely  
339 further decrease albedo and increase surface meltwater production (thereby increasing surface  
340 lowering, potentially leading to a positive cycle until debris thickens sufficiently to insulate the  
341 surface) (Kirkbride and Warren, 1999; Stokes et al., 2007). Measuring meltwater production is  
342 crucial, but difficult beneath (thin) debris layers, and often impossible where access to the ice-  
343 debris interface is not feasible. More broadly, the future evolution of debris-covered glacier  
344 surface geometry remains inadequately addressed, for example, whether meltwater will primarily  
345 be transported rapidly off the glacier in channels or stored within large systems of linked  
346 supraglacial ponds, thereby buffering diurnal proglacial discharge.

347           On a finer scale, a detailed process understanding of meltwater storage and transport  
348 through supraglacial ponds and pond systems is lacking, particularly of water circulation within,  
349 between, and out of ponds. While often just one discrete, channelised output is visible, water has  
350 also been observed to seep beneath the debris layer and emerge in unexpected locations (Miles  
351 et al., 2019). There has been little focus on how these links between ponds will change as ponds  
352 expand and eventually coalesce. Volumetric measurements of supraglacial ponds are scarce,  
353 rendering it difficult to accurately calculate how much meltwater is being stored on the glacier  
354 surface. Additionally, little attention has been paid to the effect of debris (heated by solar  
355 radiation) falling into a pond on the pond temperature and thus its subaqueous melt rate.

356           Understanding of the various pathways and rates of meltwater transport across a debris-  
357 covered glacier surface would benefit from additional focused research. For example, supraglacial  
358 streams are commonly difficult to discern in debris-covered regions of the glacier surface; this is  
359 particularly true for smaller surface streams and diffuse flows, which are less easily located and  
360 consequently remain largely unreported. On a smaller scale, the occurrence of some ice ablation



361 beneath even a thick debris layer implies that during much of the ablation season, water must  
362 exist between the ice surface and the debris layer (McCarthy et al., 2017), likely as a thin but  
363 variable film. However, the planform structure and meltwater transport of any such drainage  
364 network remain unknown, although transport must subsequently occur as a saturated surface  
365 layer or - initially at least - as small, inefficient rivulets.

366 Water storage within and below the supraglacial debris layer is likely but unexplored. Such  
367 storage would introduce temporary delays in the transport of meltwater through the system, thus  
368 affecting meltwater hydrochemistry (Tranter et al., 2002, 1993), the development of other parts  
369 of the drainage network, and proglacial discharge. However, despite its importance in contrasting  
370 with standard models of supraglacial hydrology based on research at clean-ice glaciers, small-scale  
371 meltwater storage delays remain unknown, which at least partly results from the difficulty  
372 involved in gaining access to the ice-debris interface beneath thick surface debris. Similar issues  
373 are present for the hydrology of snowpacks overlying thick debris; the extent that the snowpack  
374 delays runoff and how much snowmelt enters the hydrological system are similarly unaddressed.

### 375 3. Englacial hydrology

#### 376 3.1 Englacial zone

377 Exceptionally, englacial conduits at High Mountain Asian debris-covered glaciers have been at least  
378 as well explored by glaciospeleologists as at clean-ice glaciers. Such exploration has been carried  
379 out primarily in the Nepal Himalaya, including at Khumbu Glacier (Gulley et al., 2009a), Ngozumpa  
380 Glacier (Benn et al., 2017, 2009; Gulley and Benn, 2007), Ama Dablam and Lhotse Glaciers (Gulley  
381 and Benn, 2007), as well as several debris-covered glaciers in the Tien Shan (Narama et al., 2017).  
382 Largely on the basis of such studies, Gulley et al. (2009) proposed three formation mechanisms for  
383 englacial conduits within debris-covered glaciers:

- 384 I. Cut-and-closure type conduits appear to be particularly prevalent within High  
385 Mountain Asian debris-covered glaciers, relative to clean-ice counterparts. Since the  
386 process requires more rapid supraglacial channel incision than surface ablation, this  
387 prevalence could result from the presence of cold surface ice and/or surface debris,  
388 both impeding general surface lowering. Under such conditions, incision will continue  
389 to the hydrologic base-level of the glacier (Section 2.1.2), with englacial conduits  
390 forming by supraglacial channel closure from snow infill and, in some cases, by ice creep  
391 (Gulley et al., 2009b, 2009a). These conduits may be repeatedly abandoned and  
392 reactivated as water supply varies through the year. However, such conduits rarely  
393 close completely due to their shallow depth below the surface, and may contain  
394 sediment that provides lines of secondary permeability by which the conduit may  
395 subsequently be reactivated (Benn et al., 2009; Gulley et al., 2009a; Gulley and Benn,  
396 2007). Cut-and-closure conduits have been reported on Khumbu (Gulley et al., 2009a)  
397 and Ngozumpa Glaciers (Thompson et al., 2012).
- 398 II. Meltwater may aggregate to form englacial conduits by exploiting lines/planes of  
399 secondary permeability; for example, those left by relict cut-and-closure conduits or  
400 debris-filled and/or compressed former surface crevasses (Benn et al., 2012; Gulley et

401 al., 2009b; Gulley and Benn, 2007; E. S. Miles et al., 2018a). Along these low-  
402 permeability zones, discharge through the icy matrix leads to the development of  
403 enlarging lines of preferential flow due to viscous heat dissipation, eventually forming  
404 an englacial conduit (Benn et al., 2012).

405 **III.** Englacial conduits may also form by hydrofracturing (Benn et al., 2012, 2009; Gulley et  
406 al., 2009b), though this process is generally restricted to upper, debris-free areas where  
407 surface runoff can enter open crevasses (Benn et al., 2012). In the lower ablation area,  
408 low surface gradients, low strain, and compression reduce the capacity for crevassing.  
409 Conduit formation by hydrofracturing has been invoked in association with longitudinal  
410 crevasses on Khumbu Glacier (Benn et al., 2012, 2009), promoted by the combined  
411 effect of transverse stresses and high water pressure at the base of supraglacial lakes.  
412 Multiple stages of hydrofracture, followed by conduit closure through freeze-on, were  
413 interpreted from a series of successively lower niches eroded into pond walls (Benn et  
414 al., 2009).

415 If a stream exploits a crevasse for sufficient time, it forms a moulin, as on clean-ice glaciers.  
416 Although such instances are rare, steep-gradient moulins have been observed in the upper  
417 ablation area of some High Mountain Asian debris-covered glaciers (e.g. Southern Inylchek Glacier,  
418 Tien Shan and Baltoro Glacier, Pakistan Karakoram (Narama et al., 2017; Quincey et al., 2009)),  
419 and a shallow-gradient moulin reported in the lower ablation area of Khumbu Glacier (Figure 6E)  
420 (Miles et al., 2019). Indeed, explored englacial conduits, such as on Khumbu and Ngozumpa  
421 Glaciers, also had shallow gradients (Benn et al., 2017; Gulley et al., 2009a; Gulley and Benn, 2007),  
422 suggesting predominant formation in these instances by cut-and-closure rather than crevasse  
423 exploitation.


424 Englacial conduits have been observed at multiple elevations within High Mountain Asian  
425 debris-covered glaciers, often showing numerous levels of incision hypothesised to result from  
426 sequential supraglacial pond drainage events as the base-level has moved (Gulley et al., 2009a;  
427 Gulley and Benn, 2007). According to this model, each conduit can have varying local base-levels  
428 through time (Section 2.1.2), with elevations ultimately limited by the glacier's contemporary  
429 base-level, determined by the height at which water leaves the glacier (Gulley et al., 2009a; Gulley  
430 and Benn, 2007). Furthermore, as the surface gradient of the ablation area of debris-covered  
431 glaciers is typically very low, the hydraulic gradient (Shreve, 1972) is correspondingly low,  
432 encouraging meandering and the formation of sinuous englacial conduits (Miles et al., 2019), as  
433 observed on Khumbu and Ngozumpa Glaciers (Benn et al., 2017; Gulley and Benn, 2007).

434 Longer-distance water transport has been inferred to occur through perennial sub-  
435 marginal conduits, likely formed by cut-and-closure, located along the edge of debris-covered  
436 glaciers (Benn et al., 2017; Thompson et al., 2016). Such marginal features provide longer-distance  
437 and more hydraulically efficient pathways than conduits within the central glacier, due to the  
438 frequent presence of infilled crevasse traces that can be exploited by water flowing at the margins  
439 (Gulley and Benn, 2007). Centrally located englacial conduits may become re-exposed due to  
440 lowering of the surrounding surface, routing water back to the surface (Figure 7) (Miles et al.,

441 2019). This process may make these conduits more discontinuous, particularly when combined  
442 with the commonly hummocky topography (Miles et al., 2017a).

443 Englacial systems have been observed at shallow depths below the surfaces of High  
444 Mountain Asian debris-covered glaciers. These typically consist of short conduits (channelised,  
445 distributed or a combination), englacial reservoirs, and/or shallow moulines, primarily linking  
446 supraglacial ponds (Miles et al., 2017a, 2019; Narama et al., 2017). Such linked supraglacial-  
447 englacial systems may be created and/or maintained by supraglacial pond drainage into englacial  
448 conduits (Gulley and Benn, 2007; Narama et al., 2017). Narama et al. (2017) found that the  
449 seasonal drainage cycle of supraglacial ponds on seven Tien Shan glaciers was characterised by a  
450 connection to an established englacial drainage system later in the summer; 94% of ponds drained  
451 and connected during all three years studied. Englacial conduits may thus play an important role  
452 in the life cycles of perched ponds (Benn et al., 2017; Miles et al., 2017a).

453 Deeper englacial drainage networks can vary in efficiency in response to numerous factors,  
454 including supraglacial pond drainage events. On Dokriani Glacier, Garhwal Himalaya, englacial  
455 conduits were inferred to be efficient and active through the entire melt season, with proglacial  
456 discharge proportional to supraglacial water production (Hasnain and Thayyen, 1994). Conversely,  
457 on Khumbu Glacier, a channelised but inefficient englacial system was inferred in the pre-monsoon  
458 season (Miles et al., 2019). This system did not link to the supraglacial pond chain, but was routed  
459 to the surface close to the terminus, suggesting that deep englacial to shallow-englacial-  
460 supraglacial links are also possible. While this inefficient englacial system was characterised by  
461 slow transport velocities, previous observations of faster transit through Khumbu Glacier during  
462 the drainage of a tributary glacier's supraglacial pond implies that the system can adapt rapidly to  
463 greater meltwater inputs (E. S. Miles et al., 2018a; Miles et al., 2019).



464

*Figure 7 – A relict englacial conduit (~10 m in height) in the centre of an ice cliff on Khumbu Glacier, Nepal Himalaya, exposed after a drainage event of the associated supraglacial pond, viewed: **A)** from upglacier, and **B)** from downglacier. On the downglacier side, tens of metres of surface lowering has occurred and the previously englacial conduit is now visible from the surface, meandering and incising for ~200 m farther downglacier before flowing into a pond. Image credit for A: Evan Miles; and B: Katie Miles.*

465 The efficiency of englacial meltwater transport has also been noted to change through the  
466 melt season at High Mountain Asian debris-covered glaciers. The influx of large volumes of  
467 monsoon precipitation during the summer months may result in the reopening of englacial (and  
468 subglacial) conduits, giving potential for considerable englacial ablation (Benn et al., 2012); for a  
469 surface pond of 500 m<sup>2</sup>, sufficient energy to melt ~2,600 m<sup>3</sup> of temperate ice is released over a  
470 single monsoon season (Miles et al., 2016). This additional meltwater ultimately leads to conduit  
471 erosion (Miles et al., 2017b; Sakai et al., 2000), which may be further enhanced by pond drainage  
472 events, as the warmer drained water (Section 2.1.2) conveys large amounts of energy, adding  
473 further to total glacier mass loss (Benn et al., 2012; Miles et al., 2016; Sakai et al., 2000; Thompson  
474 et al., 2016).

475 For englacial conduits located near the surface, rapid expansion can result in conduit  
476 collapse if the ceiling is not sufficiently supported. A supraglacial, possibly relict, channel formed  
477 from englacial conduit collapse exposes new bare ice faces, including ice cliffs, which may then  
478 contribute to more rapid lowering of the glacier surface (Section 2.1.1) (Benn et al., 2017;  
479 Kraaijenbrink et al., 2016b; Miles et al., 2016; Sakai et al., 2000; Thompson et al., 2016, 2012).  
480 Ablation rates and surface subsidence can be further enhanced if the new depression becomes  
481 flooded by that increased meltwater production, supplemented by upglacier inputs, providing new  
482 depressions for supraglacial ponds to form or expand and coalesce (Section 2.1.2) (Benn et al.,  
483 2012, 2001; Kirkbride, 1993; Kraaijenbrink et al., 2016b; Miles et al., 2017a; Sakai et al., 2000;  
484 Thompson et al., 2012).

485 Meltwater may be stored englacially within debris-covered glaciers, ranging from small,  
486 shallow englacial reservoirs (Miles et al., 2019) to deeper and potentially larger reservoirs. The  
487 latter type has been inferred, for example, for glaciers feeding the Hunza river system, Central  
488 Karakoram, at the start of the melt season due to a notable lag between the initiation of glacier  
489 ablation and higher discharges observed downstream (Hewitt et al., 1989). Similarly, the initiation  
490 of an outburst flood at Lhotse Glacier was partly attributed to the release of meltwater stored  
491 within englacial conduits that became over-pressurised from greater meltwater production and  
492 transit during the transitional pre-monsoon season (Rounce et al., 2017). Other inferences have  
493 been made from supraglacial pond water-level measurements, such as at Imja Tsho, Nepal  
494 Himalaya, where the post-melt season lake level was constant despite lower air temperatures and  
495 lower precipitation, which would both serve to reduce meltwater production. This situation was  
496 explained by recharge from englacially and subglacially stored water being progressively released  
497 over time (Thakuri et al., 2016).

### 498 3.2 Englacial knowledge gaps

499 Despite relatively extensive englacial glacioclimatological exploration, numerous gaps remain in our  
500 knowledge of the englacial drainage of High Mountain Asian debris-covered glaciers. For example,  
501 as at clean-ice glaciers, the thermal regime of the glacier exerts a significant control on the location  
502 and formation of an englacial drainage system, yet the thermal regime is unknown for almost all  
503 High Mountain Asian debris-covered glaciers. A recent study suggested that the lower area of  
504 Khumbu Glacier may primarily comprise temperate ice (K. E. Miles et al., 2018) allowing the  
505 existence of a deep englacial drainage system (Miles et al., 2019). However, this research was

506 confined to a single glacier and its representativeness for other debris-covered glaciers in High  
507 Mountain Asia remains unknown.

508 Knowledge of the influence of supraglacial debris on englacial (and subglacial) drainage  
509 systems is incomplete. On Miage Glacier, the upglacier ice, which is cleaner and covered with a  
510 thin supraglacial debris layer, was shown to produce an efficient subsurface drainage system to  
511 the terminus from early in the melt season. In contrast, the heavily debris-covered lower ablation  
512 area restricted the development of supraglacial drainage, leading to an inefficient subsurface  
513 system that ultimately flowed into the efficient system driven from upglacier (Fyffe et al., 2019b).  
514 While there are similarities between the drainage system of Miage and the few High Mountain  
515 Asian debris-covered glaciers studied, the generally thicker debris layer and much greater  
516 prevalence of supraglacial ponds towards the terminus of the latter will additionally influence the  
517 hydrological system of such glaciers – an influence that remains unexplored.

518 Improved understanding is required of the links between the englacial system and other  
519 hydrological domains, such as supraglacial-to-englacial transitions (through cut-and-closure  
520 conduits, weaknesses in the ice, and supraglacial pond drainages). Research into the shallow  
521 englacial system is needed, including determining how much of a distinction there is between  
522 shallow englacial and supraglacial systems, considering the rapidly changing surface topography  
523 that is typical of High Mountain Asian debris-covered glaciers. Finally, the potential for englacial  
524 meltwater storage has received very little attention.

## 525 4. Subglacial hydrology

### 526 4.1 Subglacial zone


527 Knowledge of subglacial drainage at High Mountain Asian debris-covered glaciers is limited,  
528 although some evidence at least points to the existence of such systems. For example,  
529 glaciological investigations indicated that the proglacial stream of a retreating tributary of  
530 Khumbu Glacier reached Khumbu's bed (Benn, pers. comm., 2018). This conduit was assumed to  
531 follow the bed for some distance downglacier, similar to the perennial sub-marginal conduits  
532 present at the edge of the neighbouring Ngozumpa Glacier (Benn et al., 2017; Miles et al., 2019;  
533 Thompson et al., 2016). However, this water did not persist subglacially, and instead was  
534 documented to exit the glacier supraglacially. This upward routing of water likely occurs due to  
535 the commonly high hydrological base-level of such glaciers, possibly following the glacier's cold-  
536 temperate transition surface (K. E. Miles et al., 2018; Miles et al., 2019).

537 Beyond the studies outlined above, all other information relating to the subglacial drainage  
538 of High Mountain Asia debris-covered glaciers is inferred. For example, the presence of meltwater  
539 at the bed has been inferred from surface velocity records from remote sensing (e.g. Quincey et  
540 al., 2009) or field-based GPS (e.g. Tsutaki et al., 2019), using inferences similar to those for clean-  
541 ice glaciers. Relatively rapid surface velocities in the central areas of glaciers have been recorded  
542 during summer months, when melting and rainfall delivery are greatest (Figure 8). Such velocity  
543 increases have been interpreted as indicative of basal motion lubricated by subglacial drainage  
544 (Benn et al., 2017; Copland et al., 2009; Kääh, 2005; Kodama and Mae, 1976; Kraaijenbrink et al.,

545 2016a; Kumar and Dobhal, 1997; Mayer et al., 2006; Quincey et al., 2009). Similar remote sensing  
546 studies of surging debris-covered glaciers, particularly in the Karakoram, have inferred the  
547 presence of subglacial water that enables rapid surface velocities during surge phases (Copland et  
548 al., 2009; Quincey et al., 2011; Steiner et al., 2018a), such as the maximum velocity of  $> 250 \text{ m a}^{-1}$   
549 reported at South Skamri Glacier, Pakistan Karakoram (Copland et al., 2009).

550 The existence of channelised subglacial drainage has been inferred from the presence of  
551 proglacial outlet streams at the terminus of debris-covered glaciers. During the melt season, these  
552 channels discharge large volumes of heavily debris-laden water, implying sediment entrainment  
553 during transport along the bed (Quincey et al., 2009). This transport pathway has also been  
554 inferred from comparisons of supraglacial with proglacial solute concentrations on Lirung Glacier,  
555 where high proglacial  $\text{Ca}^{2+}$  and  $\text{SO}_4^{2-}$  concentrations indicated prolonged contact with reactive  
556 debris, inferred to occur during subglacial drainage (Bhatt et al., 2007). Similarly, a perennially  
557 active subglacial system on Dokriani Glacier was inferred to be connected with the englacial system  
558 on the basis of proglacial electrical conductivity measurements (Hasnain and Thayyen, 1994).

559 Variations in subglacial system efficiency have been inferred from studies focusing on  
560 proglacial streams. For example, the increasing efficiency and interconnection of the subglacial  
561 system of Gangotri Glacier, Garhwal Himalaya, was inferred from an increase in the net flux and  
562 size of subglacially eroded suspended particles through the melt season (Haritashya et al., 2010).  
563 On the same glacier, dye tracing experiments showed that the channelised subglacial drainage  
564 system became progressively more efficient with greater meltwater inputs through the melt  
565 season (Pottakkal et al., 2014). Dye tracing experiments have also demonstrated a transition from  
566 distributed to channelised subglacial drainage through the melt season, for example at both  
567 Dokriani Glacier and Hailuogou Glacier, Mt. Gongga, Tibet (Hasnain et al., 2001; Liu et al., 2018).  
568 On a diurnal scale, Kumar et al. (2009) found that the total ion concentration of proglacial  
569 meltwater at Gangotri Glacier increased from the afternoon onwards, interpreted as an enhanced  
570 subglacial component due to the englacial system developing through the day and transporting a  
571 greater proportion of supraglacial meltwater to the solute-rich glacier bed. Finally, substantial  
572 subglacial meltwater storage at debris-covered Lirung Glacier was inferred from its lower diurnal  
573 discharge variability relative to nearby debris-free Khimsung Glacier, Nepal Himalaya (Wilson et  
574 al., 2016).



575

Figure 8 – Surface velocity maps of Lirung Glacier, Nepal Himalaya, during summer (left) and winter (right), with three transverse velocity profiles (A-C) at the locations marked. From Kraaijenbrink et al. (2016b).

## 576 4.2 Subglacial knowledge gaps

577 Very little is known about the subglacial drainage of High Mountain Asian debris-covered glaciers,  
 578 largely due to the difficulty in accessing these systems. Furthermore, many debris-covered glaciers  
 579 in High Mountain Asia terminate in lakes (Section 5.1.1), which increases the likelihood of some  
 580 form of subglacial drainage system but reduces the likelihood of that system being channelised.  
 581 Such lakes also severely hamper direct access to any outflow streams that might be present.  
 582 Assuming the existence of such conduits, it is entirely unknown whether subglacial networks flow  
 583 directly into proglacial ponds at the bed, are routed to the surface upglacier and flow in  
 584 supraglacially (similar to the pathway of some englacial drainage at Ngozumpa Glacier (Benn et al.,  
 585 2017)), or are partially or wholly lost to groundwater. Additionally, the existence of base-level  
 586 englacial streams and a perched water table are highly likely to complicate the detection of, and  
 587 distinction between, englacial and subglacial systems, at least approaching the terminus. For  
 588 example, towards the terminus of Khumbu Glacier, it has been inferred that the high local base-  
 589 level results in the uprouting of the subglacial/deep englacial drainage system to the surface, yet,  
 590 since the ice here is temperate, some meltwater would nonetheless be expected at the bed (K. E.  
 591 Miles et al., 2018; Miles et al., 2019). However, basal ice temperatures and conditions for almost  
 592 all other High Mountain Asian debris-covered glaciers are entirely unknown.

593 Transitions between the englacial and subglacial system are important to understand, as  
 594 are discovering and tracking lost meltwater components – lost potentially to groundwater, to  
 595 short- or long-term storage within the glacier, or to evaporation from the terminal moraine. If


596 extensive subglacial drainage environments are discovered, the influence of the supraglacial debris  
597 cover on those systems should also be investigated.

## 598 5. Proglacial hydrology

### 599 5.1 Proglacial zone

#### 600 5.1.1 Proglacial lakes

601 One of the most distinctive characteristics of the proglacial zone of High Mountain Asian debris-  
602 covered glaciers is the frequent presence of a proglacial lake (Figure 9), which are far more  
603 common than at equivalent clean-ice glaciers. These lakes form by a continuation of the processes  
604 of glacier thinning and supraglacial pond growth (Section 2.1.2) facilitated by the deposition of  
605 sufficient debris by debris-covered glaciers to create high, arcuate terminal moraines. Here,  
606 perched supraglacial ponds expand both downwards, eventually cutting to base-level, and  
607 laterally, often eventually coalescing to produce one large lake above and over the terminus  
608 (Basnett et al., 2013; Kattelmann, 2003; Mertes et al., 2016; Röhl, 2008; Watanabe et al., 2009).  
609 Although less common, base-level lakes that penetrate the full glacier thickness can form farther  
610 upglacier and expand downglacier through stagnant terminus ice, for example Imja Tsho on Imja-  
611 Lhotse Shar Glacier, Nepal Himalaya (Figure 9) (Watanabe et al., 2009). The exact location of such  
612 a proglacial lake may be determined by the location of shallow englacial conduits that provide pre-  
613 existing lines of weakness as the perched ponds grow (Benn et al., 2017; Thompson et al., 2012).  
614 Proglacial lakes will therefore determine the hydrological base-level of the glacier, and are often  
615 dammed by the terminal moraine (Thompson et al., 2012).



616

*Figure 9 – Proglacial lake (Imja Tsho) with a frozen and snow-covered surface at Imja-Lhotse Shar Glacier, Nepal Himalaya. A) full length of Imja Tsho (~2.7 km in October 2018), looking upstream towards the calving front of Imja-Lhotse Shar Glacier. B) detached (stagnant) glacier ice that dams the lake. The black star and arrow in B) show the location and direction A) was taken in. Image credits: Katie Miles.*

617 The formation of moraine-dammed proglacial lakes represents a final stage in the surface  
618 lowering and overall mass loss of debris-covered glaciers. Benn et al. (2012) defined three stages  
619 in the development of debris-covered glaciers: in regime one, all parts of the glacier are



620 dynamically active; in regime two, surface lowering has begun and ice velocities decrease; in  
621 regime three, glaciers are stagnant and rapid recession may occur. The formation of a base-level  
622 lake indicates that a glacier has entered this third regime, and rapid recession may then occur  
623 through further expansion of that proglacial lake (Benn et al., 2012). A growing number of  
624 proglacial lakes of increasing size have been observed in recent decades across the Hindu Kush  
625 Himalaya (Gardelle et al., 2011; Haritashya et al., 2018b; Nie et al., 2017; Thompson et al., 2012).  
626 The pattern of proglacial lake formation varies across the region, with proglacial lake area in the  
627 western Himalaya decreasing 30–50% from 1990–2009 compared to an increase of 20–65%  
628 towards the east, where proglacial lakes are already more prevalent (Gardelle et al., 2011;  
629 Maharjan et al., 2018). This pattern at least partly results from greater glacier recession in the west  
630 over this period (Gardelle et al., 2011).

631 Proglacial lakes expand through similar mechanisms to supraglacial ponds (i.e. subaqueous  
632 melting and subaerial ice face melting; Section 2.1.2) until they are limited by substrate. Lake  
633 expansion therefore enhances glacial mass loss and meltwater production while the lake is  
634 underlain or dammed by ice (Carrivick and Tweed, 2013; Röhl, 2008). Once calving is triggered, it  
635 becomes the dominant method of subsequent lake growth (Röhl, 2008; Thompson et al., 2012).  
636 Calving into a proglacial lake progresses from notch development and roof collapse to large-scale,  
637 full-height slab calving enabled by the lake deepening to the glacier bed (Kirkbride and Warren,  
638 1997; Thompson et al., 2012). The water depth may then be sufficient to trigger extending flow in  
639 the now-unsupported ice cliff, increasing flow velocities and weakening the ice through crevasse  
640 formation and dynamically induced thinning (King et al., 2019; Kirkbride and Warren, 1999;  
641 Thompson et al., 2012; Tsutaki et al., 2019). This can result in rapid and potentially unstable  
642 calving, substantially increasing glacier mass loss, as has been observed during several kilometres  
643 of such retreat at Tasman Glacier, New Zealand (Kirkbride and Warren, 1999) and modelled for  
644 lake- and land-terminating glaciers in the Bhutan Himalaya (Tsutaki et al., 2019). Upglacier  
645 expansion of the proglacial lake (Watanabe et al., 2009) may have implications for the glacier's  
646 drainage system, such as by earlier interruption of meltwater routing (Carrivick and Tweed, 2013).

647 Very large proglacial lakes can alter a glacier's microclimate due to a lake's lower albedo  
648 and higher thermal heat capacity relative to the surrounding ice and soil, thereby producing locally  
649 cooler summer air temperatures and warmer autumn temperatures (Carrivick and Tweed, 2013).  
650 This can slow local summer ice ablation and consequently reduce the amount of meltwater being  
651 produced and transported through the glacier, with implications for the development of englacial  
652 and subglacial drainage systems. If a moraine-dammed proglacial lake is present then the  
653 overwhelming majority of water transported through a debris-covered glacier is likely to pass  
654 through that lake (Benn et al., 2017). This has implications for drainage through the glacier and for  
655 the potential occurrence of glacial lake outburst floods (GLOFs).


### 656 5.1.2 Proglacial streams

657 Proglacial runoff from debris-covered glaciers can form a significant proportion of the discharge of  
658 large rivers downstream, particularly in High Mountain Asia: the Indus, Dudh Koshi, Ganges, and  
659 Brahmaputra rivers all stem from glacial meltwaters (Pritchard, 2019; Ragettli et al., 2015; Wilson  
660 et al., 2016). In particular, glacial runoff buffers both seasonal (Bolch et al., 2019; Pritchard, 2019)

661 and annual (Pohl et al., 2017) water shortages. Loss of glacier volume due to longer, warmer melt  
662 seasons and decreased snow accumulation could result in periods of much reduced water  
663 availability, greatly influencing downstream communities and ecology (Bolch et al., 2019; Pohl et  
664 al., 2017; Pritchard, 2019).

665 Proglacial discharge measurements, estimates, and models have been run across High  
666 Mountain Asia, such as on individual glaciers in Nepal (Braun et al., 1993; Fujita and Sakai, 2014;  
667 Ragettli et al., 2015; Rana et al., 1997; Savéan et al., 2015; Soncini et al., 2016; Tangborn and Rana,  
668 2000), Tibet (Kehrwald et al., 2008), the Tien Shan (Chen and Ding, 2009; Han et al., 2010; Sorg et  
669 al., 2012), India (Hasnain, 1999, 1996; Khan et al., 2017; Singh et al., 2005, 1995; Singh and  
670 Bengtsson, 2004; Thayyen and Gergan, 2010), and for multiple catchments and entire regions  
671 (Winiger et al., 2005). However, such records are relatively short: of the studies listed above, five  
672 measured discharge for a year or less; three have 2–3 years of measurements; and only one has 6  
673 years of measurements. The others use modelling to obtain estimates of proglacial discharge.

674 The presence of surface debris can have a notable effect on a glacier's proglacial discharge,  
675 resulting in a proglacial hydrograph that is different from that of a clean-ice glacier. While no such  
676 comparison has been made for a High Mountain Asian debris-covered glacier, an example is shown  
677 from the debris-covered Dome Glacier, Canadian Rockies (Figure 10) (Mattson, 2000). Here,  
678 discharge was muted both diurnally and through the ablation season compared to the  
679 neighbouring clean-ice Athabasca Glacier (Figure 10); variation in annual discharge volume from  
680 Dome Glacier between the two years was 1%, compared to 24% from Athabasca Glacier. This is  
681 due partly to the suppression of surface melt by a debris cover (Section 2.1.1), and partly to the  
682 lags that are induced as a result of the debris layer – the additional time to conduct heat through  
683 the debris and the warmer local air temperatures due to the warming debris introduces a delay.  
684 Thus, peak melt can occur up to several hours after the maximum radiation receipt at the debris  
685 surface (Carenzo et al., 2016; Conway and Rasmussen, 2000; Evatt et al., 2015), and an exceptional  
686 case has been recorded as being up to 24 hours later for debris layers > 0.85 m thick (Fyffe et al.,  
687 2014). This lag in diurnal peak melt is thus reflected in the timing of the highest stream flow,  
688 producing a later and less pronounced peak in the diurnal pattern of a debris-covered glacier's  
689 proglacial stream (Fyffe et al., 2019a, 2014).



690

Figure 10 – Hydrographs of proglacial discharge of the clean-ice Athabasca Glacier and the adjacent debris-covered Dome Glacier, Canadian Rockies, over the ablation months of July and August 1994 and 1995. Redrawn from Mattson (2000).

691 Lags in proglacial discharge from debris-covered glaciers may also be caused by the  
 692 temporary storage of water within the surface debris layer, for example, during rainfall events.  
 693 This may influence subglacial and proglacial discharge by delaying and buffering water transfer at  
 694 the surface, potentially affecting basal water pressures and minimising peaks in proglacial  
 695 discharge (Brock et al., 2010). In the Himalaya, monsoon precipitation is thought to exert a  
 696 significant control on proglacial discharge hydrographs at high rainfall intensities. For example,  
 697 Thayyen et al. (2005) suggested such an intensity was  $> \sim 20 \text{ mm d}^{-1}$ , which occurred on 20% of  
 698 rainfall days during four years of monsoon measurements on Dokriani Glacier. Early in the melt  
 699 season, meltwater is also stored within the snowpack of debris-covered glaciers, providing a  
 700 further delay in the transport of meltwater from the surface into the subsurface drainage system  
 701 (Singh et al., 2006b). In the last two decades the amount of snowfall accumulation has decreased  
 702 across the Himalaya, and is projected to decrease a further 20–40% by 2100 (Salerno et al., 2015;  
 703 Smith and Bookhagen, 2018; Viste and Sorteberg, 2015); this buffer will thus be further reduced,  
 704 influencing the future proglacial hydrograph pattern of debris-covered glaciers.

705 Groundwater stored within high-elevation glacial catchments has been inferred to interact  
 706 with proglacial (and subglacial) stream networks, affecting their discharge patterns due to  
 707 additional water storage and subsequent release (Gremaud et al., 2009; Smart, 1996, 1988). For  
 708 example, a  $\sim 45$  day lag between precipitation and discharge was observed for 12 glacierised and  
 709 non-glacierised Himalayan catchments, indicating storage of up to two-thirds of the river discharge  
 710 in a groundwater aquifer system before the monsoon, greatly affecting the annual discharge  
 711 pattern (Andermann et al., 2012c). This has similarly been shown by much reduced river

712 suspended sediment concentrations measured post-monsoon, having been diluted as  
713 groundwater begins to be released (Andermann et al., 2012b, 2012a). Comparable processes may  
714 occur beneath the glaciers themselves, for example, at Khumbu Glacier in the pre-monsoon  
715 season, where more meltwater entered the glacier's subsurface drainage system than exited the  
716 glacier at the terminus (Miles et al., 2019). Indeed, in the Jade Dragon Snow Mountain region of  
717 southwest China, 29% of glacier meltwater was calculated to be stored in a karst aquifer (Zeng et  
718 al., 2015). Groundwater sinks of subglacial meltwater can therefore comprise a significant portion  
719 of the total glacial output, potentially resulting in underestimation of glacial ablation.

720 A range of models has been used to predict future runoff from debris-covered glaciers using  
721 various future climatic scenarios for a single glacier basin (Ragettli et al., 2015; Singh et al., 2008,  
722 2006a; Zhang et al., 2007) and multiple glacier basins (Immerzeel et al., 2012; Lowe and Collins,  
723 2001) up to a regional scale (Rees and Collins, 2006; Shea and Immerzeel, 2016). Currently, a large  
724 proportion of debris-covered glaciers worldwide, particularly in the Himalaya, have negative mass  
725 balances (Bolch et al., 2012, 2011; Käab et al., 2012; Scherler et al., 2011). A recently observed  
726 decline in Himalayan snowfall will contribute further to the decreasing mass of these glaciers by  
727 both reducing accumulation rates and exposing the glacier surface to atmospheric melting earlier  
728 in the melt season (Salerno et al., 2015). Glacier contributions to catchment discharge in many  
729 regions have been predicted to increase over the next few decades, but as the glaciers continue  
730 to shrink, peak water will be surpassed and this proportion will begin to reduce substantially due  
731 to the reduced volume of the remaining glaciers (Barnett et al., 2005; Bolch, 2017; Bolch et al.,  
732 2012; Huss, 2011; Huss and Hock, 2018; Lutz et al., 2014). Shea and Immerzeel (2016) estimated  
733 that most basins will have declining glacier contributions to streamflow by 2100, and water  
734 shortages may then be a concern for many populated areas in the Karakoram, while reduced peak  
735 flows may represent a greater concern in the eastern Himalaya.

## 736 5.2 Proglacial knowledge gaps

737 Few glacial discharge monitoring stations have been in place for longer than a decade in High  
738 Mountain Asia, leaving current and future discharge volumes unknown for most debris-covered  
739 glaciers. The volume and temporal variability of potential glacial meltwater losses to groundwater,  
740 and whether these re-join the glacial system (subglacially, proglacially, or further downstream),  
741 are also poorly understood.

742 Projections of future changes in proglacial hydrology are hampered by the absence of  
743 accurate predictions of the future geometric development of High Mountain Asian debris-covered  
744 glaciers. For example, if surface lowering remains the dominant response to climate warming,  
745 glaciers may melt entirely and/or form large proglacial lakes that then dominate mass loss  
746 processes. Conversely, the inverted mass balance regime could result in a separation of stagnant,  
747 heavily debris-covered lower glacier tongues from the upper, less debris-covered regions,  
748 potentially providing ideal conditions for a base-level lake to form in between, dammed by the  
749 detached debris-covered ice.

## 750 6. Future research themes

751 Based on the review above, we identify six hydrological research themes, including examples of  
752 appropriate techniques, that would contribute substantially to advancing our understanding of the  
753 hydrology of High Mountain Asian debris-covered glaciers.

### 754 I. Elucidating glacier-wide water balance

755 Given the importance of glaciers as a source of water in high mountain regions (Immerzeel et al.,  
756 2020), more robust quantification of water inputs into, and outputs from, the glacier system is  
757 paramount. Detailed and temporally and spatially extensive hydrological field measurements are  
758 required to better constrain numerical model parameterisations. Water inputs should be  
759 simulated and examined independently of glacier-fed river discharge, with attention to process  
760 parameterisation to facilitate improvements in efforts to close the water balance. Water storage  
761 is also an important component of the water balance, discussed further in research theme IV  
762 below.

763 The limited measurement to date of precipitation across High Mountain Asia, particularly  
764 in terms of partitioned snow and rainfall and synoptic and seasonal-to-annual variations in  
765 precipitation gradient and rainfall fraction, should be augmented by establishing a network of  
766 robust automatic weather stations over a range of catchments, surface types, and elevations.  
767 Glacier surface elevation change should be measured simultaneously by remote sensing and  
768 ground-based methods – for example, by ultrasonic rangefinders – to calibrate and validate models of  
769 melt and mass balance. These approaches would also aid in determining the impact of  
770 anthropogenic black carbon aerosols and other light-absorbing impurities on albedo, supported  
771 by remote-sensing studies of surface characteristics. Precipitation gradients should be quantified  
772 further through dedicated accumulation measurements.

773 The retention of meltwater, for example by refreezing of meltwater within supraglacial  
774 debris, firn, crevasses, or the body of the glacier, requires better characterisation. Empirical data  
775 collected from snow pits and shallow ice cores would be sufficient to quantify such mass retention  
776 over short timescales, complemented by longer-term records derived from deeper coring or visual  
777 examination of layering present in borehole walls. In the accumulation area, these methods would  
778 provide the additional bonus of historical records of local accumulation.

779 The amount of water lost through evaporation and sublimation should be assessed through  
780 comprehensive studies of eddy covariance coupled with meteorological measurements. Future  
781 research should examine these processes not only from snow-covered areas, recently shown to  
782 be a key source of water loss (Stigter et al., 2018), but also over the accumulation and debris-  
783 covered ablation areas and the terminal and lower lateral moraines, which may equally contribute  
784 to evaporation and sublimation losses. Quantifying these moisture fluxes may be possible either  
785 by direct field measurement or by remote sensing for longer timescales.

786 Other research needs include quantifying losses to groundwater and better evaluating the  
787 role of debris in driving the observed hysteretic behaviour of downstream annual hydrographs.  
788 Hydrochemical and isotopic analyses may shed light on water sources and variations therein, while  
789 catchment-scale dye or gas tracing studies tied closely to continuous measurements of discharge  
790 at various locations on and beyond the glacier could help to define the volumes of water delivered  
791 to groundwater systems (and if so, the proportion that re-joins the proglacial stream farther  
792 downvalley).

## 793 **II. Understanding hydrological processes influencing glacier mass balance**

794 The efficiency of rainfall and meltwater routing from higher elevation locations should be  
795 evaluated due to its potential effect on glacier accumulation and mass balance by englacial  
796 melting. For example, heat fluxes driven by meltwater conveyance to the englacial and subglacial  
797 environments of debris-covered glaciers (i.e. cryo-hydrologic warming (Gilbert et al., 2020; Phillips  
798 et al., 2010)), could be explored using numerical models guided by field-based measurements of  
799 supraglacial water fluxes and temperatures, along with geophysical and/or borehole-based  
800 investigations of englacial temperature fields. Specific loci and timescales of meltwater routing,  
801 storage, and release should be determined. Englacial drainage pathways are of particular  
802 importance due to their strong association with the formation of supraglacial ice cliffs, which  
803 account for a disproportionate amount of surface melt at heavily debris-covered glaciers.  
804 Investigations should map current streams and monitor changes in surface topography and  
805 hydrology (for example, the collapse and surface exposure of shallow englacial systems) both  
806 remotely, using satellite images where streams are large enough, and in the field. The latter should  
807 be supplemented by dye tracing experiments to characterise the hydraulic properties of englacial  
808 systems.

809 There is a need for accurate knowledge of spatial variations in surface debris characteristics  
810 and thicknesses, and of meltwater located at the ice-debris interface, to improve models of surface  
811 vapour fluxes. Thus, meteorological stations are required to measure water content or relative  
812 humidity. Debris thickness maps and the existence of ponded and surface water could be  
813 constructed by refining algorithms from remotely sensed data (both thermal imagery and surface  
814 lowering) or on the basis of manual field measurements of ponds and high-frequency ground-  
815 penetrating radar to identify water present at the interface between the supraglacial debris layer  
816 and the underlying ice. Future investigations of supraglacial pond expansion rates should focus on  
817 wide-scale systematic field-based bathymetry, pond-sediment stratigraphic assessment, and  
818 measurements of pond water and basal sediment temperatures at multiple depths (particularly to  
819 assess vertical heat transfer from warm supraglacial pond water to the base of the pond),  
820 combined with the development of numerical models of heat transfer by such mechanisms.

## 821 **III. Identifying the influence of drainage and meltwater storage on ice motion**

822 Meltwater present at the bed or the terminus of debris-covered glaciers can affect the velocity of  
823 both land- and lake-terminating glaciers; a better understanding and inclusion of subglacial  
824 hydrological processes into models of glacier dynamics will improve future simulations of ice flow  
825 and glacier evolution. Within subglacial hydrological processes, better quantification is needed of

826 the inputs to the system (i.e. coupling meteorological data with melt modelling), the volume of  
827 water present at the bed (for example by monitoring subglacial water pressure in deep borehole  
828 arrays), and the volumes of water lost from the system (i.e. by calculating the glacier's water  
829 balance).

830 Ice motion should be separated into its constituent components (i.e. ice deformation and  
831 basal motion), with particular focus on measurements acquired during the melt season and on an  
832 individual glacier scale. Basal water pressure, and consequently glacier sliding, should be estimated  
833 through analysis of variations in glacier surface velocity obtained, for example through combining  
834 remote-sensing data with field-based GPS studies. The recently available and constantly growing  
835 archive of rapid-repeat, high-resolution optical and radar remotely sensed imagery will help future  
836 work to improve knowledge of seasonal velocities (e.g. Dehecq et al., 2019). Deeper ice velocities  
837 and strain can be recorded within boreholes, ideally extending to the glacier bed. Such boreholes  
838 can also allow measurements of the glacier thermal regime and bed substrate, while improved  
839 mapping of glacier bed topography across High Mountain Asia is necessary to constrain estimates  
840 of ice thicknesses. Finally, in order to assess the influence of calving from proglacial lakes, the  
841 above measurements should be collected in comparative studies of both lake- and land-  
842 terminating High Mountain Asian glaciers.

#### 843 **IV. Characterising seasonal changes in hydrology**

844 Targeted research is needed to measure seasonal changes in hydrological storage components,  
845 particularly those that are specific to debris-covered glaciers. Improved understanding of storage  
846 components is needed to represent the drainage system of debris-covered glaciers appropriately  
847 in hydrological models. For example, seasonal changes in the area and volume of perched  
848 supraglacial ponds could be achieved at the glacier scale using rapid-repeat optical satellite  
849 imagery to maximise likelihood of observation and/or by using high-resolution synthetic aperture  
850 radar satellite data, which are insensitive to cloud cover. Detailed examination of the water  
851 content of the supraglacial debris layer (including the seasonal thaw dynamics of the debris layer,  
852 influencing its hydraulic transmissivity) can be made using soil moisture sensors installed at  
853 multiple depth intervals, while through-debris transmissivity and snowpack storage/release could  
854 be assessed by dye tracing experiments. These processes will aide better understanding of the role  
855 of debris, snow, and firn in transmitting meltwater to supraglacial streams and the subsurface  
856 drainage system, including seasonal storage and release from subsurface reservoirs.

857 Process-based understanding of seasonal hydrological changes could also be improved by  
858 detailed field-based studies. Glacier drainage systems respond dynamically to the seasonal  
859 production of meltwater; this is clear at clean-ice glaciers where snowline retreat stimulates the  
860 progressive upglacier transition from inefficient to efficient drainage. Research is needed at High  
861 Mountain Asian debris-covered glaciers to evaluate whether distinctive seasonal dynamics can be  
862 explained by additional storage components or specific melt-generation patterns. These  
863 phenomena can be addressed through dye tracing, glaciospeleology or bulk proglacial meltwater  
864 analysis. Such studies would also result in a better general understanding of the nature and form  
865 of englacial and subglacial drainage at High Mountain Asian debris-covered glaciers.

866 Finally, the seasonal structure and dynamics of debris-covered glacier hydrological systems  
867 should be understood in the context of projected future melt and discharge. An integrated effort  
868 to assess seasonal changes in debris-covered glacier hydrology should be coupled with melt season  
869 meteorological and ablation measurements, as well as development of a continuous discharge  
870 record through proglacial discharge monitoring stations.

## 871 **V. Evaluating hydrological hazards**

872 The growth in both number and size of supraglacial ponds is one of the clearest visual signs of  
873 debris-covered glacier decay. Future research should focus on predicting formation and growth of  
874 such ponds by combining glacier melt projections (e.g. Kraaijenbrink et al., 2017; Rounce et al.,  
875 2020) with modelled glacier bed overdeepenings (e.g. Linsbauer et al., 2016). Moraine-impounded  
876 sites (such as where base-level terminal lakes have been observed to develop) are more complex;  
877 investigations of the drainage capacity (evidence of free-drainage as opposed to impoundment),  
878 combined with remotely sensed observations of expanding and coalescing supraglacial pond  
879 chains, may provide a suitable starting point. Improved understanding of supraglacial pond  
880 expansion rates, discussed in research theme II, is also crucial, while accurately modelling the  
881 longevity of ice cliffs could be improved with high-resolution digital elevation models (obtained,  
882 for example, through Structure-from-Motion) coupled with simple numerical melt modelling.

883 Assessments of how 'dangerous' a lake is (potential for a catastrophic GLOF occurring)  
884 often disagree (e.g. Haritashya et al., 2018a; Maharjan et al., 2018; Rounce et al., 2016) and, while  
885 recent events such as the 2015 Gorkha earthquake suggest that the terminal moraines of glacial  
886 lakes may be more stable than hitherto considered, large-scale remote observations cannot assess  
887 internal or small-scale superficial damage caused by such events (Byers et al., 2017; Kargel et al.,  
888 2016). Such studies should be improved in terms of their sophistication, for example addressing a  
889 broader range of factors that contribute to the formation of a hazardous lake (e.g. Rounce et al.,  
890 2016), many of which may be site specific. Traditional magnitude-frequency relationships may no  
891 longer be relevant as the current state of mountain environments is non-stationary and beyond  
892 historic precedence. Therefore, alternative forms of event prediction are needed, such as site-  
893 specific hazard development depending on different event magnitudes.

894 Field-based measurements should be made on, and downstream of, individual proglacial  
895 lakes to determine potential hazards and the GLOF risk. Knowledge of moraine dam composition  
896 (including sediment type and the presence or absence of an ice core) and the existence of seepage  
897 or piping is needed, and could be addressed by radar, seismic studies, or drilling into moraines to  
898 characterise soil strength and composition. Flood hydrographs could be better constrained by  
899 geotechnical modelling to understand dam failure mechanisms. While predicting the timing of an  
900 outburst flood is nearly impossible, particularly those originating from englacial and subglacial  
901 sources, characterising subsurface drainage and routing and seasonal release of stored water may  
902 help to identify likely timing and locations of sudden outbursts (research theme IV). Cascading  
903 hydrological hazards, which may be triggered by very high-elevation and often hanging glaciers  
904 that are seldom studied, should also be considered. The thermal conditions and hydrology of these



905 glaciers should be investigated, for example, by surface ground-penetrating radar guided by  
906 borehole-based sensors, dye tracing and discharge monitoring.

## 907 **VI. Predicting future hydrological changes over short and long timescales**

908 Understanding the timescales over which debris-covered glaciers will lose mass, thereby  
909 influencing the amount of meltwater generated and subsequent hydrological processes, depends  
910 on developing a new generation of detailed glacier models that capture both the complex  
911 feedbacks between debris transport by ice and the processes affecting sub-debris ablation over  
912 timescales longer than a few decades (Rowan et al., 2015). Numerical model predictions need to  
913 integrate opposing processes on different scales, for example, encompassing both the glacier-scale  
914 'debris-cover anomaly' (the observed, but still unexplained, debris-covered glacier mass loss rates  
915 that are similar to those of clean-ice glaciers (Brun et al., 2019; Gardelle et al., 2012; Pellicciotti et  
916 al., 2015)) and the smaller-scale insulating effect of the debris. Field and remote-sensing data  
917 relating to mass balance and ice flow processes are required at the correct scale and resolution for  
918 use in numerical models of glacier mass change, parameterised with sufficient process-based  
919 understanding to predict how these controls will evolve over time. The inclusion of these small-  
920 scale and complex processes within regional models (e.g. Kraaijenbrink et al., 2017) to improve  
921 the accuracy of large-scale mass-loss predictions should also be explored.

922 As debris-covered glaciers get smaller, primarily by surface lowering, the debris cover will  
923 thicken and increase insulation, reducing ablation over a potentially greater area of the terminus.  
924 Debris-covered glaciers are therefore already larger and likely to decline slower than equivalent  
925 clean-ice glaciers; as a result, the meltwater of clean-ice glaciers will temporarily provide a  
926 relatively larger component of the annual hydrological budget as they lose mass preferentially.  
927 Robust dynamic glacier models are therefore needed to predict changing hydrographs and  
928 contributions to downstream water supplies, particularly as peak water passes. Supraglacial ponds  
929 play an important role in modulating the proglacial hydrograph and, in the long-term, may provide  
930 a natural water supply reservoir during periods of drought. However, sedimentation rates within  
931 ponds, and therefore their likely longevity, should be quantified by *in situ* hydrological stations.

932 The acceleration of debris-covered glacier mass loss and decrease in glacial runoff as peak  
933 water passes are likely to lead to proglacial streams becoming proportionately more sediment-  
934 laden. This may be enhanced during the melt season, particularly in regions of High Mountain Asia  
935 affected by heavy monsoon rains, which can enhance supraglacial debris erosion. Furthermore,  
936 ice within larger debris-covered glaciers is older than in smaller glaciers and will thus contain a  
937 longer legacy of environmental contaminants (e.g. Hodson, 2014; Li et al., 2017). Ultimately, this  
938 may result in more pollutants being delivered via proglacial streams to water supplies, particularly  
939 during the melt season. Discharge and water quality should therefore be monitored with  
940 hydrological monitoring stations on proglacial streams across High Mountain Asia. Combined with  
941 modelling efforts and improved hydrological understanding, this will allow mitigation strategies to  
942 be planned for the vast downstream populations that depend on that meltwater.

943

## 944 7. Summary

945 In this review, we have summarised our understanding of the hydrology of High Mountain Asian  
946 debris-covered glaciers, identified numerous knowledge gaps, and suggested six themes for future  
947 research. While research has advanced substantially in recent years, there remain many questions  
948 about how the hydrological systems of debris-covered glaciers behave, and how this varies  
949 through both space and time. This limitation is largely due to the position of debris-covered  
950 glaciers in hard-to-reach areas because of logistical difficulties and/or political instability, an  
951 inability to gather observations beneath the surface due to the reduced performance of  
952 combustion-powered equipment at high elevation, and the persistence of challenging weather  
953 conditions for fieldwork through much of the year. Consequently, large uncertainty accompanies  
954 any projections of future water supply, a concern for tens of millions of people across several  
955 countries. Closing these knowledge gaps should thus prioritise generating information that best  
956 improves robust model-based projections of water supply from High Mountain Asian debris-  
957 covered glaciers. There is an inevitable trade-off between the cost of collecting the necessary  
958 empirical data to close these gaps and the benefits returned in terms of improved model outputs.  
959 In light of these requirements and considerations, we conclude by identifying two principal  
960 priorities for scientists and two principal priorities for policymakers and funders.

961 Our first priority for scientists is to improve understanding of patterns and rates of surface  
962 melting, particularly beneath debris layers of different properties and thicknesses on High  
963 Mountain Asian debris-covered glaciers. To this end, multi-variable analytical models should be  
964 developed to generate Østrem-type relationships applicable to a variety of debris properties (such  
965 as lithology, shape, grain-size texture, and variability therein) and energy-balance regimes (thereby  
966 factoring in influences such as elevation), extending the work of, for example Evatt et al. (2015).  
967 Our second priority for scientists is to improve understanding of the basal hydrology of debris-  
968 covered glaciers across all of High Mountain Asia. Currently, little is known about the subglacial  
969 environment, including in many instances where the glacier base is, what the basal temperature  
970 field is, and whether subglacial drainage occurs at all. Yet, these properties are central to the  
971 quality of water supplied by such glaciers, as well as to their actual and modelled deformation  
972 rates and motion fields, which govern their modelled response to anticipated climate change. In  
973 order to maximise benefits relative to cost, field investigations of the subglacial environment could  
974 be undertaken at a limited number of sites to evaluate and guide larger-scale remote sensing and  
975 modelling studies.

976 Our first priority for policymakers and funders is to improve access for scientists to glaciers  
977 across High Mountain Asia. In this regard, we believe the provision of a small number of bases with  
978 effective transport infrastructure, open to international scientific teams, would facilitate a step  
979 change in research activity and output. Our second priority for policymakers and funders is to  
980 produce a better administrative environment for effective scientific collaboration. This should  
981 include, for example, the development of memorandums of understanding between countries to  
982 simplify regulations for research permitting and border crossing as part of a scientific research  
983 project. It should also involve adopting best practice in terms of ensuring a uniform approach to  
984 the quality control and homogeneity of data series, and archiving and sharing freely accessible

985 data. This would be in the interest of all involved parties, since maintaining a clean and reliable  
986 water supply is a fundamental part of building sustainable development (United Nations, 2015),  
987 which in High Mountain Asia can only be realised by improving understanding of future changes in  
988 the timing and magnitude of meltwater production from hitherto poorly studied debris-covered  
989 glaciers.

990

## 991 8. Author contributions

992 KM and BH planned the manuscript. KM led the manuscript writing and illustration. All authors  
993 contributed to the writing and editing of the manuscript.

994

## 995 9. Competing interests

996 The authors declare that they have no conflict of interest.

997

## 998 10. Acknowledgements

999 This research was supported by the ‘EverDrill’ Natural Environment Research Council Grant  
1000 awarded to Aberystwyth University (NE/P002021) and the Universities of Leeds and Sheffield  
1001 (NE/P00265X). KM is funded by an AberDoc PhD Studentship, with fieldwork costs supported by  
1002 the Mount Everest Foundation and Postgraduate Research Awards from the British Society for  
1003 Geomorphology, the Royal Geographical Society (with IBG) and Aberystwyth University  
1004 Department of Geography and Earth Sciences. TIF acknowledges the Leverhulme Trust (RF-2018-  
1005 584/4). The authors would like to thank Antony Smith (Aberystwyth University) for the initial  
1006 illustration of Figure 2 and redrawing Figure 10. Figure 4 is reproduced from the Journal of  
1007 Glaciology with permission of the International Glaciological Society. Figure 8 is reproduced  
1008 according to a Creative Commons Attribution 4.0 License and Figure 10 is reproduced with  
1009 permission of the publisher. They are also grateful to Himalayan Research Expeditions for  
1010 organising the logistics that supported fieldwork in Nepal, and in particular Mahesh Magar for  
1011 guiding and navigation. The authors thank two anonymous reviewers, and two previous reviewers  
1012 in *The Cryosphere Discussions*, all the comments of which led to a much-improved manuscript.

1013

## 1014 11. References

1015 Adhikary, S., Nakawo, M., Seko, K., Shakya, B., 2000. Dust influence on the melting process of  
1016 glacier ice: experimental results from Lirung Glacier, Nepal Himalayas, in: Nakawo, M.,  
1017 Raymond, C.F., Fountain, A. (Eds.), Debris-Covered Glaciers. International Association of  
1018 Hydrological Sciences, Oxford, pp. 43–52.

- 1019 Ageta, Y., Iwata, S., Yabuki, H., Naito, N., Sakai, A., Narama, C., Karma, 2000. Expansion of glacier  
1020 lakes in recent decades in the Bhutan Himalayas, in: Nakawo, M., Raymond, C.F., Fountain, A.  
1021 (Eds.), Debris-Covered Glaciers. International Association of Hydrological Sciences, Oxford,  
1022 pp. 165–175.
- 1023 Andermann, C., Bonnet, S., Crave, A., Davy, P., Longuevergne, L., Gloaguen, R., 2012a. Sediment  
1024 transfer and the hydrological cycle of Himalayan rivers in Nepal. *Comptes Rendus Geosci.* 344,  
1025 627–635. <https://doi.org/10.1016/j.crte.2012.10.009>
- 1026 Andermann, C., Crave, A., Gloaguen, R., Davy, P., Bonnet, S., 2012b. Connecting source and  
1027 transport: Suspended sediments in the Nepal Himalayas. *Earth Planet. Sci. Lett.* 351–352,  
1028 158–170. <https://doi.org/10.1016/j.epsl.2012.06.059>
- 1029 Andermann, C., Longuevergne, L., Bonnet, S., Crave, A., Davy, P., Gloaguen, R., 2012c. Impact of  
1030 transient groundwater storage on the discharge of Himalayan rivers. *Nat. Geosci.* 5, 127–132.  
1031 <https://doi.org/10.1038/ngeo1356>
- 1032 Anderson, R.S., 2000. A model of ablation-dominated medial moraines and the generation of  
1033 debris-mantled glacier snouts. *J. Glaciol.* 46, 459–469.
- 1034 Barnett, T.P., Adam, J.C., Lettenmaier, D.P., 2005. Potential impacts of a warming climate on water  
1035 availability in snow-dominated regions. *Nature* 438, 303–309.  
1036 <https://doi.org/10.1038/nature04141>
- 1037 Basnett, S., Kulkarni, A.V., Bolch, T., 2013. The influence of debris cover and glacial lakes on the  
1038 recession of glaciers in Sikkim Himalaya, India. *J. Glaciol.* 59, 1035–1046.  
1039 <https://doi.org/10.3189/2013JoG12J184>
- 1040 Benn, D.I., Bolch, T., Hands, K., Gulley, J.D., Luckman, A., Nicholson, L., Quincey, D.J., Thompson,  
1041 S.S., Toumi, R., Wiseman, S., 2012. Response of debris-covered glaciers in the Mount Everest  
1042 region to recent warming, and implications for outburst flood hazards. *Earth-Science Rev.*  
1043 114, 156–174. <https://doi.org/10.1016/j.earscirev.2012.03.008>
- 1044 Benn, D.I., Gulley, J.D., Luckman, A., Adamek, A., Glowacki, P.S., 2009. Englacial drainage systems  
1045 formed by hydrologically driven crevasse propagation. *J. Glaciol.* 55, 513–523.  
1046 <https://doi.org/10.3189/002214309788816669>
- 1047 Benn, D.I., Thompson, S.S., Gulley, J.D., Mertes, J.R., Luckman, A., Nicholson, L., 2017. Structure  
1048 and evolution of the drainage system of a Himalayan debris-covered glacier, and its  
1049 relationship with patterns of mass loss. *Cryosph.* 11, 2247–2264. [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-2017-29)  
1050 [2017-29](https://doi.org/10.5194/tc-2017-29)
- 1051 Benn, D.I., Wiseman, S., Hands, K.A., 2001. Growth and drainage of supraglacial lakes on debris-  
1052 mantled Ngozumpa Glacier, Khumbu Himal, Nepal. *J. Glaciol.* 47, 626–638.  
1053 <https://doi.org/10.3189/172756501781831729>
- 1054 Benn, D.I., Wiseman, S., Warren, C.R., 2000. Rapid growth of a supraglacial lake, Ngozumpa Glacier,  
1055 Khumbu Himal, Nepal. *IAHS Publ.* 264, 177–185.
- 1056 Bhatt, M.P., Masuzawa, T., Yamamoto, M., Takeuchi, N., 2007. Chemical characteristics of pond  
1057 waters within the debris area of Lirung Glacier in Nepal Himalaya. *J. Limnol.* 66, 71–80.
- 1058 Bocchiola, D., Senese, A., Mihalcea, C., Mosconi, B., D’Agata, C., Smiraglia, C., Diolaiuti, G., 2015.  
1059 An ablation model for debris-covered ice: The case study of Venerocolo glacier (Italian Alps).  
1060 *Geogr. Fis. e Din. Quat.* 38, 113–128. <https://doi.org/10.4461/GFDQ.2015.38.11>

- 1061 Bolch, T., 2017. Hydrology: Asian glaciers are a reliable water source. *Nature* 545, 161–162.  
1062 <https://doi.org/10.1038/545161a>
- 1063 Bolch, T., Kulkarni, A., Kaab, A., Huggel, C., Paul, F., Cogley, J.G., Frey, H., Kargel, J.S., Fujita, K.,  
1064 Scheel, M., Bajracharya, S., Stoffel, M., 2012. The State and Fate of Himalayan Glaciers.  
1065 *Science*. 336, 310–314. <https://doi.org/10.1126/science.1215828>
- 1066 Bolch, T., Pieczonka, T., Benn, D.I., 2011. Multi-decadal mass loss of glaciers in the Everest area  
1067 (Nepal Himalaya) derived from stereo imagery. *Cryosph.* 5, 349–358.  
1068 <https://doi.org/10.5194/tc-5-349-2011>
- 1069 Bolch, T., Shea, J.M., Liu, S., Azam, F.M., Gao, Y., Gruber, S., 2019. Status and Change of the  
1070 Cryosphere in the Extended Hindu Kush Himalaya Region, in: Wester, P., Mishra, A., Mukherji,  
1071 A., Shrestha, A.B. (Eds.), *The Hindu Kush Himalaya Assessment*. Springer, Cham, Switzerland,  
1072 pp. 209–255. [https://doi.org/doi.org/10.1007/978-3-319-92288-1\\_7](https://doi.org/doi.org/10.1007/978-3-319-92288-1_7)
- 1073 Bookhagen, B., Burbank, D.W., 2006. Topography, relief, and TRMM-derived rainfall variations  
1074 along the Himalaya. *Geophys. Res. Lett.* 33, 1–5. <https://doi.org/10.1029/2006GL026037>
- 1075 Braun, L.N., Grabs, W., Rana, B., 1993. Application of a Conceptual Precipitation- Runoff Model in  
1076 the Langtang Khola Basin , Nepal Himalaya. *Snow Glacier Hydrol. (Proceedings Kathmandu*  
1077 *Symp. Novemb. 1992)*. IAHS Publ. no. 218,1993. 218, 221–237.
- 1078 Brock, B.W., Mihalcea, C., Kirkbride, M.P., Diolaiuti, G., Cutler, M.E.J., Smiraglia, C., 2010.  
1079 Meteorology and surface energy fluxes in the 2005-2007 ablation seasons at the Miage  
1080 debris-covered glacier, Mont Blanc Massif, Italian Alps. *J. Geophys. Res. Atmos.* 115, 1–16.  
1081 <https://doi.org/10.1029/2009JD013224>
- 1082 Brun, F., Berthier, E., Wagnon, P., Käab, A., Treichler, D., 2017. A spatially resolved estimate of High  
1083 Mountain Asia glacier mass balances from 2000 to 2016. *Nat. Geosci.* 10, 668–673.  
1084 <https://doi.org/10.1038/ngeo2999>
- 1085 Brun, F., Buri, P., Miles, E.S., Wagnon, P., Steiner, J.F., Berthier, E., Ragetti, S., Kraaijenbrink, P.D.A.,  
1086 Immerzeel, W.W., Pellicciotti, F., 2016. Quantifying volume loss from ice cliffs on debris-  
1087 covered glaciers using high-resolution terrestrial and aerial photogrammetry. *J. Glaciol.* 62,  
1088 684–695. <https://doi.org/10.1017/jog.2016.54>
- 1089 Brun, F., Wagnon, P., Berthier, E., Jomelli, V., Maharjan, S.B., Shrestha, F., Kraaijenbrink, P.D.A.,  
1090 2019. Heterogeneous Influence of Glacier Morphology on the Mass Balance Variability in High  
1091 Mountain Asia. *J. Geophys. Res. Earth Surf.* 124, 1331–1345.  
1092 <https://doi.org/10.1029/2018JF004838>
- 1093 Brun, F., Wagnon, P., Berthier, E., Shea, J.M., Immerzeel, W.W., Kraaijenbrink, P.D.A., Vincent, C.,  
1094 Reverchon, C., Shrestha, D., Arnaud, Y., 2018. Ice cliff contribution to the tongue-wide  
1095 ablation of Changri Nup Glacier, Nepal, central Himalaya. *Cryosph.* 12, 3439–3457.  
1096 <https://doi.org/10.5194/tc-12-3439-2018>
- 1097 Buri, P., Miles, E.S., Steiner, J.F., Immerzeel, W.W., Wagnon, P., Pellicciotti, F., 2016a. A physically-  
1098 based 3-D model of ice cliff evolution on a debris-covered glacier. *J. Geophys. Res. Earth Surf.*  
1099 121, 2471–2493. <https://doi.org/10.1002/2016JF004039>
- 1100 Buri, P., Pellicciotti, F., Steiner, J.F., Miles, E.S., Immerzeel, W.W., 2016b. A grid-based model of  
1101 backwasting of supraglacial ice cliffs on debris-covered glaciers. *Ann. Glaciol.* 57, 199–211.  
1102 <https://doi.org/10.3189/2016AoG71A059>

- 1103 Byers, A.C., Byers, E.A., McKinney, D.C., Rounce, D.R., 2017. A field-based study of impacts of the  
1104 2015 earthquake on potentially dangerous glacial lakes in Nepal. *Himalaya* 37, 26–41.
- 1105 Careno, M., Pellicciotti, F., Mabillard, J., Reid, T., Brock, B.W., 2016. An enhanced temperature  
1106 index model for debris-covered glaciers accounting for thickness effect. *Adv. Water Resour.*  
1107 94, 457–469. <https://doi.org/10.1016/j.advwatres.2016.05.001>
- 1108 Carrivick, J.L., Tweed, F.S., 2013. Proglacial Lakes: Character, behaviour and geological importance.  
1109 *Quat. Sci. Rev.* 78, 34–52. <https://doi.org/10.1016/j.quascirev.2013.07.028>
- 1110 Casey, K.A., Käab, A., Benn, D.I., 2012. Geochemical characterization of supraglacial debris via in  
1111 situ and optical remote sensing methods: a case study in Khumbu Himalaya, Nepal. *Cryosph.*  
1112 6, 85–100. <https://doi.org/10.5194/tc-6-85-2012>
- 1113 Chaturvedi, R.K., Kulkarni, A., Karyakarte, Y., Joshi, J., Bala, G., 2014. Glacial mass balance changes  
1114 in the Karakoram and Himalaya based on CMIP5 multi-model climate projections. *Clim.*  
1115 *Change* 123, 315–328. <https://doi.org/10.1007/s10584-013-1052-5>
- 1116 Chen, C., Ding, Y., 2009. The application of artificial neural networks to simulate meltwater runoff  
1117 of Keqikaer Glacier, south slope of Mt. Tuomuer, western China. *Environ. Geol.* 57, 1839–  
1118 1845. <https://doi.org/10.1007/s00254-008-1471-1>
- 1119 Chikita, K., Jha, J., Yamada, T., 1998. The basin expansion mechanism of a supraglacial lake in the  
1120 Nepal Himalaya. *J. Hokkaido Univ. Fac. Sci. Ser. VII Geophys.* 11, 501–521.
- 1121 Chu, V.W., 2014. Greenland ice sheet hydrology: A review. *Prog. Phys. Geogr.* 38, 19–54.  
1122 <https://doi.org/10.1177/0309133313507075>
- 1123 Collier, E., Maussion, F., Nicholson, L., Mölg, T., Immerzeel, W.W., Bush, A., 2015. Impact of debris  
1124 cover on glacier ablation and atmosphere-glacier feedbacks in the Karakoram. *Cryosph.* 9,  
1125 1617–1632. <https://doi.org/10.5194/tc-9-1617-2015>
- 1126 Collier, E., Nicholson, L., Brock, B.W., Maussion, F., Essery, R.L.H., Bush, A.B.G., 2014. Representing  
1127 moisture fluxes and phase changes in glacier debris cover using a reservoir approach.  
1128 *Cryosph.* 8, 1429–1444. <https://doi.org/10.5194/tc-8-1429-2014>
- 1129 Conway, H., Rasmussen, L.A., 2000. Summer temperature profiles within supraglacial debris on  
1130 Khumbu Glacier, Nepal, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered*  
1131 *Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 89–97.
- 1132 Copland, L., Pope, S., Bishop, M.P., Shroder, J.F., Clendon, P., Bush, A., Kamp, U., Seong, Y.B., Owen,  
1133 L.A., 2009. Glacier velocities across the Karakoram Himalaya. *Ann. Glaciol.* 50, 1–18.  
1134 <https://doi.org/10.3189/172756409789624229>
- 1135 Dehecq, A., Gourmelen, N., Gardner, A.S., Brun, F., Goldberg, D., Nienow, P.W., Berthier, E.,  
1136 Vincent, C., Wagnon, P., Trouvé, E., 2019. Twenty-first century glacier slowdown driven by  
1137 mass loss in High Mountain Asia. *Nat. Geosci.* 12, 22–27. <https://doi.org/10.1038/s41561-018-0271-9>
- 1139 Dunning, S.A., Rosser, N.J., Mccoll, S.T., Reznichenko, N.V., 2015. Rapid sequestration of rock  
1140 avalanche deposits within glaciers. *Nat. Commun.* 6, 1–7.  
1141 <https://doi.org/10.1038/ncomms8964>
- 1142 Eriksson, M., Jianchu, X., Shrestha, A., Vaidya, R., Nepal, S., Sandström, K., 2009. The Changing  
1143 Himalayas: Impact of climate change on water resources and livelihoods in the greater  
1144 Himalayas. *Int. Cent. Integr. Mt. Dev.* 114, 1–28. <https://doi.org/10.1144/SP312.3>

- 1145 Evatt, G.W., Abrahams, I.D., Heil, M., Mayer, C., Kingslake, J., Mitchell, S.L., Fowler, A.C., Clark, C.D.,  
1146 2015. Glacial melt under a porous debris layer. *J. Glaciol.* 61, 825–836.  
1147 <https://doi.org/10.3189/2015JoG14J235>
- 1148 Fountain, A.G., Walder, J.S., 1998. Water flow through temperate glaciers. *Rev. Geophys.* 36, 299.  
1149 <https://doi.org/10.1029/97RG03579>
- 1150 Fujita, K., Sakai, A., 2014. Modelling runoff from a Himalayan debris-covered glacier. *Hydrol. Earth*  
1151 *Syst. Sci.* 18, 2679–2694. <https://doi.org/10.5194/hess-18-2679-2014>
- 1152 Fyffe, C.L., Brock, B.W., Kirkbride, M.P., Black, A.R., Smiraglia, C., Diolaiuti, G., 2019a. The influence  
1153 of supraglacial debris on proglacial runoff fluctuations and water chemistry. *J. Hydrol.* 576,  
1154 41–57. <https://doi.org/10.1016/j.jhydrol.2019.06.023>
- 1155 Fyffe, C.L., Brock, B.W., Kirkbride, M.P., Mair, D.W.F., Arnold, N.S., Smiraglia, C., Diolaiuti, G., Diotri,  
1156 F., 2019b. Do debris-covered glaciers demonstrate distinctive hydrological behaviour  
1157 compared to clean glaciers? *J. Hydrol.* 570, 584–597.  
1158 <https://doi.org/10.1016/j.jhydrol.2018.12.069>
- 1159 Fyffe, C.L., Reid, T.D., Brock, B.W., Kirkbride, M.P., Diolaiuti, G., Smiraglia, C., Diotri, F., 2014. A  
1160 distributed energy-balance melt model of an alpine debris-covered glacier. *J. Glaciol.* 60, 587–  
1161 602. <https://doi.org/10.3189/2014JoG13J148>
- 1162 Gardelle, J., Arnaud, Y., Berthier, E., 2011. Contrasted evolution of glacial lakes along the Hindu  
1163 Kush Himalaya mountain range between 1990 and 2009. *Glob. Planet. Change* 75, 47–55.  
1164 <https://doi.org/10.1016/j.gloplacha.2010.10.003>
- 1165 Gardelle, J., Berthier, E., Arnaud, Y., 2012. Slight mass gain of Karakoram glaciers in the early  
1166 twenty-first century. *Nat. Geosci.* 5, 322–325. <https://doi.org/10.1038/ngeo1450>
- 1167 Gibson, M.J., Glasser, N.F., Quincey, D.J., Mayer, C., Rowan, A.V., Irvine-Fynn, T.D.L., 2017a.  
1168 Temporal variations in supraglacial debris distribution on Baltoro Glacier, Karakoram between  
1169 2001 and 2012. *Geomorphology* 295, 572–585.  
1170 <https://doi.org/10.1016/j.geomorph.2017.08.012>
- 1171 Gibson, M.J., Glasser, N.F., Quincey, D.J., Rowan, A.V., Irvine-Fynn, T.D.L., 2017b. Changes in glacier  
1172 surface cover on Baltoro glacier, Karakoram, north Pakistan, 2001–2012. *J. Maps* 13, 100–108.  
1173 <https://doi.org/10.1080/17445647.2016.1264319>
- 1174 Gilbert, A., Sinisalo, A., Gurung, T.R., Fujita, K., Maharjan, S.B., Sherpa, T.C., Fukuda, T., 2020. The  
1175 influence of water percolation through crevasses on the thermal regime of a Himalayan  
1176 mountain glacier. *Cryosph.* 14, 1273–1288. <https://doi.org/10.5194/tc-14-1273-2020>
- 1177 Gremaud, V., Goldscheider, N., Savoy, L., Favre, G., Masson, H., 2009. Geological structure,  
1178 recharge processes and underground drainage of a glacierised karst aquifer system,  
1179 Tsanfleuron-Sanetsch, Swiss Alps. *Hydrogeol. J.* 17, 1833–1848.  
1180 <https://doi.org/10.1007/s10040-009-0485-4>
- 1181 Gulley, J.D., Benn, D.I., 2007. Structural control of englacial drainage systems in Himalayan debris-  
1182 covered glaciers. *J. Glaciol.* 53, 399–412. <https://doi.org/10.3189/002214307783258378>
- 1183 Gulley, J.D., Benn, D.I., Müller, D., Luckman, A., 2009a. A cut-and-closure origin for englacial  
1184 conduits in uncrevassed regions of polythermal glaciers. *J. Glaciol.* 55, 66–80.  
1185 <https://doi.org/10.3189/002214309788608930>
- 1186 Gulley, J.D., Benn, D.I., Screatton, E., Martin, J., 2009b. Mechanisms of englacial conduit formation

- 1187 and their implications for subglacial recharge. *Quat. Sci. Rev.* 28, 1984–1999.  
1188 <https://doi.org/10.1016/j.quascirev.2009.04.002>
- 1189 Hambrey, M.J., Quincey, D.J., Glasser, N.F., Reynolds, J.M., Richardson, S.J., Clemmens, S., 2008.  
1190 Sedimentological, geomorphological and dynamic context of debris-mantled glaciers, Mount  
1191 Everest (Sagarmatha) region, Nepal. *Quat. Sci. Rev.* 28, 1084.  
1192 <https://doi.org/10.1016/j.quascirev.2009.04.009>
- 1193 Han, H., Wang, J., Wei, J., Liu, S., 2010. Backwasting rate on debris-covered Koxkar glacier,  
1194 Tuomuer mountain, China. *J. Glaciol.* 56, 287–296.  
1195 <https://doi.org/10.3189/002214310791968430>
- 1196 Hannah, D.M., Kansakar, S.R., Gerrard, A.J., Rees, G., 2005. Flow regimes of Himalayan rivers of  
1197 Nepal: Nature and spatial patterns. *J. Hydrol.* 308, 18–32.  
1198 <https://doi.org/10.1016/j.jhydrol.2004.10.018>
- 1199 Haritashya, U.K., Kargel, J.S., Shugar, D.H., Leonard, G.J., Strattman, K., Watson, C.S., Shean, D.,  
1200 Harrison, S., Mandli, K.T., Regmi, D., 2018a. Evolution and controls of large glacial lakes in the  
1201 Nepal Himalaya. *Remote Sens.* 10, 1–31. <https://doi.org/10.3390/rs10050798>
- 1202 Haritashya, U.K., Kargel, J.S., Shugar, D.H., Leonard, G.J., Strattman, K., Watson, C.S., Shean, D.E.,  
1203 Harrison, S., Mandli, K.T., Regmi, D., 2018b. Evolution and controls of large glacial lakes in the  
1204 Nepal Himalaya. *Remote Sens.* 10, 1–31. <https://doi.org/10.3390/rs10050798>
- 1205 Haritashya, U.K., Kumar, A., Singh, P., 2010. Particle size characteristics of suspended sediment  
1206 transported in meltwater from the Gangotri Glacier, central Himalaya - An indicator of  
1207 subglacial sediment evacuation. *Geomorphology* 122, 140–152.  
1208 <https://doi.org/10.1016/j.geomorph.2010.06.006>
- 1209 Hasnain, S.I., 1999. Runoff characteristics of a glacierized catchment, Garhwal Himalaya, India.  
1210 *Hydrol. Sci. J.* 44, 847–854. <https://doi.org/10.1080/02626669909492284>
- 1211 Hasnain, S.I., 1996. Factors controlling suspended sediment transport in Himalayan glacier  
1212 meltwaters. *J. Hydrol.* 181, 49–62. [https://doi.org/10.1016/0022-1694\(95\)02917-6](https://doi.org/10.1016/0022-1694(95)02917-6)
- 1213 Hasnain, S.I., Jose, P.G., Ahmad, S., Negi, D.C., 2001. Character of the subglacial drainage system  
1214 in the ablation area of Dokriani glacier, India, as revealed by dye-tracer studies. *J. Hydrol.* 248,  
1215 216–223. [https://doi.org/10.1016/S0022-1694\(01\)00404-8](https://doi.org/10.1016/S0022-1694(01)00404-8)
- 1216 Hasnain, S.I., Thayyen, R.J., 1994. Hydrograph separation of bulk meltwaters of Dokriani Bamak  
1217 glacier basin, based on electrical conductivity. *Curr. Sci.* 67, 189–193.  
1218 <https://doi.org/24095811>
- 1219 Hewitt, K., Wake, C.P., Young, G.J., David, C., 1989. Hydrological investigations at Biafo Glacier,  
1220 Karakoram range, Himalaya: an important source of water for the Indus River. *Ann. Glaciol.*  
1221 13, 103–108. <https://doi.org/10.3189/S0260305500007710>
- 1222 Hock, R., Rasul, G., Adler, C., Caceres, S., Gruber, Y., Hirabayashi, Y., Jackson, M., Käab, A., Kang, S.,  
1223 Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., Steltzer, H., 2019. High Mountain  
1224 Areas, in: Pörtner, H.O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska,  
1225 E., Mintenbeck, K., Nicolai, M., Okem, A., Petzold, J., Rama, B., Weyer, N. (Eds.), IPCC Special  
1226 Report on the Ocean and Cryosphere in a Changing Climate.
- 1227 Hodson, A.J., 2014. Understanding the dynamics of black carbon and associated contaminants in  
1228 glacial systems. *WIREs Water* 1, 141–149. <https://doi.org/10.1002/wat2.1016>



- 1229 Horodyskyj, U.N., 2015. Contributing factors to ice mass loss on Himalayan debris-covered glaciers.  
1230 University of Colorado, Boulder.
- 1231 Hubbard, B., Nienow, P.W., 1997. Alpine subglacial hydrology. *Quat. Sci. Rev.* 16, 939–955.  
1232 [https://doi.org/10.1016/S0277-3791\(97\)00031-0](https://doi.org/10.1016/S0277-3791(97)00031-0)
- 1233 Huss, M., 2011. Present and future contribution of glacier storage change to runoff from  
1234 macroscale drainage basins in Europe. *Water Resour. Res.* 47, 1–14.  
1235 <https://doi.org/10.1029/2010WR010299>
- 1236 Huss, M., Hock, R., 2018. Global-scale hydrological response to future glacier mass loss. *Nat. Clim.*  
1237 *Chang.* 8, 135–140. <https://doi.org/10.1038/s41558-017-0049-x>
- 1238 Immerzeel, W.W., Kraaijenbrink, P.D.A., Shea, J.M., Shrestha, A.B., Pellicciotti, F., Bierkens, M.F.P.,  
1239 De Jong, S.M., 2014. High-resolution monitoring of Himalayan glacier dynamics using  
1240 unmanned aerial vehicles. *Remote Sens. Environ.* 150, 93–103.  
1241 <https://doi.org/10.1016/j.rse.2014.04.025>
- 1242 Immerzeel, W.W., Lutz, A.F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S.,  
1243 Davies, B.J., Elmore, A.C., Emmer, A., Feng, M., Fernandez, A., Haritashya, U., Kargel, J.S.,  
1244 Hoppes, M., Kraaijenbrink, P.D.A., Kulkarni, A.V., Mayewski, P., Nepal, S., Pacheco, P., Painter,  
1245 T.H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A.B., Viviroli, D., Wada, Y.,  
1246 Xiao, C., Yao, T., Bailie, J.E.M., 2020. Importance and vulnerability of the world’s water towers.  
1247 *Nature* 577, 364–369. <https://doi.org/10.1038/s41586-019-1822-y>
- 1248 Immerzeel, W.W., van Beek, L.P.H., Bierkens, M.F.P., 2010. Climate change will affect the Asian  
1249 water towers. *Science*. 328, 1382–1385. <https://doi.org/10.1126/science.1183188>
- 1250 Immerzeel, W.W., van Beek, L.P.H., Konz, M., Shrestha, A.B., Bierkens, M.F.P., 2012. Hydrological  
1251 response to climate change in a glacierized catchment in the Himalayas. *Clim. Change* 110,  
1252 721–736. <https://doi.org/10.1007/s10584-011-0143-4>
- 1253 Inoue, J., Yoshida, M., 1980. Ablation and Heat Exchange over the Khumbu Glacier. *J. Japanese Soc.*  
1254 *Snow Ice* 41, 26–33. [https://doi.org/10.5331/seppyo.41.Special\\_26](https://doi.org/10.5331/seppyo.41.Special_26)
- 1255 Irvine-Fynn, T.D.L., Hodson, A.J., Moorman, B.J., Vatne, G., Hubbard, A.L., 2011. Polythermal  
1256 Glacier Hydrology: A Review. *Rev. Geophys.* 49, 1–37.  
1257 <https://doi.org/10.1029/2010RG000350>
- 1258 Irvine-Fynn, T.D.L., Porter, P.R., Rowan, A.V., Quincey, D.J., Gibson, M.J., Bridge, J.W., Watson, C.S.,  
1259 Hubbard, A.L., Glasser, N.F., 2017. Supraglacial Ponds Regulate Runoff From Himalayan  
1260 Debris-Covered Glaciers. *Geophys. Res. Lett.* 44, 11,894–11,904.  
1261 <https://doi.org/10.1002/2017GL075398>
- 1262 Iwata, S., Watanabe, O., Fushimi, H., 1980. Surface morphology in the ablation area of the Khumbu  
1263 Glacier. *J. Japanese Soc. Snow Ice* 41, 9–17. [https://doi.org/10.5331/seppyo.41.Special\\_9](https://doi.org/10.5331/seppyo.41.Special_9)
- 1264 Jansson, P., Hock, R., Schneider, T., 2003. The concept of glacier storage: a review. *J. Hydrol.* 282,  
1265 116–129. [https://doi.org/10.1016/S0022-1694\(03\)00258-0](https://doi.org/10.1016/S0022-1694(03)00258-0)
- 1266 Jarosch, A.H., Gudmundsson, M.T., 2012. A numerical model for meltwater channel evolution in  
1267 glaciers. *Cryosph.* 6, 493–503. <https://doi.org/10.5194/tc-6-493-2012>
- 1268 Juen, M., Mayer, C., Lambrecht, A., Han, H., Liu, S., 2014. Impact of varying debris cover thickness  
1269 on ablation: A case study for Koxkar Glacier in the Tien Shan. *Cryosphere* 8, 377–386.  
1270 <https://doi.org/10.5194/tc-8-377-2014>

- 1271 Kääb, A., 2005. Combination of SRTM3 and repeat ASTER data for deriving alpine glacier flow  
1272 velocities in the Bhutan Himalaya. *Remote Sens. Environ.* 94, 463–474.  
1273 <https://doi.org/10.1016/j.rse.2004.11.003>
- 1274 Kääb, A., Berthier, E., Nuth, C., Gardelle, J., Arnaud, Y., 2012. Contrasting patterns of early twenty-  
1275 first-century glacier mass change in the Himalayas. *Nature* 488, 495–498.  
1276 <https://doi.org/10.1038/nature11324>
- 1277 Kargel, J.S., Leonard, G.J., Shugar, D.H., Haritashya, U.K., Bevington, A., Fielding, E.J., Fujita, K.,  
1278 Geertsema, M., Miles, E.S., Steiner, J., Anderson, E., Bajracharya, S., Bawden, G.W.,  
1279 Breashears, D.F., Byers, A., Collins, B., Dhital, M.R., Donnellan, A., Evans, T.L., Geai, M.L.,  
1280 Glasscoe, M.T., Green, D., Gurung, D.R., Heijenk, R., Hilborn, A., Hudnut, K., Huyck, C.,  
1281 Immerzeel, W.W., Jiang, L., Jibson, R., Kääb, A., Khanal, N.R., Kirschbaum, D., Kraaijenbrink,  
1282 P.D.A., Lamsal, D., Liu, S., Lv, M., McKinney, D., Nahirnick, N.K., Nan, Z., Ojha, S., Olsenholler,  
1283 J., Painter, T.H., Pleasants, M., Pratima, K.C., Yuan, Q.I., Raup, B.H., Regmi, D., Rounce, D.R.,  
1284 Sakai, A., Shangguan, D., Shea, J.M., Shrestha, A.B., Shukla, A., Stumm, D., Van Der Kooij, M.,  
1285 Voss, K., Wang, X., Weihs, B., Wolfe, D., Wu, L., Yao, X., Yoder, M.R., Young, N., 2016.  
1286 Geomorphic and geologic controls of geohazards induced by Nepal’s 2015 Gorkha  
1287 earthquake. *Science*. 351. <https://doi.org/10.1126/science.aac8353>
- 1288 Kattelmann, R., 2003. Glacial lake outburst floods in the Nepal Himalaya: A manageable hazard?  
1289 *Nat. Hazards* 28, 145–154. <https://doi.org/10.1023/A:1021130101283>
- 1290 Kayastha, R.B., Takeuchi, Y., Nakawo, M., Ageta, Y., 2000. Practical prediction of ice melting  
1291 beneath various thickness of debris cover on Khumbu Glacier, Nepal, using a positive degree-  
1292 day factor, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered Glaciers*.  
1293 International Association of Hydrological Sciences, Oxford, pp. 71–81.
- 1294 Kehrwald, N.M., Thompson, L.G., Tandong, Y., Mosley-Thompson, E., Schotterer, U., Alifimov, V.,  
1295 Beer, J., Eikenberg, J., Davis, M.E., 2008. Mass loss on Himalayan glacier endangers water  
1296 resources. *Geophys. Res. Lett.* 35, 2–7. <https://doi.org/10.1029/2008GL035556>
- 1297 Khan, A.A., Pant, N.C., Sarkar, A., Tandon, S.K., Thamban, M., Mahalinganathan, K., 2017. The  
1298 Himalayan cryosphere: A critical assessment and evaluation of glacial melt fraction in the  
1299 Bhagirathi basin. *Geosci. Front.* 8, 107–115. <https://doi.org/10.1016/j.gsf.2015.12.009>
- 1300 King, O., Bhattacharya, A., Bhambri, R., Bolch, T., 2019. Glacial lakes exacerbate Himalayan glacier  
1301 mass loss. *Sci. Rep.* 9, 1–9. <https://doi.org/10.1038/s41598-019-53733-x>
- 1302 King, O., Quincey, D.J., Carrivick, J.L., Rowan, A.V., 2017. Spatial variability in mass loss of glaciers  
1303 in the Everest region, central Himalaya, between 2000 and 2015. *Cryosph.* 11, 407–426.  
1304 <https://doi.org/10.5194/tc-2016-99>
- 1305 Kirkbride, M.P., 2011. Debris-Covered Glaciers, in: Singh, V., Singh, P., Haritashya, U. (Eds.),  
1306 *Encyclopedia of Snow, Ice and Glaciers*. Springer Netherlands, pp. 180–182.  
1307 [https://doi.org/10.1007/978-90-481-2642-2\\_622](https://doi.org/10.1007/978-90-481-2642-2_622)
- 1308 Kirkbride, M.P., 1993. The temporal significance of transitions from melting to calving termini in  
1309 the glaciers of the central Southern Alps of New Zealand. *The Holocene* 3, 232–240.
- 1310 Kirkbride, M.P., Deline, P., 2013. The formation of supraglacial debris covers by primary dispersal  
1311 from transverse englacial debris bands. *Earth Surf. Process. Landforms* 38, 1779–1792.  
1312 <https://doi.org/10.1002/esp.3416>
- 1313 Kirkbride, M.P., Warren, C.R., 1999. Tasman Glacier, New Zealand: 20th-century thinning and

- 1314 predicted calving retreat. *Glob. Planet. Change* 22, 11–28. <https://doi.org/10.1016/S0921->  
 1315 8181(99)00021-1
- 1316 Kirkbride, M.P., Warren, C.R., 1997. Calving processes at a grounded ice cliff. *Ann. Glaciol.* 24, 116–  
 1317 121.
- 1318 Kodama, H., Mae, S., 1976. The Flow of Glaciers in the Khumbu Region. *J. Japanese Soc. Snow Ice*  
 1319 38, 31–36.
- 1320 Komori, J., 2008. Recent expansions of glacial lakes in the Bhutan Himalayas. *Quat. Int.* 184, 177–  
 1321 186. <https://doi.org/10.1016/j.quaint.2007.09.012>
- 1322 Kraaijenbrink, P.D.A., Bierkens, M.F.P., Lutz, A.F., Immerzeel, W.W., 2017. Impact of a global  
 1323 temperature rise of 1.5 degrees Celsius on Asia's glaciers. *Nature* 549, 257–260.  
 1324 <https://doi.org/10.1038/nature23878>
- 1325 Kraaijenbrink, P.D.A., Meijer, S.W., Shea, J.M., Pellicciotti, F., De Jong, S.M., Immerzeel, W.W.,  
 1326 2016a. Seasonal surface velocities of a Himalayan glacier derived by automated correlation  
 1327 of unmanned aerial vehicle imagery. *Ann. Glaciol.* 57, 103–113.  
 1328 <https://doi.org/10.3189/2016AoG71A072>
- 1329 Kraaijenbrink, P.D.A., Shea, J., Pellicciotti, F., de Jong, S., Immerzeel, W., 2016b. Object-based  
 1330 analysis of unmanned aerial vehicle imagery to map and characterise surface features on a  
 1331 debris-covered glacier. *Remote Sens. Environ.* 186, 581–595.  
 1332 <https://doi.org/10.1016/j.rse.2016.09.013>
- 1333 Kumar, K., Miral, M.S., Joshi, S., Pant, N., Joshi, V., Joshi, L.M., 2009. Solute dynamics of meltwater  
 1334 of Gangotri glacier, Garhwal Himalaya, India. *Environ. Geol.* 58, 1151–1159.  
 1335 <https://doi.org/10.1007/s00254-008-1592-6>
- 1336 Kumar, S., Dobhal, D.P., 1997. Climatic effects and bedrock control on rapid fluctuations of Chhota  
 1337 Shigri glacier, northwest Himalaya, India. *J. Glaciol.* 43, 467–472.
- 1338 Lejeune, Y., Bertrand, J., Wagnon, P., Morin, S., 2013. A physically based model of the year-round  
 1339 surface energy and mass balance of debris-covered glaciers. *J. Glaciol.* 59, 327–344.  
 1340 <https://doi.org/10.3189/2013JoG12J149>
- 1341 Li, J., Yuan, G., Wu, M., Sun, Y., Han, P., Wang, G., 2017. Evidence for persistent organic pollutants  
 1342 released from melting glacier in the central Tibetan Plateau, China. *Environ. Pollut.* 220, 178–  
 1343 185. <https://doi.org/10.1016/j.envpol.2016.09.037>
- 1344 Linsbauer, A., Frey, H., Haeberli, W., Machguth, H., Azam, M.F., Allen, S., 2016. Modelling glacier-  
 1345 bed overdeepenings and possible future lakes for the glaciers in the Himalaya-Karakoram  
 1346 region. *Ann. Glaciol.* 57, 119–130. <https://doi.org/10.3189/2016AoG71A627>
- 1347 Liu, Q., Liu, S., Cao, W., 2018. Seasonal Variation of Drainage System in the Lower Ablation Area of  
 1348 a Monsoonal Temperate Debris-Covered Glacier in Mt. Gongga, South-Eastern Tibet. *Water*  
 1349 10, 1–11. <https://doi.org/10.3390/w10081050>
- 1350 Liu, Q., Mayer, C., Liu, S., 2015. Distribution and interannual variability of supraglacial lakes on  
 1351 debris-covered glaciers in the Khan Tengri-Tumor Mountains, Central Asia. *Environ. Res. Lett.*  
 1352 10, 1–10. <https://doi.org/10.1088/1748-9326/10/1/014014>
- 1353 Lowe, A.T., Collins, D.N., 2001. Modelling runoff from large glacierized basins in the Karakoram  
 1354 Himalaya using remote sensing of the transient snowline. *Remote Sens. Hydrol.* 99–104.

- 1355 Lutz, A.F., Immerzeel, W.W., Shrestha, A.B., Bierkens, M.F.P., 2014. Consistent increase in High  
1356 Asia's runoff due to increasing glacier melt and precipitation. *Nat. Clim. Chang.* 4, 587–592.  
1357 <https://doi.org/10.1038/nclimate2237>
- 1358 Maharjan, S.B., Mool, P.K., Lizong, W., Xiao, G., Shrestha, F., Shrestha, R.B., Khanal, N.R.,  
1359 Bajracharya, S.R., Joshi, S., Shai, S., Baral, P., 2018. The status of glacial lakes in the Hindu Kush  
1360 Himalaya, ICIMOD Res. ed. ICIMOD, Kathmandu.
- 1361 Marston, R., 1983. Supraglacial Stream Dynamics on the Juneau Icefield. *Ann. Assoc. Am. Geogr.*  
1362 73, 597–608.
- 1363 Mattson, L.E., 2000. The influence of a debris cover on the midsummer discharge of Dome Glacier,  
1364 Canadian Rocky Mountains, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-  
1365 Covered Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 25–33.
- 1366 Mayer, C., Lambrecht, A., Belò, M., Smiraglia, C., Diolaiuti, G., 2006. Glaciological characteristics of  
1367 the ablation zone of Baltoro glacier, Karakoram, Pakistan. *Ann. Glaciol.* 43, 123–131.  
1368 <https://doi.org/10.3189/172756406781812087>
- 1369 McCarthy, M., Pritchard, H., Willis, I.C., King, E., 2017. Ground-penetrating radar measurements of  
1370 debris thickness on Lirung Glacier, Nepal. *J. Glaciol.* 63, 543–555.  
1371 <https://doi.org/10.1017/jog.2017.18>
- 1372 Mertes, J.R., Thompson, S.S., Booth, A.D., Gulley, J.D., Benn, D.I., 2016. A conceptual model of  
1373 supraglacial lake formation on debris-covered glaciers based on GPR facies analysis. *Earth  
1374 Surf. Process. Landforms* 42, 903–914. <https://doi.org/10.1002/esp.4068>
- 1375 Mihalcea, C., Mayer, C., Diolaiuti, G., Lambrecht, A., Smiraglia, C., Tartari, G., 2006. Ice ablation  
1376 and meteorological conditions on the debris-covered area of Baltoro glacier, Karakoram,  
1377 Pakistan. *Ann. Glaciol.* 43, 292–300. <https://doi.org/10.3189/172756406781812104>
- 1378 Miles, E.S., Pellicciotti, F., Willis, I.C., Steiner, J.F., Buri, P., Arnold, N.S., 2016. Refined energy-  
1379 balance modelling of a supraglacial pond, Langtang Khola, Nepal. *Ann. Glaciol.* 57, 29–40.  
1380 <https://doi.org/10.3189/2016AoG71A421>
- 1381 Miles, E.S., Steiner, J.F., Willis, I.C., Buri, P., Immerzeel, W.W., Chesnokova, A., Pellicciotti, F.,  
1382 2017a. Pond dynamics and supraglacial-englacial connectivity on debris-covered Lirung  
1383 Glacier. *Front. Earth Sci.* 5. <https://doi.org/10.3389/feart.2017.00069>
- 1384 Miles, E.S., Watson, C.S., Brun, F., Berthier, E., Esteves, M., Quincey, D.J., Miles, K.E., Hubbard, B.,  
1385 Wagnon, P., 2018a. Glacial and geomorphic effects of a supraglacial lake drainage and  
1386 outburst event, Nepal Himalaya. *Cryosph.* 12, 3891–3905. [https://doi.org/10.5194/tc-2018-  
152](https://doi.org/10.5194/tc-2018-<br/>1387 152)
- 1388 Miles, E.S., Willis, I.C., Arnold, N.S., Steiner, J.F., Pellicciotti, F., 2017b. Spatial, seasonal, and  
1389 interannual variability of supraglacial ponds in the Langtang Valley of Nepal, 1999 to 2013. *J.  
1390 Glaciol.* 63, 88–105. <https://doi.org/10.1017/jog.2016.120>
- 1391 Miles, E.S., Willis, I.C., Buri, P., Steiner, J.F., Arnold, N.S., Pellicciotti, F., 2018b. Surface pond energy  
1392 absorption across four Himalayan glaciers accounts for 1/8 of total catchment ice loss.  
1393 *Geophys. Res. Lett.* 45. <https://doi.org/10.1029/2018GL079678>
- 1394 Miles, K.E., Hubbard, B., Quincey, D.J., Miles, E.S., Irvine-Fynn, T.D.L., Rowan, A.V., 2019. Surface  
1395 and subsurface hydrology of debris-covered Khumbu Glacier, Nepal, revealed by dye tracing.  
1396 *Earth Planet. Sci. Lett.* 513, 176–186. <https://doi.org/doi.org/10.1016/j.epsl.2019.02.020>

- 1397 Miles, K.E., Hubbard, B., Quincey, D.J., Miles, E.S., Sherpa, T.C., Rowan, A.V., Doyle, S.H., 2018.  
1398 Polythermal structure of a Himalayan debris-covered glacier revealed by borehole  
1399 thermometry. *Sci. Rep.* 8, 1–9. <https://doi.org/10.1038/s41598-018-34327-5>
- 1400 Narama, C., Daiyrov, M., Tadono, T., Yamamoto, M., Kääh, A., Morita, R., Jinro, U., 2017. Seasonal  
1401 drainage of supraglacial lakes on debris-covered glaciers in the Tien Shan Mountains, Central  
1402 Asia. *Geomorphology* 286, 133–142. <https://doi.org/10.1016/j.geomorph.2017.03.002>
- 1403 Nicholson, L., Benn, D.I., 2013. Properties of natural supraglacial debris in relation to modelling  
1404 sub-debris ice ablation. *Earth Surf. Process. Landforms* 38, 490–501.  
1405 <https://doi.org/10.1002/esp.3299>
- 1406 Nicholson, L., Benn, D.I., 2006. Calculating ice melt beneath a debris layer using meteorological  
1407 data. *J. Glaciol.* 52, 463–470. <https://doi.org/10.3189/172756506781828584>
- 1408 Nicholson, L.I., McCarthy, M., Pritchard, H.D., Willis, I., 2018. Supraglacial debris thickness  
1409 variability: Impact on ablation and relation to terrain properties. *Cryosph.* 12, 3719–3734.  
1410 <https://doi.org/10.5194/tc-12-3719-2018>
- 1411 Nie, Y., Sheng, Y., Liu, Q., Liu, L., Liu, S., Zhang, Y., Song, C., 2017. A regional-scale assessment of  
1412 Himalayan glacial lake changes using satellite observations from 1990 to 2015. *Remote Sens.*  
1413 *Environ.* 189, 1–13. <https://doi.org/10.1016/j.rse.2016.11.008>
- 1414 Nuimura, T., Fujita, K., Yamaguchi, S., Sharma, R.R., 2012. Elevation changes of glaciers revealed  
1415 by multitemporal digital elevation models calibrated by GPS survey in the Khumbu region,  
1416 Nepal Himalaya, 1992–2008. *J. Glaciol.* 58, 648–656. <https://doi.org/10.3189/2012JoG11J061>
- 1417 Østrem, G., 1959. Ice Melting under a Thin Layer of Moraine, and the Existence of Ice Cores in  
1418 Moraine Ridges. *Geogr. Ann.* 41, 228–230.  
1419 <https://doi.org/10.1080/20014422.1959.11907953>
- 1420 Pellicciotti, F., Stephan, C., Miles, E.S., Herreid, S., Immerzeel, W.W., Bolch, T., 2015. Mass-balance  
1421 changes of the debris-covered glaciers in the Langtang Himal, Nepal, from 1974 to 1999. *J.*  
1422 *Glaciol.* 61, 373–386. <https://doi.org/10.3189/2015JoG13J237>
- 1423 Pelto, M.S., 2000. Mass balance of adjacent debris-covered and clean glacier ice in the North  
1424 Cascades, Washington, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered*  
1425 *Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 35–42.
- 1426 Phillips, T., Rajaram, H., Steffen, K., 2010. Cryo-hydrologic warming: A potential mechanism for  
1427 rapid thermal response of ice sheets. *Geophys. Res. Lett.* 37, 1–5.  
1428 <https://doi.org/10.1029/2010GL044397>
- 1429 Pieczonka, T., Bolch, T., Junfeng, W., Shiyin, L., 2013. Heterogeneous mass loss of glaciers in the  
1430 Aksu-Tarim Catchment (Central Tien Shan) revealed by 1976 KH-9 Hexagon and 2009 SPOT-5  
1431 stereo imagery. *Remote Sens. Environ.* 130, 233–244.  
1432 <https://doi.org/10.1016/j.rse.2012.11.020>
- 1433 Pohl, E., Gloaguen, R., Andermann, C., Knoche, M., 2017. Glacier melt buffers river runoff in the  
1434 Pamir Mountains. *Water Resour. Res.* 53, 2467–2489. <https://doi.org/10.1002/2016WR019431>
- 1436 Pottakkal, J.G., Ramanathan, A., Singh, V.B., Sharma, P., Azam, M.F., Linda, A., 2014.  
1437 Characterization of subglacial pathways draining two tributary meltwater streams through  
1438 the lower ablation zone of Gangotri glacier system, Garhwal Himalaya, India. *Curr. Sci.* 107,

- 1439 613–621. <https://doi.org/24103533>
- 1440 Pritchard, H.D., 2019. Asia's shrinking glaciers protect large populations from drought stress.  
1441 Nature 569, 649–654. <https://doi.org/10.1038/s41586-019-1240-1>
- 1442 Quincey, D.J., Braun, M., Glasser, N.F., Bishop, M.P., Hewitt, K., Luckman, A., 2011. Karakoram  
1443 glacier surge dynamics. Geophys. Res. Lett. 38, 2–7. <https://doi.org/10.1029/2011GL049004>
- 1444 Quincey, D.J., Copland, L., Mayer, C., Bishop, M., Luckman, A., Belò, M., 2009. Ice velocity and  
1445 climate variations for Baltoro Glacier, Pakistan. J. Glaciol. 55, 1061–1071.  
1446 <https://doi.org/10.3189/002214309790794913>
- 1447 Quincey, D.J., Richardson, S.D., Luckman, A., Lucas, R.M., Reynolds, J.M., Hambrey, M.J., Glasser,  
1448 N.F., 2007. Early recognition of glacial lake hazards in the Himalaya using remote sensing  
1449 datasets. Glob. Planet. Change 56, 137–152.  
1450 <https://doi.org/10.1016/j.gloplacha.2006.07.013>
- 1451 Radić, V., Bliss, A., Beedlow, A.C., Hock, R., Miles, E.S., Cogley, J.G., 2014. Regional and global  
1452 projections of twenty-first century glacier mass changes in response to climate scenarios from  
1453 global climate models. Clim. Dyn. 42, 37–58. <https://doi.org/10.1007/s00382-013-1719-7>
- 1454 Ragettli, S., Bolch, T., Pellicciotti, F., 2016a. Heterogeneous glacier thinning patterns over the last  
1455 40 years in Langtang Himal, Nepal. Cryosph. 10, 2075–2097. <https://doi.org/10.5194/tc-10-2075-2016>
- 1457 Ragettli, S., Immerzeel, W.W., Pellicciotti, F., 2016b. Contrasting climate change impact on river  
1458 flows from high-altitude catchments in the Himalayan and Andes Mountains. Proc. Natl. Acad.  
1459 Sci. U. S. A. 113, 9222–9227. <https://doi.org/10.1073/pnas.1606526113>
- 1460 Ragettli, S., Pellicciotti, F., Immerzeel, W.W., Miles, E.S., Petersen, L., Heynen, M., Shea, J.M.,  
1461 Stumm, D., Joshi, S., Shrestha, A., 2015. Unraveling the hydrology of a Himalayan catchment  
1462 through integration of high resolution in situ data and remote sensing with an advanced  
1463 simulation model. Adv. Water Resour. 78, 94–111.  
1464 <https://doi.org/10.1016/j.advwatres.2015.01.013>
- 1465 Rana, B., Nakawo, M., Fukushima, Y., Ageta, Y., 1997. Application of a conceptual precipitation-  
1466 runoff model (HYCYMODEL) in a debris-covered glacierized basin in the Langtang Valley,  
1467 Nepal Himalaya. Ann. Glaciol. 25, 226–231.
- 1468 Raymond, C.F., Nolan, M., 2000. Drainage of a glacial lake through an ice spillway, in: Nakawo, M.,  
1469 Raymond, C.F., Fountain, A. (Eds.), Symposium: Debris-Covered Glaciers. International  
1470 Association of Hydrological Sciences, Oxford, pp. 199–210.
- 1471 Rees, G., Collins, D., 2006. Regional differences in response of flow in glacier-fed Himalayan rivers  
1472 to climatic warming. Hydrol. Process. 20, 2157–2169. <https://doi.org/10.1002/hyp>
- 1473 Reid, T.D., Brock, B.W., 2014. Assessing ice-cliff backwasting and its contribution to total ablation  
1474 of debris-covered Miage glacier, Mont Blanc massif, Italy. J. Glaciol. 60, 3–13.  
1475 <https://doi.org/10.3189/2014JoG13J045>
- 1476 Reid, T.D., Brock, B.W., 2010. An energy-balance model for debris-covered glaciers including heat  
1477 conduction through the debris layer. J. Glaciol. 56, 903–916.
- 1478 Reynolds, J.M., 2000. On the formation of supraglacial lakes on debris-covered glaciers. Debris-  
1479 covered glaciers 264, 153–161.

- 1480 Röhl, K., 2008. Characteristics and evolution of supraglacial ponds on debris-covered Tasman  
1481 Glacier, New Zealand. *J. Glaciol.* 54, 867–880. <https://doi.org/10.3189/002214308787779861>
- 1482 Röhl, K., 2006. Thermo-erosional notch development at fresh-water-calving Tasman Glacier, New  
1483 Zealand. *J. Glaciol.* 52, 203–213. <https://doi.org/10.3189/172756506781828773>
- 1484 Rounce, D.R., Byers, A.C., Byers, E.A., McKinney, D.C., 2017. Brief Communications: Observations  
1485 of a glacier outburst flood from Lhotse Glacier, Everest area, Nepal. *Cryosph.* 11, 443–449.  
1486 <https://doi.org/10.5194/tc-2016-239>
- 1487 Rounce, D.R., Hock, R., Shean, D.E., 2020. Glacier Mass Change in High Mountain Asia Through  
1488 2100 Using the Open-Source Python Glacier Evolution Model (PyGEM). *Front. Earth Sci.* 7, 1–  
1489 20. <https://doi.org/10.3389/feart.2019.00331>
- 1490 Rounce, D.R., McKinney, D.C., Lala, J.M., Byers, A.C., Watson, C.S., 2016. A New Remote Hazard  
1491 and Risk Assessment Framework for Glacial Lakes in the Nepal Himalaya. *Hydrol. Earth Syst.*  
1492 *Sci.* 3455–3475. <https://doi.org/10.5194/hess-2016-161>
- 1493 Rowan, A.V., Egholm, D.L., Quincey, D.J., Glasser, N.F., 2015. Modelling the feedbacks between  
1494 mass balance, ice flow and debris transport to predict the response to climate change of  
1495 debris-covered glaciers in the Himalaya. *Earth Planet. Sci. Lett.* 430, 427–438.  
1496 <https://doi.org/10.1016/j.epsl.2015.09.004>
- 1497 Sakai, A., 2012. Glacial Lakes in the Himalayas : A Review on Formation and Expansion Processes.  
1498 *Glob. Environ. Res.* 16, 23–30.
- 1499 Sakai, A., Fujita, K., 2010. Formation conditions of supraglacial lakes on debris-covered glaciers in  
1500 the Himalaya. *J. Glaciol.* 56, 9249–9251.
- 1501 Sakai, A., Nakawo, M., Fujita, K., 2002. Distribution Characteristics and Energy Balance of Ice Cliffs  
1502 on Debris-Covered Glaciers, Nepal Himalaya. *Arctic, Antarct. Alp. Res.* 34, 12–19.
- 1503 Sakai, A., Nakawo, M., Fujita, K., 1998. Melt rate of ice cliffs on the Lirung Glacier, Nepal Himalayas,  
1504 1996. *Bull. Glacier Res.* 16, 57–66.
- 1505 Sakai, A., Nishimura, K., Kadota, T., Takeuchi, N., 2009. Onset of calving at supraglacial lakes on  
1506 debris-covered glaciers of the Nepal Himalaya. *J. Glaciol.* 55, 909–917.  
1507 <https://doi.org/10.3189/002214309790152555>
- 1508 Sakai, A., Takeuchi, N., Fujita, K., Nakawo, M., 2000. Role of supraglacial ponds in the ablation  
1509 process of a debris-covered glacier in the Nepal Himalayas, in: Nakawo, M., Raymond, C.F.,  
1510 Fountain, A. (Eds.), *Debris-Covered Glaciers*. International Association of Hydrological  
1511 Sciences, Oxford, pp. 119–130.
- 1512 Salerno, F., Guyennon, N., Thakuri, S., Viviano, G., Romano, E., Vuillermoz, E., Cristofanelli, P.,  
1513 Stocchi, P., Agrillo, G., Ma, Y., Tartari, G., 2015. Weak precipitation, warm winters and springs  
1514 impact glaciers of south slopes of Mt. Everest (central Himalaya) in the last 2 decades (1994-  
1515 2013). *Cryosph.* 9, 1229–1247. <https://doi.org/10.5194/tc-9-1229-2015>
- 1516 Salerno, F., Thakuri, S., D'Agata, C., Smiraglia, C., Manfredi, E.C., Viviano, G., Tartari, G., 2012.  
1517 Glacial lake distribution in the Mount Everest region: Uncertainty of measurement and  
1518 conditions of formation. *Glob. Planet. Change* 92–93, 30–39.  
1519 <https://doi.org/10.1016/j.gloplacha.2012.04.001>
- 1520 Savéan, M., Delclaux, F., Chevallier, P., Wagnon, P., Gonga-Saholiariliva, N., Sharma, R., Neppel, L.,  
1521 Arnaud, Y., 2015. Water budget on the Dudh Koshi River (Nepal): Uncertainties on

- 1522 precipitation. *J. Hydrol.* 531, 850–862. <https://doi.org/10.1016/j.jhydrol.2015.10.040>
- 1523 Scherler, D., Bookhagen, B., Strecker, M.R., 2011. Spatially variable response of Himalayan glaciers  
1524 to climate change affected by debris cover. *Nat. Geosci.* 4, 156–159.  
1525 <https://doi.org/10.1038/ngeo1068>
- 1526 Scherler, D., Wulf, H., Gorelick, N., 2018. Global Assessment of Supraglacial Debris-Cover Extents.  
1527 *Geophys. Res. Lett.* 45, 11,798–11,805. <https://doi.org/10.1029/2018GL080158>
- 1528 Scott, C.A., Zhang, F., Mukherji, A., Immerzeel, W.W., Mustafa, D., Bharati, L., 2019. Water in the  
1529 Hindu Kush Himalaya, in: Wester, P., Mishra, A., Mukherji, A., Shrestha, A. (Eds.), *The Hindu*  
1530 *Kush Himalaya Assessment*. Springer, Cham, Switzerland, pp. 257–299.  
1531 [https://doi.org/doi.org/10.1007/978-3-319-92288-1\\_8](https://doi.org/doi.org/10.1007/978-3-319-92288-1_8)
- 1532 Shea, J.M., Immerzeel, W.W., 2016. An assessment of basin-scale glaciological and hydrological  
1533 sensitivities in the Hindu Kush-Himalaya. *Ann. Glaciol.* 57, 308–318.  
1534 <https://doi.org/10.3189/2016AoG71A073>
- 1535 Shea, J.M., Immerzeel, W.W., Wagnon, P., Vincent, C., Bajracharya, S., 2015. Modelling glacier  
1536 change in the Everest region, Nepal Himalaya. *Cryosph.* 9, 1105–1128.  
1537 <https://doi.org/10.5194/tc-9-1105-2015>
- 1538 Shean, D., 2017. High Mountain Asia 8-meter DEM Mosaics Derived from Optical Imagery, Version  
1539 1. Tile 677. <https://doi.org/https://doi.org/10.5067/KXOVQ9L172S2>
- 1540 Shreve, R.L., 1972. Movement of Water in Glaciers. *J. Glaciol.* 11, 205–214.  
1541 <https://doi.org/10.3189/S002214300002219X>
- 1542 Singh, P., Arora, M., Goel, N.K., 2006a. Effect of climate change on runoff of a glacierized Himalayan  
1543 basin. *Hydrol. Process.* 20, 1979–1992. <https://doi.org/10.1002/hyp.5991>
- 1544 Singh, P., Bengtsson, L., 2004. Hydrological sensitivity of a large Himalayan basin to climate change.  
1545 *Hydrol. Process.* 18, 2363–2385. <https://doi.org/10.1002/hyp.1468>
- 1546 Singh, P., Haritashya, U.K., Kumar, N., 2008. Modelling and estimation of different components of  
1547 streamflow for Gangotri Glacier basin, Himalayas. *Hydrol. Sci. J.* 53, 309–322.  
1548 <https://doi.org/10.1623/hysj.53.2.309>
- 1549 Singh, P., Haritashya, U.K., Kumar, N., Singh, Y., 2006b. Hydrological characteristics of the Gangotri  
1550 Glacier, central Himalayas, India. *J. Hydrol.* 327, 55–67.  
1551 <https://doi.org/10.1016/j.jhydrol.2005.11.060>
- 1552 Singh, P., Haritashya, U.K., Ramasastri, K.S., Kumar, N., 2005. Diurnal variations in discharge and  
1553 suspended sediment concentration, including runoff-delaying characteristics, of the Gangotri  
1554 Glacier in the Garhwal Himalayas. *Hydrol. Process.* 19, 1445–1457.  
1555 <https://doi.org/10.1002/hyp.5583>
- 1556 Singh, P., Kumar, N., Ramasastri, K.S., Singh, Y., 2000. Influence of a fine debris layer on the melting  
1557 of snow and ice on a Himalayan glacier, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.),  
1558 *Debris-Covered Glaciers*. International Association of Hydrological Sciences, Oxford, pp. 63–  
1559 69.
- 1560 Singh, P., Ramasastri, K.S., Singh, U.K., Gergan, J.T., Dobhal, D.P., 1995. Hydrological characteristics  
1561 of the Dokriani Glacier in the Garhwal Himalayas. *Hydrol. Sci. J.* 40, 243–257.  
1562 <https://doi.org/10.1080/02626669509491407>



- 1563 Smart, C.C., 1996. Statistical evaluation of glacier boreholes as indicators of basal drainage  
1564 systems. *Hydrol. Process.* 10, 599–613. [https://doi.org/10.1002/\(sici\)1099-](https://doi.org/10.1002/(sici)1099-)  
1565 1085(199604)10:4<599::aid-hyp394>3.0.co;2-8
- 1566 Smart, C.C., 1988. Artificial Tracer Techniques for the Determination of the Structure of Conduit  
1567 Aquifers. *Groundwater.* <https://doi.org/10.1111/j.1745-6584.1988.tb00411.x>
- 1568 Smith, T., Bookhagen, B., 2018. Changes in seasonal snow water equivalent distribution in High  
1569 Mountain Asia (1987 to 2009). *Sci. Adv.* 4. <https://doi.org/10.1126/sciadv.1701550>
- 1570 Soncini, A., Bocchiola, D., Confortola, G., Minora, U., Vuillermoz, E., Salerno, F., Viviano, G.,  
1571 Shrestha, D., Senese, A., Smiraglia, C., Diolaiuti, G., 2016. Future hydrological regimes and  
1572 glacier cover in the Everest region: The case study of the upper Dudh Koshi basin. *Sci. Total*  
1573 *Environ.* 1084–1101. <https://doi.org/10.1016/j.scitotenv.2016.05.138>
- 1574 Sorg, A., Bolch, T., Stoffel, M., Solomina, O., Beniston, M., 2012. Climate change impacts on glaciers  
1575 and runoff in Tien Shan (Central Asia). *Nat. Clim. Chang.* 2, 725–731.  
1576 <https://doi.org/10.1038/nclimate1592>
- 1577 Steiner, J.F., Buri, P., Miles, E.S., Ragetti, S., Pellicciotti, F., 2019. Supraglacial ice cliffs and ponds  
1578 on debris-covered glaciers: Spatio-temporal distribution and characteristics. *J. Glaciol.*  
1579 *COMPLETE*, 1–16. <https://doi.org/10.1017/jog.2019.40>
- 1580 Steiner, J.F., Kraaijenbrink, P.D.A., Jiduc, S.G., Immerzeel, W.W., 2018a. Brief communication: The  
1581 Khurdopin glacier surge revisited - extreme flow velocities and formation of a dammed lake  
1582 in 2017. *Cryosph.* 12, 95–101. <https://doi.org/10.5194/tc-12-95-2018>
- 1583 Steiner, J.F., Litt, M., Stigter, E.E., Shea, J.M., Bierkens, M.F.P., Immerzeel, W.W., 2018b. The  
1584 Importance of Turbulent Fluxes in the Surface Energy Balance of a Debris-Covered Glacier in  
1585 the Himalayas. *Front. Earth Sci.* 6, 1–18. <https://doi.org/10.3389/feart.2018.00144>
- 1586 Stigter, E.E., Litt, M., Steiner, J.F., Bonekamp, P.N.J., Shea, J.M., Bierkens, M.F.P., Immerzeel, W.W.,  
1587 2018. The Importance of Snow Sublimation on a Himalayan Glacier. *Front. Earth Sci.* 6, 1–16.  
1588 <https://doi.org/10.3389/feart.2018.00108>
- 1589 Stokes, C.R., Popovnin, V., Aleynikov, A., Gurney, S.D., Shahgedanova, M., 2007. Recent glacier  
1590 retreat in the Caucasus Mountains, Russia, and associated increase in supraglacial debris  
1591 cover and supra-/proglacial lake development. *Ann. Glaciol.* 46, 195–203.  
1592 <https://doi.org/10.3189/172756407782871468>
- 1593 Sundal, A.V., Shepherd, A., Nienow, P.W., Hanna, E., Palmer, S., Huybrechts, P., 2009. Evolution of  
1594 supra-glacial lakes across the Greenland Ice Sheet. *Remote Sens. Environ.* 113, 2164–2171.  
1595 <https://doi.org/10.1016/j.rse.2009.05.018>
- 1596 Takeuchi, N., Sakai, A., Kohshima, S., Fujita, K., Nakawo, M., 2012. Variation in Suspended  
1597 Sediment Concentration of Supraglacial Lakes on Debris-covered Area of the Lirung Glacier in  
1598 the Nepal Himalayas. *Glob. Environ. Res.* 16, 95–104.
- 1599 Takeuchi, Y., Kayastha, R.B., Nakawo, M., 2000. Characteristics of ablation and heat balance in  
1600 debris-free and debris-covered areas on Khumbu Glacier, Nepal Himalayas, in the pre-  
1601 monsoon season, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered Glaciers.*  
1602 *International Association of Hydrological Sciences, Oxford*, pp. 53–61.
- 1603 Tangborn, W., Rana, B., 2000. Mass balance and runoff of the partially debris- covered Langtang  
1604 Glacier, Nepal, in: Nakawo, M., Raymond, C.F., Fountain, A. (Eds.), *Debris-Covered Glaciers.*

- 1605 International Association of Hydrological Sciences, Oxford, pp. 99–108.
- 1606 Thakuri, S., Salerno, F., Bolch, T., Guyennon, N., Tartari, G., 2016. Factors controlling the  
1607 accelerated expansion of Imja Lake, Mount Everest region, Nepal. *Ann. Glaciol.* 57, 245–257.  
1608 <https://doi.org/10.3189/2016AoG71A063>
- 1609 Thakuri, S., Salerno, F., Smiraglia, C., Bolch, T., D’Agata, C., Viviano, G., Tartari, G., 2014. Tracing  
1610 glacier changes since the 1960s on the south slope of Mt. Everest (central Southern Himalaya)  
1611 using optical satellite imagery. *Cryosph.* 8, 1297–1315. [https://doi.org/10.5194/tc-8-1297-](https://doi.org/10.5194/tc-8-1297-2014)  
1612 2014
- 1613 Thayyen, R.J., Gergan, J.T., 2010. Role of glaciers in watershed hydrology : a preliminary study of a  
1614 “Himalayan catchment.” *Cryosph.* 4, 115–128. <https://doi.org/10.5194/tcd-3-443-2009>
- 1615 Thayyen, R.J., Gergan, J.T., Dobhal, D.P., 2005. Monsoonal control on glacier discharge and  
1616 hydrograph characteristics, a case study of Dokriani Glacier, Garhwal Himalaya, India. *J.*  
1617 *Hydrol.* 306, 37–49. <https://doi.org/10.1016/j.jhydrol.2004.08.034>
- 1618 Thompson, S.S., Benn, D.I., Dennis, K., Luckman, A., 2012. A rapidly growing moraine-dammed  
1619 glacial lake on Ngozumpa Glacier, Nepal. *Geomorphology* 145–146, 1–11.  
1620 <https://doi.org/10.1016/j.geomorph.2011.08.015>
- 1621 Thompson, S.S., Benn, D.I., Mertes, J.R., Luckman, A., 2016. Stagnation and mass loss on a  
1622 Himalayan debris-covered glacier: Processes, patterns and rates. *J. Glaciol.* 62, 467–485.  
1623 <https://doi.org/10.1017/jog.2016.37>
- 1624 Tranter, M., Brown, G.H., Raiswell, R., Sharp, M.J., Gurnell, A., 1993. A conceptual model of solute  
1625 acquisition by Alpine glacial meltwaters. *J. Glaciol.* 39, 573–581.  
1626 <https://doi.org/10.3198/1993JoG39-133-573-581>
- 1627 Tranter, M., Sharp, M.J., Lamb, H.R., Brown, G.H., Hubbard, B., Willis, I.C., 2002. Geochemical  
1628 weathering at the bed of Haut glacier d’Arolla, Switzerland - a new model. *Hydrol. Process.*  
1629 16, 959–993. <https://doi.org/10.1002/hyp.309>
- 1630 Tsutaki, S., Fujita, K., Nuimura, T., Sakai, A., Sugiyama, S., Komori, J., 2019. Contrasting thinning  
1631 patterns between lake- and land-terminating glaciers in the Bhutanese Himalaya. *Cryosph.*  
1632 13, 2733–2750.
- 1633 United Nations, 2015. Transforming our world: the 2030 agenda for sustainable development,  
1634 *A/RES/70/1*. <https://doi.org/10.1201/b20466-7>
- 1635 van Woerkom, T., Steiner, J.F., Kraaijenbrink, P.D.A., Miles, E.S., Immerzeel, W.W., 2019. Sediment  
1636 supply from lateral moraines to a debris-covered glacier in the Himalaya. *Earth Surf. Dyn.* 7,  
1637 411–427. <https://doi.org/10.5194/esurf-7-411-2019>
- 1638 Vaughan, D.G., Comiso, J.C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray, T., Paul,  
1639 F., Ren, J., Rignot, E., Solomina, O., Steffen, K., Zhang, T., 2013. Observations: Cryosphere, in:  
1640 Stocker, T.F., D. Qin, G.-K., Plattner, M., Tignor, S., Allen, K., Boschung, J., A.Nauels, Xia, Y.,  
1641 Bex, V., Midgley, P.M. (Eds.), *Climate Change 2013: The Physical Science Basis. Contribution*  
1642 *of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
1643 *Change.* Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp.  
1644 317–382.
- 1645 Viste, E., Sorteberg, A., 2015. Snowfall in the Himalayas: an uncertain future from a little-known  
1646 past. *Cryosph.* 9, 1147–1167. <https://doi.org/10.5194/tc-9-1147-2015>

- 1647 Wang, X., Ding, Y., Liu, S., Jiang, L., Wu, K., Jiang, Z., Guo, W., 2013. Changes of glacial lakes and  
1648 implications in Tian Shan, central Asia, based on remote sensing data from 1990 to 2010.  
1649 *Environ. Res. Lett.* 8, 044052. <https://doi.org/10.1088/1748-9326/8/4/044052>
- 1650 Wang, X., Liu, S., Han, H., Wang, J., Liu, Q., 2012. Thermal regime of a supraglacial lake on the  
1651 debris-covered Koxkar Glacier, southwest Tianshan, China. *Environ. Earth Sci.* 67, 175–183.  
1652 <https://doi.org/10.1007/s12665-011-1490-1>
- 1653 Watanabe, T., Lamsal, D., Ives, J.D., 2009. Evaluating the growth characteristics of a glacial lake  
1654 and its degree of danger of outburst flooding: Imja Glacier, Khumbu Himal, Nepal. *Nor. Geogr.*  
1655 *Tidsskr. - Nor. J. Geogr.* 63, 255–267. <https://doi.org/10.1080/00291950903368367>
- 1656 Watson, C.S., Quincey, D.J., Carrivick, J.L., Smith, M.W., 2017. Ice cliff dynamics in the Everest  
1657 region of the Central Himalaya. *Geomorphology* 278, 238–251.  
1658 <https://doi.org/10.1016/j.gloplacha.2016.04.008>
- 1659 Watson, C.S., Quincey, D.J., Carrivick, J.L., Smith, M.W., 2016. The dynamics of supraglacial ponds  
1660 in the Everest region, central Himalaya. *Glob. Planet. Change* 142, 14–27.  
1661 <https://doi.org/10.1016/j.gloplacha.2016.04.008>
- 1662 Watson, C.S., Quincey, D.J., Carrivick, J.L., Smith, M.W., Rowan, A.V., Richardson, R., 2018.  
1663 Heterogeneous water storage and thermal regime of supraglacial ponds on debris-covered  
1664 glaciers. *Earth Surf. Process. Landforms* 43, 229–241. <https://doi.org/10.1002/esp.4236>
- 1665 Wessels, R.L., Kargel, J.S., Kieffer, H.H., 2002. ASTER measurement of supraglacial lakes in the  
1666 Mount Everest region of the Himalaya. *Ann. Glaciol.* 34, 399–408.  
1667 <https://doi.org/10.3189/172756402781817545>
- 1668 Wilson, A.M., Williams, M.W., Kayastha, R.B., Racoviteanu, A., 2016. Use of a hydrologic mixing  
1669 model to examine the roles of meltwater, precipitation and groundwater in the Langtang  
1670 River basin, Nepal. *Ann. Glaciol.* 57, 155–168. <https://doi.org/10.3189/2016AoG71A067>
- 1671 Winiger, M., Gumpert, M., Yamout, H., 2005. Karakorum-Hindukush-western Himalaya: Assessing  
1672 high-altitude water resources. *Hydrol. Process.* 19, 2329–2338.  
1673 <https://doi.org/10.1002/hyp.5887>
- 1674 Zeng, C., Liu, Z., Yang, J., Yang, R., 2015. A groundwater conceptual model and karst-related carbon  
1675 sink for a glacierized alpine karst aquifer, Southwestern China. *J. Hydrol.* 529, 120–133.  
1676 <https://doi.org/10.1016/j.jhydrol.2015.07.027>
- 1677 Zhang, Y., Liu, S., Ding, Y., 2007. Glacier meltwater and runoff modelling, Keqicar Baqi glacier,  
1678 southwestern Tien Shan, China. *J. Glaciol.* 53, 91–98.  
1679 <https://doi.org/10.3189/172756507781833956>
- 1680 Zhao, L., Ding, R., Moore, J.C., 2014. Glacier volume and area change by 2050 in high mountain  
1681 Asia. *Glob. Planet. Change* 122, 197–207. <https://doi.org/10.1016/j.gloplacha.2014.08.006>
- 1682