


# Journal Pre-proof



Rapid Systematic Review: The Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in the Context of COVID-19

Maria Elizabeth Loades, DClinPsy, Eleanor Chatburn, MA, Nina Higson-Sweeney, BSc, Shirley Reynolds, PhD, Roz Shafran, PhD, Amberly Brigden, MSc, Catherine Linney, MA, Megan Niamh McManus, BSc, candidate, Catherine Borwick, MSc, Esther Crawley, PhD

PII: S0890-8567(20)30337-3

DOI: <https://doi.org/10.1016/j.jaac.2020.05.009>

Reference: JAAC 2970

To appear in: *Journal of the American Academy of Child & Adolescent Psychiatry*

Received Date: 30 April 2020

Revised Date: 27 May 2020

Accepted Date: 28 May 2020

Please cite this article as: Loades ME, Chatburn E, Higson-Sweeney N, Reynolds S, Shafran R, Brigden A, Linney C, McManus MN, Borwick C, Crawley E, Rapid Systematic Review: The Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in the Context of COVID-19, *Journal of the American Academy of Child & Adolescent Psychiatry* (2020), doi: <https://doi.org/10.1016/j.jaac.2020.05.009>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc. on behalf of the American Academy of Child and Adolescent Psychiatry.

Rapid Systematic Review: The Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in the Context of COVID-19  
RH = Loneliness Mental Health Rapid Review

Maria Elizabeth Loades, DCLinPsy, Eleanor Chatburn, MA, Nina Higson-Sweeney, BSc, Shirley Reynolds, PhD, Roz Shafran, PhD, Amberly Brigden, MSc, Catherine Linney, MA, Megan Niamh McManus, BSc candidate, Catherine Borwick, MSc, Esther Crawley, PhD

Editorial  
Clinical Guidance  
Supplemental Material

Accepted May 28, 2020

Dr. Loades and Mss. Chatburn, Higson-Sweeney, and Borwick are with the University of Bath, Claverton Down, UK. Prof Reynolds is with the School of Psychology and Clinical Language Sciences, University of Reading, UK. Prof Sharan is with UCL Great Ormond Street Institute of Child Health, London, UK. Mss. Brigden and Linney are PhD students at the University of Bristol, UK. Prof Crawley is with the University of Bristol, UK. Ms. McManus is with the University of Edinburgh, UK.

The authors have reported no funding for this work. All research at Great Ormond Street Hospital NHS Foundation Trust and UCL Great Ormond Street Institute of Child Health is made possible by the NIHR Great Ormond Street Hospital Biomedical Research Centre. This report is independent research. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

*Conceptualization:* Loades, Chatburn, Reynolds, Shafran, Borwick, Crawley

*Data curation:* Loades, Chatburn, Higson-Sweeney, Brigden, Linney, McManus

*Formal analysis:* Loades, Chatburn, Reynolds

*Methodology:* Loades, Chatburn, Reynolds

*Project administration:* Loades, Chatburn, Higson-Sweeney, Borwick

*Supervision:* Reynolds, Shafran, Crawley

*Writing – original draft:* Loades

*Writing – review and editing:* Loades, Chatburn, Higson-Sweeney, Reynolds, Shafran, Brigden, Linney, McManus, Borwick, Crawley

ORCID:

Maria Elizabeth Loades, DCLinPsy: 0000-0002-0839-3190

Eleanor Chatburn, MA: 0000-0002-6745-6737

Nina Higson-Sweeney, BSc: 0000-0002-6926-0463

Shirley Reynolds, PhD: 0000-0001-9975-2023

Roz Shafran, PhD: 0000-0003-2729-4961

Amberly Brigden, MSc: 0000-0002-7958-7881

Catherine Linney, MA: 0000-0002-3873-3686

Megan Niamh McManus, BSc candidate: 0000-0002-7206-3444

Catherine Borwick, MSc: 0000-0002-5423-7279

Esther Crawley, PhD: 0000-0002-2521-0747

The authors very much appreciate the valuable comments made by the anonymous peer reviewers.

Disclosure: Dr. Loades has received funding from the National Institute for Health Research (NIHR Doctoral Research Fellowship, DRF-2016-09-021). Ms. Brigden has received funding from the National Institute for Health Research (NIHR Doctoral Research Fellowship, DRF-2017-10-169). Drs. Reynolds, Shafran, Crawley and Mss. Chatburn, Higson-Sweeney, Linney, McManus, and Borwick have reported no biomedical financial interests or potential conflicts of interest.

Correspondence to Maria Loades, Department of Psychology, University of Bath, Bath, BA2 7AY, England; e-mail: [m.e.loades@bath.ac.uk](mailto:m.e.loades@bath.ac.uk).

**Abstract**

**Objective:** Disease containment of COVID-19 has necessitated widespread social isolation. We aimed to establish what is known about how loneliness and disease containment measures impact on the mental health in children and adolescents.

**Method:** For this rapid review, we searched MEDLINE, PSYCHINFO, and Web of Science for articles published between 01/01/1946 and 03/29/2020. 20% of articles were double screened using pre-defined criteria and 20% of data was double extracted for quality assurance.

**Results:** 83 articles (80 studies) met inclusion criteria. Of these, 63 studies reported on the impact of social isolation and loneliness on the mental health of previously healthy children and adolescents (n=51,576; mean age 15.3) 61 studies were observational; 18 were longitudinal and 43 cross sectional studies assessing self-reported loneliness in healthy children and adolescents. One of these studies was a retrospective investigation after a pandemic. Two studies evaluated interventions. Studies had a high risk of bias although longitudinal studies were of better methodological quality. Social isolation and loneliness increased the risk of depression, and possibly anxiety at the time loneliness was measured and between 0.25 to 9 years later. Duration of loneliness was more strongly correlated with mental health symptoms than intensity of loneliness.

**Conclusion:** Children and adolescents are probably more likely to experience high rates of depression and probably anxiety during and after enforced isolation ends. This may increase as enforced isolation continues. Clinical services should offer preventative support and early intervention where possible and be prepared for an increase in mental health problems.

**Key words:** loneliness, pandemic, COVID-19, disease containment, mental health

## Introduction

The COVID-19 pandemic has resulted in governments implementing disease containment measures such as school closures, social distancing and home quarantine. Children and adolescents are experiencing a prolonged state of physical isolation from their peers, teachers, extended family and community networks. Quarantine in adults generally has negative psychological effects including confusion, anger, and post-traumatic distress.<sup>1,2</sup> Duration of quarantine, infection fears, boredom, frustration, lack of necessary supplies, lack of information, financial loss, and stigma appear to increase the risk of negative psychological outcomes.<sup>1</sup> Social distancing and school closures may therefore increase mental health problems in children and adolescents, already at higher risk of developing mental health problems compared to adults<sup>3</sup> at a time when they are also experiencing anxiety over a health threat and threats to family employment/income.

Social distancing and school closures are likely to result in increased loneliness in children and adolescents whose usual social contacts are curtailed by the disease containment measures. Loneliness is the painful emotional experience of a discrepancy between actual and desired social contact<sup>4</sup> Although social isolation is not necessarily synonymous with loneliness, early indications in the COVID-19 context indicate that more than one third of adolescents report high levels of loneliness<sup>5,6</sup> and almost half of 18-24-year olds are lonely during lockdown.<sup>7</sup> There are well established links between loneliness and mental health.<sup>8</sup> The purpose of this review was to establish what is known about the relationship between loneliness and mental health problems in healthy children and adolescents and to establish whether disease containment measures including quarantine and social isolation are predictive of future mental health problems. We included cross sectional, observational, retrospective and case control studies if studies included mainly children and adolescents, who had experienced loneliness or had used validated measures of social isolation and mental health problems. To capture the possible effects of social isolation and the expected

mediator (loneliness) on mental health problems, we included search terms to capture these two areas.

## **Method**

We conducted a rapid review to provide timely evidence synthesis to inform urgent healthcare policy decision-making.<sup>9</sup> A rapid review adheres to the essential principles of systematic reviews, including scientific rigour, transparency, and reproducibility.<sup>9,10</sup> It uses “abbreviated” systemic review methodology including: limiting search criteria, faster data extraction, and using narrative synthesis methods.<sup>11,12</sup>

*Search strategy and selection criteria [see Table S1, Table S2, Table S3 for full search strategy]*

We searched MEDLINE, PSYCHINFO, Web of Science and the Cochrane Library. Our search terms were informed by recent rapid reviews in the COVID-19 context<sup>1</sup> and included definitions of loneliness and social isolation to capture the impact of social distancing and school closures. Terms captured ‘children’ or ‘adolescents’ AND ‘quarantine’ or ‘social isolation’ or ‘loneliness’ AND ‘mental health’ with a focus on the most common mental health problems in this age group: depression and anxiety.

Peer reviewed studies were selected if they were published (1946 to 03/29/2020); reporting primary research; included predominantly children/adolescents (mean age < 21)<sup>13</sup>; published in English (web of science only); participants had experienced either social isolation or loneliness; valid assessment of depression, anxiety, trauma, OCD, mental health, or mental wellbeing.

*Study selection and data collection*

We checked 20% of all study eligibility results (both included and excluded) to ensure adherence to the eligibility criteria. Data were extracted into a purpose-designed database: A random 20% of the data was double entered to ensure accuracy.

A truncated quality assessment was conducted by one author (SR) using criteria adapted from the NIH<sup>14</sup> (see table 1).

[insert table 1 here]

### *Data Synthesis*

We conducted a narrative synthesis within the following categories: (1) the impact of loneliness on mental health in healthy populations (further divided into cross-sectional and longitudinal evidence), (2) pandemic-specific findings, and (3) intervention studies.

### **Results**

We located 4531 articles (see Figure 1) of which 83 articles (80 studies) met the inclusion criteria. Of these, 18 articles (17 studies) reported on the impact of loneliness in those with a variety of health conditions including mental health problems (12 studies), physical health problems (1 study) and neurodevelopmental conditions (4 studies). The remaining 65 articles reported on 63 studies which examined the impact of loneliness or disease containment measures on healthy children and adolescents. For the purposes of this rapid review, we will focus our analyses on these 63 studies.

[insert figure 1 here]

Figure 1. Flow diagram showing search results.<sup>15</sup>

The 63 studies were mainly from the USA, China, Europe and Australia. Included studies were also conducted in India, Malaysia, Korea, Thailand, Israel, Iran, and Russia. 61 studies were observational and 2 studies reported on interventions. Of the 61 observational studies, 43 studies were cross-sectional only, 6 longitudinal only and 12 reported both cross sectional and longitudinal findings. 1 study was a retrospective study after a pandemic. In cross sectional studies, likely confounders (e.g. adversity, SES) were rarely controlled, meaning that the association between loneliness and mental health outcomes in these studies is very likely to be inflated<sup>16</sup>. Four longitudinal studies used multi-informant approaches including self-report and parent and/or teacher report to assess mental health outcomes. Importantly, they typically assessed and controlled for confounds and could assess the most plausible direction of causality between loneliness/social isolation and mental health.

#### *The impact of loneliness on mental health*

Tables 2 and 3 describe the 60 studies which examined the impact of loneliness on mental health. 53 studies stated that they measured the impact of loneliness on mental health. 7 studies stated that they measured the impact of social isolation<sup>17-23</sup> on mental health, but the social isolation measures used were either subscales or questions from loneliness scales, or strongly overlapped with the construct of loneliness. Therefore, we have considered them together with studies that measured loneliness. Participants were mainly school or university students or taking part in longitudinal cohort studies.

Forty-five studies examined the cross-sectional relationship between depressive symptoms and loneliness and/or social isolation.<sup>17,20,21,23-66</sup> The majority were conducted in adolescent (N = 23) and young adult (N = 16) samples, although six studies included children under the age of 10. Most reported moderate to large correlations ( $0.12 \leq r \leq 0.81$ ) and most included a measure of depressive symptoms. Two studies reported odds ratios, with those who were lonely 5.8<sup>45</sup> to 40 times<sup>49</sup> more likely to score above clinical cut-offs for depression. The associations were stronger in older



participants<sup>37</sup> and in female participants.<sup>46</sup> However, the strength and direction of the associations did not differ by age of the sample. Fewer studies (N = 23) examined symptoms of anxiety. Those that did found small to moderate associations between anxiety and loneliness/social isolation ( $0.18 \leq r \leq 0.54$ ). The duration of loneliness was more strongly associated with anxiety than intensity of loneliness.<sup>42,67</sup> Social anxiety was moderately to strongly associated with loneliness/social isolation ( $0.33 \leq r \leq 0.72$ ) and there were moderate associations between generalized anxiety and loneliness/social isolation ( $r = 0.37, 0.40$ ).<sup>21,34</sup> One study found a small association between panic and loneliness ( $r = 0.13$ ).<sup>61,62</sup> In the single study which reported odds ratios, being lonely was associated with increased odds of being anxious by 1.63 to 5.49 times.<sup>49</sup> Positive associations were also reported between social isolation/loneliness and suicidal ideation,<sup>24,27,28</sup> self-harm,<sup>24</sup> and eating disorder risk behaviour.<sup>24</sup> Negative associations were reported between social isolation/loneliness and wellbeing<sup>68,69</sup> and mental health.<sup>22</sup>

Eighteen studies followed participants over time (see table 3).<sup>17-19,55-58,60-63,65,70-75</sup> Several of these were conducted in childhood (N = 6), or adolescence (N = 8), although three were in university students. Most (N = 12) had only one follow up time point usually between 1 and 3 years.

12 of the 15 studies found that loneliness is associated with depression and explained a significant amount of the variance in severity of depression symptoms several months to several years later.<sup>55,57-63,71,73,74</sup> Two studies found that loneliness in childhood at age 5 was not associated with depression several years later<sup>59,60</sup> although other studies which assessed loneliness during childhood found evidence that it is associated with subsequent depression<sup>55,72</sup> One large study of adolescents (n=3088) found that loneliness was not associated with depression one year later.<sup>56</sup> There were mixed findings in another large study of adolescents (n = 541) which found a significant association between loneliness and subsequent depression, although this did not hold in a cross-lagged model<sup>17</sup> suggesting a possible bidirectional relationships between the variables. A study of university

students found evidence of a gender difference, with loneliness being associated with later depression in female participants but not in male participants.<sup>18</sup> In a large longitudinal cohort of vulnerable young people, aged 11 to 17, after controlling for caregiver neglect and other relevant covariates, a substantial increase in self-reported peer isolation (1 S.D.) was associated with an increase in depression symptoms (0.49 S.D.).<sup>71</sup> Duration of peer loneliness rather than the intensity of peer loneliness is associated with depression 8 years later (i.e. from age 5 to age 13); in contrast family related loneliness was not independently associated with subsequent depression.<sup>59</sup>

Three of the four studies which examined the longitudinal effect of loneliness on anxiety found that loneliness was associated with later anxiety.<sup>56,64,75</sup> Two of these studies assessed social anxiety, and one measured anxiety as a broad construct. One study did not find that loneliness/social isolation at age 5 was associated with anxiety at age 12.<sup>19</sup> One study of young adolescents found differences by gender, with loneliness being associated with later social anxiety in male participants but not female participants.<sup>75</sup> None of these studies measured loneliness during childhood.

Other mental health outcomes reported over time included internalizing symptoms which were associated with prior loneliness in primary school age children,<sup>72</sup> and suicidal ideation during adolescence, which was not associated with prior loneliness during childhood.<sup>60</sup>

#### *The impact of social isolation in an infectious disease context*

One study<sup>76</sup> reported on mental health and social isolation in the context of different infections including H1N1, SARS, and avian flu (see table 2). This retrospective study included 398 parents of exposed children from the USA, Canada and Mexico, of whom 20.9% experienced social isolation and a further 3.8% had been quarantined. Parents of children reported on their child's experience of trauma and on their current mental health. One third of parents whose children had been subject to disease containment measures said their child had needed mental health service input because of

their pandemic related experiences. The most reported diagnoses were acute stress disorder (16.7%), adjustment disorder (16.7%), grief (16.7%), and PTSD (6.2%). Two different parent-reported measures of PTSD symptoms found that those children exposed to disease containment measures scored significantly higher for PTSD symptoms post-pandemic. On the PTSD Checklist Civilian Version, 28% of children who had experienced isolation/quarantine scored about the cut-off for PTSD, compared to 5.8% of those who had not experienced isolation/quarantine. Similarly, on the UCLA PTSD Reaction Index, 30% of children who experienced isolation/quarantine scored about the cut-off for PTSD, compared to 1.1% of those who had not experienced isolation/quarantine (effect size: Cramer's  $V = 0.449$ ). Mean scores were 4 times higher in the isolated/quarantined group than in those who had not been isolated/quarantined. The most common trauma symptoms in the quarantined/isolated group were avoidance/numbing (57.8%), re-experiencing (57.8%), and arousal (62.5%).

### *Interventions*

Two randomised control trials measured loneliness and mental health outcomes following an intervention aimed at the general population (peer mentoring<sup>77</sup> and classroom based,<sup>78</sup> see table 4). In both instances the comparator was no intervention/with follow-up and education as usual. A relatively intensive peer mentor program, with an adult mentor, 4-6 hours per month for 4 months on average, reduced loneliness and mental health problems (small to medium effects) for victims of bullying and victimization. However, a brief (two session) universal classroom-based program delivered in schools including psychosocial support through peer mentors and a staff mental health support team did not reduce loneliness. Neither intervention specifically addressed mental health problems which had developed in the context of loneliness; therefore we are unable to answer our second review question which was what interventions are effective for those who have developed mental health problems as a result of social isolation or loneliness.

[insert tables 2 – 4 here]

## Discussion

This rapid systematic review of 63 studies of 51, 576 participants found a clear association between loneliness and mental health problems in children and adolescents. Loneliness was associated with future mental health problems up to 9 years later. The strongest association was with depression. These findings were consistent across studies of children, adolescents, and young adults. There may also be gender differences with some research indicating that loneliness was more strongly associated with elevated depression symptoms in girls and with elevated social anxiety in boys.<sup>18,75</sup> The length of loneliness appears to be a predictor of future mental health problems<sup>59</sup>. This is of particular relevance in the COVID-19 context as politicians in different countries consider the length of time that schools should remain closed, and the implementation of social distancing within schools.

Furthermore, in the one study that examined mental health problems after enforced isolation and quarantine in previous pandemics, children who had experienced enforced isolation or quarantine were five times more likely to require mental health service input and experienced higher levels of post-traumatic stress. This suggests that the current social distancing measures enforced on children because of COVID-19 could lead to an increase in mental health problems, as well as possible post-traumatic stress. These results are consistent with preliminary, unpublished data emerging from China during the COVID-19 pandemic where children aged 3 to 18 are commonly displaying behavioural manifestations of anxiety including: clinginess, distraction, fear of asking questions about the pandemic, and irritability<sup>79</sup> Furthermore, a large survey of young adult students in China has reported that around one in four are experiencing at least mild anxiety symptoms<sup>80</sup> In the UK, early results from the Co-SPACE (COVID-19 Supporting Parents, Adolescents and Children in Epidemics) online survey of over 1500 parents suggest high levels of COVID-19-related worries and

fears, with younger children (age four to 10) significantly more worried than older children (age 11 to 16).<sup>81,82</sup>

In addition to the more direct effects of enforced isolation and quarantine, loneliness as an unintended consequence of disease containment measures seems to be particularly problematic for young people<sup>5,7</sup>. This may be because of the particular importance of the peer group for identity and support during this developmental stage.<sup>83,84</sup> This propensity to experience loneliness may make young people particularly vulnerable to loneliness in the COVID-19 context, which, based on our findings, may further exacerbate the mental health impacts of the disease containment measures. More studies have examined the relationship between loneliness and depression than loneliness and anxiety. Losing links to other people and feeling excluded can result in an affective response of depression.<sup>85</sup> Social anxiety was more strongly associated with loneliness than other anxiety subtypes. This may be because social anxiety is triggered by a perceived threat to social relationships or status.<sup>86</sup>

It is difficult to predict the effect COVID-19 will have on the mental health of children and young people. The subjective social isolation experienced by participants did not mirror the current features of social isolation experienced by many children and adolescents worldwide. Social isolation was not enforced upon the participants, nor was social isolation almost ubiquitous across their peer groups and across the communities in which they live. As loneliness involves social comparison,<sup>87</sup> it is possible that the shared experience of social isolation imposed by disease containment measures may mitigate the negative effects. The studies were also not in the context of an uncertain but dangerous threat to health. These features limit the extent to which we can extrapolate from existing evidence to the current context. In order to make evidence based decisions on how to mitigate the impact of a second wave, we need further research on the mental health impacts of social isolation in the disease containment context of a global pandemic. In this context, to more

specifically understand the impacts of loneliness, measures such as the Loneliness and Aloneness Scale for Children and Adolescents (LACA) that assess the duration and the intensity of loneliness, and that separate peer-related loneliness from parent-related loneliness could be elucidating.

This rapid systematic review was conducted rapidly, in 3 weeks, to inform our response to COVID-19. We double screened 20% of all articles and data extracted. In line with Cochrane rapid review guidance,<sup>10</sup> grey literature, and trial registry databases were not searched, hand-search strategies were not employed, and only English language publications were included, meaning that some relevant studies may have been missed. During the rapid data extraction phase, there was no scope to contact authors to request any missing information. The main limitation from this review is the lack of high-quality studies investigating mental health problems after enforced isolation. All but one study investigated social isolation that was not enforced on young people and was not common across a peer group. The effect of widespread social distancing could mitigate against the social isolation described with increased use of internet mediated relationships which can be beneficial to adolescents.<sup>88</sup> Most studies were cross-sectional, and therefore the direction of the association cannot be inferred. Few studies used independent (i.e. not self-report) measures of mental health or social isolation/loneliness, increasing the risk of bias. Furthermore, the studies were mainly observational and did not consistently control for potential confounders. The majority of studies focused on depression and anxiety, and other mental health problems are important to measure in future research.

However, we used all available evidence on social isolation and loneliness to inform the likely outcome for healthy children and adolescents subjected to social isolation. The results were consistent across all study methodology for depression, (but less so for anxiety) suggesting these results are reliable. The results are also consistent with one study investigating mental health problems in children<sup>76</sup> after pandemics improving our confidence in the results. However, the post

pandemic study has several limitations in that the sample was self-selecting, and the demographics of the children and the time elapsed since the experience were not reported. There is little evidence pertaining to interventions. We have focused on healthy populations in this review and will report on those with pre-existing conditions including mental health problems elsewhere.

#### *Implications for Policy and Practice*

The review indicates that felt loneliness is associated with adverse mental health in children and adolescents. There is limited evidence that indicates specific interventions to prevent loneliness or to reduce its effects on mental health and well-being. However, there are well-established practical and psychological strategies that may help promote child and adolescent mental health in the context of involuntary social isolation e.g. during the COVID-19 pandemic. Reducing the impact of enforced physical distancing by maintaining the structure, quality, and quantity of social networks, and helping children and adolescents to experience social rewards, feel part of a group, and know that there are others they can look to for support is likely to be important.<sup>8</sup> Finding ways to give children and adolescents a sense of belonging within the family and to feel that they are part of a wider community should be a priority. Therefore, providing accurate information about the relative risks and benefits of social media and networking to parents who overestimate the dangers of allowing their children too much screen time may help young people access the benefits of virtual social contact.

However, simply increasing the frequency of contact may not address young people's subjective experience of loneliness.<sup>20</sup> Helping young people to identify valued alternative activities and build structure and purpose into periods of involuntary social isolation may help to provide a wider range of rewards.<sup>89</sup> Addressing negative thoughts about social encounters (e.g. self-blame, self-devaluation) may also be effective.<sup>34,90</sup> During periods of prolonged social isolation digital technology

that provides evidence-based interventions to help young people to reappraise their thoughts and change their behaviour within the confines of the home setting may be particularly welcome.

Whilst this review did not provide evidence on interventions to improve social isolation or loneliness in healthy children and adolescents, given social distancing, digital interventions may be appropriate. Computerized Cognitive Behaviour Therapy (CBT) based self-help program, BRAVE-TA, was shown to be effective for anxiety following the Christchurch earthquake in New Zealand.<sup>91</sup> Furthermore, computerized CBT, such as MoodGym, SPARX, and 'Think, Feel, Do' generally have small but positive effects on mental health.<sup>92,93</sup> Although mobile applications for mental health have been found to be generally acceptable to children and adolescents, there is a lack of convincing evidence of effectiveness on intended mental health outcomes<sup>94</sup> and few mobile health apps have been thoroughly tested.<sup>93</sup> Self-help interventions including bibliotherapy<sup>95</sup> and computerized therapy<sup>96</sup> have shown a moderate positive effect size when compared to control groups although they are generally less effective than face to face therapies.<sup>97</sup> Importantly, reviews have tended to conclude that effects are better if there is some therapist input<sup>93,97</sup> and if parents are involved especially for younger children.<sup>92,93</sup>

The rapid review suggests that loneliness that may result from disease containment measures in the COVID-19 context could be associated with subsequent mental health problems in young people. Strategies to prevent the development of such problems should be an international priority.



## REFERENCES

1. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395(10227):912-920.
2. Hossain MS, A;Purohit,N;. Mental health outcomes of Quarantine and isolation for infection prevention: A systematic umbrella review of the global evidence. psyarxiv.com/dz5v2/. Published 2020. Accessed April 10 2020.
3. Deighton J, Lereya ST, Casey P, Patalay P, Humphrey N, Wolpert M. Prevalence of mental health problems in schools: poverty and other risk factors among 28 000 adolescents in England. *The British journal of psychiatry : the journal of mental science*. 2019:1-3.
4. Perlman D, Peplau LA. Toward a social psychology of loneliness. *Personal relationships*. 1981;3:31-56.
5. Study OA. Achieving Resilience During COVID-19 Weekly Report 2. Published 2020. Accessed 22/05/2020.
6. Abstracts from the Poster and Oral Presentations from the 18th Neonatal and Paediatric Pharmacists Group, NPPG Annual Conference. *Archives of disease in childhood*. 2013;98 (6).
7. Foundation MH. Coping with Loneliness. Published 2020. Accessed 22/05/2020.
8. Wang J, Lloyd-Evans B, Giacco D, et al. Social isolation in mental health: a conceptual and methodological review. *Soc Psychiatry Psychiatr Epidemiol*. 2017;52(12):1451-1461.
9. World Health Organisation AfHPaSR. *Rapid reviews to strengthen ealth policy and systems: a practical guide*. 2017.
10. Garritty CG, G;Kamel,C;King,V.J;Nussbaumer-Streit,B;Stevens,A;Hamel,C;Affengruber,L;. *Cochrane Rapid Reviews. Interim Guidance from teh Cochrane Rapid Review Methods Group*. 2020.
11. Ganann R, Ciliska D, Thomas H. Expediting systematic reviews: methods and implications of rapid reviews. *Implement Sci*. 2010;5:56.
12. Tricco AC, Antony J, Zarin W, et al. A scoping review of rapid review methods. *BMC Med*. 2015;13:224.
13. Manuell ME, Cukor J. Mother Nature versus human nature: public compliance with evacuation and quarantine. *Disasters*. 2011;35(2):417-442.
14. Health Nio. Study Assessment Tools. Published 2018. Accessed.
15. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
16. Westfall J, Yarkoni T. Statistically Controlling for Confounding Constructs Is Harder than You Think. *PLoS one*. 2016;11(3):e0152719-e0152719.
17. Lasgaard M, Goossens L, Elklit A. Loneliness, depressive symptomatology, and suicide ideation in adolescence: cross-sectional and longitudinal analyses. *J Abnorm Child Psychol*. 2011;39(1):137-150.
18. Liu H, Zhang M, Yang Q, Yu B. Gender differences in the influence of social isolation and loneliness on depressive symptoms in college students: a longitudinal study. *Soc Psychiatry Psychiatr Epidemiol*. 2020;55(2):251-257.
19. Matthews T, Danese A, Wertz J, et al. Social isolation and mental health at primary and secondary school entry: a longitudinal cohort study. *J Am Acad Child Adolesc Psychiatry*. 2015;54(3):225-232.
20. Matthews T, Danese A, Wertz J, et al. Social isolation, loneliness and depression in young adulthood: a behavioural genetic analysis. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51(3):339-348.
21. Richardson C, Oar E, Fardouly J, et al. The Moderating Role of Sleep in the Relationship Between Social Isolation and Internalising Problems in Early Adolescence. *Child Psychiatry Hum Dev*. 2019;50(6):1011-1020.

22. Stacciarini JM, Smith R, Garvan CW, Wiens B, Cottler LB. Rural Latinos' mental wellbeing: a mixed-methods pilot study of family, environment and social isolation factors. *Community Ment Health J.* 2015;51(4):404-413.
23. Yadegarfar MM-B, M.E;Ho,R;. Family Rejection, Social Isolation, and Loneliness as Predictors of Negative Health Outcomes (Depression, Suicidal Ideation, and Sexual Risk Behaviour) Among Thai Male-to-female Transgender Adolescents *Journal of LGBT Youth.* 2014;11(4):347-363.
24. Lasgaard MG, L;Bramsen,R.H;Trillingsgaard,T;Elklit,A;. Different sources of loneliness are associated with different forms of psychopathology in adolescence. *Journal of Research in Personality.* 2011;45(2):233-237.
25. Alpaslan AH, Kocak U, Avci K. Gender-Related Factors for Depressive Symptoms in Turkish Adolescents. *J Child Adolesc Psychiatr Nurs.* 2016;29(1):23-29.
26. Baskin TWW, B.E;Quintana,S.M;Enright,R.D;. Belongingness as a Protective Factor Against Loneliness and Potential Depression in a Multicultural Middle School. *Couns Psychol.* 2010;38(5):626-651.
27. Brage D, Campbell-Grossman C, Dunkel J. Psychological correlates of adolescent depression. *J Child Adolesc Psychiatr Nurs.* 1995;8(4):23-30.
28. Brage D, Meredith W, Woodward J. Correlates of loneliness among midwestern adolescents. *Adolescence.* 1993;28(111):685-693.
29. Chang EC, Wan L, Li P, et al. Loneliness and Suicidal Risk in Young Adults: Does Believing in a Changeable Future Help Minimize Suicidal Risk Among the Lonely? *J Psychol.* 2017;151(5):453-463.
30. Doman LCHL, A;. The relationship between loneliness and psychological well-being among third-year students: a cross-cultural investigation. *International Journal of Culture and Mental Health.* 2012;5(3):153-168.
31. Erdur-Baker OEB, A;. Mediator and Moderator Role of Loneliness in the Relationship between Peer Victimization and Depressive Symptoms *Australian Journal of Guidance and Counselling.* 2011;21(2):175-185.
32. Hudson DB, Elek SM, Campbell-Grossman C. Depression, self-esteem, loneliness, and social support among adolescent mothers participating in the new parents project. *Adolescence.* 2000;35(139):445-453.
33. Hutcherson STE, C.C;. Differentiating parent- and peer-related interpersonal correlates of depressive symptoms and social anxiety in preadolescent girls. *Journal of Social and Personal Relationships.* 2009;26(7):875-897.
34. Jackson J, Cochran SD. Loneliness and psychological distress. *J Psychol.* 1991;125(3):257-262.
35. Kim O. Sex differences in social support, loneliness, and depression among Korean college students. *Psychol Rep.* 2001;88(2):521-526.
36. Koenig LJI, A.M;Schwartz,J.A.J;. Sex differences in adolescent depression and loneliness: why are boys lonelier if girls are more depressed? . *J Res Personal.* 1994;28(1):27-43.
37. Lau S, Chan DW, Lau PS. Facets of loneliness and depression among Chinese children and adolescents. *J Soc Psychol.* 1999;139(6):713-729.
38. Mahon NE, Yarcheski A, Yarcheski TJ. Mental health variables and positive health practices in early adolescents. *Psychol Rep.* 2001;88(3 Pt 2):1023-1030.
39. MajdAra ET, S;Rezaei,A.M;. A Structural Model of Depression Based on Interpersonal Relationships: The Mediating Role of Coping Strategies and Loneliness. *Noro Psikiyatrs Ars.* 2017;54(2):125-130.
40. Markovic A, Bowker JC. Social surrogacy and adjustment: exploring the correlates of having a "social helper" for shy and non-shy young adolescents. *J Genet Psychol.* 2015;176(1-2):110-129.

41. McIntyre JCW, J;Corcoran,R;HarrisonWoods,P;Bentall,R.P;. Academic and non-academic predictors of student psychological distress: the role of social identity and loneliness. *J Ment Health*. 2018;27(3):230-239.
42. Moore D, Schultz NR, Jr. Loneliness at adolescence: Correlates, attributions, and coping. *J Youth Adolesc*. 1983;12(2):95-100.
43. Mounts NSV, D;Anderson,K.L;Boswell,M.K;. Shyness, Sociability, and Parental Support for the College Transition: Relation to Adolescents' Adjustment. *Journal of Youth and Adolescence*. 2006;35(1):68-77.
44. Purwono UF, D.C;. Depression and its relation to loneliness and religiosity in Indonesian Muslim adolescents. *Mental Health, Religion & Culture*. 2016;19(3):218-228.
45. Roberts RE, Chen YW. Depressive symptoms and suicidal ideation among Mexican-origin and Anglo adolescents. *J Am Acad Child Adolesc Psychiatry*. 1995;34(1):81-90.
46. Singhvi MK, S;Sehgal,R;Kumari,N;. Psychological correlates of loneliness among adolescents. *Indian J Psychol Sci*. 2011;2(2):8.
47. Spithoven AW, Lodder GM, Goossens L, et al. Adolescents' Loneliness and Depression Associated with Friendship Experiences and Well-Being: A Person-Centered Approach. *J Youth Adolesc*. 2017;46(2):429-441.
48. Stednitz JN, Epkins CC. Girls' and mothers' social anxiety, social skills, and loneliness: associations after accounting for depressive symptoms. *J Clin Child Adolesc Psychol*. 2006;35(1):148-154.
49. Stickley A, Koyanagi A, Kuposov R, et al. Loneliness and its association with psychological and somatic health problems among Czech, Russian and U.S. adolescents. *BMC Psychiatry*. 2016;16:128.
50. Swami V, Chamorro-Premuzic T, Sinniah D, et al. General health mediates the relationship between loneliness, life satisfaction and depression. A study with Malaysian medical students. *Soc Psychiatry Psychiatr Epidemiol*. 2007;42(2):161-166.
51. Thomas KKB, J.C;. Rejection Sensitivity and Adjustment during Adolescence: Do Friendship Self-Silencing and Parent Support Matter? *Journal of Child and Family Studies*. 2013;24:608-616.
52. Tu YZ, S;. Loneliness and Subjective Well-Being Among Chinese Undergraduates: The Mediating Role of Self-Efficacy. *Soc Indic Res*. 2014;124:963-980.
53. Vanhalst J, Luyckx K, Raes F, Goossens L. Loneliness and depressive symptoms: the mediating and moderating role of uncontrollable ruminative thoughts. *J Psychol*. 2012;146(1-2):259-276.
54. Xu J, Chen P. The rural children's loneliness and depression in Henan, China: the mediation effect of self-concept. *Soc Psychiatry Psychiatr Epidemiol*. 2019;54(9):1101-1109.
55. Boivin MH, S;Bukowski,WM. THE ROLES OF SOCIAL WITHDRAWAL, PEER REJECTION, AND VICTIMIZATION BY PEERS IN PREDICTING LONELINESS AND DEPRESSED MOOD IN CHILDHOOD. *Development and Psychopathology*. 1995;7(4):765-785.
56. Danneel S, Nelemans S, Spithoven A, et al. Internalizing Problems in Adolescence: Linking Loneliness, Social Anxiety Symptoms, and Depressive Symptoms Over Time. *J Abnorm Child Psychol*. 2019;47(10):1691-1705.
57. Ladd GW, Ettekal I. Peer-related loneliness across early to late adolescence: normative trends, intra-individual trajectories, and links with depressive symptoms. *J Adolesc*. 2013;36(6):1269-1282.
58. Lapierre MA, Zhao P, Custer BE. Short-Term Longitudinal Relationships Between Smartphone Use/Dependency and Psychological Well-Being Among Late Adolescents. *J Adolesc Health*. 2019;65(5):607-612.
59. Qualter P, Brown SL, Munn P, Rotenberg KJ. Childhood loneliness as a predictor of adolescent depressive symptoms: an 8-year longitudinal study. *Eur Child Adolesc Psychiatry*. 2010;19(6):493-501.

60. Schinka KC, van Dulmen MH, Mata AD, Bossarte R, Swahn M. Psychosocial predictors and outcomes of loneliness trajectories from childhood to early adolescence. *J Adolesc.* 2013;36(6):1251-1260.
61. Vanhalst J, Goossens L, Luyckx K, Scholte RH, Engels RC. The development of loneliness from mid- to late adolescence: trajectory classes, personality traits, and psychosocial functioning. *J Adolesc.* 2013;36(6):1305-1312.
62. Vanhalst J, Klimstra TA, Luyckx K, Scholte RH, Engels RC, Goossens L. The interplay of loneliness and depressive symptoms across adolescence: exploring the role of personality traits. *J Youth Adolesc.* 2012;41(6):776-787.
63. Vanhalst JL, K;Teppers,E;Goossens,L;. Disentangling the Longitudinal Relation Between Loneliness and Depressive Symptoms: Prospective Effects and the Intervening Role of Coping. *Journal of Social and Clinical Psychology.* 2012;31(8):810-834.
64. Wang YT, L;Guo,L;Huebner,S.E;. Family dysfunction and Adolescents' anxiety and depression: A multiple mediation model. *Journal of Applied Developmental Psychology.* 2020;66.
65. Zhou J, Li X, Tian L, Huebner ES. Longitudinal association between low self-esteem and depression in early adolescents: The role of rejection sensitivity and loneliness. *Psychol Psychother.* 2020;93(1):54-71.
66. Uba IY, S.N;Juhari,R;Talib,M.A;. Does self-esteem mediate the relationship between loneliness and depression among Malaysian teenagers? *Pertanika J Soc Sci Humanit.* 2012.
67. Ginter EJ, Lufi D, Dwinell PL. Loneliness, perceived social support, and anxiety among Israeli adolescents. *Psychol Rep.* 1996;79(1):335-341.
68. Heredia DS, M.L;osner,C.M;He,X;Castillo,L.G;Ojeda,L;. The Influence of Loneliness and Interpersonal Relations on Latina/o Middle School Students' Wellbeing. *Journal of Latinos and Education.* 2017;16(4):338-348.
69. Houghton SH, J;Carroll,A;Wood,L;Baffour,B;. It Hurts To Be Lonely! Loneliness and Positive Mental Wellbeing in Australian Rural and Urban Adolescents. *Journal of Psychologists and Counsellors in Schools.* 2016;26(1):52-67.
70. Wang G, Zhang Y, Zhao J, Zhang J, Jiang F. Mitigate the effects of home confinement on children during the COVID-19 outbreak. *Lancet.* 2020;395(10228):945-947.
71. Christ SL, Kwak YY, Lu T. The joint impact of parental psychological neglect and peer isolation on adolescents' depression. *Child Abuse Negl.* 2017;69:151-162.
72. Fontaine RG, Yang C, Burks VS, et al. Loneliness as a partial mediator of the relation between low social preference in childhood and anxious/depressed symptoms in adolescence. *Dev Psychopathol.* 2009;21(2):479-491.
73. Jones AC, Schinka KC, van Dulmen MH, Bossarte RM, Swahn MH. Changes in loneliness during middle childhood predict risk for adolescent suicidality indirectly through mental health problems. *J Clin Child Adolesc Psychol.* 2011;40(6):818-824.
74. Lalayants MP, J.D;. Loneliness and Depression or Depression-Related Factors Among Child Welfare-Involved Adolescent Females. *Child and Adolescent Social Work Journal volume.* 2015;32(2):167-176.
75. Mak HW, Fosco GM, Feinberg ME. The Role of Family for Youth Friendships: Examining a Social Anxiety Mechanism. *J Youth Adolesc.* 2018;47(2):306-320.
76. Sprang G, Silman M. Posttraumatic stress disorder in parents and youth after health-related disasters. *Disaster Med Public Health Prep.* 2013;7(1):105-110.
77. King CA, Gipson PY, Arango A, et al. LET's CONNECT community mentorship program for youths with peer social problems: Preliminary findings from a randomized effectiveness trial. *J Community Psychol.* 2018;46(7):885-902.
78. Larsen TBU, H;Tobro,M;Ardal,E;Waldahl,R.H;Djuedal,I;Holsen,I;. Promoting Mental Health and Preventing Loneliness in Upper Secondary School in Norway: Effects of a Randomized Controlled Trial. *Scandinavian Journal of Educational Research.* 2019:1-14.

79. Jiao WYW, L.N.;Liu,J;Fang,S.F;Jiao,F.Y;Pettoello-Mantovani,M;Somekh,E;. Behavioral and Emotional Disorders in Children during the COVID-19 Epidemic. *The journal of Pediatrics* 2020.
80. Cao W, Fang Z, Hou G, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry research*. 2020:112934.
81. Oxford Uo. COVID-19's impact on youth mental health the focus of new research <http://www.ox.ac.uk/news/2020-03-30-covid-19s-impact-youth-mental-health-focus-new-research>. Published 2020. Accessed April 13, 2020.
82. Weale S. Fifth of primary children afraid to leave house over Covid-19, survey finds. <https://www.theguardian.com/world/2020/apr/10/fifth-of-primary-children-afraid-to-leave-house-because-of-covid-19-survey-finds>. Published 2020. Accessed April 13, 2020.
83. Bradford Brown BE, S.A; Petrie,S. The importance of peer group ('crowd') affiliation in adolescence. *J Adolescence*. 1986;9:73-96.
84. Meeus W, Dekoviic M. Identity development, parental and peer support in adolescence: results of a national Dutch survey. *Adolescence*. 1995;30(120):931-944.
85. Beck ATC, D.A. Anxiety and depression: An information processing perspective. *anxiety Research*. 1988;88(1).
86. Leary MR. Responses to social exclusion: social anxiety, jealousy, loneliness, depression and low self-esteem. . *Journal of Social and Clinical Psychology*. 1990;9(2):8.
87. Wiseman H. On failed intersubjectivity: Recollections of loneliness experiences in offspring of Holocaust survivors. *American Journal of Orthopsychiatry*. 2008;78(3):350.
88. Przybylski AK, Orben A, Weinstein N. How Much Is Too Much? Examining the Relationship Between Digital Screen Engagement and Psychosocial Functioning in a Confirmatory Cohort Study. *J Am Acad Child Adolesc Psychiatry*. 2019.
89. Pass L, Lejuez CW, Reynolds S. Brief Behavioural Activation (Brief BA) for Adolescent Depression: A Pilot Study. *Behav Cogn Psychother*. 2018;46(2):182-194.
90. Masi CMC, H.Y;Hawkey,L.C;Capioppo,J.T;. A meta-analysis of interventions to reduce loneliness. 2011. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3865701/>. Accessed April 9 2020.
91. Moor S, Williman J, Drummond S, et al. 'E' therapy in the community: Examination of the uptake and effectiveness of BRAVE (a self-help computer programme for anxiety in children and adolescents) in primary care. *Internet Interv*. 2019;18:100249.
92. Pennant ME, Loucas CE, Whittington C, et al. Computerised therapies for anxiety and depression in children and young people: a systematic review and meta-analysis. *Behaviour research and therapy*. 2015;67:1-18.
93. Grist R, Croker A, Denne M, Stallard P. Technology Delivered Interventions for Depression and Anxiety in Children and Adolescents: A Systematic Review and Meta-analysis. *Clin Child Fam Psychol Rev*. 2019;22(2):147-171.
94. Grist R, Porter J, Stallard P. Mental Health Mobile Apps for Preadolescents and Adolescents: A Systematic Review. *J Med Internet Res*. 2017;19(5):e176.
95. Creswell C, Violato M, Fairbanks H, et al. Clinical outcomes and cost-effectiveness of brief guided parent-delivered cognitive behavioural therapy and solution-focused brief therapy for treatment of childhood anxiety disorders: a randomised controlled trial. *Lancet Psychiatry*. 2017;4(7):529-539.
96. Lillevoll KR, Vangberg HC, Griffiths KM, Waterloo K, Eisemann MR. Uptake and adherence of a self-directed internet-based mental health intervention with tailored e-mail reminders in senior high schools in Norway. *BMC Psychiatry*. 2014;14:14.
97. Bennett SD, Cuijpers P, Ebert DD, et al. Practitioner Review: Unguided and guided self-help interventions for common mental health disorders in children and adolescents: a systematic review and meta-analysis. *Journal of child psychology and psychiatry, and allied disciplines*. 2019;60(8):828-847.

98. Arslan G. *Australian Journal of Psychology*. 2020.
99. Johnson HDL, J.L.;Mahoney,M;. Interparental Conflict and Family Cohesion: Predictors of Loneliness, Social Anxiety, and Social Avoidance in Late Adolescence. 2001. doi:<https://doi.org/10.1177/0743558401163004>. Published May 1 2001.
100. Neto F, Barros J. Psychosocial concomitants of loneliness among students of Cape Verde and Portugal. *J Psychol*. 2000;134(5):503-514.
101. Wang L, Yao J. Life satisfaction and social anxiety among left-behind children in rural China: The mediating role of loneliness. *J Community Psychol*. 2020;48(2):258-266.

Journal Pre-proof

**Table 1.** Quality Assessment Tool Adapted From National Institutes of Health<sup>16</sup>

Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	Yes: 1 No: 0
Was the exposure measure objective (ie, not self-report)	Yes: 1 No: 0
Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	Yes: 1 No: 0
Was the outcome assessed objectively?	Yes or by blinded assessors: 2 By another individual, eg, parent: 1 No, ie, self-report: 0
Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposure(s) and outcome(s)?	No or unclear: 0 Some attempt, eg, SES, demographics: 1 Reasonable or comprehensive, eg, baseline depression for longitudinal studies, other exposure to stress or adversity, negative affectivity: 2
Is a longitudinal design with exposure measured before outcome?	Yes: 1 No: 0
<b>Longitudinal only</b>	
Was loss to follow up after base line 20% or less?	Yes: 1 No: 0
Were the exposure(s) assessed more than once over time?	Yes: 1 No: 0

*Note: Exposure measures indicates independent variables.*

**Table 2.** Cross-sectional Studies Examining Social Isolation/Loneliness

Author (year), Country	Sample	Total N (% male participants)	Child (≤11)/ Adolescent (12-18)/ Young adult (≥19)	Age range at baseline (years)	Mean age (SD)	Social isolation/loneliness measure	Mental Health Measure(s)	Associations between social isolation/loneliness (lon) and mental health - r (p) unless otherwise stated		
								Depression (dep)	Anxiety (anx)	Other mental health
<b>Social isolation/loneliness and concurrent mental health symptoms</b>										
Alpaslan <i>et al.</i> (2016), <sup>25</sup> Turkey	School students	487 (41·7)	Adolescent	14 to 19	16·07 (1·05)	UCLA Loneliness Scale	CDI, SDQ	Male participants: OR 1·21 Female participants: OR 1·05		
Arslan (2020), <sup>98</sup> Turkey	School students	244 (47·5)	Adolescent	14 to 18	16·27 (1·02)	8-item UCLA Loneliness Scale-Short Form	Youth Internalizing and Externalizing Behavior Screeners			Lon - mental health problems 0·41 (<·001), $\beta = \cdot 22$ (<·01).
Baskin <i>et al.</i> (2010), <sup>26</sup> USA	School students	294 (NS)	Adolescent	NS Estimated 13-14	13·11 (0·469)	Children's Loneliness Scale (CLS)	BDI-Y	R <sup>2</sup> = ·28 (<·001). Moderated by Belongingness		
Brage <i>et al.</i> (1993), <sup>28</sup> Brage <i>et al.</i> (1995), <sup>27</sup> USA	School students	156 (39·7)	Adolescent	11 to 18	14 (1·56)	Loneliness Inventory Short Form	CES-D (child version)	0·646, (<·001).		
Chang <i>et al.</i> (2017), <sup>29</sup> USA	University students	228 (23·7)	Young adult	18 to 28	19·69 (1·38)	Revised UCLA Loneliness scale	BDI, Frequency of Suicidal Ideation	0·69 (<·001). Regressions: 47% shared variance.		Lon - suicidal ideation 0·52 (<·001). Lon R <sup>2</sup> = 26·9%



							Inventory.			variance in suicidal ideation
Doman and Le Roux (2012), <sup>30</sup> South Africa	University students	275 (42·3)	Young adult	19 to 34	20·92 (NS)	Le Roux Loneliness Questionnaire	Psychological General Well-Being Index: anxiety + depressed mood	0·517 (<·01). 26·7% shared variance.	Anx: 0·365, (<·01)	
Erdur-Baker and Bugay (2011), <sup>31</sup> Turkey	School students	144 (54·2)	Adolescent	11 to 15	12·5 (1·61)	LSDQ	CDI	0·51 (NS)		
Ginter <i>et al.</i> (1996), <sup>67</sup> Israel	School students	144 (45·1)	Adolescent	11 to 16	13·90 (1·5)	The Loneliness Rating Scale (subscales for Frequency, Intensity, Duration) + additional 2 questions	Revised Children's Manifest Anxiety Scale (RCMAS)		Not lonely group: Frequency of lon – anx 0·33 (<·001), Intensity of lon – anx 0·18 (<·05). Lonely group > anx t=3·81, (<·001),	
Heredia <i>et al.</i> (2017), <sup>68</sup> USA	School students	394 (50·2)	Adolescent	12 to 15	13·52 (0·63)	LSDQ	Wellbeing - World Health Organisation Well-being Index (WHO-5)			Lon - wellbeing 0·111, (<·05). Hierarchical linear regression - loneliness accounted for 1·3% of variance in

										wellbeing
Houghton et al (2016), <sup>69</sup> Australia	School students	1143 (46·3)	Adolescent	10·1 to 16	13·20 (1·2)	Perth Aloneness Scale (includes friendship-related loneliness subscale)	Warwick-Edinburgh Mental Well Being Scale (WEMWBS)			Friendship related lon-wellbeing 0·36 (< ·001).
Hudson et al. (2000), <sup>32</sup> USA	Adolescent mothers post-partum recruited from primary health care practices	21 (0)	Adolescent	16 to 19	18 (1·14)	Revised UCLA Loneliness Scale	CES-D (child version)	0·53 (<·05).		
Hutcherson and Epkins (2009), <sup>33</sup> USA	Female school students (and their mothers)	100 (0)	Child	9 to 12	10·52 (1·04)	Loneliness Scale (LS)	Social Anxiety Scale for Children-Revised (SASC-R), CDI.	0·62 (< ·001). Controlling for soc anx 0·36 (< ·001).	Social anx: 0·65 (< ·001). Controlling for dep 0·49 (< ·001).	
Jackson and Cochran (1991), <sup>34</sup> USA	University students	293 (49·8)	Young adult	17 to 26	Median 19	Revised UCLA Loneliness Scale	Symptom Checklist-90 (SCL-90)	0·54 (< ·001). Controlling for overall symptoms 0·23 (<·01).	General anx: 0·37 (< ·001).	Obsessive Compulsiveness 0·40 (< ·001).
Johnson et al (2001), <sup>9997</sup> USA	University students	124 (43·5)	Young adult	17 to 21	Male participants 19·41 (NS) Female participants 19·69 (NS)	UCLA Loneliness Scale (Revised)	Franke and Hymel Social Anxiety and Social Avoidance Scale		Soc anx: F(6, 115) = 4·23 (< ·05). $\beta$ = ·24, p < ·01, $R^2$ = ·31, p < ·01.	
Kim (2001), <sup>35</sup> Korea	University students	452 (44·7)	Young adult	18 to 25	20·9 (2·0)	Revised UCLA Loneliness Scale	BDI	Male participants: $\beta$ = ·49 (< ·01). 24% shared variance.		
Koenig et	School students	397 (38·3)	Adolescent	14 to 18	NS	Revised UCLA	BDI	Male		

<i>al. (1994),</i> <sup>36</sup> USA						Loneliness Scale		participants: 0.55 (<.001). Female participants: 0.49 (<.001).		
Lasgaard, Goosens <i>et al.</i> (2011), <sup>24</sup> Denmark	School students	1009 (43)	Adolescent	NS	17.11 (1.11)	SELSA– SF (3 subscales: social lon, family-related lon, romantic lon)	BAI-Y, BDI-Y, Social Interaction Anxiety Scale (SIAS), Suicide Ideation subscale from the Suicide Probability Scale, Deliberate self-harm (DSH), Risk Behavior related to Eating Disorders (RiBED-8)	23% of the variance. Peer-related lon – dep $\beta =$ 0.26, $r^2 =$ 0.076; family- related lon – dep $\beta =$ 0.29, $r^2 =$ 0.089.	Anx: 14% shared variance. Peer-related lon $\beta =$ .21, $r^2 =$ .045. Family- related lon $\beta =$ .21, $r^2 =$ .045 Social anx: 21% shared variance. Peer-related lon $\beta =$ .33, $r^2 =$ .109. Romantic lon $\beta =$ .19, $r^2 =$ .040.	Suicidal ideation (SI): 14% shared variance. Peer- related lon – SI $\beta =$ .17, $r^2 =$ .027. Family- related lon – SI $\beta =$ .26, $r^2 =$ .061. Self-harm: 10% shared variance. Family-related lon $\beta =$ .31, $r^2 =$ .081. Eating Disorder (ED): risk behaviour: 6% shared variance. Family related lon – ED $\beta =$ .22, $r^2 =$ .041.
Lau <i>et al.</i> (1999), <sup>37</sup> Hong Kong	School students	6,356 (NS estimated 48)	Child/ adolescent	9 to 14	NS	Marcoen and Brumagne’s Loneliness Scale (3 subscales: Peer-Related Lon, Parent-Related Lon, and Aloneness)	CDI, RCADS	Primary school students: 0.71 (<.001). Peer-related lon 0.67 (<.001),		

								parent-related lon 0.49 (<.001), aloneness – 0.65 (<.001). 46% shared variance. Secondary school students: 0.81 (<.001). Peer-related lon 0.77, (<.001), parent-related lon 0.56 (<.001), aloneness – dep 0.72 (<.001). 65% shared variance.		
Majd Ara <i>et al.</i> (2017), <sup>39</sup> Iran	Female school students	301 (0)	Adolescent	15 to 18	16.6 (1.1)	Children's Loneliness Scale	DASS-21	0.66 (NS).		
Mahon <i>et al.</i> (2001), <sup>38</sup> USA	School students	127 (43.3)	Adolescent	12 to 14	12.9 (0.63)	Revised UCLA Loneliness Scale	Profile of Mood States - Depression-Dejection subscale	0.57 (<.001).		
Markovic and Bowker (2015), <sup>40</sup> USA	School students	157 (45)	Adolescent	NS	13.84 (.75)	LSDQ	YSR	0.39 (<.001)	Anx: 0.35, (<.001)	
Matthews <i>et al.</i>	Twin birth cohort	2066 (49)	Young adult	18	18.4 (0.36)	Multidimensional Scale of Perceived	Diagnostic Interview	0.21 (<.001)		

(2016), <sup>20</sup> UK						Social Support (MSPSS)	Schedule			
McIntyre <i>et al.</i> (2018), <sup>41</sup> UK	University students	1135	Young adult	NS	20.78 (4.35)	UCLA Loneliness Scale	PHQ-9, GAD-7, Self-harm (4 items)	0.58 (<.001) $\beta = 0.52$ (<.001)	Anx: 0.54 (<.001) $\beta =$ 0.50 (<.001)	
Moore and Schultz (1983), <sup>42</sup> USA	School students	99 (45)	Adolescent	14 to 19	17 (0.98)	UCLA Loneliness Scale (ULS) + frequency, duration, characteristics and perceived causes of loneliness	SDS, STAI	0.66 (<.001). Lon duration 0.46, (<.001). Lon frequency -dep 0.70 (<.001).	State anx: 0.48 (<.001), Lon duration 0.37 (<.001) Lon Frequency 0.48 (<.001)	
Mounts <i>et al.</i> (2006), <sup>43</sup> USA	University students – ethnically diverse sample	350 (36)	Young adult	18 to 19	NS	Revised UCLA Loneliness Scale	BDI, BAI	$\beta = .51$ , (<.001).	Anx: $\beta = .30$ (<.001)	
Neto and Barros (2000), <sup>100</sup> Portugal	School students	487 (39.3)	Adolescent	NS (estimated 15 to 18)	Cape Verde 17.5 (1.2): Portugal 17.8 (1.0).	Revised UCLA Loneliness Scale	Social Anxiety subscale		Soc anx: 0.33-0.35 (<.001)	
Purwono and French (2016), <sup>44</sup> Indonesia	Muslim school students	453 (45.9)	Adolescent	13 to 16	7th grade: 13.57 (0.44) 10th grade: 16.47 (0.43)	10 items from UCLA Loneliness Scale - modified	CES-D	0.59 (< .01).		
Richardson <i>et al.</i> (2019), <sup>21</sup> Australia	Community	528 (51)	Child/ adolescent	10 - 12	11.18 (0.56)	3 items from School Belonging and Isolation Scale	SCAS-C– subscales generalized anx, social anx and	0.46 (< .001).	Social anx: 0.50 (< .001). Generalized anx: 0.42 (< .001).	

							separation anx 3 item SMFQ		Separation anx: 0.41 (< .001).	
Roberts and Chen (1995), <sup>45</sup> USA	School students	2614 (n.s)	Adolescent	11 to 14	NS (NS)	8 item UCLA Loneliness Scale	CES-D, 4 suicide items from Oregon Adolescent Depression Project.	OR = 5.8 (< .001).		Suicidal ideation: OR 5.0
Singhvi <i>et al.</i> (2011), <sup>46</sup> India	School students	300 (50)	Adolescent	15 to 17	NS	Revised UCLA Loneliness Scale	SDS, Cohen's Perceived Stress Scale	Male participants: 0.461 (<.001) Female participants: 0.683 (<.001). Male participants: lon associated with dep [t=6.32, p<0.005, β=.461]. Female participants: lon associated with dep [t=11.38, p<.005, β=.683].	Male participants: lon associated with perceived stress [t=1.50, p<.01, β=- .108]	
Spithoven <i>et al.</i> (2017), <sup>47</sup> Belgium and	NS	Sample 1: 417 (48.4) Sample 2: 1140 (48.7)	Adolescent	NS	Sample 1: 12.47 (1.89) Sample 2: 12.81	LACA – peer-related loneliness subscale	Sample 1: CDI. Sample 2: Iowa short form of CES-	Sample 1: 0.48 (<.001). Sample 2: 0.54 (<.001).		

Netherlands					(0.42).		D.			
Stednitz and Epkins (2006), <sup>48</sup> USA	Community sample	102 (0)	Child	9 to 12	10.46 (1)	LSDQ	CDI, Social Anxiety Scale for Children – Revised (child and parent versions)	0.63 (< .001)	Social anx: self-rated 0.72 (< .001)· mother rated 0.36 (< .001).	
Stacciarini <i>et al.</i> (2015), <sup>22</sup> USA	Church and community (Latina/o immigrants)	31 (42)	Adolescent	11 to 18	13.0 (2.0)	Short version of PROMIS Health Organisation Social Isolation	SF12 Health survey			Mental health (r = -.38, p < .05)
Stickley <i>et al.</i> (2016), <sup>49</sup> Czech, Russia and USA	School students	Sample 1: 2205 (NS) Sample 2: 1995 (NS) Sample 3: 2050 (NS)	Adolescent	13 to 15	NS	Lon item from CES-D	CES-D (minus lon item), 12 statement anxiety scale	ORs: 8.04-40.13.	Anx: ORs: 1.63 - 5.49.	
Swami <i>et al.</i> (2007), <sup>50</sup> Malaysia	University students	172 (41.8)	Young adult	18 to 24	20.3 (1.25)	Revised UCLA Loneliness Scale	BDI	0.38 (< 0.01).		
Thomas and Bowker (2015), <sup>51</sup> USA	School students	103 (51.4)	Child/ Adolescent	NS (estimated 10-13)	13.73 (0.82)	LSDQ	YSR	0.42 (< 0.1)		
Tu and Zhang (2014), <sup>52</sup> China	University students	444 (38.4)	Young adult	NS	19.02 (1.26)	Revised UCLA Loneliness Scale	CES-D (7 item version), Perceived Stress Scale	$\gamma = .517$ , (< .001). $\beta = .833$ (< .001).	Stress: $\gamma = .381$ , (< .001), $\beta = .297$ (< .001)	
Uba <i>et al.</i> (2012), <sup>66</sup> Malaysia	School students	242 (49.2)	Adolescent	13 to 16	14.67 (1.27)	Revised UCLA Loneliness Scale	CDI	0.493 (< .01).		
Vanhalst, Luyckx,	University students	370 (16.5)	Young adult	NS	18.22 (1.21)	LACA	CES-D	Peer-related lon 0.58		

Raes (2012), <sup>53</sup> Belgium								(.001). Parent-related lon 0.23 (< .001).		
Wang and Yao (2020), <sup>101</sup> China	Schools (left behind children in rural China)	442 (54)	Child/Adolescent	8 to 16	11.5 (2.098)	UCLA Loneliness Scale	Social Anxiety Subscale		Soc anx: 0.332 (< .001)	
Xu and Chen (2019), <sup>54</sup> China	School students	724 (59.5)	Child/Adolescent	6 to 14	9.15 (1.79)	LSDQ	CES-D	0.492 (< 0.01).		
Yadegarfar <i>et al.</i> (2014), <sup>2325</sup> Thailand	Transgender association and university (male Transgender and cis gender)	260 (100)	Adolescent/Young adult	15 to 25	20 (NS)	SSA	DASS-21 (short version), Positive and Negative Suicide Inventory	Transgender: Soc support-dep. (B = -0.01) Lower soc support associated with higher negative risk factors related to suicidal behaviour (B = .13). Cisgender: Soc support-dep. (B = .23). Lower soc support associated with higher negative risk factors related to		



								suicidal behaviour (B = .15).	
<b>Social isolation/quarantine in the context of infectious disease</b>									
Sprang and Silman (2013), <sup>76</sup> USA, Canada and Mexico	Parents of children (who experienced H1N1/SARS/avian flu pandemics)	398 (NS)	Child	NS	NS	Children experienced pandemic – 20·9% social isolation and 3·8% quarantine	PTSD-RI; PCL-C	PTSD-RI: Children who experienced isolation/quarantine were more likely to meet cut-off score for PTSD (30%) than those who had not been in isolation or quarantine (1·1%; $X^2 = 49·56$ , $P < ·001$ , Cramer V = ·449). Mean scores in isolated/quarantined group (22·3) were 4 x general group (5·5), ( $t = 6·59$ , $P = ·000$ ). PCL-CL: Children who experienced isolation/quarantine were more likely to meet cut-off score for PTSD (28%) ( $X^2 = 31·44$ , $P < ·001$ )	

Note: Anx = Anxiety; BAI = Beck Anxiety Inventory; BAI-Y = Beck Anxiety Inventory for Youth; BDI = Beck Depression Inventory; BDI-Y = Beck Depression Inventory for Youth; CBCL = Child Behaviour Checklist; CDI = Children's Depression Inventory; CES-D = Center for Epidemiologic Studies Depression Scale; DASS-21 Depression, Anxiety, and Stress Scale, Dep = depression; GAD-7 = Generalized Anxiety Disorder – 7; Lon = Loneliness; LSDQ = Loneliness and Social Dissatisfaction Questionnaire; LACA = Loneliness and Aloneness Scale for Children and Adolescents; OR = Odds Ratio; PCL-C = PTSD Checklist Civilian Version; PHQ-9 = Patient Health Questionnaire; PTSD-RI = UCLA Posttraumatic Stress Disorder Reaction Index; RCADS = Revised Children's Anxiety and Depression Scale; SAS-A = Social Anxiety Scale for Adolescents; SCAS-C = Spence Children's Anxiety Scale- Child; SDS = Zung Self-rating Depression Scale; SDQ = Strengths and Difficulties Questionnaire; SELSA = Social and Emotional Loneliness Scale for Adults; SMFQ = Short Mood and Feelings Questionnaire-Child; SSA = Social Support Appraisals scale; STAI = State Trait Anxiety Inventory; TRF = Teacher Rating Form; YSR = Youth Self-Report Form.

**Table 3.** Longitudinal Studies Examining Social Isolation/Loneliness and Subsequent Mental Health Outcomes

Author (year), Country	Sample (selection criteria)	Total N (% male participants)	Child (≤11)/ Adolescent (12-18)/ Young adult (≥19)	Age range (years)	Mean age (S.D.) at T1	Social isolation/loneliness measure	Mental Health Measures	Cross-sectional associations r (p)	Length of follow-up (years)	Is social isolation/loneliness associated with later mental health?	
										Depression	Anxiety
Boivin <i>et al.</i> (1995), <sup>55</sup> Canada	School students	774 (51·8)	Child	9 to 12	10·8 (NS)	LSDQ	CDI	Lon-dep 0·53 (<·001)	1	T1 Lon – T2 Dep: r = 0·36 (p < ·01) T1 Lon accounted for 8·3% of the variance in T2 Dep.	
Christ <i>et al.</i> (2017), <sup>71</sup> USA	National Survey of Child and Adolescent Well-being (child welfare cohort)	2776 (47)	Adolescent	11 to 17	13·5 (NS)	LDSQ 7 peer isolation items	4 items from YSR	NS	7	Controlling for caregiver neglect and covariates, a 1 S.D. increase in peer Isolation was associated with a 0·49 S.D. increase in depression	
Danneel <i>et al.</i> (2019), <sup>56</sup> Belgium	Longitudinal cohorts	Sample 1: 1116 (51·1), Sample 2: 1423 (47·6), Sample 3: 549 (37·33)	Adolescent	Sample 1: 11 to 17 Sample 2: 11 to 18 Sample 3: 12 to 17	Sample 1: 13·79 (0·94) Sample 2: 13·59 (0·98) Sample 3: 14·82 (0·79)	LACA peer-related loneliness subscale	Samples 1 and 3 – SAS-A; CES-D. Sample 2 - CDI	Lon-Social anxiety 0·58 ≤ r ≤ 0·67. Lon-Dep 0·48 ≤ r ≤ 0·56, (all <·01).	1	Not significant	Lon --> Social Anxiety (β = 0·10, p < 0·001).
Fontaine <i>et</i>	School students	NS (52)	Child	NS	NS	LSDQ (T2)	Internalizing items	NS	2-3	T2 Lon --> Anx/Dep symptoms at	

<i>al.</i> (2009), <sup>72</sup> USA	(longitudinal cohort)			Estimated 5 to 9			from: CBCL (mother T1 and T3); TRF (teacher T1 and T2); YSR (self T2 and T3)			T3 ( $\gamma^2 = .18$ , $z = 2.60$ , $p < .01$ ).
Jones <i>et al.</i> (2011), <sup>73</sup> USA	Longitudinal cohort	889 (50)	Child	6	NS	LSDQ	CDI short form	NS	9	Indirect effects T1 Lon --> T2 Suicidal Thoughts through Dep ( $\beta = .06$ , $p < .001$ )
Ladd and Ettekal (2013), <sup>57</sup> USA	School students (longitudinal cohort)	478 (50)	Adolescent	12 to 18	12.0 (n.s)	LSDQ – revised - 3 items	Depression items CBCL (parent); TRF (teacher); YSR (self)	Lon-Dep 0.19 ( $< .01$ ) (parent), 0.38 ( $< .001$ ) (teacher) 0.62 ( $< .001$ ) (self)	7	Changes in Lon associated with changes in dep reported by teachers ( $r = 0.63$ , $p < .001$ ) and adolescents ( $r = 0.65$ , $p < .001$ ), but not parents ( $r = 0.18$ , $p = 0.13$ )
Lalayants and Prince (2015), <sup>74</sup> 83 countries	National Survey of Child and Adolescent Wellbeing (child welfare cohort)	356 (0)	Adolescent	11 to 12	NS	LSDQ	CDI	NS	1.5	T1 Lon --> T2 Dep AOR=2.93, CI=1.74-4.91, $p < .001$ . T1 lonely female participants were 5.09 times more likely (CI 2.24-11.56, $p < .001$ ) to be depressed at T2.
Lapierre <i>et al.</i> (2019), <sup>58</sup> USA	College Students	346 (33.6)	Young adult	17 to 20	19.11 (0.75)	UCLA Loneliness Scale	10 item CES-D	Lon-Dep 0.628 (T1), 0.666 (T2) ( $< .001$ )	0.25	T1 Lon – T2 Dep $r = 0.524$ , $p < .001$ T1 Lon --> T2 Dep ( $b = .21$ , $SE = .05$ , $p < .001$ ),

Lasgaard <i>et al.</i> (2011b), <sup>17</sup> Denmark	School students	T1: 1009 (43) T2: 541 (40)	Adolescent/ Young adult	15 to 26	17·11 (1·11)	SELSA-short form; MSPSS	BAI-Y, BDI-Y	Lon-Dep 0·61 (<·0005) Lon-Anx 0·51 (<·0005). Soc support - dep r = -0·12, -0·18, -0·28 (all p < ·0005)	1	T1 Lon--> T2 Dep r = 0·37, p < ·0005. Cross lagged structural equation modelling found T1 Lon did not predict dep at T2.	
Liu <i>et al.</i> (2020), <sup>18</sup> China	College students	741 (28·3)	Young adult	NS (estimated 18-20)	18·47 (0·87)	6 item index of social isolation based on only child status, number of friends, frequency of contact with friends and family; UCLA Loneliness Scale	SDS	NS	3	Female participants: T1 isolation associated with increased dep ( $\beta = 0·22$ , p < 0·001). Lon associated with increased dep. ( $\beta = 0·23$ , p < 0·001). Male participants: T1 isolation associated with increased dep. ( $\beta = 0·25$ , p < 0·01). Lon did not predict dep. ( $\beta = 0·14$ , p > 0·05)	
Mak <i>et al.</i> (2018), <sup>75</sup> USA	School students (Randomised trial)	687 (47·7)	Adolescent	NS (estimated 11 to 14)	11·27 (0·49)	LSDQ	SAS-A	Lon-social anxiety 0·41-0·45 (< ·01).	1·5 (T2), 3 (T3)		T1 Lon --> T2 Social Anxiety ( $\beta = -0·09$ , p < ·05). T2 Lon --> T3 Social Anxiety ( $\beta = -0·12$ , p < ·01)

											By gender: T2 Lon --> T3 Soc Anxiety: Boys ( $\beta = \cdot22$ , $p < \cdot001$ ). Girls ( $\beta = \cdot01$ , $p = \cdot79$ )
Matthews <i>et al.</i> (2015), <sup>19</sup> UK	Twin birth cohort	2232 (NS)	Child	5	NS	6 items from CBCL (parent) and TRF (teacher)	MASC	NS	7		T1 social isolation failed to predict T2 anx, controlling for T1 anx.
Qualter <i>et al.</i> (2010), <sup>59</sup> UK	School students	296 (49.3)	Child	5	NS	T1 and T2: Peer and Parent subscales LACA	T1: T-CARS T2 and T3: DDPCA	T1 Peer Lon-internalizing symptoms 0.32 ( $< \cdot01$ ) Parent Lon-Internalizing Symptoms 0.09.  T2 Peer Lon-Dep 0.13 ( $< \cdot05$ ) Parent Lon-Dep 0.12 ( $< \cdot05$ )	8	T1 Peer Lon-T2 Dep $r = 0.07$ . T1 Peer Lon-T3 Dep $r = 0.06$ . T2 Peer Lon – T3 Dep $r = 0.12$ , $p < \cdot05$ T1 Parent Lon – T2 Dep $r = 0.19$ , $p < \cdot01$ T1 Parent Lon-T3 Dep $r = 0.13$ , $p < \cdot05$ T2 Parent Lon-T3 Dep $r = 0.08$ Structural model: Duration of Peer Lon --> T3 dep. T1 and T2 Peer Lon, Parent Lon (T1, T2, and duration), did not independently	

										predict T3 Dep.	
Schinka <i>et al.</i> (2013), <sup>60</sup> USA	Longitudinal cohort study	832 (53)	Child	9	NS	LDSQ	T1: CBCL (mother)  T3: CDI -Short form; Suicide items from CBCL and YSR	T3 Lon- Dep - 0-10 (< .01) Lon- Suicidal Ideation r = 0-02 Lon- Suicidal Attempt r = 0-4	2 (T2), 6 (T3)	T1 Lon-T3 Dep r = 0-01 T2 Lon-T3 Dep r = -0-01 T1 Lon- T3 Suicidal Ideation r = 0-00 T2 Lon-T3 Suicidal Ideation r = 0-03 T1 Lon- T3 Suicidal Attempt r = 0-02 T2 Lon-T3 Suicidal Attempt r = -0-01	
Vanhalst, Goosens <i>et al.</i> (2013) <sup>61</sup> and Vanhalst, Klimstra <i>et al.</i> (2012), <sup>62</sup> Netherlands	Community sample via municipality registers	389 (53)	Adolescents	15	15-22 (0-60)	LACA Peer-related loneliness subscale	6 item depression questionnaire; SCARED generalized anxiety, panic and social anxiety subscales.	Lon- Dep - 0-34 -0-50 (<.001). Lon- Perceived Stress 0-23, (< .001). Lon- Generalized Anx 0-40 (< .001), Lon-Panic 0-13 (p < .05), Lon- Social Phobia 0-47 (< .001).	5	T1 Lon --> T2 Dep Symptoms (B = .13, p < .001)	
Vanhalst, Luyckx <i>et al.</i> (2012), <sup>63</sup> Belgium	University students	Sample 1: 514 (10-9) Sample 2: 437 (17)	Young adults	Sample: 19-62 (0-62) Sample 2:	NS	Sample 1: 8-item revised UCLA Loneliness	Sample 1: 12-item CES-D Sample 2: 20-item CES-D	Sample 1: Lon – dep 0-49-0-52 (< .001).	2	Sample 1: T1 lon – T2 dep r= 0-35, p < .001. T1 lon – T3 dep r=	

				18-22 (1-21)		Scale. Sample 2: LACA Peer- related loneliness subscale		Sample 2: Lon – dep r= 0.40-0.60 (< .001).		0.36, p < .001. Lon --> associated with dep across both time intervals. Sample 2: cross- lagged path from lon --> associated with dep (b = .12, p < .05)	
Wang <i>et al.</i> (2020), <sup>64</sup> China	School students	921 (48.3)	Adolescents	12 to 15	12-98 (0.66)	Revised UCLA Loneliness Scale (T1 and T2)	SCARED; DSRSC (T1 and T3)	T1 Lon- Anx 0.40 p<.001, Lon-Dep 0.57, p<.001,	1	T1 Lon-T3 Dep 0.36, p<.001. T2 Lon-T3 Dep 0.46, p<.001.	T1 Lon-T3 Anx 0.29, p<.001. T2 Lon-T3 Anx 0.36, p<.001.
Zhou <i>et al.</i> (2020), <sup>65</sup> China	School students	866 (49)	Adolescents	11 to 15	12-98 (0.67)	UCLA Loneliness Scale (T1 and T2)	DSRSC (T3)	T1 Lon-Dep r = 0.56, p < .001	2	T1 Lon-T3 Dep r = 0.38, p < .001 Controlling for age, gender and SES, T2 Lon - T3 Dep adj. b= 0.34 p <.001.	

Note: Anx = Anxiety; BAI-Y = Beck Anxiety Inventory for Youth; BDI-Y = Beck Depression Inventory for Youth; CBCL = Child Behaviour Checklist; CDI = Children's Depression Inventory; CES-D = Center for Epidemiologic Studies Depression Scale; DDPCA = Depression profile for children and adolescents; Dep = depression; DSRSC = Birlson Depression Self-Rating Scale for Children; Lon = Loneliness; LSDQ = Loneliness and Social Dissatisfaction Questionnaire; LACA = Loneliness and Aloneness Scale for Children and Adolescents; MASC = Multidimensional Anxiety Scale for Children; MSPSS = Multidimensional Scale of Perceived Social Support; NS = not specified; SAS-A = Social Anxiety Scale for Adolescents; SCARED = Scale for Child Anxiety Related Emotional Disorders; SDS = Zung Self-rating Depression Scale; SELSA = Social and Emotional Loneliness Scale for Adults; T-CARS = Teacher-Classroom Adjustment Rating Scale; T1 = Time 1; T2 = Time 2; T3 = Time 3; TRF = Teacher Rating Form; YSR = Youth Self-Report Form.

**Table 4.** Study Description and Relevant Findings: Intervention Studies

Author (year),	Sample	Total N (% male)	Age range at baseline (years)	Mean age (S.D.)	Loneliness measure	Mental Health Measures	Intervention	Comparison condition	Main findings
----------------	--------	------------------	-------------------------------	-----------------	--------------------	------------------------	--------------	----------------------	---------------


Country		participants)							
King <i>et al.</i> (2018), <sup>77</sup> USA	Experienced bullying/ Victimization, recruited via paediatric medical emergency services	218 (33·5)	12 to 15	13·50 (1·1)	Revised UCLA Loneliness Scale	Reynolds Adolescent Depression Scale - 2 short; Columbia Suicide Severity Rating Scale	LET'S CONNECT (LC) mentorship program – strengths-based approach. Mentorship lasted an average of 120·32 days (SD = 69·69), 4-6 hours/month.	No treatment	At 6 months, loneliness decreased more in the LC intervention group than to the control group ( $p < .01$ ), ES = .4.
Larsen <i>et al.</i> (2019), <sup>78</sup> Norway	School students	2254 (NS estimate 53).	15 to 19	16·82 (NS)	Loneliness Scale (modified)	Symptom Checklist	Dream School Program – aimed to change psychosocial environment of classroom, including through peer mentors and a staff mental health support team. 2 classes over 2 semesters.	Education as usual.	No significant effects on mental health or loneliness for either intervention group.

Note: ES = effect size; NS = not specified.



Figure 1: PRISMA Diagram

Journal Pre-proof



## Rapid Systematic Review: The Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in the Context of COVID-19

Maria Elizabeth Loades, DClinPsy, Eleanor Chatburn, MA, Nina Higson-Sweeney, BSc, Shirley Reynolds, PhD, Roz Shafran, PhD, Amberly Brigden, MSc, Catherine Linney, MA, Megan Niamh McManus, BSc candidate, Catherine Borwick, MSc, Esther Crawley, PhD

The authors have reported no funding for this work. All research at Great Ormond Street Hospital NHS Foundation Trust and UCL Great Ormond Street Institute of Child Health is made possible by the NIHR Great Ormond Street Hospital Biomedical Research Centre. This report is independent research. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

### Author Contributions:

E. Chatburn and M. Loades conceived the review, with input from R. Shafran, S. Reynolds and E. Crawley on refining the review questions and focus. C. Borwick developed the search strategy, with input from M. Loades, N. Higson-Sweeney, R. Shafran and S. Reynolds. N. Higson-Sweeney, A. Bridgen, C. Linney and M. McManus undertook the searches and screening. E. Chatburn and M. Loades completed the data extraction with input from A. Brigden and C. Linney. S. Reynolds conducted the quality assessment. M. Loades led on synthesising the findings, and wrote the first draft of the manuscript with input from E. Chatburn, R. Shafran, E. Crawley and S. Reynolds. All co-authors reviewed and approved the final draft of the manuscript.

The authors very much appreciate the valuable comments made by the anonymous peer reviewers.

Disclosure: Dr. Loades has received funding from the National Institute for Health Research (NIHR Doctoral Research Fellowship, DRF-2016-09-021). Ms. Brigden has received funding from the National Institute for Health Research (NIHR Doctoral Research Fellowship, DRF-2017-10-169). Drs. Reynolds, Shafran, Crawley and Mss. Chatburn, Higson-Sweeney, Linney, McManus, and Borwick have reported no biomedical financial interests or potential conflicts of interest.

## Appendix. Database searches –03/29/2020

**Table S1. Ovid MEDLINE (R)**

1	exp Adolescent/ or exp Child/ or exp Child, Preschool/ or exp Infant/ or exp Minors/ or exp Pediatrics/	3533050
2	(adolesc* or preadolesc* or pre-adolesc* or boy* or girl* or child* or infan* or preschool* or pre-school* or juvenil* or minor* or pe?diatri* or pubescen* or pre-pubescen* or prepubescen* or puberty or teen* or young* or youth* or school* or high-school* or highschool* or schoolchild* or school child*).tw,kf.	2951684
3	1 or 2	4748091
4	quarantine*.tw,kf.	4350
5	exp Quarantine/	2093
6	Quarantine.tw,kf.	3975
7	exp social isolation/	17148
8	(isolation and (infect* or SARS or influenza or flu or MERS or ebola or COVID-19)).tw,kf.	34141
9	exp Loneliness/	3552
10	4 or 5 or 6 or 7 or 8 or 9	56227
11	anxiet*/ or anxious*/ or "anxiety disorder*".tw,kf.	29320
12	depress*/ or "internal* disord*" or "low mood".tw,kf.	737
13	depressive disorder/	72188
14	exp depression/	115922
15	depress*.tw,kf.	445459
16	exp adjustment disorders/	4197
17	adjustment disorder*.tw,kf.	1642
18	low mood.tw,kf.	737
19	obsessive-compulsive disorder.tw,kf.	12336
20	stress disorders, traumatic/	672
21	stress disorders, post-traumatic/	31840
22	trauma*.tw,kf.	353295
23	((post-trauma* or posttrauma*) adj stress) or PTSD).tw,kf.	35040
24	11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23	853134
25	3 and 10 and 24	1277

Full references saved as Medline 290320 v1

**Table S2. Ovid PsycINFO**

1	(adolescent or child or child, preschool or infant or minor or pediatrics).ti,ab,id.	425212
2	(adolesc* or preadolesc* or pre-adolesc* or boy* or girl* or child* or infan* or preschool* or pre-school* or juvenil* or minor* or pe?diatri* or pubescen* or pre-pubescen* or prepubescen* or puberty or teen* or youth* or school* or high-school* or highschool* or schoolchild* or school child*).ti,ab,id.	1227549
3	1 or 2	1227549
4	quarantine.ti,ab,id.	179
5	exp *Social Isolation/	5944
6	(isolation and (infect* or SARS or influenza or flu or MERS or ebola or COVID-19)).ti,ab,id.	437
7	Disease containment*.ti,ab,id.	5
8	Lonel*.ti,ab,id.	10569
9	exp *loneliness/	3642
10	4 or 5 or 6 or 7 or 8 or 9	16688
11	anxiet*/ or anxious*/ or "anxiety disorder*".ti,ab,id.	33786
12	depress*/ or "internal* disord*"/ or "low mood".ti,ab,id.	673
13	exp *depression/	19678
14	depress*.ti,ab,id.	301583
15	exp adjustment disorders/	719
16	adjustment disorder*.ti,ab,id.	1851
17	obsessive-compulsive disorder.ti,ab,id.	15268
18	post-traumatic stress disorder.ti,ab,id.	10195
19	trauma*.ti,ab,id.	107899
20	((post-trauma* or posttrauma*) adj stress) or PTSD.ti,ab,id.	44403
21	11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20	431601
22	3 and 10 and 21	1303

Full references saved as PsycINFO 290320 v1

**Table S3. Web of Science Core Collection**

# 22	<u>3,211</u>	#21 AND #10 AND #3
# 21	<u>1,173,555</u>	#20 OR #19 OR #18 OR #17 OR #16 OR #15 OR #14 OR #13 OR #12 OR #11
# 20	<u>64,185</u>	TS=(((post-trauma* or posttrauma*) NEAR stress) or PTSD)
# 19	<u>387,085</u>	TS=trauma*
# 18	<u>15,994</u>	TS=post traumatic stress disorder
# 17	<u>25,733</u>	TS=obsessive compulsive disorder
# 16	<u>22,119</u>	TS=adjustment disorder*
# 15	<u>22,104</u>	TS=adjustment disorders
# 14	<u>627,349</u>	TS=depress*
# 13	<u>494,240</u>	TS=depression
# 12	<u>628,267</u>	TS=(depress* OR " internal* disord* " OR " low mood ")
# 11	<u>283,559</u>	TS=(anxiet* OR anxious* OR " anxiety disorder* ")
# 10	<u>77,296</u>	#9 OR #8 OR #7 OR #6 OR #5 OR #4
# 9	<u>12,570</u>	TS=loneliness
# 8	<u>15,420</u>	TS=Lonel*
# 7	<u>2,586</u>	TS=Disease containment*
# 6	<u>35,721</u>	TS=(isolation and (infect* or SARS or influenza or flu or MERS or ebola or COVID-19))
# 5	<u>17,794</u>	TS=social isolation
# 4	<u>8,759</u>	TS=quarantine
# 3	<u>3,591,598</u>	#2 OR #1
# 2	<u>3,581,837</u>	TS=(adolesc* or preadolesc* or pre-adolesc* or boy* or girl* or child* or infan* or preschool* or pre-school* or juvenil* or minor* or pe?diatri* or pubescen* or pre-pubescen* or prepubescen* or puberty or teen* or youth* or school* or high-school* or highschool* or schoolchild* or school child*)
# 1	<u>2,450,709</u>	TS=(adolescent OR child OR child, preschool OR infant OR minor OR pediatrics)

Applied 'English language' limit = 3012