



You have downloaded a document from  
**RE-BUŚ**  
repository of the University of Silesia in Katowice

**Title:** The expansion of some distributions into the Wiener series

**Author:** Anna Cichocka

**Citation style:** Cichocka Anna. (2000). The expansion of some distributions into the Wiener series. "Annales Mathematicae Silesianae" (Nr 14 (2000), s. 7-21).



Uznanie autorstwa - Użycie niekomercyjne - Bez utworów zależnych Polska - Licencja ta zezwala na rozpowszechnianie, przedstawianie i wykonywanie utworu jedynie w celach niekomercyjnych oraz pod warunkiem zachowania go w oryginalnej postaci (nie tworzenia utworów zależnych).



UNIwersYTET ŚLĄSKI  
W KATOWICACH



Biblioteka  
Uniwersytetu Śląskiego



Ministerstwo Nauki  
i Szkolnictwa Wyższego

# THE EXPANSION OF SOME DISTRIBUTIONS INTO THE WIENER SERIES

ANNA CICHOCKA

## 1. Introduction

The aim of this paper is to investigate a discrete integral transform on the real line, which seems to be better adapted for some applications than the Hermite transform (see for example [6]). Another complete orthonormal system (CON) of functions on the real line, which was introduced by Wiener is more appropriate for nonlinear differential equations of mathematical physics. The reasons are that there exist linearization formulas with respect to the argument as well as with respect to the index and that the functions tend to zero as  $|x|$  tends to infinity as quickly as  $|x|^{-1}$ .

**Notations:** Let  $\mathbb{Z}$  be the set of integers,  $\mathbb{N}$  the set of positive integers,  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ ,  $\mathbb{R}$ —the field of real numbers,  $\mathbb{C}$ —the field of complex numbers. Instead of  $\sum_{n=-\infty}^{+\infty} a_n$ , we will write  $\sum a_n$ . For abbreviation we denote  $L_2(\mathbb{R}) = L_2$ ,  $C^\infty(\mathbb{R}) = C^\infty$ . By  $\| \cdot \|$  we will denote the norm and by  $(\cdot, \cdot)$  the inner product of  $L_2$ .

## 2. The spaces $A$ and $A_k$ and their characterizations

The set of functions  $\{\varrho_n\}_{n=-\infty}^{+\infty}$ , where

$$(1) \quad \varrho_n(x) = \frac{1}{\sqrt{2\pi}} \frac{(-ix - \frac{1}{2})^n}{(-ix + \frac{1}{2})^{n+1}} \quad \text{for } n \in \mathbb{Z}, x \in \mathbb{R}$$

forms CON in  $L_2$ . These functions were introduced by N. Wiener in 1949. For details we refer to [6], [9].

Obviously  $\varrho_n \in C^\infty$ . Let  $S$  be the differential operator defined by

$$(2) \quad (Su)(x) = i(x - \frac{1}{2}i) \frac{d}{dx} [(x + \frac{1}{2}i)u(x)].$$

We can easily check that, the functions  $\varrho_n$  are eigenfunctions of the differential operator  $S$  belonging to the eigenvalues  $\lambda_n = -n$ ;

$$(3) \quad S\varrho_n = -n\varrho_n.$$

DEFINITION 1. We consider the space  $A$  defined by:

$$(4) \quad A = \{\varphi \in C^\infty : \|S^k \varphi\| < \infty, k \in \mathbb{N}_0\}.$$

By integration by parts it is easy to verify that  $(S\varphi, \varrho_n) = (\varphi, S\varrho_n)$  for  $\varphi \in A$ ,  $n \in \mathbb{Z}$ . Notice that from this definition  $A \subset L_2$ . A. H. Zemanian [10] proves that  $A$  is a complete countable multinormed space with the system of semi-norms  $\{\alpha_k\}_{k=0}^\infty$ , where  $\alpha_k(\varphi) := \|S^k \varphi\|$ . Notice also that functions  $\varrho_n$ ,  $n \in \mathbb{Z}$  belong to  $A$ .  $S$  is a continuous linear operator of  $A$  into itself.

THEOREM 1. For each element  $\varphi \in A$  we have the representation

$$(5) \quad \varphi = \sum (\varphi, \varrho_n) \varrho_n.$$

We have the following characterization of the space  $A$ :

THEOREM 2. If  $\varphi$  is in  $A$ , then  $\varphi = \sum a_n \varrho_n$  and  $\sum |n|^{2k} |a_n|^2$  is convergent for each  $k \in \mathbb{N}_0$ , where  $a_n = (\varphi, \varrho_n)$ . Conversely if  $\sum |n|^{2k} |a_n|^2$  is convergent, then the series  $\sum a_n \varrho_n$  converges to some  $\varphi$  in  $A$ .

For the proof see [10], p. 312–313 or [11], p. 268.

DEFINITION 2. For each  $k \in \mathbb{N}$  and  $\varphi \in A$  we define

$$(6) \quad |\varphi|_k^2 := ((S^2 + I)^k \varphi, \varphi).$$

Immediately we see that the following theorem holds.

THEOREM 3. We have following properties of  $|\cdot|_k$ :

$$(7) \quad |\varphi|_k^2 = \sum_{l=0}^k \binom{k}{l} \|S^l \varphi\|^2 \quad \text{for each } k \in \mathbb{N} \text{ and } \varphi \in A,$$

$$(8) \quad \|\varphi\|_{k+1} \geq \|\varphi\|_k \geq \|\varphi\| \quad \text{for each } k \in \mathbb{N} \text{ and } \varphi \in A.$$

Notice, that  $\|\cdot\|_k$  is the norm in  $A$  for each  $k \in \mathbb{N}$ .

DEFINITION 3. By  $A_k$  we denote the completion of  $A$  in the norm  $\|\cdot\|_k$ , for each  $k \in \mathbb{N}$ .

THEOREM 4. *The norms  $\|\cdot\|_k$  are compatible in the following sense: If a sequence  $\{\varphi_n\}$  of elements of  $A$  converges to zero with respect to the norm  $\|\cdot\|_m$  and is a Cauchy sequence in the norm  $\|\cdot\|_k$  ( $m \leq k$ ), then it also converges to zero in the norm  $\|\cdot\|_k$ . (Compare [7], p. 14.)*

Since the norms  $\|\cdot\|_k$  are compatible,  $A$  is complete and the relation (8) holds, so we have

$$(9) \quad A \subset \dots \subset A_{k+1} \subset A_k \subset \dots \subset A_2 \subset A_1 \subset L_2,$$

$$(10) \quad A = \bigcap_{k=1}^{\infty} A_k.$$

For the proof see [4] or [7].

We have also a characterization of  $A_k$ :

THEOREM 5. *The function  $\varphi$  belonging to  $L_2$  is in  $A_k$  if and only if the series  $\sum n^{2k} |a_n|^2$  is convergent, where  $a_n = (\varphi, \varrho_n)$ .*

For the proof see [7], p. 34.

### 3. The dual spaces $A'$ and $A'_k$ and their characterizations

DEFINITION 4. By  $A'_k$  and  $A'$  we denote the dual space of  $A_k$  and  $A$ , respectively.

It means that  $A'(A'_k)$  is the space of all continuous linear functionals on  $A(A_k)$ .

For  $f \in A'$  and  $\varphi \in A$ , we use the following notation:

$$(f, \varphi) = f(\overline{\varphi}).$$

THEOREM 6. *If  $f \in A'$  then  $f = \sum (f, \varrho_n) \varrho_n$ , where the series is convergent in  $A'$ .*

For the proof see [10] or [11].

**THEOREM 7.** *The following inclusions take place:*

$$(11) \quad L_2 \subset A'_1 \subset A'_2 \subset \dots \subset A'_k \subset A'_{k+1} \subset \dots \subset A' \quad \text{and} \quad A' = \bigcup_{k=1}^{\infty} A'_k.$$

For the proof see [4].

We have also characterizations of  $A'_k$  and  $A'$ .

**THEOREM 8.** *If  $f$  is in  $A'$ , then the series  $\sum a_n \varrho_n$ ,  $a_n = f(\varrho_n)$  converges to  $f$  in  $A'$  and there exists an integer  $k \in \mathbb{N}_0$  such that the series  $\sum_{n \neq 0} |n|^{-2k} |a_n|^2$  is convergent.*

*Conversely if the series  $\sum_{n \neq 0} |n|^{-2k} |a_n|^2$  is convergent, then there exists a continuous linear functional  $f$  in  $A'$  such that  $f(\varrho_n) = a_n$ .*

For the proof see [10], p. 323–324 or [11], p. 271–272.

**THEOREM 9.** *If  $f$  is in  $A'$ , then  $f$  belongs to  $A'_k$  if and only if the series  $\sum_{n \neq 0} |n|^{-2k} |a_n|^2$  is convergent, where  $a_n = f(\varrho_n)$ .*

For the proof see [7], p. 36, theorem 4.9.

#### 4. Connection between $A_k$ and the Sobolev space $W_{k,2}$

**DEFINITION 5.** We shall denote by  $W_{m,2}$  for  $m \in \mathbb{N}$ , the subspaces of  $L_2$  defined by

$$\{f \in L_2 : f^{(\nu)} \in L_2 \quad \text{for } \nu \in \{0, 1, \dots, m\}, \text{ where } f^{(\nu)} \text{ is the weak derivative of } f\}.$$

The vector space  $W_{m,2}$  equipped with the norm:

$$\|f\|_{m,2} = \left( \sum_{\nu=0}^m \|f^{(\nu)}\|^2 \right)^{1/2}$$

is called Sobolev space and it is Banach space.

**DEFINITION 6.** By  $D_{L_2}$  we shall denote the space of all smooth functions  $\varphi$  such that  $\varphi^{(\nu)} \in L_2$  for  $\nu \in \mathbb{N}$ .

The convergence in  $D_{L_2}$  may be defined by the family of norms  $\|\cdot\|_{m,2}$  for  $m = 0, 1, 2, \dots$

Let  $\Lambda$  be a linear functional on  $D_{L_2}$  continuous with respect to the norm  $\|\cdot\|_{m,2}$ . It is known that such functional takes the following form (see [8], p. 201)

$$(12) \quad \Lambda(\varphi) = \sum_{\nu=0}^m \int_R f_\nu(x) \varphi^{(\nu)}(x) dx \quad \text{for } \varphi \in D_{L_2},$$

where  $f_\nu$  for  $\nu \in \{0, 1, \dots, m\}$  are fixed elements of  $L_2$ . We shall now show that the functional  $\Lambda$  can be extended on  $W_{m,2}$  by continuity and formula (12) also holds.

Let be  $\varphi \in W_{m,2}$ . Then by virtue of density of  $D_{L_2}$  in  $W_{m,2}$  (see for example [1]), there exists a sequence  $\{\varphi_n\}_{n \in \mathbb{N}}$ ,  $\varphi_n \in D_{L_2}$  for  $n \in \mathbb{N}$ , such that:

$$(13) \quad \left\| \varphi_n^{(\nu)} - \varphi^{(\nu)} \right\| \xrightarrow{n \rightarrow \infty} 0 \quad \text{for } \nu = 0, 1, \dots, m.$$

From Hölder's inequality we have:

$$(14) \quad \left| \int_R [f_\nu(x) \varphi_n^{(\nu)}(x) - f_\nu(x) \varphi^{(\nu)}(x)] dx \right| \leq \|f_\nu\| \|\varphi_n^{(\nu)} - \varphi^{(\nu)}\|.$$

Since (13) and (14), therefore

$$\Lambda(\varphi) = \lim_{n \rightarrow \infty} \Lambda(\varphi_n) = \sum_{\nu=0}^m \int_R f_\nu(x) \varphi^{(\nu)}(x) dx \quad \text{for } \varphi \in W_{m,2}.$$

**THEOREM 10.** *Let  $\Lambda : D_{L_2} \rightarrow \mathbb{C}$  be a linear functional continuous with respect to the norm  $\|\cdot\|_{m,2}$  and  $g \in D_{L_1}$ ,  $\int_R g(x) dx = 1$  and  $g_\varepsilon(x) = \varepsilon^{-1} g(\frac{x}{\varepsilon})$  and  $\Lambda_{g_\varepsilon}(x) = \Lambda(g_\varepsilon(x - \cdot))$ .*

Then

$$\int_R \Lambda_{g_\varepsilon}(x) \varphi(x) dx \quad \text{tends to } \Lambda(\varphi) \text{ as } \varepsilon \rightarrow 0 \text{ for } \varphi \in W_{m,2}.$$

**PROOF.** An easy computation shows that:

$$(15) \quad \int_R \Lambda_{g_\varepsilon}(x) \varphi_n(x) dx - \Lambda(\varphi_n) =$$

$$= \sum_{\nu=0}^m \int_R \int_R [f_\nu(x - \varepsilon y) - f_\nu(x)] g(y) dy \varphi_n^{(\nu)}(x) dx \quad \text{for } \varphi_n \in D_{L_2}.$$

Assume that a sequence  $\{\varphi_n\}_{n \in \mathbb{N}}$ ,  $\varphi_n \in D_{L_2}$  for  $n \in \mathbb{N}$ , converges to  $\varphi$  in  $W_{m,2}$ , then

$$a) \quad \int_R \Lambda_{g_\varepsilon}(x) \varphi_n(x) dx \xrightarrow{n \rightarrow \infty} \int_R \Lambda_{g_\varepsilon}(x) \varphi(x) dx$$

$$b) \quad \Lambda(\varphi_n) \xrightarrow{n \rightarrow \infty} \Lambda(\varphi)$$

$$c) \quad \sum_{\nu=0}^m \int_R \int_R [f_\nu(x - \varepsilon y) - f_\nu(x)] g(y) dy \varphi_n^{(\nu)}(x) dx \rightarrow \\ \sum_{\nu=0}^m \int_R \int_R [f_\nu(x - \varepsilon y) - f_\nu(x)] g(y) dy \varphi^{(\nu)}(x) dx.$$

From a) b) c) it follows:

$$(15') \quad \int_R \Lambda_{g_\varepsilon}(x) \varphi(x) dx - \Lambda(\varphi) = \sum_{\nu=0}^m \int_R \int_R [f_\nu(x - \varepsilon y) - f_\nu(x)] \\ g(y) dy \varphi^{(\nu)}(x) dx \quad \text{for } \varphi \in W_{m,2}.$$

The rest part of the proof is as in the proof of Lemma 1 in [2].

**THEOREM 11.**

$$(16) \quad A_k \subset W_{k,2} \quad \text{for } k = 1, 2, \dots$$

**PROOF.** (Compare Th. 6 in [2]). Let  $\varphi$  be in  $A_k$ . In accordance with Definition 3  $\varphi$  is in  $L_2$ .

It is easy to see that

$$(17) \quad \varrho'_n = -ni\varrho_{n-1} + (2n+1)i\varrho_n - (n+1)i\varrho_{n+1}.$$

According to Theorem 5 the function  $\varphi$  can be represented as follows

$$\varphi = \sum a_n \varrho_n, \quad \text{where } a_n = (\varphi, \varrho_n).$$

Formally differentiating this series we obtain

$$\sum a_n \varrho'_n = -i \sum n a_n \varrho_{n-1} + i \sum (2n+1) a_n \varrho_n \\ - i \sum (n+1) a_n \varrho_{n+1}.$$

By virtue of Theorem 5 it follows that the series  $\sum |n|^2 |a_n|^2$ ,  $\sum |2n+1|^2 |a_n|^2$  and  $\sum |n+1|^2 |a_n|^2$  are convergent so  $\sum n a_n \varrho_{n-1}$ ,  $\sum (2n+1) a_n \varrho_n$  and  $\sum (n+1) a_n \varrho_{n+1}$  are convergent in  $L_2$ . Hence  $\varphi'$  is in  $L_2$ .

Analogously we will prove theorem 14. First using the formulæ (26) and (27) it can be shown that  $\varphi^{(\nu)} \in L_2$  for  $\nu \in \{0, 1, \dots, k\}$  in the case  $k > 1$ .

REMARK 1. *Remark 1* If  $\Lambda \in D'_{L_2}$  is functional continuous with respect to the norm  $\|\cdot\|_{m,2}$ , then we will write that

$$(18) \quad \Lambda \in W_{-m,2} := (W_{m,2})'$$

THEOREM 12. *If  $\Lambda$  belongs to  $W_{-k,2}$ , then the restriction  $\tilde{\Lambda}$  of the functional  $\Lambda$  to  $A_k$  belongs to  $A'_k$ .*

PROOF. By virtue of Theorem 10

$$(19) \quad \int_R \Lambda_{g_\varepsilon}(x) \varphi(x) dx$$

tends to  $\Lambda(\varphi)$  as  $\varepsilon \rightarrow 0$  for  $\varphi \in W_{k,2}$ .

Therefore, in view of Theorem 11 we infer that the expression (19) converges to  $\Lambda(\varphi)$  when  $\varepsilon \rightarrow 0$  for  $\varphi \in A_k$ . Taking into account that  $A_k$  is complete we conclude, by the Banach–Steinhaus theorem that  $\tilde{\Lambda}$  belongs to  $A'_k$ .

COROLLARY 1. *If  $\Lambda$  is in  $W_{-k,2}$ , then  $\tilde{\Lambda} = \sum a_n \varrho_n$  in  $A'_k$ , where  $a_n = \tilde{\Lambda}(\bar{\varrho}_n)$ .*

### 5. Some functionals from the space $A'_k$

Now we are going to show the examples of functionals in  $A'_k$ .

DEFINITION 7. For  $k \in \mathbb{N}$  and  $\varphi \in A$  we define the functionals:

$$(20) \quad \frac{1}{(\cdot + 0i)^k}(\varphi) = \lim_{\varepsilon \rightarrow 0^+} \int_{-\infty}^{+\infty} \frac{1}{(x + \varepsilon i)^k} \cdot \varphi(x) dx$$

$$(21) \quad \frac{1}{(\cdot - 0i)^k}(\varphi) = \lim_{\varepsilon \rightarrow 0^+} \int_{-\infty}^{+\infty} \frac{1}{(x - \varepsilon i)^k} \cdot \varphi(x) dx$$



DEFINITION 8. For  $k = 1$  we define functional

$$(22) \quad \frac{1}{(\cdot)}(\varphi) := \int_{-\infty}^{+\infty} (\varphi(x) - \varphi(-x)) \frac{1}{2x} dx \quad \text{for } \varphi \in A.$$

For  $k > 1$  we define functionals depending on evenness and oddness of  $k$ :  
for even  $k = 2m$

$$(23) \quad \frac{1}{(\cdot)^{2m}}(\varphi) := \int_{-\infty}^{+\infty} x^{-2m} \left[ (\varphi(x) + \varphi(-x)) \frac{1}{2} - \left( \varphi(0) + \varphi^{(2)}(0) \frac{1}{2!} x^2 + \dots + \right. \right. \\ \left. \left. + \varphi^{(2m-2)}(0) \frac{1}{(2m-2)!} x^{2m-2} \right) \right] dx \quad \text{for } m = 1, 2, \dots$$

for odd  $k = 2m + 1$ :

$$(24) \quad \frac{1}{(\cdot)^{2m+1}}(\varphi) := \int_{-\infty}^{+\infty} x^{-(2m+1)} \left[ (\varphi(x) - \varphi(-x)) \frac{1}{2} \right. \\ \left. - \left( \varphi'(0) \frac{1}{1!} x + \dots + \varphi^{(2m-1)}(0) \frac{1}{(2m-1)!} x^{2m-1} \right) \right] dx \\ \text{for } m = 1, 2, \dots \text{ and } \varphi \in A.$$

Compare the definition in [5], p. 199.

Notice that these functionals belongs to  $W_{-k,2}$  for appropriate  $k$ .

THEOREM 13. Functionals  $\frac{1}{(\cdot)}$ ,  $\frac{1}{\cdot+0i}$ ,  $\frac{1}{\cdot-0i}$ ,  $\delta \in A'_1$ .

THEOREM 14. Functionals  $\frac{1}{(\cdot)^k}$ ,  $\frac{1}{(\cdot+0i)^k}$ ,  $\frac{1}{(\cdot-0i)^k}$ ,  $\delta^{(k-1)} \in A'_k \setminus A'_{k-1}$ ,  
for  $k > 1$ .

We base on the following result (see for example [3]):

**Cauchy's representation theorem.** *If a simple closed curve  $C$ , positively oriented and lying in the region  $R$ , contains only points  $c$  of  $R$  in its interior, then a function  $f(z)$  analytic in  $R$  can be represented for points  $c$  interior to  $C$  by*

$$(25) \quad f(c) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z-c} dz,$$

and it's  $n$ -th derivative

$$(26) \quad f^{(n)}(c) = \frac{n!}{2\pi i} \int_C \frac{f(z)}{(z-c)^{n+1}} dz,$$

Moreover,

$$(27) \quad \int_C f(z) dz = 0.$$

(A region is defined as an open set which is arcwise connected.)

We denote by  $\Gamma_R^+$  and  $\Gamma_R^-$  the parts of circle

$$(28) \quad \Gamma_R^+ = \{Re^{it} : 0 \leq t \leq \pi\} \quad \text{and} \quad \Gamma_R^- = \{Re^{-it} : 0 \leq t \leq \pi\}$$

for constant  $R > 0$

and we use the following property of rational functions:

REMARK 2. Remark 2 If

- a)  $W(z)$  is rational function,
- b)  $W$  has no poles on the real line,
- c)  $\lim_{|z| \rightarrow +\infty} zW(z) = 0$

then:

$$(29) \quad \int_{-\infty}^{+\infty} W(x) dx = \lim_{R \rightarrow +\infty} \int_{[-R, R] \cup \Gamma_R^+} W(z) dz,$$

and

$$(30) \quad \int_{-\infty}^{+\infty} W(x) dx = \lim_{R \rightarrow +\infty} \int_{[-R, R] \cup \Gamma_R^-} W(z) dz.$$

Now using the Cauchy's representation theorem and remark 2 we can calculate the integrals:

LEMMA 1. For each  $\varepsilon > 0$  we have

$$(31) \quad \int_{-\infty}^{+\infty} \frac{1}{x + \varepsilon i} \bar{\varrho}_n(x) dx = \begin{cases} 0, & \text{for } n < 0 \\ i\sqrt{2\pi}(-1)^{n+1} \frac{(\frac{1}{2}-\varepsilon)^n}{(\frac{1}{2}+\varepsilon)^{n+1}}, & \text{for } n \geq 0 \end{cases}$$

$$(32) \quad \int_{-\infty}^{+\infty} \frac{1}{x - \varepsilon i} \bar{\varrho}_n(x) dx = \begin{cases} i\sqrt{2\pi}(-1)^n \frac{(\frac{1}{2}-\varepsilon)^{-n-1}}{(\frac{1}{2}+\varepsilon)^{-n}}, & \text{for } n < 0 \\ 0, & \text{for } n \geq 0 \end{cases}$$

PROOF. We want to calculate the integral

$$(33) \quad \int_{-\infty}^{+\infty} \frac{1}{x + \varepsilon i} \bar{\varrho}_n(x) dx = \frac{1}{i\sqrt{2\pi}} \int_{-\infty}^{+\infty} \frac{1}{x + \varepsilon i} \frac{(x + \frac{1}{2}i)^n}{(x - \frac{1}{2}i)^{n+1}} dx.$$

For  $n \geq 0$  the function  $f(z) = \frac{(z + \frac{1}{2}i)^n}{(z - \frac{1}{2}i)^{n+1}}$  is analytic in the half-plane  $\text{Im } z < \frac{1}{2}$ , so by the Cauchy's representation theorem we have

$$(34) \quad \int_{[-R, R] \cup \Gamma_R^-} \frac{1}{z + \varepsilon i} f(z) dz = -2\pi i f(-\varepsilon i) \quad \text{for } R > \varepsilon.$$

Using the remark 2 we receive

$$(35) \quad \int_{-\infty}^{+\infty} \frac{1}{x + \varepsilon i} \frac{(x + \frac{1}{2}i)^n}{(x - \frac{1}{2}i)^{n+1}} dx = -2\pi i \frac{(-\varepsilon i + \frac{1}{2}i)^n}{(-\varepsilon i - \frac{1}{2}i)^{n+1}} = 2\pi (-1)^n \frac{(\frac{1}{2} - \varepsilon)^n}{(\frac{1}{2} + \varepsilon)^{n+1}}.$$

So from (33) and (35) we have (31) for  $n \geq 0$ .

For  $n < 0$

$$(36) \quad \int_{-\infty}^{+\infty} \frac{1}{x + \varepsilon i} \bar{\varrho}_n(x) dx = \frac{1}{i\sqrt{2\pi}} \int_{-\infty}^{+\infty} \frac{1}{x + \varepsilon i} \frac{(x - \frac{1}{2}i)^{p-1}}{(x + \frac{1}{2}i)^p} dx \quad \text{where } p = -n.$$

Since the function  $f(z) = \frac{1}{z + \varepsilon i} \frac{(z - \frac{1}{2}i)^{p-1}}{(z + \frac{1}{2}i)^p}$  is analytic in the half-plane  $\text{Im } z \geq 0$  so  $\int_{[-R, R] \cup \Gamma_R^+} f(z) dz = 0$  for each  $R > 0$ , and by the remark 2

we have (31) for  $n < 0$ .

It means that:

$$(37) \quad \frac{1}{x + \varepsilon i} = i\sqrt{2\pi} \sum_{n=0}^{\infty} (-1)^{n+1} \frac{(\frac{1}{2} - \varepsilon)^n}{(\frac{1}{2} + \varepsilon)^{n+1}} \varrho_n(x) \quad \text{in } L_2$$

Similarily we have

$$(38) \quad \frac{1}{x - \varepsilon i} = -i\sqrt{2\pi} \sum_{n=-\infty}^{-1} (-1)^n \frac{(\frac{1}{2} - \varepsilon)^{-n-1}}{(\frac{1}{2} + \varepsilon)^{-n}} \varrho_n(x) \quad \text{in } L_2.$$

By definition 7 we have Fourier representation:

$$(39) \quad \frac{1}{\cdot + 0i}(\varphi) = i2\sqrt{2\pi} \sum_{n=0}^{+\infty} (-1)^{n+1} \varrho_n$$

and from theorem 9  $\frac{1}{\cdot + 0i} \in A'_1$ .

Analogously we can receive

$$(40) \quad \frac{1}{\cdot - 0i} = i2\sqrt{2\pi} \sum_{n=-\infty}^{-1} (-1)^n \varrho_n \quad \text{in } A'_1.$$

We can also easily calculate that:

$$(41) \quad \delta = \frac{2}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} (-1)^n \cdot \varrho_n.$$

Moreover, from the Sohotzki formulas:

$$(42) \quad (-1)\pi i\delta + \frac{1}{(\cdot)} = \frac{1}{\cdot + 0i}$$

or

$$(43) \quad \pi i\delta + \frac{1}{(\cdot)} = \frac{1}{\cdot - 0i},$$

we have:

$$(44) \quad \frac{1}{(\cdot)} = i\sqrt{2\pi} \sum_{n=-\infty}^{-1} (-1)^n \cdot \varrho_n + i\sqrt{2\pi} \sum_{n=0}^{\infty} (-1)^{n+1} \cdot \varrho_n.$$

From the above statement and theorem 9, theorem 13 follows.

Analogously we will prove theorem 14. First using the formulas (26) and (27) of the Cauchy's representation theorem and remark 2, we calculate the integrals.

LEMMA 2. For  $k > 1, k \in \mathbb{N}$  and  $\varepsilon > 0$  we have:

$$(45) \quad \int_{-\infty}^{+\infty} \frac{1}{(x + \varepsilon i)^k} \bar{\varrho}_n(x) dx = \begin{cases} 0, & \text{for } n < 0 \\ \frac{\sqrt{2\pi}}{i^k} (-1)^n \\ \left\{ \sum_{l=0}^{\min\{k-1, n\}} \frac{1}{l!(k-1-l)!} (n-l+1) \dots (n+k-1-l) \right. \\ \left. \left( \frac{1}{2} - \varepsilon \right)^{n-l} \left( \frac{1}{2} + \varepsilon \right)^{-n-k+l} \right\} & \text{for } n \geq 0 \end{cases}$$

$$(46) \quad \int_{-\infty}^{+\infty} \frac{1}{(x - \varepsilon i)^k} \bar{\varrho}_n(x) dx = \begin{cases} \frac{\sqrt{2\pi}}{i^k} (-1)^{n+1} \\ \left\{ \sum_{l=0}^{\min\{k-1, -n-1\}} \frac{1}{l!(k-1-l)!} (n+l) \dots (n+l-k+2) \right. \\ \left. \left( \frac{1}{2} - \varepsilon \right)^{-n-1-l} \left( \frac{1}{2} + \varepsilon \right)^{n-k+1+l} \right\} & \text{for } n < 0 \\ 0, & \text{for } n \geq 0 \end{cases}$$

This implies from definition 7

$$(47) \quad \frac{1}{(\cdot + 0i)^k} (\bar{\varrho}_n) = \begin{cases} 0, & \text{for } n < 0 \\ \frac{\sqrt{2\pi} \cdot 2^k}{i^k} (-1)^n \\ \left\{ \sum_{l=0}^{\min\{k-1, n\}} \frac{1}{l!(k-1-l)!} (n+1-l) \dots (n+k-1-l) \right\} \\ \text{for } n \geq 0 \end{cases}$$

$$(48) \quad \frac{1}{(\cdot - 0i)^k} (\bar{\varrho}_n) = \begin{cases} \frac{\sqrt{2\pi}}{i^k} (-1)^{n+1} \\ \left\{ \sum_{l=0}^{\min\{k-1, -n-1\}} \frac{1}{l!(k-1-l)!} (n+l) \dots (n+l-k+2) \right\} \\ \text{for } n < 0 \\ 0, & \text{for } n \geq 0 \end{cases}$$

Notice that, since  $\bar{\varrho}_n(x) = \varrho_n(-x)$ , for  $k > 1, k \in \mathbb{N}$  we have:

$$(49) \quad \delta^{(k-1)}(\bar{\varrho}_n) = (-1)^{k-1} \delta \left( \bar{\varrho}_n^{(k-1)} \right) = \varrho_n^{(k-1)}(0)$$

and

(50)

$$\varrho_n^{(k-1)}(0) = \begin{cases} \frac{i}{\sqrt{2\pi}} (-1)^n \frac{2^k}{i^k} \\ \left\{ \sum_{l=0}^{\min\{k-1, -n-1\}} \binom{k-1}{l} (-n-l) \dots (-n+k-l-2) \right\} \\ \text{for } n < 0 \\ \frac{i}{\sqrt{2\pi}} (-1)^{n+k-1} \frac{2^k}{i^k} \\ \left\{ \sum_{l=0}^{\min\{k-1, n\}} \binom{k-1}{l} (n-l+1) \dots (n+k-1-l) \right\} \\ \text{for } n \geq 0 \end{cases}$$

From Sochotzki formulas:

$$(51) \quad \frac{1}{(\cdot)^k} = \frac{1}{(\cdot + 0i)^k} - \frac{\pi i (-1)^k}{(k-1)!} \delta^{(k-1)}$$

or

$$(52) \quad \frac{1}{(\cdot)^k} = \frac{1}{(\cdot - 0i)^k} + \frac{\pi i (-1)^k}{(k-1)!} \delta^{(k-1)}$$

we can receive for  $k \in \mathbb{N}, k > 1$ :

(53)

$$\frac{1}{(\cdot)^k} (\bar{\varrho}_n) = \begin{cases} \frac{\sqrt{2\pi} 2^{k-1}}{i^k} (-1)^{n+k} \\ \left\{ \sum_{l=0}^{\min\{k-1, -n-1\}} \frac{1}{l!(k-1-l)!} (-n-l) \dots (-n+k-l-2) \right\} \\ \text{for } n < 0 \\ \frac{\sqrt{2\pi} 2^{k-1}}{i^k} (-1)^n \\ \left\{ \sum_{l=0}^{\min\{k-1, n\}} \frac{1}{l!(k-1-l)!} (n-l+1) \dots (n+k-1-l) \right\} \\ \text{for } n \geq 0 \end{cases}$$

From the previous statement and theorem 9, theorem 14 follows.

## REFERENCES

- [1] R. A. Adams, *Sobolev spaces*, Academic Press, New York 1975.
- [2] A. Cichocka, W. Kierat, *An application of the Wiener functions to the Dirichlet problem of Laplace equation*, to appear in *Integral Transforms and Special Functions*.
- [3] H. Cohn, *Conformal Mapping on Riemann Surfaces*, Dover Publications, Inc. New York 1980.
- [4] I. M. Gelfand, G. E. Shilov, *Generalized Functions*, vol. 2, Academic Press, New York and London 1968.
- [5] I. M. Gelfand, G. E. Shilov, *Generalized Functions*, vol. 1, Academic Press, New York and London 1964.
- [6] H. J. Glaeske, *On the Wiener-Laguerre transform of generalized functions*, in: *Generalized functions and convergence*, eds. P. Antosik, A. Kamiński.
- [7] W. Kierat, *Applications of quadratic forms generated by Sturm-Liouville operators in the theory of generalized functions*, Prace Naukowe Uniwersytetu Śląskiego nr 388, Katowice 1980.
- [8] L. Schwartz, *Théorie des distributions*, Hermann, Paris 1966.
- [9] N. Wiener, *Extrapolation, Interpolation and Smoothing of Stationary Time Series*, Technological Press of MIT and J. Wiley, New York 1949.
- [10] A. H. Zemanian, *Generalized Integral Transformations*, Intersci. Publ., New York 1968.
- [11] *Orthonormal Series Expansions of Certain Distributions and Distributional Transform Calculus*, J. Math. Anal. Appl. 14 (1966), 263-271.

INSTYTUT MATEMATYKI  
UNIwersytet ŚLĄSKI  
BANKOWA 14  
40 007 KATOWICE  
POLAND